


Erratum to: Ultrapure dialysis water obtained with additional ultrafilter may reduce inflammation in patients on hemodialysis

Biagio Di Iorio¹ · Lucia Di Micco¹ · Dario Bruzzese² · Luca Nardone¹ · Luigi Russo³ · Pietro Formisano⁴ · Vittoria D'Esposito⁴ · Domenico Russo³ 

Published online: 6 October 2017
© Italian Society of Nephrology 2017

Erratum to: J Nephrol
DOI [10.1007/s40620-017-0422-x](https://doi.org/10.1007/s40620-017-0422-x)

In original publication, the Table 4 was incorrect. The correct Table 4 has been given below:

Table 4 Concentrations of hemoglobin, ESA doses and erythropoietin resistance index (ERI) at Control and Study phase

Variable	Number	Control-phase	Study-phase	Percent variation study versus control phase
Hemoglobin (g/dl)	32	11.2 ± 0.9	11.9 ± 1.3 [•]	+6.25
Epoietin alpha (units per week)	19	11,313 ± 650	8670 ± 554 [•]	-23.4
Darboepoietin alpha (units per week)	10	80 ± 11	61 ± 12 ⁺	-23.7
Mircera (unit per month)	3	233 ± 130	152 ± 67	-34.8
ERI: EPO (week/BW)/Hb	29	13.1 ± 7.5	9.5 ± 6.0 [•]	

Numbers are mean ± SD

[•]*p* < 0.01

⁺*p* < 0.05

The online version of the original article can be found under doi: [10.1007/s40620-017-0422-x](https://doi.org/10.1007/s40620-017-0422-x).

✉ Domenico Russo
domenicorusso51@hotmail.com

¹ Department of Nephrology, “A. Landolfi” Hospital (Solofra, Avellino), Solofra, Italy

² Department of Statistics, University of Naples “FEDERICO II”, Naples, Italy

³ Department of Public Health, University of Naples “FEDERICO II”, Naples, Italy

⁴ Department of Genetics, University of Naples “FEDERICO II”, Naples, Italy