

Erratum to: Utilization of bioimpedance spectroscopy in the prevention of chronic breast cancer-related lymphedema

David I. Kaufman¹ · Chirag Shah² · Frank A. Vicini³ · Marisa Rizzi¹

Published online: 20 September 2017
© Springer Science+Business Media, LLC 2017

Erratum to: Breast Cancer Res Treat DOI 10.1007/s10549-017-4451-x

In the original publication of the article, under the heading, Study limitations in the Discussion section, the second paragraph, fifth sentence was published incorrectly as “Patients with an L-Dex increase of ≥ 5.5 undergo...sleeve for 4 weeks”.

The corrected sentence should read as below:

Patients with an L-Dex increase of ≥ 7 undergo circumference arm measurements and then treatment with an OTC compression sleeve for 4 weeks, while those with a volume increase of 5–10 cc will undergo L-Dex testing and compression sleeve for 4 weeks.

The online version of the original article can be found under doi:[10.1007/s10549-017-4451-x](https://doi.org/10.1007/s10549-017-4451-x).

✉ David I. Kaufman
dkat727@aol.com

- ¹ Breast Cancer Specialists, Bethpage, NY, USA
- ² Department of Radiation Oncology, Cleveland Clinic, Taussig Cancer Institute, Cleveland, OH, USA
- ³ Michigan Healthcare Professionals, 21st Century Oncology, Farmington Hills, MI, USA