



Correction to: Clinically meaningful and lasting HbA_{1c} improvement rarely occurs after 5 years of type 1 diabetes: an argument for early, targeted and aggressive intervention following diagnosis

Krishnarajah Nirantharakumar^{1,2} · Nuredin Mohammed² · Konstantinos A. Toulis² · G. Neil Thomas² · Parth Narendran^{1,3}

Received: 28 August 2017 / Accepted: 20 December 2017 / Published online: 25 April 2018
© Springer-Verlag GmbH Germany, part of Springer Nature 2018

Correction to: Diabetologia

<https://doi.org/10.1007/s00125-018-4574-6>

In Table 1 the data for age group, sex and Townsend index were incorrectly identified as mean ± SD instead of *n* (%). The table is corrected here.

The online version of the original article can be found at <https://doi.org/10.1007/s00125-018-4574-6>

✉ G. Neil Thomas
g.n.thomas@bham.ac.uk

✉ Parth Narendran
p.narendran@bham.ac.uk

¹ Department of Diabetes, University Hospitals Birmingham NHS Foundation Trust, Birmingham, UK

² Public Health, Epidemiology and Biostatistics, Institute of Applied Health Research, College of Medical and Dental Sciences, University of Birmingham, Edgbaston, Birmingham B15 2TT, UK

³ Institute of Immunology and Immunotherapy, Research College of Medical and Dental Sciences, University of Birmingham, Edgbaston, Birmingham B15 2TT, UK

Table 1 Baseline characteristics of participants and HbA_{1c} summary statistics

Characteristics	All participants <i>N</i> = 4525	Participants contributing for 10 years and above <i>n</i> = 938
Age group (years)		
0–10	1294 (28.6)	293 (31.3)
10–20	1737 (38.4)	324 (34.5)
20–30	798 (17.6)	138 (14.7)
30–40	696 (15.4)	183 (19.5)
Sex		
Male	2743 (60.6)	574 (61.2)
Female	1782 (39.4)	364 (38.8)
Townsend index		
1	1020 (22.5)	229 (24.4)
2	869 (19.2)	178 (19.0)
3	923 (20.4)	190 (20.3)
4	828 (18.3)	175 (18.7)
5	650 (14.4)	137 (14.6)
Missing	235 (5.2)	29 (3.0)
Follow-up period (years)		
Mean (SD)	6.0 ± 4.6	13.2 ± 2.4
Median (25th–75th percentile)	5.0 (2.1–9.2)	13.2 (11.2–14.8)
HbA _{1c} (mmol/mol) ^a		
Mean (SD)	72.6 ± 20.6	73.7 ± 19.5
Median (25th–75th percentile)	69.4 (58.5–82.5)	70.5 (60.7–83.6)
HbA _{1c} (%) ^a		
Mean (SD)	8.8 ± 4.0	8.9 ± 3.9
Median (25th–75th percentile)	8.5 (7.5–9.7)	8.6 (7.7–9.8)

Data presented as *n* (%), mean ± SD or median (25th–75th percentile)

^a Number of measurements in study period: 41,583 for all participants group, 16,989 for participants contributing for 10 years and above