



Codon bias confers stability to human mRNAs

Fabian Hia¹, Sheng Fan Yang¹, Yuichi Shichino², Masanori Yoshinaga¹, Yasuhiro Murakawa^{3,4}, Alexis Vandenbon⁵, Akira Fukao⁶, Toshinobu Fujiwara⁶, Markus Landthaler^{7,8}, Tohru Natsume⁹, Shungo Adachi⁹ , Shintaro Iwasaki^{2,10} & Osamu Takeuchi^{1,*} 

Abstract

Codon bias has been implicated as one of the major factors contributing to mRNA stability in several model organisms. However, the molecular mechanisms of codon bias on mRNA stability remain unclear in humans. Here, we show that human cells possess a mechanism to modulate RNA stability through a unique codon bias. Bioinformatics analysis showed that codons could be clustered into two distinct groups—codons with G or C at the third base position (GC3) and codons with either A or T at the third base position (AT3): the former stabilizing while the latter destabilizing mRNA. Quantification of codon bias showed that increased GC3-content entails proportionately higher GC-content. Through bioinformatics, ribosome profiling, and *in vitro* analysis, we show that decoupling the effects of codon bias reveals two modes of mRNA regulation, one GC3- and one GC-content dependent. Employing an immunoprecipitation-based strategy, we identify ILF2 and ILF3 as RNA-binding proteins that differentially regulate global mRNA abundances based on codon bias. Our results demonstrate that codon bias is a two-pronged system that governs mRNA abundance.

Keywords codon bias; codon optimality; GC-content; mRNA stability; translation efficiency

Subject Category RNA Biology

DOI 10.15252/embr.201948220 | Received 3 April 2019 | Revised 8 August 2019 | Accepted 19 August 2019 | Published online 3 September 2019

EMBO Reports (2019) 20: e48220

Introduction

Messenger RNA (mRNA) regulation represents an essential part of regulating a myriad of physiological processes in cells, being indicated in the maintenance of cellular homeostasis to immune

responses [1–3]. In addition to transcription regulation, post-transcriptional regulation of mRNA stability is vital to the fine-tuning of mRNA abundance. To date, several mRNA-intrinsic properties, often in 5' or 3' untranslated regions (UTRs), have been shown to affect mRNA stability [4,5]. Due to the recent advances in technology, the contribution of mRNA stability to gene expression has been suggested [6]. However, the regulation of mRNA stability, which is possibly governed by mRNA-intrinsic features, has not been fully elucidated.

One of the most crucial mRNA-intrinsic features is codon bias. To scrutinize this bias in usage of redundant codons, several metrics to measure how efficiently codons are decoded by ribosomes (codon optimality) have been proposed. In a classical metric called the codon Adaptation Index (cAI), gene optimality is calculated by comparison between codon usage bias of a target gene and reference genes, which are highly expressed [7,8]. Another index termed the tRNA Adaption Index (tAI) gauges how efficiently tRNA is utilized by the translating ribosome [9,10]. More recently, the normalized translation efficiency (nTE), which takes into consideration not only the availability of tRNA but also demand, was also proposed [11]. In addition to these, there are estimators of codon ribosome translation speed [12] as well as calculators of species-specific tAI [13].

Recently, Presnyak and colleagues showed that mRNA half-lives are correlated with optimal codon content based on a metric, the codon stabilization coefficient (CSC), which was calculated from the correlations between the codon frequencies in mRNAs and stabilities of mRNAs. Additionally, they showed that the substitutions of codons with their synonymous optimal and non-optimal counterparts resulted in significant increases and decreases in mRNA stability in yeast [14]. This effect was brought by an RNA-binding protein (RBP) Dhh1p (mammalian ortholog DDX6), which senses ribosome elongation speed [14–16]. In yeast, these differences in ribosome elongation speed in turn are influenced by tRNA availability and demand [11,17,18]. Taken together, codons can be designated into optimal and non-optimal categories: the former hypothesized to be

1 Department of Medical Chemistry, Graduate School of Medicine, Kyoto University, Kyoto, Japan

2 RNA Systems Biochemistry Laboratory, RIKEN Cluster for Pioneering Research, Wako, Japan

3 Division of Genomic Technologies, RIKEN Center for Life Science Technologies, Yokohama, Japan

4 RIKEN Preventive Medicine and Diagnosis Innovation Program, Yokohama, Japan

5 Laboratory of Infection and Prevention, Institute for Frontier Life and Medical Sciences, Kyoto University, Kyoto, Japan

6 Laboratory of Biochemistry, Department of Pharmacy, Kindai University, Higashiosaka City, Japan

7 RNA Biology and Posttranscriptional Regulation, Max Delbrück Center for Molecular Medicine Berlin, Berlin Institute for Molecular Systems Biology, Berlin, Germany

8 IRI Life Sciences, Institut für Biologie, Humboldt-Universität zu Berlin, Berlin, Germany

9 Molecular Profiling Research Center for Drug Discovery (molprof), National Institute of Advanced Industrial Science and Technology (AIST), Tokyo, Japan

10 Department of Computational Biology and Medical Sciences, Graduate School of Frontier Sciences, The University of Tokyo, Kashiwa, Japan

*Corresponding author. Tel: +81 75 753 9500; Fax: +81 75 753 9502; E-mail: otake@mfour.med.kyoto-u.ac.jp

decoded efficiently and accurately [19,20], while the latter slow ribosome elongation, resulting in decreased mRNA stability [14–16]. It is also important to make the distinction that common and rare codons do not necessarily imply optimal and non-optimal codons.

At present, codon optimality-mediated decay has been extensively studied and established particularly in *Saccharomyces cerevisiae* as well as other model organisms such as *Schizosaccharomyces pombe*, *Drosophila melanogaster*, *Danio rerio*, *Escherichia coli*, *Trypanosoma brucei*, and *Neurospora crassa* [21–27]. At present, the molecular mechanisms of this system of codon optimality in humans are under intense scrutiny [28, preprint: 29].

In this study, we show that codon bias-mediated decay exists in humans. Principal component analysis (PCA) showed that codons could be clustered into two distinct groups: codons with A or T at the third base position (AT3) and codons with either G or C at the third base position (GC3). This clustering was associated with mRNA half-lives enabling us to determine GC3 and AT3 codons as stabilizing and non-stabilizing codons, respectively. In this regard, the increased usage of GC3 codons entails an inevitable increase GC-content. We then developed an algorithm to quantify the codon bias of GC3 codons. With ribosome profiling, we show that codon bias-derived occupancy scores agreed with ribosome occupancy. Additionally, bioinformatics analysis revealed that frameshifts abrogate this GC3-AT3 delineation. We then verified our results *in vitro* using optimized and deoptimized reporter constructs. Here, we propose that GC3 codons and AT3 codons are optimized and deoptimized codons, respectively. Importantly, frameshifted optimized transcripts retain a certain level of stability, suggesting that overall the overall GC-content of transcripts is an additional determinant of stability. Finally, employing a ribonucleoprotein immunoprecipitation strategy, we identified RNA-binding proteins, which were bound to transcripts with low or high GC3-content. We propose that interleukin enhancer-binding factor 2 (ILF2) mediates mRNA stability of transcripts via codon bias.

Results

Codons in *Homo sapiens* can be categorized into GC3 and AT3 codons

To examine whether a system of codon bias exists in humans, we first compared codon frequencies in *Homo sapiens* and other model organisms. Hierarchical clustering analysis of codon frequency data obtained from Ensembl database [30] showed a difference between lower eukaryotes such as *Saccharomyces cerevisiae* and *Caenorhabditis elegans*, and higher eukaryotes such as *Homo sapiens* and *Mus musculus* (Fig 1A). To investigate codon bias in humans, we downloaded human coding sequence (CDS) data from the Ensembl BioMart database and calculated the codon counts for each coding sequence. For each CDS, we calculated the codon frequencies by expressing the codon counts as a fraction of the total number of codons in the CDS. We then performed a principal component analysis (PCA) on the CDS codon frequencies. The first principal component (PC1) of the PCA, which accounted for 22.85% of the total variance, divided codons into two clusters: codons with either G or C at the third base position (GC3) and codons with either A or T at the third base position (AT3) (Fig 1B). Interestingly, the

division within the second principal component (PC2) appeared to be split along the number of G/C or A/T bases in codons. We repeated our analysis on the CDS sequences from *S. cerevisiae* and found no such clustering (Fig EV1A). However, we discovered that the factor loading scores of the codons along the first principal component of our analysis in yeast corresponded to the CSC metric [14], albeit differences in the order (Fig EV1B). The above-mentioned results therefore raised the possibility that the PCA method might have identified optimal and non-optimal codons; GC3 and AT3 codons in humans may have a valid effect on mRNA stability. To investigate the agreement between the PCA method and CSC in humans, we calculated the CSC scores in humans using published datasets of global mRNA decay rates in physiologically growing HEK293 cells (GSE69153) [Data ref: 31,32] and compared them to the PC1 factor loading scores of the codons (Fig EV1C). We observed a correlation of $R^2 = 0.58$ between the two outputs, indicating a moderately strong agreement despite the methodologies being different.

We then tested the link between mRNA stability and GC3-AT3 codons using the above-mentioned mRNA stability data (GSE69153) [Data ref: 31,32]. Briefly, we divided the transcripts equally into quartiles based on their half-lives and averaged the codon frequencies within the quartiles. Strikingly, genes with short half-lives were associated with AT3 codons, while genes with longer half-lives were associated with GC3 codons (Fig 1C), suggesting a connection between third base of codons and the stability of mRNAs.

Broadly, the codon bias in mRNA can predict the stability of the mRNA. Classification by GC3-content might potentially implicate GC-content as a factor, which might affect the stability of mRNA. By summing the GC3 frequencies and GC bases of CDS sequences, we could determine the GC3- and GC-content of a gene (Dataset EV1). We then visualized the genome-wide GC3 and GC landscape by plotting the corresponding values via a histogram (Fig 1D). GC3-content was represented as a bimodal distribution with a range of values from the minimum of 24.1% to the maximum of 100%, while GC-content appeared similarly as a bimodal distribution with a range of values from a minimum of 27.6% to the maximum of 79.7%. A Pearson correlation analysis ($R^2 = 0.869$) between gene GC-content and GC3-content (Fig EV1D) reflected an enrichment of GC-content with increased GC3-content. Indeed, higher GC3-content was generally associated with better stability (Fig 1E top and Fig EV1E). To further verify the impact of GC3-content on mRNA stability, we plot the GC3-content data in Fig 1E (top) in the form of cumulative distribution functions and found these distributions to be significantly different from the genome average (Fig EV1F). As with our analysis with GC3-content, we grouped the half-life data by GC-content (Fig 1E, bottom) and observed a similar increase in half-lives even with the GC-content grouping. Interestingly, we also noted a decrease in half-life beyond a GC-content of 60%; this decrease also coincides with the decrease in half-lives in the GC3-content grouping (Fig EV1D). While we are currently unable to explain the associated decrease in both plots at extreme GC3- and GC-content, it would be interesting to investigate this particular drop-off in stability in the future.

Additionally, we noted that the codon bias *per se* was different between yeast and humans (Figs 1B and EV1A) [14]. We also observed this difference in *Xenopus*, zebrafish, and *Drosophila*, when compared to humans [24,33]. We repeated our analysis, this




Figure 1.

Figure 1. Bioinformatics analysis reveals that biased codons can be categorized into GC3 and AT3 codons, respectively.

- A Hierarchical clustering analysis of model organisms and their average CDS codon frequencies.
- B Principal component analysis of the CDS codon frequencies of 9,666 protein-coding genes. PC1 and PC2 indicate the first and second principal components.
- C Heatmap of half-lives of mRNA and their CDS codon frequencies. The transcripts were ranked according to their half-lives and divided equally into quartiles. The respective codon frequencies of each group were then averaged.
- D Histogram illustrating the distribution of genes and their respective GC3- and GC-content.
- E Comparison of average transcript mRNA half-lives across their respective GC3- and GC-content ranges. Number of transcripts within each gene optimality range is indicated above their respective points.

Data information: In (E), error bars represent the 95% confidence intervals.

time grouping the half-life dataset by their respective cAI (Fig EV1G). With the cAI dataset, we were able to observe increased half-life with an associated increased in cAI albeit only from the range of 0.75–0.95. In contrast, the PCA-derived GC3-content method was better able to recapitulate this increase in half-life compared to the cAI metric. Taken together, our analysis allowed us to designate GC3 and AT3 codons as stabilizing and destabilizing codons, respectively. Additionally, high GC3-content in transcripts inevitably results in high GC-content, which is a feature of stable mRNAs.

We then asked about the biological relevance associated with codon bias. Taking the 5% of lowest and highest ranked genes into account, we observed that genes with high GC3-content were enriched in developmental processes, while genes with low GC3-content were enriched in cellular division processes (Fig EV1H and I), suggesting the importance of codon bias-mediated mRNA decay across dynamic cellular processes in humans.

GC3-AT3 codon bias can explain ribosome occupancy to a certain extent

Given that GC3-AT3 codons were associated with high and low stability, respectively, we wondered whether these two groups were synonymous with optimal and non-optimal codons. It has been proposed that slower ribosome elongation rate modulated by low codon optimality affects the stability of mRNAs in yeast [14]. This led us to examine whether decelerated ribosomes could be observed especially in regions where optimality was low. From the PCA, PC1 factor loadings of the codons were indicative of how much a particular codon contributed to the AT3-GC3 grouping, i.e., instability–stability (Fig EV2A). Therefore, as a measure of estimating ribosome occupancy, the factor loading scores of the codons from the first principal component were utilized to derive codon bias-derived occupancy scores (refer to Materials and Methods for details on the calculation of scores). Because we speculated that a single codon would be insufficient in eliciting any noticeable effects on the speed of the ribosome, we divided each CDS into 25 bins from start codon to stop codon and summed up the codon bias-derived occupancy scores. We then compared these scores with corresponding ribosome occupancies derived from ribosome profiling [34]. Ribosome occupancy obtained from HEK293 cells growing under physiological conditions generally coincided with codon bias-derived occupancy (Fig 2A). These measurements were highly reproducible between replicates of ribosome profiling experiments across the transcriptome ($R^2 = 0.750$, 16,423 transcripts) (Fig EV2B). We observed a significantly better prediction of ribosome occupancy by codon bias-derived occupancy scores than that

derived from scrambled codon bias-derived occupancy scores (Fig 2B). Unfortunately, at the individual codon level, we only observed a weak but positive correlation ($R^2 = 0.13$) between ribosome occupancy and codon bias-derived scores (Fig EV2C). We believe that this difference in both calculations can be attributed to the binning of the ribosome occupancy data, which ensures that any reasonable slowing of ribosomes in regions of low optimality could be accurately manifested. Indeed, representative transcripts showed a good correlation between our binned codon bias-derived occupancy scores and ribosome occupancy as exemplified by EIF2B2, DYNC1L2, and IDH3G transcripts (Fig EV2D).

Although translation elongation and initiation are distinct steps, previous literature has suggested that optimal codons are also enriched in mRNAs with high translation [35]. Ribosome footprint reads normalized by mRNA abundances from RNA-seq enable the calculation of translation efficiency, which in turn is also generally regarded as the translation initiation rate [36]. Therefore, to establish the link between translation status and codon bias, we calculated the translation efficiency (TE)—ribosome footprints normalized by mRNA abundance. Indeed, our results showed that mRNAs with high GC3-content generally possessed high TE (Fig 2C). This phenomena also coincide with known research in zebrafish and yeast in that optimal genes generally have high TE [33,37]. To exclude the effect of mRNA abundances on TE, we grouped mRNA of similar abundances into separate groups and repeated our analysis (Fig EV2E). Within these groups, we still observed a general increase in TE within each of the groups, albeit a decrease in TE at a GC-content of 70–80% across all ranges of mRNA abundances (similar to Fig 2C).

To verify whether GC3 and AT3 codons were indeed associated with stability and instability respectively, we performed PCA on +1 and –1 frameshifted CDS sequences genome-wide and show that the GC3-AT3 demarcation was abolished (Fig 2D and E). Interestingly, we found that GC-rich (two or three G/C bases) and AT-rich (two or three A/T bases) codons contributed strongly to PC1 of the frameshifted data, showing that GC/AT-content is a natural consequence GC3-AT3 usage (Figs 1D and EV1D).

Thus far, we show that GC3 and AT3 codons are associated with mRNA stability, ribosome translation speed, and efficiency, thereby suggesting that the former and latter can be designated into optimal and non-optimal codons, respectively.

Codon bias affects mRNA stability

We then experimentally validated our bioinformatics observations of GC3 and AT3 codons in human cells. We developed a scheme based on the PC1 factor loadings in which we previously utilized in




Figure 2. GC3-AT3 codon bias can explain ribosome occupancy to a certain extent.

A Average ribosome occupancy and their respective codon bias-derived occupancy scores across the CDS of transcript (in 25 bins). Ribosome occupancy for 16,423 transcripts and their respective codon bias-derived occupancy were firstly binned into 25 bins, and the mean occupancy was calculated for each bin.

B Cumulative distribution plots showing the distributions of correlations between ribosome occupancy and codon bias-derived occupancy. Correlations obtained from the ribosome occupancy and scrambled codon bias-derived occupancy served as the control. A Kolmogorov–Smirnov test was performed between the codon bias-derived occupancy and the control group.

C Comparison of average transcript translation efficiencies (TEs) across their respective GC3-content ranges. Number of transcripts within each gene optimality range is indicated above their respective points.

D, E Principal component analysis of CDS codon frequencies of protein-coding genes derived from a +1 frameshift (D) and a –1 frameshift (E). Shaded ellipses indicate codons which are GC-rich (orange) and AT-rich (blue).

Data information: In (A, C), error bars represent the 95% confidence intervals.

our ribosome profiling analysis (Fig EV2A). Based on this scheme, codons could be optimized and deoptimized with regard to GC3-content within their codon boxes, *i.e.*, synonymous substitutions (Fig EV3A). Single box codons such as TGG (Trp) and ATG (Met) would remain unchanged. We synthesized two independent genes (*REL* and *IL6*) with differential GC3-content (Fig EV3B, Dataset EV2) and examined the stability of these reporter RNAs in HEK293 cells utilizing the Tet-off system (Fig 3A). As expected, the optimized transcripts of *REL* and *IL6* were more stable than their wild-type counterparts. Additionally, the decay rate of the deoptimized *IL6* reporter was faster, confirming that low GC3-content transcripts were unstable.

In addition to the RNA stability, higher GC3-content was also associated with higher translation efficiency (Fig 2C), thereby increasing protein production. Indeed, the protein abundance of the optimized *REL* reporter was higher than *REL-WT* even after normalization of protein abundance by steady-state mRNA levels (Figs 3B and EV3C). Using enzyme-linked immunosorbent assay (ELISA), we observed that expression of *IL6-OPT* resulted in a 1.5-fold and twofold significantly higher level of IL6 compared to its WT and *IL6-DE*, respectively (Fig 3C). In a similar fashion, normalization of IL6 protein abundance by mRNA levels revealed that translation efficiency of the optimized *IL6* reporter was higher than its WT and deoptimized reporter counterparts (Fig EV3D). We tested our *REL* reporters in HeLa cells and show that the high protein abundance of *REL-OPT* could also be observed (Fig EV3E). Similarly, actinomycin-based stability measurements of the *REL* reporters in HeLa cells revealed a similar increase in mRNA stability in the *REL-OPT* transcript (Fig EV3F). Moreover, polysome fractionation and subsequent qPCR analysis revealed that within the polysome fractions, *REL-OPT* transcript amounts were proportionately higher than *REL-WT* transcripts, suggesting that *REL-OPT* was translated more efficiently than *REL-WT* (Fig 3D). Thus far, our results validate the bioinformatics analyses and show that GC3 and AT3 codons can be designated as optimal and non-optimal codons.

GC-content as an additional determinant of stability

We then hypothesized that if the effect on mRNA stability was entirely the result of translational elongation, blocking translational elongation would restore stability to transcripts possessing low optimality to levels similar to that of their high optimality counterparts. We therefore treated cells expressing the *REL* reporters with a translation inhibitor, cycloheximide (CHX), and assayed the mRNA decay rates via the Tet-off system (Fig 4A). Treatment with CHX improved the stability of both *REL-OPT* and *REL-WT* transcripts compared to the control group. Interestingly, the stability of CHX-treated *REL-WT* transcripts was still significantly lower than that of CHX-treated *REL-OPT* transcripts. We repeated our experiments using the *IL6* reporters and found that in a similar fashion, CHX-treated *IL6-DE* transcripts were stabilized, albeit, not to the same extent as CHX-treated *IL6-OPT* (Fig 4B). Following this, we repeated our experiments using a different translation inhibitor, anisomycin (ANI), and obtained similar results (Fig 4C and D), suggesting that a translation-independent mRNA degradation pathway could also be present. It should be noted that an important caveat to the use of global translation inhibitors, CHX in particular, is that they have been

reported to potentially distort mRNA level measurements as well as translation efficiency [38–40].

We then synthesized a +1 frameshifted version of the *REL-OPT* transcript, removing any potential stop codons, which would have resulted in premature termination of transcription, and measured its stability via the Tet-off system (Fig 4E). This frameshifted version, while retaining a high GC-content (similar to *REL-OPT*), possessed a lower GC3-content than its in-frame counterpart (Fig EV3B). Surprisingly, the frameshifted version was still more stable than the WT form, yet less stable compared to its in-frame optimized counterpart, suggesting that high GC/low AU-content was able to retain a significant amount of transcript stability. To verify our findings, we similarly synthesized a +1 frameshifted version of the *IL6-OPT* transcript, which had a high GC-content (similar to *IL6-OPT*) but a GC3-content of 39.15%, the GC3-content falling between its WT and DE counterparts (Fig EV3B). This frameshifted version of *IL6* was relatively more stable compared to the DE transcript (Fig 4F). Taken together, our results reinforce the notion that in addition to GC3-content, GC-content could be an additional determinant of stability. Taken together, our results show that codon bias encompasses two modes of mRNA regulation, GC3- and GC-content dependent.

RNA-binding proteins differentially bind to transcripts of varying degrees of codon bias

Having shown that high optimality content inevitably accords high GC-content that in turn promotes mRNA stability, we wondered whether there were RNA-binding proteins (RBPs), which scrutinize, discriminate, or even affect an mRNA's fate. To identify RBPs that were either bound to transcripts bearing high or low optimality, we performed a ribonucleoprotein immunoprecipitation-based approach termed ISRIM (*In vitro* Specificity-based RNA Regulatory protein Identification Method) [41]. Lysates of HEK293 cells were mixed with FLAG peptide-conjugated *REL* and *IL6* transcripts of high and low optimality, and their interacting proteins were determined using mass spectrometry. We then calculated the fold changes based on the abundance of RBPs bound to *REL-WT* with respect to *REL-OPT* (Fig 5A).

As *IL6* transcripts possessed three levels of GC3-content (*OPT*, *WT*, and *DE*), we defined high GC3-content binding RBPs based on the RBP enrichment of *IL6-DE* to *IL6-WT* (Fig 5B) as well as *IL6-WT* compared to *IL6-OPT* (Fig 5C). Similarly, we defined low GC3-content binding RBPs based on the RBP enrichment of *IL6-DE* compared to *IL6-WT* (Fig 5B) as well as *IL6-WT* to *IL6-OPT* (Fig 5C). By selecting common RBPs belonging to each group, we defined a set of RBPs, which bound differentially to high GC3 and low GC3 *IL6* transcripts, respectively (Fig EV4A). We then selected RBP candidates, which were specifically enriched with either low or high GC3 transcripts common to both *REL* and *IL6* ISRIM experiments (Fig EV4B, Dataset EV3). In all, we show that RBPs can differentiate between transcripts of high GC3- and low GC3-content.

ILF2 regulates the stability of low GC3/high AT3 transcripts

We investigated the role of RBPs in modulating the stability of transcripts with different codon bias. Of interest were ILF2 and ILF3, RBPs identified from the list of RBPs interacting exclusively with low optimality transcripts. ILF2 and ILF3, also known as NF45 and




Figure 3. GC3-content of transcripts determines their fate.

A HEK293 Tet-off experiments showing the degradation of *REL-OPT* and *REL-WT* transcripts (left) and *IL6-OPT*, *IL6-WT*, and *IL6-DE* transcripts (right), post-doxycycline addition.

B Representative immunoblot of FLAG-tagged *REL-OPT* and *REL-WT* in HEK293T cells transfected with either empty plasmids, or plasmids bearing *REL-OPT* or *REL-WT*. The immunoblot is representative of three independent experiments. ACTB is shown as the loading controls.

C ELISA of secreted IL6 concentrations of *IL6-OPT*, *IL6-WT*, and *IL6-DE* from HEK293T cells transfected with plasmids bearing *IL6-OPT*, *IL6-WT*, and *IL6-DE*.

D Fold changes of *REL-OPT* and *REL-WT* transcript levels (top) relative to their abundances from fraction 1 as detected by qPCR across polysome fractions (below). Data represent the mean ± SD for three biological replicates.

Data information: In (A, D), data are representative of three independent experiments each with three replicates. The data represent the mean ± SD for three replicates. A two-way ANOVA with the Holm–Sidak multiple comparisons was performed. *P*-values are denoted as follows: *P* < 0.05 (*), *P* < 0.01 (**), and *P* < 0.001 (***). The half-lives of the respective transcripts are indicated in brackets. In (C), the data are representative of three independent experiments each with three replicates. The data represent the mean ± SD for three replicates. A one-way ANOVA with Tukey’s multiple comparisons was performed between samples where *P* < 0.01 (**) and *P* < 0.001 (***).




Figure 4.

Figure 4. GC-content as an additional determinant of stability.

- A, B HEK293 Tet-off experiments showing the degradation of *REL-OPT* and *REL-WT* transcripts (A) and *IL6-OPT*, *IL6-WT*, and *IL6-DE* transcripts (B), under vehicle (DMSO) and cycloheximide (CHX) treatment, post-doxycycline addition.
- C, D HEK293 Tet-off experiments showing the degradation of *REL-OPT* and *REL-WT* transcripts (C) and *IL6-OPT*, *IL6-WT*, and *IL6-DE* transcripts (D), under vehicle (PBS) and anisomycin (ANI) treatment, post-doxycycline addition.
- E, F HEK293 Tet-off experiments showing the degradation of *REL-OPT*, *REL-OPT* (+1 Frameshift), and *REL-WT* transcripts (E) and *IL6-OPT*, *IL6-WT*, *IL6-DE*, and *IL6-OPT* (+1 Frameshift) transcripts (F), post-doxycycline addition.

Data information: In (A–F), data are representative of three independent experiments each with three replicates. The data represent the mean \pm SD for three replicates. A two-way ANOVA with the Holm-Sidak multiple comparisons was performed. *P*-values are denoted as follows: *P* < 0.05 (*), *P* < 0.01 (**), and *P* < 0.001 (***).

NF90/NF110, respectively, are well known to function dominantly as heterodimers, which bind double-stranded RNA. ILF3 has been extensively studied, having shown to bind to AU-rich sequences in 3' UTR of target RNA to repress its translation [42]. We hypothesize that the binding of ILF2 and ILF3 as a heterodimer to their targets occurs as low optimality transcripts are inadvertently AU-rich. Here, we focused on the effects of these RBPs on low optimality transcripts. Firstly, using published RIP-seq data of ILF2 in two multiple myeloma cell lines, H929 and JLN3, we observed that ILF2 interacts with low optimality transcripts (Fig EV5A) [Data ref: 43,44]. Additionally, we analyzed RNA-seq data obtained from the ENCODE project of K562 cells treated by CRISPR interference targeting ILF2 [Data ref: 45]. Strikingly, we observed that transcripts that possessed low optimality scores were upregulated, whereas transcripts that possessed high optimality scores were downregulated (Figs 6A and EV5B). The abundance changes in representative mRNAs by ILF2 knockdown were antiparallel to their GC3-content (Fig EV5C).

However, differences in mRNA levels do not necessarily imply a difference in mRNA stability. To confirm whether mRNA stability was indeed affected, we examined the stability of FLAG-tagged versions of *REL-OPT* and *REL-WT* in the Tet-off system after ILF2 and ILF3 knockdown via siRNA (Fig 6B and C). Interestingly, we observed that the optimized reporter was more unstable under the knockdown of both ILF2 and ILF3, whereas the WT reporter was more stable with the knockdown of ILF2 and a combination of both ILF2 and ILF3 knockdown. In agreement with this, we found a significant increase in protein levels of *REL-WT* when cells were treated with ILF2- and ILF3-targeting siRNA (Figs 6D and EV5D). However, despite seeing a decrease in stability of the GC3-optimized reporter under both ILF2 and ILF3 knockdown, we were unable to observe this change at the protein level. Focusing our attention on ILF2, we expressed FLAG-tagged versions of *REL-OPT* and *REL-WT*, along with the two isoforms of ILF2, and detected the reporter protein levels via Western blot. A significant decrease in band intensity was observed for the *REL-WT* bands when both isoforms of ILF2 were expressed, whereas the amount of *REL-OPT* was not changed (Figs 6E and EV5E). Taken together, our results suggest that ILF2 and ILF3 affect mRNA transcripts with low GC3-content (and inadvertently low GC-content) to induce their decay.

Next, we sought to identify possible motifs, which are enriched in ILF2/3 targets. Based on the RIP-seq data in JLN3 and H929 [Data ref: 43,44], we identified common transcripts, which were more than fivefold differentially upregulated, and subjected their cDNA sequences to *de novo* motif identification via the MEME (Multiple EM for Motif Elicitation) software [46]. Our analysis

identified AU-rich motifs of about 6–7 nt long (Fig EV5F) as well as their distributions mainly in the CDS and 3'UTR along target transcripts. It should be noted that that these motifs are enriched in mRNA targets and may not necessarily imply *bona fide* binding motifs of ILF2/3. Therefore, we performed an additional motif search on a recently identified and experimentally validated ILF3 motif from RNA Bind-n-seq experiments by Dotu and colleagues [47] and found a similar distribution of motifs in the CDS and 3'UTR of targets (Fig EV5F).

Discussion

This study provides a framework describing codon bias-mediated RNA decay in humans. We first show that GC3 codons are associated with stability and AT3 codons with instability. We quantified codon bias by calculating the GC3-content within the CDS of genes and showed that GC3-content is strongly correlated with RNA stability and amount of protein expressed. In general, the use of optimal GC3 codons correlated with higher GC-content at a genome-wide level. We then show a modest agreement between codon bias-derived scores and ribosome occupancy as determined by ribosome profiling. Using GC3-optimized and GC3-deoptimized reporters, we validate our bioinformatics observations *in vitro*. Screening of RNA-binding proteins and further *in vitro* analysis suggests a role of ILF2, possibly in complex with ILF3, in the codon-mediated regulation of mRNA. Taken together, we conclude that gene expression can be shaped by codon bias and inevitably by GC/AU-content through the modulation of mRNA stability in human cells.

Investigating the system of codon bias in humans

Since translation elongation is affected by tRNA availability, the tRNA Adaptation Index (tAI), which is based on genomic tRNA copy number, has been used as a surrogate for codon optimality. However, in contrast to yeast, tRNA copy number in the genome is not always correlated with tRNA abundance in higher eukaryotes [48]. Hence, this metric is less suitable for quantifying codon optimality in humans. Independent of tRNA-based metrics, we addressed these challenges by utilizing an unsupervised learning algorithm, PCA, to identify features in that were mRNA-intrinsic. In the PCA of both yeast and humans, we demonstrated that the first principal component mirrored optimal/non-optimal assignments. We also show that the codon bias is different between these two organisms (Figs 1B and EV1A). In humans, the classification of codons into AT3 and GC3 groups was striking, but the percentage by which it accounts for its variation, however, was modest.




Figure 5. RNA-binding proteins bind differentially to transcripts with different levels of GC3-content.

A–C Volcano plots showing the enrichment of RBPs, which bind to *REL-WT* relative to *REL-OPT* transcripts (A), *IL6-DE* relative to *IL6-WT* transcripts (B), and *IL6-WT* relative to *IL6-OPT* transcripts (C).

Data information: In (A–C), vertical dotted lines indicate a 1.5-fold enrichment, while horizontal dashed lines indicate the *P*-value cut-off of 0.05. Points shaded in blue indicate RBPs, which have a differential fold change of more than 1.5 and *P* < 0.05.

From the PCA, the first and second principal components only explain a quarter of total variance in codon frequencies (Fig 1B), implying that other factors that explain bias of codon frequency possibly remain in human cells. The limitation of this method is reflected in the use of codon frequencies as our input data for the PCA. This approach might have neglected other factors of stability or instability, which might be codon-independent or which might be inherent at the nucleotide level. Assuming that evolution drives the selection of codons, synonymous codon usage in different organisms must be fine-tuned over time to achieve precise expression levels of mRNA and eventually proteins in essential physiological processes. Indeed, similar to our findings, a study by Bazzini *et al* [33] showed that a system of codon optimality is conserved among vertebrates, *Xenopus* and Zebrafish. In addition, they demonstrated that in Zebrafish embryos, low codon optimality was associated with shorter poly(A) tail length in addition to lower levels of translation. Our data together with recently published work by Wu and colleagues [28] indicate that a system of codon optimality exists in humans.

Our investigations show that high GC3/AT3-content or GC/AT-content in mRNA is selected for to modulate transcript stability in essential physiological processes, but is subject to constraints by amino sequence. Indeed, we show that transcripts with high and low GC3-content were linked to particular physiological and cellular processes (Fig EV1H and I). In a particular study, Gingold and colleagues argue that tRNA abundances vary in proliferating and differentiating cell types [49]. Interestingly, they showed that codons preferred by cell cycling genes were AT3 codons, while pattern specification preferred codons tended to be GC3 codons—in agreement with our GO analyses. In *Drosophila*, the correlation between codon optimality and mRNA stability has been demonstrated to be attenuated in neural development, possibly allowing the effect of trans-acting factors to dominate development [24].

Our results show that the codon bias we have identified affects ribosome occupancy to a significant but limited extent (Fig 2B). At the level of individual codon occupancies, we only observed a weak but positive correlation ($R^2 = 0.13$) between ribosome occupancy and codon optimality-derived scores (Fig EV2C). These results, however, are not surprising given that studies based on ribosome profiling data found no correlations between ribosome occupancy and rare codons [50,51]. In view of this, we binned the CDS into 25 evenly spaced groups to ensure that any reasonable slowing of ribosomes in regions of low optimality could be accurately represented by the GC3-AT3 bias. However, we acknowledge that our metric is only able to demonstrate a prediction to a limited extent. There are many factors that can affect ribosome profiling results such as




Figure 6.

Figure 6. ILF2 regulates the stability of low GC3/high AT3 transcripts.

- A Cumulative distribution plots showing the difference in distribution of transcript optimality between upregulated and downregulated transcripts in K562 cells subject to ILF2 CRISPR interference targeting ILF2. Transcript quantities are indicated in the figure legend.
- B, C HEK293 Tet-off experiments showing the degradation of *REL-OPT* (B) and *REL-WT* (C) transcripts with ILF2 and ILF3 siRNA and control (CTR) siRNA treatment, post-doxycycline addition.
- D Representative immunoblot of FLAG-tagged *REL-OPT* and *REL-WT* expressed in HEK293T cells under ILF2 and ILF3 siRNA treatment. The immunoblot is representative of three independent experiments. ACTB is shown as loading controls.
- E Representative immunoblot of FLAG-tagged *REL-OPT* and *REL-WT* in HEK293T cells co-expressed with two different isoforms of ILF2. The immunoblot is representative of three independent experiments. ACTB is shown as loading controls.

Data information: In (A), Wilcoxon signed rank tests were performed on the upregulated and downregulated groups against the control group. *P*-values are denoted (right). In (B, C), data are representative of three independent experiments in which the data represent the mean \pm SD for three biological replicates. A two-way ANOVA with the Holm–Sidak multiple comparisons was performed. *P*-values are denoted as follows: $P < 0.05$ (*) and $P < 0.01$ (**).

growth conditions, coverage, cloning and sequencing biases, methods of bioinformatics analysis, and experimental noise [18,52,53].

Taking into account our *in vitro* experiment results together with the ribosomal profiling results, we suggest that GC3 and AT3 codons are synonymous with optimal and non-optimal codons. Additionally, our study along with others' suggests that slower elongation of ribosome is a key feature of mRNA stability. However, it should be noted that in our analysis methodology, the assumption that stability is solely a function of ribosome speed might only hold true to a limited extent. There is evidence to show that mRNA-intrinsic features that have the propensity to regulate ribosome velocity are essential in maintaining the function and correct expression of proteins, the failure of which may result in degradation of the mRNA and protein: Although codon optimality is a dominant factor in general, other factors may also be involved in decelerated ribosomes, such as secondary structures [54,55]. These obstacles for ribosome elongation are reversible and dynamically regulated by RNA helicases [56,57]. Importantly, these structures may serve to reduce ribosome speed when the nascent peptide requires additional time to fold to its correct conformation [58]. Furthermore, it has been shown in *Neurospora* that codon usage can regulate co-translational protein folding and, subsequently, its function [59].

As such, while we have shown that the optimizations of transcripts lead to increases in protein production, further studies are required to investigate protein folding dynamics and determine whether the produced protein still retains its functionality. Furthermore, in a study of two model organisms, *E. coli* and *S. cerevisiae* by Tuller *et al* [60], the rate of translation elongation was shown to be determined by the folding energy, codon bias, and amino acid charge at the beginning of the CDS. It is likely that these factors may also affect the local speed of the ribosome further down the CDS and, by extension, the stability of the mRNA. Further studies will be required to elucidate the role of RNA secondary structures and helicases and their relevance to codon bias, protein folding, and mRNA stability decay.

In attempts to quantify the effect of ribosomal density on mRNA stability, several studies have demonstrated that in general, increased ribosomal density results in increased mRNA stability of a transcript [61,62]. This phenomenon has been attributed to competition between the initiation complex and decay factors as well as ribosomes sterically excluding decay factors from accessing the mRNA [63,64]. To this effect, reduction in translation initiation has been shown to decrease ribosomal density and, subsequently, mRNA stability [65]. On the other hand, inhibiting translation elongation causes an increase in ribosome density and, consequently,

mRNA stability [66]. Here, we show that optimized transcripts are highly polysome-bound as opposed to their WT counterparts suggesting increased rates of translation initiation (Fig 3D). This is corroborated by our ribosome profiling findings that high GC3-containing transcripts have higher TE (Fig 2C), possibly protecting transcripts from decay factors.

In this regard, transcripts with high optimality have higher translation initiation rates, causing them to be highly polysome-bound. Additionally, optimized codons allow for efficient decoding and, thus, smoother ribosome traffic. On the other hand, transcripts with low optimality tend to be less polysome-bound with frequent ribosome deceleration and/or stalling. Our ribosome profiling analyses in Fig 2B, however, is tailored to comparing the relative ribosome densities (in bins) within an individual transcript, against the codon bias optimality scores. While we show relative accumulation of ribosomes in low optimality regions locally within a transcript, this particular analysis can neither be extended to comparing total ribosome densities across the transcriptome nor compared to the polysome profiling results.

Interestingly, in a separate study in *Neurospora*, gene expression modulated by codon usage was shown to be due to the effects of transcription rather than translation [67]. In a follow-up study, the group also demonstrated C/G bias is able to promote gene expression by suppressing premature transcription termination [68]. In addition, several other studies have demonstrated that in mammalian cells, GC-rich genes are transcribed with increased efficiency, resulting in higher levels of transcripts independent of mRNA degradation [69,70]. Next, a study by Fu *et al*, which investigated the effects of codon usage bias on two proto-oncogenes with similar amino acid identity, but differing levels of optimality, KRAS and HRAS, showed that codon usage can affect both transcription efficiency and translation efficiency, suggesting that the effect of codon bias is multilevel [71]. In this and another study, changing the rare codons of KRAS to common ones increased its enrichment in the polysome fractions [72]. Likewise, *REL-OPT* transcripts were enriched in the polysome fractions compared to *REL-WT* transcripts. Nevertheless, our investigations also show that steady-state transcript copy number of the optimized reporter transcripts was significantly higher than that of the WT (and DE versions) (Fig EV3C and D). In addition to this, however, we also show increased translation efficiency in mRNA that contains a higher proportion of optimized codons. In our study and several other vertebrates, however, translation is the predominant effector of gene expression [33].

At the time of writing this manuscript, a study was published by Wu and colleagues, which demonstrated that translation is indeed a

determinant of mRNA stability in human cells [28]. While paper by Wu *et al* had assigned optimal and non-optimal designations to codons via the calculation of the CSC derived from ORFeome and SLAM-seq experiments, we noted that some of the findings paralleled ours. Indeed, the codon designations of optimal and non-optimal codons also showed modest delineation of codons into GC3 and AT3 codons, respectively. In another article published in the bioRxiv preprint server, Forrest and colleagues utilized a combination of endogenous and human ORFeome collection mRNAs in human cells to derive the CSC for human cells [preprint: 29]. Similar to the study by Wu and colleagues, the codon designations of optimal and non-optimal codons also showed a modest division of codons into GC3 and AT3 codons, respectively. Similarly, we also show that the use of optimal and non-optimal codons can affect both mRNA stability and translation initiation to a large extent (Figs 1–3), albeit transcription to a limited extent. However, we have yet to identify an RBP that is involved in direct co-translational decay of mRNAs in humans as with that in yeast. Moreover, DDX6, the mammalian ortholog of DHH1, was recently demonstrated in humans to be involved in miRNA-driven translational repression, not mRNA destabilization as previously shown in yeast [73]. DDX6 aside, it would certainly be exciting for future experiments to uncover the nature of this elusive RBP.

c-Rel, a protein encoded by the *REL* gene and a canonical nuclear factor κ B (NF- κ B) subunit, is expressed abundantly in differentiated lymphoid cells and has been shown to be vital in thymic regulatory T-cell development in addition to controlling cancer via activated regulatory T cells [74,75]. Given the inherent low optimality and associated instability of *REL* in its WT form (Fig 3A), we wonder whether besides transcriptional control of *REL*, could there be other post-transcriptional regulation systems at play. Further studies would be necessary to investigate whether codon optimality or codon optimality-associated RBPs modulate *REL* gene expression.

In our investigation, mRNA stability can be affected by GC3- and GC-content. It is important to note that the latter of which is also implicated in several processes such as miRNA binding, mRNA folding, and splicing, which in turn can affect mRNA stability. It is thus plausible that GC-content can also affect gene expression independent of RBP association. A study of transcriptome miRNA-binding sites has shown that effective miRNA-binding sites tend to dwell in G-poor and U-rich environments [76]. In addition, while our analyses are CDS-based, it has been shown that GC-content of both introns and exons is important in splicing via RNA structures [77–79]. Taken together, we propose that codon bias is able to exert its effects at multiple levels, consequently effecting gene and protein expression.

The stability of mRNA can be modulated by RBPs, which bind AU-rich sequences

Whereas AU-rich elements (AREs) in the 3'UTR have been traditionally targeted by RBPs, we found that coding regions are also targeted by ARE-recognizing RBPs. The identification of the heterodimeric complex consisting of ILF2 and ILF3 among others shows that a wide array of RBPs recognizes low optimality (AU-rich) sequences (Fig 5). However, the binding of ILF2/3 to target RNA presents as a challenge when trying to identify its target motif. Studies have shown that the RNA-binding portion of the ILF2/3 complex, ILF3, in

particular is a promiscuous RBP, binding to RNA with no obvious sequence specificity [80]. It is interesting to note that several binding motifs, all of which are AU-rich have been proposed for ILF3. Analysis of ILF3 RNA Bind-n-seq measurements identified a 9 nt AU-rich motif that is bound to by ILF3 [47]. Kuwano and colleagues show that NF90, the shorter isoform of ILF3, specifically targets a 30 nt AU-rich sequence in mRNA 3'UTRs and represses their translation, not stability [42]. This state of promiscuousness was compounded by a recent study by Wu and colleagues, in which in almost all genes where ILF3 occupancy was detected on the genome by ChIP-seq, ILF3 occupancy was on the corresponding transcript. Indeed, ILF3 is a multifunctional protein, affecting several biological processes. In addition to ours, other studies have shown that ILF3 can contribute to splicing [81], stabilization, nuclear export [82], and, as mentioned, translation [42].

ILF2 on the other hand has been less scrutinized compared to its partner. From our experiments, we find that the longer isoform of ILF2 is predominantly and highly expressed, while the shorter isoform is low in expression. Additionally, we observed that overexpression of the longer isoform appeared to upregulate the expression of the shorter isoform albeit to a small extent. From the literature, it is known that ILF2 stabilizes ILF3 in the heterodimeric form [83]. We postulate that it is possible that the ILF2/3 heterodimer represses translation of mRNA with AU-rich sequences at a steady state in both CDS and 3'UTR. Knockdown of ILF2/3 relieves the repression on translation initiation allowing an increase in bound (translating) ribosomes, which sterically exclude decay factors from accessing the mRNA, thereby increasing stability. Indeed, the knockdown of ILF2, which is critical in maintaining the stability of the heterodimeric complex, results in a stabilization of mRNA possibly due to increased ribosome traffic. At the protein level, while the knockdown of ILF2 results in an increased protein expression of target mRNA, the combined effect of both ILF3 and ILF2 knockdown results in a higher increase in target mRNA expression as compared to the ILF2-only knockdown. Unfortunately, in the case of the ILF2/3 siRNA experiments (Fig 6D), we were unable to achieve a complete knockdown of ILF2 due to the very high and constitutive production of ILF2. However, we still noted a small reduction in ILF3 protein levels, hinting that ILF2 stabilizes ILF3 in the heterodimer form. In addition, taking into consideration reports that ILF2 and ILF3 can function independently of each other [84–86], it is also possible that ILF2 and ILF3 regulate the fate of mRNA differently, ILF2 being able to dimerize with other binding partners such as ZFR and SPNR. It is unknown, however, how optimized transcripts are affected. Whereas our screens revealed that ILF2/3 bind exclusively to low optimality targets, we noted from our analysis of ILF2 knockdown data from the ENCODE database [Data ref: 45] as well as tests from our reporter constructs that high optimality transcripts are being regulated. Given this, we postulate that ILF2/3 might not interact directly with high optimality targets. Instead, ILF2/3 may be indirectly (de)antagonizing certain transcripts, which may code for other regulators of high optimality genes. Further investigations will be required to assess how high optimality transcripts are antagonized.

Our screens also detected HNRNPD/AUF1, which destabilizes transcripts via recognition of AU-rich motifs [87], binding to low optimality mRNAs (Dataset EV3). These observations emphasize the importance of AU-content, which is strongly connected with low

optimality, in RNA destabilization. However, it is possible that these factors induced the degradation of AU-rich transcripts different from the model proposed by Presnyak and Radhakrishnan [14,15] as our RBP identification method was not fully reflective of the active translational status required for co-translational degradation of mRNA transcripts. Further studies would be necessary to discern whether these or other factors act as sensors of codon optimality during translation.

In conclusion, in human cells, the redundancy of the genetic code allows the choice between alternative codons for the same amino acid, which may exert dramatic effects on the process of translation and mRNA stability. In our experiments, we show that two modes of mRNA regulation exist—GC3- and GC-content dependent. This system potentially confers freedom for calibrating protein and mRNA abundances without altering protein sequence. Beginning from our exploratory analysis, we have developed an approach to quantify codon bias and demonstrate that beneath the redundancy of codons, exists a system, which modulates mRNA and, consequently, protein abundance.

Materials and Methods

Cell cultures, growth, and transfection conditions

HEK293T cells were maintained in Dulbecco's modified Eagle's medium (DMEM) (Nacalai Tesque), supplemented with 10% (v/v) fetal bovine serum. HEK293 Tet-off cells were maintained in Minimum Essential Medium Eagle—Alpha Modification (α -MEM) (Nacalai Tesque), supplemented with 10% (v/v) Tet-off system-approved fetal bovine serum (Takara Bio) and 100 μ g/ml of G418 (Nacalai Tesque). For REL and IL6 overexpression experiments, plasmids were transfected using PEI MAX (Polysciences, Inc). For co-transfection of ILF2 siRNA with REL plasmids, Lipofectamine 2000 was used as per the manufacturer's protocol. ILF2 siRNA that targeted ILF2 at exons 8 and 9 was Silencer Select siRNA, S7399 (Ambion, Life Technologies). Actinomycin D-based stability assays in HeLa cells were performed by adding actinomycin D to the transfected cells to a final concentration of 2 μ g/ml.

Plasmid construction

Codon-optimized *REL* (*REL-OPT*) and *IL6* (*IL6-OPT*) and codon-deoptimized *IL6* (*IL6-DE*) sequences were synthesized as gBlocks Gene Fragments (Integrated DNA Technologies) (Dataset EV2). The *REL-OPT* (+1 frameshift) sequence was constructed by adding a +1 frameshift just after the start codon. Resulting stop codons were removed to ensure no premature termination. These sequences and corresponding WT sequences were polymerase chain reaction (PCR) amplified (with the inclusion of a FLAG tag for *REL* sequences) and inserted into the pcDNA3.1(+) vector (Invitrogen) and pTRE-TIGHT vector (Takara Bio). The sequences were confirmed via restriction enzyme digest and sequencing.

Tet-off assay

HEK293 Tet-off cells (Clontech) were transfected with pTRE-TIGHT plasmids bearing the (de)optimized and WT sequences and

incubated overnight at 37°C. Transcriptional shut-off for the indicated plasmids was achieved by the addition of doxycycline (LKT Laboratories Inc.) to a final concentration of 1 μ g/ml. Cycloheximide-based stability assays in HEK293 Tet-off cells were performed by adding actinomycin D to the transfected cells to a final concentration of 50 μ g/ml. Anisomycin-based stability assays in HEK293 Tet-off cells were performed by adding anisomycin to the transfected cells to a final concentration of 20 μ g/ml. Samples were harvested at the indicated time points after the addition of doxycycline (and cycloheximide/anisomycin).

RNA extraction, reverse transcription-PCR, and quantitative real-time PCR

Total RNA was isolated from cells using TRIzol reagent (Invitrogen) as per the manufacturer's instructions. Reverse transcription was performed using the ReverTra Ace qPCR RT Master Mix with gDNA Removal Kit (Toyobo) as per the manufacturer's instructions. cDNA was amplified with PowerUp SYBR Green Master Mix (Applied Biosystems), and quantitative real-time PCR (qPCR) was performed on the StepOne Real-Time PCR System (Applied Biosystems). To quantify transcript abundance of the REL reporters, pTRE-TIGHT plasmids bearing the (de)optimized and WT reporter sequences were used as standards. Human GAPDH abundance was used for normalization. The list of qPCR primers can be found in Dataset EV2.

Sucrose gradient centrifugation (polysome profiling)

HEK293T were transfected with equal concentrations of *REL-OPT* and *REL-WT* plasmids. Cells were lysed the next day in polysome buffer [20 mM 4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid (HEPES-KOH) (pH 7.5), 100 mM KCl, 5 mM MgCl₂, 0.25% (v/v) Nonidet P-40, 10 μ g/ml cycloheximide, 100 units/ml RNase inhibitor, and protease inhibitor cocktail (Roche)]. Lysates were loaded on top of a linear 15–60% sucrose gradient [15–60% sucrose, 20 mM HEPES-KOH [pH 7.5], 100 mM KCl, 5 mM MgCl₂, 10 μ g/ml cycloheximide, 100 units/ml RNase inhibitor, and protease inhibitor cocktail (Roche)]. After ultracentrifugation at 256,800 g for 2.5 h at 4°C in a HITACHI P40ST rotor, fractions were collected from the top of the gradient and subjected to UV-densitometric analysis. The absorbance profiles of the gradients were determined at 254 nm. For disassociation of ribosome and polysome, EDTA was added to Mg²⁺-free polysome buffer and 15–60% sucrose gradient at concentrations of 50 and 20 mM, respectively. For RNA analysis, RNA from each fraction was extracted via the High Pure RNA Isolation Kit (Roche) and subject to reverse transcription and qPCR.

Immunoblot analysis

Samples were lysed in RIPA buffer (20 mM Tris-HCl [pH 8], 150 mM NaCl, 10 mM EDTA, 1% Nonidet-P40, 0.1% SDS, 1% sodium deoxycholate, and cOmplete Mini EDTA-free Protease Inhibitor Cocktail [Roche]). Protein concentration was determined by the BCA Protein Assay (Thermo Fisher). Whole-cell lysates were resolved by SDS-PAGE and transferred onto PVDF membranes (Bio-Rad). The following antibodies were used for immunoblot

analysis: mouse monoclonal anti-FLAG (F3165, Sigma), mouse monoclonal anti-ILF2 (sc-365283, Santa Cruz Biotechnology), mouse anti- β -actin (sc-47778, Santa Cruz), and mouse IgG HRP-linked F(ab')₂ fragment (NA9310, GE Healthcare). Luminescence was detected with a luminescent image analyzer (Amersham Imager 600; GE Healthcare).

ELISA

HEK293T cells were transfected with pcDNA3.1(+) plasmids bearing the (de)optimized and WT sequences and incubated overnight at 37°C. Cell supernatant was aspirated, and the cell monolayer was washed with 1× PBS (pre-warmed at 37°C). Pre-warmed DMEM was added to the monolayer, and the cells were incubated for 2 h at 37°C. Thereafter, the cell supernatant was harvested and centrifuged at 300 ×g to pellet residual cells. The resulting supernatant was decanted, and the concentration of secreted IL6 was measured by the Human IL6 ELISA Kit (Invitrogen) according to the manufacturer's instructions.

ISIRM (In vitro specificity-based RNA regulatory protein identification method)

Preparation of bait RNAs

T7-tagged cDNA template was PCR-amplified and subjected to *in vitro* transcription using a MEGascript T7 Kit (Applied Biosystems). Amplified cRNA was purified with an RNeasy Mini Kit (Qiagen) and then subjected to FLAG conjugation as described (10) with some modifications. Briefly, 60 μ l of freshly prepared 0.1 M NaIO₄ was added to 60 μ l of 250 pmol cRNA, and the mixture was incubated at 0°C for 10 min. The 3' dialdehyde RNA was precipitated with 1 ml of 2% LiClO₄ in acetone followed by washing with 1 ml acetone. The pellet was dissolved in 10 μ l of 0.1 M sodium acetate, pH 5.2, and then mixed with 12 μ l of 30 mM hydrazide-FLAG peptide. The reaction solution was mixed at room temperature for 30 min. The resulting imine moiety of the cRNA was reduced by adding 12 μ l of 1 M NaCNBH₃ and then incubated at room temperature for 30 min. The RNA was purified with an RNeasy Mini Kit (Qiagen).

Purification and analysis of RNA-binding proteins

Purification and analysis of RNA-binding protein (RBP) were carried out as described [41] with some modifications. Briefly, HEK293T cells were lysed with lysis buffer [10 mM HEPES (pH 7.5), 150 mM NaCl, 50 mM NaF, 1 mM Na₃VO₄, 5 μ g/ml leupeptin, 5 μ g/ml aprotinin, 3 μ g/ml pepstatin A, 1 mM phenylmethylsulfonyl fluoride (PMSF), and 1 mg/ml digitonin] and cleared by centrifugation. The cleared lysate was incubated with indicated amounts of FLAG-tagged bait RNA, antisense oligos, and FLAG-M2-conjugated agarose for 1 h. The agarose resin was then washed three times with wash buffer [10 mM HEPES (pH 7.5), 150 mM NaCl, and 0.1% Triton X-100], and co-immunoprecipitated RNA and proteins were eluted with FLAG elution buffer [0.5 mg/ml FLAG peptide, 10 mM HEPES (pH 7.5), 150 mM NaCl, and 0.05% Triton X-100]. The bait RNA-associated proteins were digested with lysyl endopeptidase and trypsin. Digested peptide mixture was applied to a Mightysil-PR-18 (Kanto Chemical) frit-less column (45 \times 3 0.150 mm ID) and separated using a 0–40% gradient of acetonitrile containing 0.1% formic

acid for 80 min at a flow rate of 100 nl/min. Eluted peptides were sprayed directly into a mass spectrometer (Triple TOF 5600+; AB Sciex). MS and MS/MS spectra were obtained using the information-dependent mode. Up to 25 precursor ions above an intensity threshold of 50 counts/s were selected for MS/MS analyses from each survey scan. All MS/MS spectra were searched against protein sequences of RefSeq (NCBI) human protein database using the Protein Pilot software package (AB Sciex), and its decoy sequences then selected the peptides (FDR < 1%). Ion intensity of peptide peaks was obtained using Progenesis Q1 for proteomics software (version 3 Nonlinear Dynamics, UK) according to the manufacturer's instructions.

Ribosome profiling and RNA-seq

Ribosome profiling was performed according to the method previously described with following modifications [34]. RNA concentration of naïve HEK293T lysate was measured by Qubit RNA BR Assay Kit (Thermo Fisher Scientific). The lysate containing 10 μ g RNA was treated with 20 U of RNase I (Lucigen) for 45 min at 25°C. After ribosomes were recovered by ultracentrifugation, RNA fragments corresponding to 26–34 nt were excised from footprint fragment purification gel. Library length distribution was checked using a microchip electrophoresis system (MultiNA, MCE-202, Shimadzu).

For RNA-seq, total RNA was extracted from the lysate using TRIzol LS reagent (Thermo Fisher Scientific) and Direct-zol RNA Kit (Zymo Research). Ribosomal RNA was depleted using the Ribo-Zero Gold rRNA Removal Kit (Human/Mouse/Rat) (Illumina), and the RNA-seq library was prepared using TruSeq Stranded mRNA Library Prep Kit (Illumina) according to the manufacturer's instructions.

The libraries were sequenced on a HiSeq 4000 (Illumina) with a single-end 50 bp sequencing run. Reads were aligned to human hg38 genome as described [34,88]. The offsets of A site from the 5' end of ribosome footprints were determined empirically as 15 for 25–30 nt, 16 for 31–32 nt, and 17 for 33 nt. For RNA-seq, offsets were set to 15 for all mRNA fragments. For calculation of the ribosome occupancies, mRNAs with lower than one footprint per codon were excluded. For calculation of the translation efficiencies (TEs), we counted the number of reads within each CDS, and ribosome profiling counts were normalized by RNA-seq counts using the DESeq package [89]. Reads corresponding to the first and last five codons of each CDS were omitted from the analysis of TEs. The Custom R Scripts will be available upon requests.

Bioinformatics and computational analyses

Principal component analysis

To calculate the codon frequencies of individual genes from *H. sapiens*, we first downloaded coding sequences (CDS) data (human genes, GRCh38p12) from the Ensembl BioMart Database. For each CDS, we tabulated the occurrences of each codon – excluding the stop codons. We then expressed the codon counts as a percentage of the total number of codons in its CDS to obtain the codon frequencies for each CDS. The codon frequencies for all 9,666 CDS were used as the input for the PCA using

the Python 3.4 environment via the factextra program [90]. Finally, the data were trimmed to remove truncated sequences as well as sequences with non-canonical start codons to a final of 9,898 genes.

Hierarchical clustering analysis

mRNA transcripts ranked in order of their half-lives divided equally into four groups, and their average half-lives within each group were calculated. The corresponding codon frequencies of transcripts within each group were averaged. Hierarchical clustering was performed using the average linkage method to cluster the codon frequencies in R using the ggplot2 program [91].

Quantification of GC3-content

To quantify GC3-content, we summed up the codon frequencies of GC3 codons and expressed the frequencies on a percentage scale.

Calculation of cAI and CSC

cAI values were calculated using the standalone CAIcal program [92] in which the human mean codon usage dataset obtained from the Kazusa Codon Usage Database [93] was used as the reference set. The CSC was calculated as described by Presnyak and colleagues [14] using the HEK293 mRNA stability dataset (GSE69153) [Data ref: 31,32].

Binning of ribosomal occupancy frequencies and calculation of codon bias-derived occupancy scores

To quantify codon bias for ribosome profiling, the factor loading scores of the codons from the first principal component were normalized linearly on a percentage scale from 0 to 1, where 0 corresponded to the codon with the lowest score (AAT) and 1 to the codon with the highest score (GCC) (Fig EV2A). Binning of the ribosome occupancies was performed in the R environment via a custom script. To calculate the corresponding codon bias-derived occupancy scores, we substituted the codon sequences of mRNA transcripts with their respective codon scores and, in a similar fashion, binned the data into 25 bins. As the scores of codons should inversely reflect the ribosome occupancy (i.e., higher ribosome occupancy associated with lower codon scores), we calculated the reciprocal of the binned codon scores within each bin for all 25 bins to derive the codon bias-derived occupancy scores. Both ribosome occupancy and codon bias-derived occupancy scores were normalized on a linear scale, and a Pearson correlation was performed on each transcript. To exclude the possibility that the correlations were due to chance, we shuffled the bins for the codon bias-derived occupancy scores within each individual transcripts and calculated the Pearson correlation between shuffled and ribosomal occupancy data.

De novo motif discovery

Common transcripts that were more than fivefold differentially upregulated between the RIP-seq data [Data ref: 43,44] in JLN3 and H929 cells were firstly identified. The corresponding cDNA sequences of the transcripts were downloaded from the UCSC table browser, with the option of masking repeats in the sequences [94]. The sequences were subject to *de novo* motif discovery via the MEME (Multiple EM for Motif Elicitation) software under the MEME tool suite of programs [46].

Data availability

Ribosome profiling and RNA-seq results of HEK293 cells have been deposited at GEO and can be accessed under dataset GSE126298 (<https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE126298>).

Expanded View for this article is available online.

Acknowledgements

The authors express their gratitude to all members of the laboratory of Medical Chemistry, Kyoto University, for their kind advice and discussions. DNA libraries were sequenced by the Vincent J. Coates Genomics Sequencing Laboratory at UC Berkeley, supported by NIH S10 OD018174 Instrumentation Grant. Computations were supported by Manabu Ishii, Itoshi Nikaido, and the Bioinformatics Analysis Environment Service on RIKEN Cloud at RIKEN ACCC. This work was supported by the JSPS KAKENHI (18H05278), AMED-CREST from Japan Agency for Medical Research and Development, and the JSPS through Core-to-Core Program. This work was supported by Joint Usage/Research Center program of Institute for Frontier Life and Medical Sciences, Takeda Science Foundation, the Uehara Memorial Foundation. S.I. was supported by Grant-in-Aid for Scientific Research on Innovative Areas "nascent chain biology" (JP17H05679) and Grant-in-Aid for Young Scientists (A) (JP17H04998) from JSPS, the Pioneering Projects ("Cellular Evolution") and the Aging Project from RIKEN, and Takeda Science Foundation.

Author contributions

FH wrote the manuscript and together with SFY performed the experiments and analyzed the data. MY provided insightful comments and proofreading for the manuscript. YM and ML performed the mRNA decay experiments. YS and SI performed the ribosomal profiling and proofreading of the manuscript. SA and TN performed the ISRM experiments. AV provided advice and bioinformatics expertise. AF and TF performed the polysome profiling experiments. OT supervised and designed the experiments.

Conflict of interest

The authors declare that they have no conflict of interest.

References

- Huang L, Lou C-H, Chan W, Shum EY, Shao A, Stone E, Karam R, Song H-W, Wilkinson MF (2011) RNA homeostasis governed by cell type-specific and branched feedback loops acting on NMD. *Mol Cell* 43: 950–961
- Mino T, Murakawa Y, Fukao A, Vandenbon A, Wessels H-H, Ori D, Uehata T, Tarte S, Akira S, Suzuki Y *et al* (2015) Regnase-1 and roquin regulate a common element in inflammatory mRNAs by spatiotemporally distinct mechanisms. *Cell* 161: 1058–1073
- Yoshinaga M, Nakatsuka Y, Vandenbon A, Ori D, Uehata T, Tsujimura T, Suzuki Y, Mino T, Takeuchi O (2017) Regnase-1 maintains iron homeostasis via the degradation of transferrin receptor 1 and prolyl-hydroxylase-domain-containing protein 3 mRNAs. *Cell Rep* 19: 1614–1630
- Leppik K, Das R, Barna M (2018) Functional 5' UTR mRNA structures in eukaryotic translation regulation and how to find them. *Nat Rev Mol Cell Biol* 19: 158–174
- Cheng J, Maier KC, Avsec Ž, Rus P, Gagneur J (2017) Cis-regulatory elements explain most of the mRNA stability variation across genes in yeast. *RNA* 23: 1648–1659

6. Vogel C, Marcotte EM (2012) Insights into the regulation of protein abundance from proteomic and transcriptomic analyses. *Nat Rev Genet* 13: 227–232
7. Zhou T, Weems M, Wilke CO (2009) Translationally optimal codons associate with structurally sensitive sites in proteins. *Mol Biol Evol* 26: 1571–1580
8. Sharp PM, Li WH (1987) The codon adaptation index—a measure of directional synonymous codon usage bias, and its potential applications. *Nucleic Acids Res* 15: 1281–1295
9. dos Reis M, Savva R, Wernisch L (2004) Solving the riddle of codon usage preferences: a test for translational selection. *Nucleic Acids Res* 32: 5036–5044
10. dos Reis M, Wernisch L, Savva R (2003) Unexpected correlations between gene expression and codon usage bias from microarray data for the whole *Escherichia coli* K-12 genome. *Nucleic Acids Res* 31: 6976–6985
11. Pechmann S, Frydman J (2013) Evolutionary conservation of codon optimality reveals hidden signatures of co-translational folding. *Nat Struct Mol Biol* 20: 237–243
12. Dana A, Tuller T (2014) Mean of the typical decoding rates: a new translation efficiency index based on the analysis of ribosome profiling data. *G3* 5: 73–80
13. Sabi R, Volvovitch Daniel R, Tuller T (2017) stAlcalc: tRNA adaptation index calculator based on species-specific weights. *Bioinformatics* 33: 589–591
14. Presnyak V, Alhusaini N, Chen Y-H, Martin S, Morris N, Kline N, Olson S, Weinberg D, Baker KE, Graveley BR et al (2015) Codon optimality is a major determinant of mRNA stability. *Cell* 160: 1111–1124
15. Radhakrishnan A, Chen Y-H, Martin S, Alhusaini N, Green R, Collier J (2016) The DEAD-box protein Dhh1p couples mRNA decay and translation by monitoring codon optimality. *Cell* 167: 122–132.e9
16. Sweet T, Kovalak C, Collier J (2012) The DEAD-box protein Dhh1 promotes decapping by slowing ribosome movement. *PLoS Biol* 10: e1001342
17. Dana A, Tuller T (2014) The effect of tRNA levels on decoding times of mRNA codons. *Nucleic Acids Res* 42: 9171–9181
18. Gardin J, Yeasmin R, Yurovsky A, Cai Y, Skiena S, Futcher B (2014) Measurement of average decoding rates of the 61 sense codons *in vivo*. *Elife* 3: e03735
19. Drummond DA, Wilke CO (2008) Mistranslation-induced protein misfolding as a dominant constraint on coding-sequence evolution. *Cell* 134: 341–352
20. Akashi H (1994) Synonymous codon usage in *Drosophila melanogaster*: natural selection and translational accuracy. *Genetics* 136: 927–935
21. Harigaya Y, Parker R (2016) Analysis of the association between codon optimality and mRNA stability in *Schizosaccharomyces pombe*. *BMC Genom* 17: 895
22. Lee Y, Zhou T, Tartaglia GG, Vendruscolo M, Wilke CO (2010) Translationally optimal codons associate with aggregation-prone sites in proteins. *Proteomics* 10: 4163–4171
23. Mishima Y, Tomari Y (2016) Codon usage and 3'UTR length determine maternal mRNA stability in zebrafish. *Mol Cell* 61: 874–885
24. Burow DA, Martin S, Quail JF, Alhusaini N, Collier J, Cleary MD (2018) Attenuated codon optimality contributes to neural-specific mRNA decay in *Drosophila*. *Cell Rep* 24: 1704–1712
25. Boël G, Letso R, Neely H, Price WN, Wong K-H, Su M, Luff J, Valecha M, Everett JK, Acton TB et al (2016) Codon influence on protein expression in *E. coli* correlates with mRNA levels. *Nature* 529: 358–363
26. Jeacock L, Faria J, Horn D (2018) Codon usage bias controls mRNA and protein abundance in trypanosomatids. *Elife* 7: e32496
27. de Freitas Nascimento J, Kelly S, Sunter J, Carrington M (2018) Codon choice directs constitutive mRNA levels in trypanosomes. *Elife* 7: e32467
28. Wu Q, Medina SG, Kushawah G, DeVore ML, Castellano LA, Hand JM, Wright M, Bazzini AA (2019) Translation affects mRNA stability in a codon-dependent manner in human cells. *Elife* 8: e45396
29. Forrest ME, Narula A, Sweet TJ, Arango D, Hanson G, Ellis J, Oberdoerffer S, Collier J, Rissland OS (2018) Codon usage and amino acid identity are major determinants of mRNA stability in humans. *bioRxiv* <https://doi.org/10.1101/488676> [PREPRINT]
30. Zerbino DR, Achuthan P, Akanni W, Amode MR, Barrell D, Bhai J, Billis K, Cummins C, Gall A, Girón CG et al (2018) Ensembl 2018. *Nucleic Acids Res* 46: D754–D761
31. Murakawa Y, Hinz M, Mothes J, Schuetz A, Uhl M, Wyler E, Yasuda T, Mastrobuoni G, Friedel CC, Dölken L et al (2015) Gene Expression Omnibus GSE69153 (<https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE69153>). [DATASET]
32. Murakawa Y, Hinz M, Mothes J, Schuetz A, Uhl M, Wyler E, Yasuda T, Mastrobuoni G, Friedel CC, Dölken L et al (2015) RC3H1 post-transcriptionally regulates A20 mRNA and modulates the activity of the IKK/NF- κ B pathway. *Nat Commun* 6: 7367
33. Bazzini AA, Del Viso F, Moreno-Mateos MA, Johnstone TG, Vejnar CE, Qin Y, Yao J, Khokha MK, Giraldez AJ (2016) Codon identity regulates mRNA stability and translation efficiency during the maternal-to-zygotic transition. *EMBO J* 35: 2087–2103
34. McGlincy NJ, Ingolia NT (2017) Transcriptome-wide measurement of translation by ribosome profiling. *Methods* 126: 112–129
35. Tuller T, Kupiec M, Ruppin E (2007) Determinants of protein abundance and translation efficiency in *S. cerevisiae*. *PLoS Comput Biol* 3: e248
36. Iwasaki S, Ingolia NT (2016) Seeing translation. *Science* 352: 1391–1392
37. Ingolia NT, Ghaemmaghami S, Newman JRS, Weissman JS (2009) Genome-wide analysis *in vivo* of translation with nucleotide resolution using ribosome profiling. *Science* 324: 218–223
38. Duncan CDS, Mata J (2017) Effects of cycloheximide on the interpretation of ribosome profiling experiments in *Schizosaccharomyces pombe*. *Sci Rep* 7: 1–11
39. Gerashchenko MV, Gladyshev VN (2014) Translation inhibitors cause abnormalities in ribosome profiling experiments. *Nucleic Acids Res* 42: e134
40. Santos DA, Shi L, Tu BP, Weissman JS (2019) Cycloheximide can distort measurements of mRNA levels and translation efficiency. *Nucleic Acids Res* 47: 4974–4985
41. Adachi S, Homoto M, Tanaka R, Hioki Y, Murakami H, Suga H, Matsumoto M, Nakayama KI, Hatta T, Iemura S et al (2014) ZFP36L1 and ZFP36L2 control LDLR mRNA stability via the ERK–RSK pathway. *Nucleic Acids Res* 42: 10037–10049
42. Kuwano Y, Pullmann R, Marasa BS, Abdelmohsen K, Lee EK, Yang X, Martindale JL, Zhan M, Gorospe M (2010) NF90 selectively represses the translation of target mRNAs bearing an AU-rich signature motif. *Nucleic Acids Res* 38: 225–238
43. Marchesini M, Ogoti Y, Fiorini E, Aktas Samur A, Nezi L, D'Anca M, Storti P, Samur MK, Ganan-Gomez I, Fulciniti MT et al (2017) Gene Expression Omnibus GSE83662 (<https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE83662>). [DATASET]
44. Marchesini M, Ogoti Y, Fiorini E, Aktas Samur A, Nezi L, D'Anca M, Storti P, Samur MK, Ganan-Gomez I, Fulciniti MT et al (2017) ILF2 is a regulator of RNA splicing and DNA damage response in 1q21-amplified multiple myeloma. *Cancer Cell* 32: 88–100.e6

45. Snyder M (2017) ENCODE Project Experiment ENCSR073QLQ (<https://www.encodeproject.org/experiments/ENCSR073QLQ/>) [DATASET]
46. Bailey TL, Boden M, Buske FA, Frith M, Grant CE, Clementi L, Ren J, Li WW, Noble WS (2009) MEME SUITE: tools for motif discovery and searching. *Nucleic Acids Res* 37: W202–W208
47. Dotu I, Adamson SI, Coleman B, Fournier C, Ricart-Altimiras E, Eyraas E, Chuang JH (2018) SARNAClust: semi-automatic detection of RNA protein binding motifs from immunoprecipitation data. *PLoS Comput Biol* 14: e1006078
48. Zheng G, Qin Y, Clark WC, Dai Q, Yi C, He C, Lambowitz AM, Pan T (2015) Efficient and quantitative high-throughput transfer RNA sequencing. *Nat Methods* 12: 835–837
49. Gingold H, Tehler D, Christoffersen NR, Nielsen MM, Asmar F, Kooistra SM, Christophersen NS, Christensen LL, Borre M, Sørensen KD et al (2014) A dual program for translation regulation in cellular proliferation and differentiation. *Cell* 158: 1281–1292
50. Charneski CA, Hurst LD (2013) Positively charged residues are the major determinants of ribosomal velocity. *PLoS Biol* 11: e1001508
51. Ingolia NT, Lareau LF, Weissman JS (2011) Ribosome profiling of mouse embryonic stem cells reveals the complexity and dynamics of mammalian proteomes. *Cell* 147: 789–802
52. Artieri CG, Fraser HB (2014) Accounting for biases in riboprofiling data indicates a major role for proline in stalling translation. *Genome Res* 24: 2011–2021
53. Lareau LF, Hite DH, Hogan GJ, Brown PO (2014) Distinct stages of the translation elongation cycle revealed by sequencing ribosome-protected mRNA fragments. *Elife* 3: e01257
54. Pop C, Rouskin S, Ingolia NT, Han L, Phizicky EM, Weissman JS, Koller D (2014) Causal signals between codon bias, mRNA structure, and the efficiency of translation and elongation. *Mol Syst Biol* 10: 770
55. Endoh T, Sugimoto N (2016) Mechanical insights into ribosomal progression overcoming RNA G-quadruplex from periodical translation suppression in cells. *Sci Rep* 6: 22719
56. Thandapani P, Song J, Gandin V, Cai Y, Rouleau SG, Garant J-M, Boisvert F-M, Yu Z, Perreault J-P, Topisirovic I et al (2015) Aven recognition of RNA G-quadruplexes regulates translation of the mixed lineage leukemia protooncogenes. *Elife* 4: e06234
57. Pan L, Li Y, Zhang H-Y, Zheng Y, Liu X-L, Hu Z, Wang Y, Wang J, Cai Y-H, Liu Q et al (2017) DHX15 is associated with poor prognosis in acute myeloid leukemia (AML) and regulates cell apoptosis via the NF- κ B signaling pathway. *Oncotarget* 8: 89643–89654
58. Faure G, Ogurtsov AY, Shabalina SA, Koonin EV (2016) Role of mRNA structure in the control of protein folding. *Nucleic Acids Res* 44: 10898–10911
59. Yu C-H, Dang Y, Zhou Z, Wu C, Zhao F, Sachs MS, Liu Y (2015) Codon usage influences the local rate of translation elongation to regulate co-translational protein folding. *Mol Cell* 59: 744–754
60. Tuller T, Veksler-Lublinsky I, Gazit N, Kupiec M, Ruppel E, Ziv-Ukelson M (2011) Composite effects of gene determinants on the translation speed and density of ribosomes. *Genome Biol* 12: R110
61. Edri S, Tuller T (2014) Quantifying the effect of ribosomal density on mRNA stability. *PLoS ONE* 9: e102308
62. Neymotin B, Ettore V, Gresham D (2016) Multiple transcript properties related to translation affect mRNA degradation rates in *Saccharomyces cerevisiae*. *G3* 6: 3475–3483
63. Schwartz DC, Parker R (2000) mRNA decapping in yeast requires dissociation of the cap binding protein, eukaryotic translation initiation factor 4E. *Mol Cell Biol* 20: 7933–7942
64. Chan LY, Mugler CF, Heinrich S, Vallotton P, Weis K (2018) Non-invasive measurement of mRNA decay reveals translation initiation as the major determinant of mRNA stability. *Elife* 7: e32536
65. Schwartz DC, Parker R (1999) Mutations in translation initiation factors lead to increased rates of deadenylation and decapping of mRNAs in *Saccharomyces cerevisiae*. *Mol Cell Biol* 19: 5247–5256
66. Saini P, Eyler DE, Green R, Dever TE (2009) Hypusine-containing protein eIF5A promotes translation elongation. *Nature* 459: 118–121
67. Zhou Z, Dang Y, Zhou M, Li L, Yu C, Fu J, Chen S, Liu Y (2016) Codon usage is an important determinant of gene expression levels largely through its effects on transcription. *Proc Natl Acad Sci USA* 113: E6117–E6125
68. Zhou Z, Dang Y, Zhou M, Yuan H, Liu Y (2018) Codon usage biases co-evolve with transcription termination machinery to suppress premature cleavage and polyadenylation. *Elife* 7: e33569
69. Kudla G, Lipinski L, Caffin F, Helwak A, Zylicz M (2006) High guanine and cytosine content increases mRNA levels in mammalian cells. *PLoS Biol* 4: e180
70. Newman ZR, Young JM, Ingolia NT, Barton GM (2016) Differences in codon bias and GC content contribute to the balanced expression of TLR7 and TLR9. *Proc Natl Acad Sci USA* 113: E1362–E1371
71. Fu J, Dang Y, Counter C, Liu Y (2018) Codon usage regulates human KRAS expression at both transcriptional and translational levels. *J Biol Chem* 293: 17929–17940
72. Lampson BL, Pershing NLK, Prinz JA, Lacsina JR, Marzluff WF, Nicchitta CV, MacAlpine DM, Counter CM (2013) Rare codons regulate KRas oncogenesis. *Curr Biol* 23: 70–75
73. Freimer JW, Hu TJ, Billeloch R (2018) Decoupling the impact of microRNAs on translational repression versus RNA degradation in embryonic stem cells. *Elife* 7: e38014
74. Grinberg-Bleyer Y, Oh H, Desrichard A, Bhatt DM, Caron R, Chan TA, Schmid RM, Hayden MS, Klein U, Ghosh S (2017) NF- κ B c-Rel is crucial for the regulatory T cell immune checkpoint in cancer. *Cell* 170: 1096–1108.e13
75. Oh H, Grinberg-Bleyer Y, Liao W, Maloney D, Wang P, Wu Z, Wang J, Bhatt DM, Heise N, Schmid RM et al (2017) An NF- κ B transcription-factor-dependent lineage-specific transcriptional program promotes regulatory T cell identity and function. *Immunity* 47: 450–465.e5
76. Gumienny R, Zavolan M (2015) Accurate transcriptome-wide prediction of microRNA targets and small interfering RNA off-targets with MIRZA-G. *Nucleic Acids Res* 43: 9095
77. Zafrir Z, Tuller T (2015) Nucleotide sequence composition adjacent to intronic splice sites improves splicing efficiency via its effect on pre-mRNA local folding in fungi. *RNA* 21: 1704–1718
78. Amit M, Donyo M, Hollander D, Goren A, Kim E, Gelfman S, Lev-Maor G, Burstein D, Schwartz S, Postolsky B et al (2012) Differential GC content between exons and introns establishes distinct strategies of splice-site recognition. *Cell Rep* 1: 543–556
79. Zhang J, Kuo CCJ, Chen L (2011) GC content around splice sites affects splicing through pre-mRNA secondary structures. *BMC Genom* 12: 90
80. Parrott AM, Walsh MR, Mathews MB (2007) Analysis of RNA: protein interactions *in vivo*: identification of RNA-binding partners of nuclear factor 90. *Meth Enzymol* 429: 243–260
81. Zhou Z, Licklider LJ, Gygi SP, Reed R (2002) Comprehensive proteomic analysis of the human spliceosome. *Nature* 419: 182–185
82. Pfeifer I, Elsby R, Fernandez M, Faria PA, Nussenzweig DR, Lossos IS, Fontoura BMA, Martin WD, Barber GN (2008) NFAR-1 and -2 modulate

- translation and are required for efficient host defense. *Proc Natl Acad Sci USA* 105: 4173–4178
83. Guan D, Altan-Bonnet N, Parrott AM, Arrigo CJ, Li Q, Khaleduzzaman M, Li H, Lee C-G, Pe'ery T, Mathews MB (2008) Nuclear factor 45 (NF45) is a regulatory subunit of complexes with NF90/110 involved in mitotic control. *Mol Cell Biol* 28: 4629–4641
 84. Harashima A, Guettouche T, Barber GN (2010) Phosphorylation of the NFAR proteins by the dsRNA-dependent protein kinase PKR constitutes a novel mechanism of translational regulation and cellular defense. *Genes Dev* 24: 2640–2653
 85. Wolkowicz UM, Cook AG (2012) NF45 dimerizes with NF90, Zfr and SPNR via a conserved domain that has a nucleotidyltransferase fold. *Nucleic Acids Res* 40: 9356–9368
 86. Graber T, Baird S, Kao P, Mathews M, Holcik M (2010) NF45 functions as an IRES trans-acting factor that is required for translation of cIAP1 during the unfolded protein response. *Cell Death Differ* 17: 719–729
 87. Gratacós FM, Brewer G (2010) The role of AUF1 in regulated mRNA decay. *Wiley Interdiscip Rev RNA* 1: 457–473
 88. Akichika S, Hirano S, Shichino Y, Suzuki T, Nishimasu H, Ishitani R, Sugita A, Hirose Y, Iwasaki S, Nureki O et al (2019) Cap-specific terminal N6-methylation of RNA by an RNA polymerase II-associated methyltransferase. *Science* 363: eaav0080
 89. Anders S, Huber W (2010) Differential expression analysis for sequence count data. *Genome Biol* 11: R106
 90. Kassambara A, Mundt F (2017) factoextra: extract and visualize the results of multivariate data analyses.
 91. Wickham H, Chang W, Henry L, Pedersen TL, Takahashi K, Wilke C, Woo K, RStudio (2018) ggplot2: create elegant data visualisations using the grammar of graphics
 92. Puigbò P, Bravo IG, Garcia-Vallve S (2008) CAIcal: a combined set of tools to assess codon usage adaptation. *Biol Direct* 3: 38
 93. Nakamura Y, Gojobori T, Ikemura T (2000) Codon usage tabulated from international DNA sequence databases: status for the year 2000. *Nucleic Acids Res* 28: 292
 94. Karolchik D, Hinrichs AS, Furey TS, Roskin KM, Sugnet CW, Haussler D, Kent WJ (2004) The UCSC table browser data retrieval tool. *Nucleic Acids Res* 32: D493–D496