

POINT Do next-generation sequencing results drive diagnostic and therapeutic decisions in MDS?

Felicitas Thol¹ and Uwe Platzbecker²

¹Department of Hematology, Hemostasis, Oncology, and Stem Cell Transplantation, Hannover Medical School, Hannover, Germany; and ²Medical Clinic and Policlinic 1, Hematology and Cellular Therapy, University Hospital Leipzig, Leipzig, Germany

This article has a companion Counterpoint by Sanz et al.

Pro (but with a “little but”)

Over the past 10 years, next-generation sequencing (NGS) has allowed major advances in unraveling the complex and diverse molecular signature of myelodysplastic syndromes (MDS).¹⁻⁴ As a result, over 50 recurrent mutations have been identified in MDS.^{1,2} In the majority of cases, at least 1 mutation (frequently 2-4) can be identified with targeted sequencing in an individual MDS patient.^{1,2} There is not one single mutation that is 100% pathognomonic for MDS, and we have learned that mutations in MDS cluster in functional genetic groups. While NGS has undoubtedly played a relevant role in the identification of novel mutations in MDS, it is legitimate to ask whether and how NGS can be beneficial for an individual MDS patient. Should we perform NGS for our MDS patients in routine clinical practice, and if so, when and at which technical level? Our answer is yes (in the vast majority of patients) and includes some caveats that will be discussed.

Diagnosing MDS

Conventional diagnosis of MDS is based on cytology, histology, and cytogenetics. So far, NGS cannot replace this diagnostic workup and should not serve as a “stand-alone” diagnostic tool. However, the mutational status of *SF3B1* is now included as a parameter of the revised World Health Organization (WHO) classification of myeloid neoplasms and is therefore recommended in all low-risk cases given the excellent prognosis.⁵ In case a *SF3B1* mutation is present, only 5% of ring sideroblasts (RS) instead of 15% need to be detected in order to diagnose MDS with RS.⁵ Appropriate conventional morphological and cytogenetic workup is mandatory for patients with cytopenia of unknown origin. Nevertheless, NGS can add helpful diagnostic information (Figure 1). For example, if a bone marrow aspirate or biopsy specimen reveals mild cytopenia and dysplasia, it can be challenging even for experienced morphologists to distinguish reactive changes from true MDS.⁶ Here, NGS including a large panel of genes can help to prove or exclude a clonal disease including clonal cytopenia of undetermined significance, which can have a significant impact on subsequent surveillance strategies.⁷ Additionally, clonal hematopoiesis of indeterminate potential (CHIP), which describes a novel entity in which somatic mutations associated with myeloid malignancies are found in blood or bone marrow cells of people, where other criteria for hematologic neoplasia (eg, cytopenia or dysplasia) are not met, needs to be clearly distinguished from MDS.⁸⁻¹⁰ Besides MDS defining criteria (eg, cytopenia or dysplasia), a greater number of mutations with a higher variant allelic frequency (VAF) are more molecularly supportive for the diagnosis of MDS than CHIP.¹¹ The discrimination of hypoplastic MDS from aplastic anemia (AA) can be difficult (eg, mutations in *PIGA* are relatively frequent in AA but very rare in MDS).^{12,13} Here, a greater number of mutations with a higher VAF are rather supportive for the diagnosis of MDS. *CEBPa*, *NPM1*, and *FLT3* mutations are acute myeloid leukemia (AML)-defining mutations and might be important to look for in cases with excess of blasts, especially when close to the 20% threshold, where intensive chemotherapy is an option. Identification of a *KIT* mutation can be helpful in suspected cases with systemic mastocytosis with associated hematological disease (SM-AHD) (Figure 2). Similarly, *STAT3* mutational screening can be useful for the detection of large granular lymphocyte leukemia associated cytopenia.¹⁴ It is likely that more molecular markers will find their way into the WHO classification of MDS in the future.

Prognostication of MDS

While the International Prognostic Scoring System (IPSS) and the revised IPSS (IPSS-R) are the gold standard for prognostication in MDS,^{15,16} the presence of some molecular markers incl. *SF3B1* (as noted above) or *TP53*¹ are known to have an independent prognostic effect. In fact, the presence of *TP53* adds to the prognosis of patients with deletion of chromosome 5q (del5q) in low-risk MDS and




Figure 1. Schematic overview of the NGS workup and potential clinical consequences in case of confirmed MDS by conventional techniques. AML-defining mutations include *CEBPA*, *NPM1*, and *FLT3-ITD*. HSCTx, allogeneic hematopoietic stem cell transplantation (possibly with preceding chemotherapy); *SF3B1*, mutations in *SF3B1*; *TP53*, mutations in *TP53*. *Currently not approved.

complex karyotype in high-risk MDS.^{17,18} Therefore, we recommend routine testing of the latter in patients with low-risk MDS (*SF3B1* all patients and *TP53* in del5q only) or high-risk disease (*TP53* in complex karyotype). Routine testing of other poor-risk mutations like *RUNX1* or *ASXL1* might help to further identify patients with a worse prognosis of all stages. We perform testing regularly in

lower-risk cases where an intensification of disease surveillance followed by putative allogeneic hematopoietic stem cell transplantation (HSCT) might be an option. NGS techniques can analyze a large number of genes, almost at the same cost irrespective of the gene number. Therefore, this is often the preferred sequencing method, as it gives additional information about the number of




Figure 2. Schematic overview of mutations in MDS as potential therapeutic targets. CMML, chronic myelomonocytic leukemia; sAML, secondary acute myeloid leukemia; TKIs, tyrosine kinase inhibitors.

mutated genes, which has been shown to be prognostic (ie, the more mutated genes a patient has, the worse the outcome).^{1,2} Ongoing efforts by an international MDS consortium try to integrate molecular data within the IPSS and IPSS-R. Furthermore, NGS has been used to predict the AML risk in healthy individuals with clonal hematopoiesis.¹⁹

Prediction of response and outcome to conventional therapy

Currently, treatment options in MDS are limited and include growth factors, lenalidomide, hypomethylating agents (HMAs), and HSCT. The number of mutations has been linked to response to erythropoiesis-stimulating agent (ESA) in low-risk MDS.²⁰ Luspatercept is a novel agent used to treat anemia in low-risk MDS patients failing ESA treatment.²¹ The drug directly targets the transforming growth factor β pathway. In a phase 2 trial, *SF3B1*-mutated patients were more likely to respond to luspatercept.²¹ While lenalidomide is standard for patients with isolated del5q, there is evidence that patients with del5q who also harbor a *TP53* mutation (up to 20%) are less likely to respond to lenalidomide and have a worse prognosis.²²

Interestingly, retrospective data in higher-risk MDS and AML patients treated with dose-intense decitabine (20 mg/m² over 10 days) suggest that patients with *TP53* mutations were more likely to respond to this intensive epigenetic treatment than patients with other mutations.²³ Furthermore, *TET2* mutational status has been identified to be predictive for response to standard HMAs, although the data are conflicting.²⁴⁻²⁶ This is also because the higher response rate did not translate into a survival benefit for *TET2*-mutated patients,^{24,25} and again, prospective data are missing. Therefore, we would not recommend routine testing before treatment with HMAs only. Last, but not least, the outcome of HSCT can be impacted by the molecular profile of a patient. While retrospective data of smaller cohorts have led to conflicting results, data from larger cohorts revealed *TP53* and *RAS* mutations as independent adverse markers in HSCT.²⁷⁻²⁹ Patients with a high-risk mutational profile might benefit the most from intensified posttransplant surveillance strategies.³⁰

Novel targeted therapy

Precision oncology, defined as molecular profiling of tumors to identify targetable alterations, is very appealing, as it combines the hopes for a more effective treatment with less toxicity compared with conventional treatment. Due to the molecular heterogeneity of MDS, a potential target needs to be identified individually in each MDS patient. In MDS, targeted therapies are just beginning to emerge (eg, *IDH1/2* inhibitors).³¹ In a phase 1/2 trial, enasidenib (AG221), a selective *IDH2* mutant inhibitor, was used as monotherapy in refractory/relapsed AML patients, leading to an overall response rate of 38%.³² In MDS, ~4% to 5% of patients carry an *IDH2* mutation, and 2% to 3% carry an *IDH1* mutation.^{1,2,33} *IDH1* and *IDH2* inhibitors are currently not approved for AML or MDS patients in Europe, but enasidenib has received approval by the US Food and Drug Administration for elderly relapsed/refractory AML patients. It is likely that the response data can be transferred to patients with MDS, especially when progressing to sAML. However, several trials for *IDH1*- and *IDH2*-mutant AML and MDS patients are currently ongoing, making molecular screening with NGS necessary for patient enrollment. Other examples of potential targeted therapies are FLT3-ITD inhibitors in sAML or so-called

splicing modulators in lower-risk MDS. Splicing gene mutations are among the most frequent mutations found in MDS (eg, *SF3B1*, *SRSF2*, *U2AF1*, and *ZRSR2* mutations).^{1,2,34} Splicing modulators are intended to induce “splicesomal sickness” and consequently cell death. Currently, treatment with splicing modulators is only possible within clinical trials, and it will be interesting to see the outcome of these trials. Other ongoing “targeted approaches” involve JAK-2 inhibitors in MDS/myeloproliferative neoplasm cases and as well as KIT modulation in SM-AHD. It will be interesting to see whether mutated *TP53* proves to be a promising target, as early results with Prima-1Met (APR-246), an activator of *TP53* function, suggest.³⁵ In general, clinical trials with targeted therapies are currently still rare and only possible with the availability of NGS.

Further aspects

Over the past decade, we have gained novel insights into the genetic origin of inherited myeloid malignancies.^{36,37} Mutations in *GATA2*, *RUNX1*, *DDX41*, and *ETV6* are examples of germline mutations associated with MDS and AML.^{36,37} The detection of germline mutations has a significant impact not only for treatment decisions but also for the patient’s family. Obtaining a detailed family history, testing germline material (eg, skin fibroblasts), and genetic counseling of patients and families should go hand in hand. Last, but not least, NGS has evolved as a very attractive tool for measurable residual disease (MRD) monitoring after chemotherapy or HSCT.³⁸⁻⁴⁰ We acknowledge that access to NGS is limited in some countries and is also associated with a clear economic burden. Thus, a targeted sequencing approach that accounts for the diagnostic/clinical scenario can be sensible, keeping the potential therapeutic implications in mind (Figures 1 and 2). Besides reporting the type and location of the mutation, the VAF should also be included in NGS report. Is NGS necessary for every MDS patient? We believe that NGS should be performed in the vast majority of MDS patients also because we as clinicians have to learn the impact and relevance of this clinical information on a patient-by-patient basis. However, NGS testing outside clinical trial options is not helpful for patients who have a very short life expectancy related to another underlying critical disease and/or very advanced disease.

In summary, we would advocate for NGS (commonly with targeted approach) in the vast majority of MDS patients, as NGS results can be helpful with the diagnosis, prognostication, and treatment response prediction in MDS patients (Figures 1 and 2). In addition, NGS opens the opportunity of targeted therapies (in clinical trials). The relevance of NGS is likely to increase in the next few years with further implications such as MRD monitoring. With NGS becoming cheaper over the years, our “little but” will become even smaller.

Acknowledgments

U.P. received support from Deutsche Jose-Carreras Leukämienstiftung and Boll-Stiftung.

Authorship

Contribution: U.P. and F.T. both performed a literature search and wrote the paper.

Conflict-of-interest disclosure: The authors declare no competing financial interests.

Correspondence: Uwe Platzbecker, Medical Clinic and Polyclinic 1, Hematology and Cellular Therapy, Leipzig University

References

1. Papaemmanuil E, Gerstung M, Malcovati L, et al; Chronic Myeloid Disorders Working Group of the International Cancer Genome Consortium. Clinical and biological implications of driver mutations in myelodysplastic syndromes. *Blood*. 2013; 122(22):3616-3627, quiz 3699.
2. Haferlach T, Nagata Y, Grossmann V, et al. Landscape of genetic lesions in 944 patients with myelodysplastic syndromes. *Leukemia*. 2014;28(2):241-247.
3. Papaemmanuil E, Cazzola M, Boulton J, et al; Chronic Myeloid Disorders Working Group of the International Cancer Genome Consortium. Somatic SF3B1 mutation in myelodysplasia with ring sideroblasts. *N Engl J Med*. 2011;365(15): 1384-1395.
4. Yoshida K, Sanada M, Shiraishi Y, et al. Frequent pathway mutations of splicing machinery in myelodysplasia. *Nature*. 2011;478(7367):64-69.
5. Arber DA, Orazi A, Hasserjian R, et al. The 2016 revision to the World Health Organization classification of myeloid neoplasms and acute leukemia. *Blood*. 2016;127(20):2391-2405.
6. Mufti GJ, Bennett JM, Goasguen J, et al; International Working Group on Morphology of Myelodysplastic Syndrome. Diagnosis and classification of myelodysplastic syndrome: International Working Group on Morphology of myelodysplastic syndrome (IWGM-MDS) consensus proposals for the definition and enumeration of myeloblasts and ring sideroblasts. *Haematologica*. 2008;93(11):1712-1717.
7. Malcovati L, Galli A, Travaglino E, et al. Clinical significance of somatic mutation in unexplained blood cytopenia. *Blood*. 2017;129(25):3371-3378.
8. Jaiswal S, Fontanillas P, Flannick J, et al. Age-related clonal hematopoiesis associated with adverse outcomes. *N Engl J Med*. 2014;371(26):2488-2498.
9. Xie M, Lu C, Wang J, et al. Age-related mutations associated with clonal hematopoietic expansion and malignancies. *Nat Med*. 2014;20(12):1472-1478.
10. Genovese G, Kähler AK, Handsaker RE, et al. Clonal hematopoiesis and blood-cancer risk inferred from blood DNA sequence. *N Engl J Med*. 2014;371(26):2477-2487.
11. Steensma DP, Bejar R, Jaiswal S, et al. Clonal hematopoiesis of indeterminate potential and its distinction from myelodysplastic syndromes. *Blood*. 2015;126(1):9-16.
12. Yoshizato T, Dumitriu B, Hosokawa K, et al. Somatic mutations and clonal hematopoiesis in aplastic anemia. *N Engl J Med*. 2015;373(1):35-47.
13. Heuser M, Schlarman C, Dobbernack V, et al. Genetic characterization of acquired aplastic anemia by targeted sequencing. *Haematologica*. 2014;99(9):e165-e167.
14. Morgan EA, Lee MN, DeAngelo DJ, et al. Systematic *STAT3* sequencing in patients with unexplained cytopenias identifies unsuspected large granular lymphocytic leukemia. *Blood Adv*. 2017;1(21):1786-1789.
15. Greenberg P, Cox C, LeBeau MM, et al. International scoring system for evaluating prognosis in myelodysplastic syndromes. *Blood*. 1997;89(6):2079-2088.
16. Greenberg PL, Tuechler H, Schanz J, et al. Revised international prognostic scoring system for myelodysplastic syndromes. *Blood*. 2012;120(12):2454-2465.
17. Jädersten M, Saft L, Smith A, et al. TP53 mutations in low-risk myelodysplastic syndromes with del(5q) predict disease progression. *J Clin Oncol*. 2011;29(15):1971-1979.
18. Bejar R, Stevenson K, Abdel-Wahab O, et al. Clinical effect of point mutations in myelodysplastic syndromes. *N Engl J Med*. 2011;364(26):2496-2506.
19. Abelson S, Collord G, Ng SWK, et al. Prediction of acute myeloid leukaemia risk in healthy individuals. *Nature*. 2018; 559(7714):400-404.
20. Kosmider O, Passet M, Santini V, et al; GFM, FISM AND D-MDS. Are somatic mutations predictive of response to erythropoiesis stimulating agents in lower risk myelodysplastic syndromes? *Haematologica*. 2016;101(7):e280-e283.
21. Platzbecker U, Germing U, Götze KS, et al. Luspatercept for the treatment of anaemia in patients with lower-risk myelodysplastic syndromes (PACE-MDS): a multicentre, open-label phase 2 dose-finding study with long-term extension study. *Lancet Oncol*. 2017;18(10):1338-1347.
22. Jädersten M, Saft L, Pellagatti A, et al. Clonal heterogeneity in the 5q- syndrome: p53 expressing progenitors prevail during lenalidomide treatment and expand at disease progression. *Haematologica*. 2009;94(12):1762-1766.
23. Welch JS, Petti AA, Miller CA, et al. TP53 and decitabine in acute myeloid leukemia and myelodysplastic syndromes. *N Engl J Med*. 2016;375(21):2023-2036.
24. Bejar R, Lord A, Stevenson K, et al. TET2 mutations predict response to hypomethylating agents in myelodysplastic syndrome patients. *Blood*. 2014;124(17):2705-2712.
25. Itzykson R, Kosmider O, Cluzeau T, et al; Groupe Francophone des Myelodysplasies (GFM). Impact of TET2 mutations on response rate to azacitidine in myelodysplastic syndromes and low blast count acute myeloid leukemias. *Leukemia*. 2011; 25(7):1147-1152.
26. Traina F, Visconte V, Elson P, et al. Impact of molecular mutations on treatment response to DNMT inhibitors in myelodysplasia and related neoplasms. *Leukemia*. 2014; 28(1):78-87.
27. Lindsley RC, Saber W, Mar BG, et al. Prognostic mutations in myelodysplastic syndrome after stem-cell transplantation. *N Engl J Med*. 2017;376(6):536-547.
28. Yoshizato T, Nannya Y, Atsuta Y, et al. Genetic abnormalities in myelodysplasia and secondary acute myeloid leukemia: impact on outcome of stem cell transplantation. *Blood*. 2017; 129(17):2347-2358.
29. Heuser M, Gabdoulline R, Löffeld P, et al. Individual outcome prediction for myelodysplastic syndrome (MDS) and

- secondary acute myeloid leukemia from MDS after allogeneic hematopoietic cell transplantation. *Ann Hematol.* 2017;96(8): 1361-1372.
30. Platzbecker U, Middeke J, Sockel K, et al. Measurable-residual disease guided treatment with azacitidine to prevent haematological relapse in patients with MDS and AML – results from the multicenter phase 2 RELAZA2 trial. *Lancet Oncol.* 2018;19(12):1668-1679.
 31. Stein EM. IDH2 inhibition in AML: finally progress? *Best Pract Res Clin Haematol.* 2015;28(2-3):112-115.
 32. Stein EM, DiNardo CD, Pollyea DA, et al. Enasidenib in mutant *IDH2* relapsed or refractory acute myeloid leukemia. *Blood.* 2017;130(6):722-731.
 33. Thol F, Weissinger EM, Krauter J, et al. IDH1 mutations in patients with myelodysplastic syndromes are associated with an unfavorable prognosis. *Haematologica.* 2010;95(10): 1668-1674.
 34. Thol F, Kade S, Schlarman C, et al. Frequency and prognostic impact of mutations in *SRSF2*, *U2AF1*, and *ZRSR2* in patients with myelodysplastic syndromes. *Blood.* 2012;119(15): 3578-3584.
 35. Maslah N, Verger E, Partouche N, et al. Synergistic effects of PRIMA-1MET (APR-246) and azacitidine (AZA) in TP53-mutated myelodysplastic syndromes. Paper presented at Annual Congress of the European Haematology Association. 15 June 2018. Stockholm, Sweden.
 36. Feurstein S, Drazer MW, Godley LA. Genetic predisposition to leukemia and other hematologic malignancies. *Semin Oncol.* 2016;43(5):598-608.
 37. DiNardo CD, Bannon SA, Routbort M, et al. Evaluation of patients and families with concern for predispositions to hematologic malignancies within the Hereditary Hematologic Malignancy Clinic (HHMC). *Clin Lymphoma Myeloma Leuk.* 2016;16(7):417-428.e412.
 38. Jongen-Lavrencic M, Grob T, Hanekamp D, et al. Molecular minimal residual disease in acute myeloid leukemia. *N Engl J Med.* 2018;378(13):1189-1199.
 39. Thol F, Gabdoulline R, Liebich A, et al. Measurable residual disease (MRD) monitoring by next-generation sequencing before allogeneic hematopoietic cell transplantation in AML. *Blood.* 2018;132(16):1703-1713.
 40. Duncavage EJ, Jacoby MA, Chang GS, et al. Mutation clearance after transplantation for myelodysplastic syndrome. *N Engl J Med.* 2018;379(11):1028-1041.

DOI 10.1182/bloodadvances.2018022434
 © 2019 by The American Society of Hematology