

Overexpression of *WT1* and *PRAME* predicts poor outcomes of patients with myelodysplastic syndromes with thrombocytopenia

Qiu-Sha Huang,¹⁻³ Jing-Zhi Wang,¹⁻³ Ya-Zhen Qin,¹⁻³ Qiao-Zhu Zeng,¹⁻³ Qian Jiang,¹⁻³ Hao Jiang,¹⁻³ Jin Lu,¹⁻³ Hui-Xin Liu,⁴ Yi Liu,⁵ Jing-Bo Wang,⁶ Li Su,⁷ Hong-Yu Zhang,⁸ Zhen-Ling Li,⁹ Su-Jun Gao,¹⁰ Bo Huang,^{11,12} Yu-Ying Liu,¹¹ Yan-Rong Liu,¹⁻³ Lan-Ping Xu,^{1-3,13} Xiao-Jun Huang,^{1-3,13} and Xiao-Hui Zhang^{1-3,13}

¹Peking University People's Hospital, Peking University Institute of Hematology, Beijing, China; ²National Clinical Research Center for Hematologic Disease, Beijing, China; ³Beijing Key Laboratory of Hematopoietic Stem Cell Transplantation, Beijing, China; ⁴Department of Clinical Epidemiology, Peking University People's Hospital, Beijing, China; ⁵The Sixth Medical Center of People's Liberation Army of China General Hospital, Beijing, China; ⁶Beijing Aerospace General Hospital, Beijing, China; ⁷Xuanwu Hospital Capital Medical University, Beijing, China; ⁸Peking University Shenzhen Hospital, Shenzhen, China; ⁹China-Japan Friendship Hospital, Beijing, China; ¹⁰The First Bethune Hospital of Jilin University, Jilin, China; ¹¹Department of Immunology & National Key Laboratory of Medical Molecular Biology, Institute of Basic Medical Sciences, Chinese Academy of Medical Sciences and Peking Union Medical College, Beijing, China; ¹²Department of Biochemistry & Molecular Biology, Tongji Medical College, Huazhong University of Science & Technology, Wuhan, China; and ¹³Collaborative Innovation Center of Hematology, Peking University, Beijing, China

Key Points

- Overexpression of *WT1* and *PRAME* are common in MDS patients with thrombocytopenia. Both are independent poor prognostic factor for outcome.
- The evaluation of *WT1/PRAME* transcript analysis can better risk-stratify the patients, thus guiding individualized treatment.

Thrombocytopenia is associated with life-threatening bleeding and is common in myelodysplastic syndromes (MDS). Robust molecular prognostic biomarkers need to be developed to improve clinical decision making for patients with MDS with thrombocytopenia. Wilms tumor 1 (*WT1*) and preferentially expressed antigen in melanoma (*PRAME*) are promising immunogenic antigen candidates for immunotherapy, and their clinical effects on patients with MDS with thrombocytopenia are still not well understood. We performed a multicenter observational study of adult patients with MDS with thrombocytopenia from 7 different tertiary medical centers in China. We examined bone marrow samples collected at diagnosis for *WT1* and *PRAME* transcript levels and then analyzed their prognostic effect for patients with MDS with thrombocytopenia. In total, we enrolled 1110 patients diagnosed with MDS with thrombocytopenia. Overexpression of *WT1* and *PRAME* was associated with elevated blast percentage, worse cytogenetics, and higher Revised International Prognostic Scoring System (IPSS-R) risk. Further, both *WT1* and *PRAME* overexpression were independent poor prognostic factors for acute myeloid leukemia evolution, overall survival, and progression-free survival. Together, the 2 genes overexpression identified a population of patients with MDS with substantially worse survival. On the basis of *WT1* and *PRAME* transcript levels, patients with MDS with IPSS-R low risk were classified into 2 significantly divergent prognostic risk groups: a low-favorable group and a low-adverse group. The low-adverse group had survival similar to that of patients in the intermediate-risk group. Our study demonstrates that the evaluation of *WT1/PRAME* transcript analysis may improve the prognostication precision and better risk-stratify the patients.

Introduction

The myelodysplastic syndromes (MDS) comprise a group of clonal bone marrow stem cell disorders characterized by ineffective hematopoiesis, peripheral cytopenias, and propensity to acute myeloid leukemia (AML) transformation.¹⁻³ Thrombocytopenia is common in MDS and leads to significant mortality because of life-threatening bleeding risk and increased risk for transformation to AML.⁴ The initial cytopenia

Submitted 10 June 2019; accepted 11 September 2019. DOI 10.1182/bloodadvances.2019000564.

Data can be shared by request sent to the corresponding author at zhangxh100@sina.com.

The full-text version of this article contains a data supplement.

© 2019 by The American Society of Hematology

presentation of patients with MDS with thrombocytopenia can be variable, including isolated thrombocytopenia, bicytopenia, and pancytopenia. At this time, there is no information available on the differences in clinical profiles and prognosis among different cytopenia subtypes. Especially for patients with MDS with isolated thrombocytopenia, the diagnosis of immune thrombocytopenia rather than MDS may initially occur,⁵⁻⁸ as different kinds of causes, from normal variants to severe congenital and acquired disorders, can be responsible for thrombocytopenia. Given the difficulty of diagnosis coupled with the fact that MDS with thrombocytopenia subtype is poorly described, proper workup of both patient and disease features is essential to provide diagnostic and therapeutic strategy recommendations.

MDS are a group of heterogeneous disorders for which customized treatment plans are tailored to the predicted prognosis, which makes the precise prediction of the prognosis necessary for the management of patients. Current prognostic scoring systems stratify patients with MDS into different risk groups on the basis of clinical/hematological parameters, but not of molecular genetic characteristics. The current scoring systems of MDS need to be improved in the future because of the poor interobserver concordance of morphologic evaluation variables and the fact that approximately 50% of patients lack informative cytogenetic abnormalities.⁹ It is essential that more multicenter prospective studies are devoted to developing robust prognostic factors. Recently, some somatic mutations have been linked to the prognosis of patients with MDS. Although our burgeoning knowledge of gene sequencing could provide novel insights into the prognostic effects of somatic mutations in patients with MDS, no specific mutation has been detected in some patients with MDS, and this approach with low mutation frequency and interpatient variation has been complicated by reports of frequent somatic mutations in the aging healthy population.¹⁰ Significantly, the assessment of associated genes can be accomplished through detecting transcript levels, with the advantages of simplicity and wide applicability in clinical practice.

Wilms tumor 1 (*WT1*) and preferentially expressed antigen in melanoma (*PRAME*) are promising immunogenic antigens candidates for specific immunotherapy.^{11,12} *WT1* is a transcription factor that plays a critical role in regulating myeloid differentiation in hematopoiesis. During hematopoiesis, abnormal expression of *WT1* retards cell proliferation and/or differentiation.¹³⁻¹⁵ *PRAME* encodes a tumor-associated antigen that is preferentially expressed in human melanoma.¹⁶ *PRAME* has been described to be a repressor of retinoic acid (RA) signaling, capable of inhibiting RA-induced hematopoietic differentiation, cell cycle arrest, and apoptosis.¹⁷⁻²⁰ Accumulating evidence has suggested that *WT1* and *PRAME* are overexpressed in many malignant neoplasms, including leukemia and MDS.²¹⁻²⁶ Both *WT1* and *PRAME* have been shown to be associated with MDS clone.^{21,23,27-29} Previous studies have demonstrated the prognostic significance and the usefulness of monitoring minimal residual disease of *WT1* and *PRAME* expression level in patients with acute leukemia.^{23,25,26,30,31} However, the prognostic significance of *WT1* and *PRAME* is not yet well understood in patients with MDS with thrombocytopenia. A combination of current prognostic scoring systems with *WT1* and *PRAME* transcript levels may further guide the design of individualized treatment of patients with MDS. To address the question, transcript levels of *WT1* and *PRAME* were detected from a large

set of samples of patients with MDS with thrombocytopenia to investigate whether *WT1* and *PRAME* can predict long-term prognosis in our study.

Materials and methods

Patients and study design

We conducted a retrospective multicenter analysis in adult patients presenting with thrombocytopenia at the onset of MDS. We have reviewed all the patients who were diagnosed with MDS between July 2007 and February 2018 from 7 different tertiary medical centers in China. The diagnosis of MDS was confirmed on the basis of standard criteria.^{1-3,32,33} Two experts separately assessed all cytogenetic studies to identify the abnormalities. We included 1110 patients with MDS with thrombocytopenia from 6582 patients with MDS. According the number of cytopenia lines at the onset of the disease, 191 patients with isolated thrombocytopenia (group A), 389 patients with bicytopenia (group B), and 530 patients with pancytopenia (group C) were enrolled in this study (as shown in Figure 1). All samples and clinical data were collected with patient consent under an institutional review board-approved protocol in accordance with the Declaration of Helsinki. Patient clinical profiles are shown in Table 1.

RNA extraction, complementary DNA synthesis, and RQ-PCR

We analyzed pretreatment bone marrow samples from adult patients with MDS with thrombocytopenia enrolled in our study. Samples for evaluation were available for 1042 patients for *WT1* transcript levels and 529 patients for *PRAME*. Total RNA extraction, reverse transcription of RNA, and TaqMan-based RQ-PCR technology measurement of *WT1* and *PRAME* transcript were performed as we previously described.^{23,25,26,30,31,34} The house-keeping gene Abelson (ABL) was selected as a control gene for RNA expression. All samples were analyzed in duplicate. The transcript levels were calculated as *WT1* or *PRAME* transcript copy number/ABL copy number in percentage.

Treatment regimen

The regimen included allogeneic hematopoietic stem cell transplantation (allo-HSCT), chemotherapy, hypomethylating agents (HMA), and supportive care, as previously published.³⁵⁻³⁷


Pretransplant conditioning therapy consisted of simustine, busulfan, cyclophosphamide, cytarabine, and rabbit antithymocyte globulin. Short-term methotrexate, mycophenolate mofetil, and cyclosporine A were used for graft-versus-host disease prophylaxis. All HSCT recipients were treated with granulocyte colony stimulating factor-mobilized, fresh, unmanipulated bone marrow cells, and peripheral blood stem cells. Stem cell harvesting, HLA typing, and donor selection were performed as described previously.³⁸

Different intensive therapies were given to 174 patients with *WT1* or *PRAME* overexpression. A total of 124 patients underwent allo-HSCT, and 50 patients only received chemotherapy or HMA.

Definitions

Cytopenia in MDS was defined as: hemoglobin lower than 10 g/dL, platelet count lower than $100 \times 10^9/L$, and absolute neutrophil count lower than $1.8 \times 10^9/L$.^{1-3,32,33,39} Severe thrombocytopenia was

Figure 1. Algorithm of study design and patient selection.



defined as PLT lower than $20 \times 10^9/L$, moderate thrombocytopenia was defined as PLT 20 to $50 \times 10^9/L$, and mild thrombocytopenia was defined as PLT 50 to $100 \times 10^9/L$.^{4,6} Constant cytopenia was defined as at least 6 months unless cytogenetic studies revealed MDS.^{1-3,33} Bicytopenia in our study was defined as blood cell count that satisfied platelet count lower than $100 \times 10^9/L$ and 1 of 2 followed the following criteria: hemoglobin lower than 10 g/dL and absolute neutrophil count lower than $1.8 \times 10^9/L$. Pancytopenia was defined as hemoglobin lower than 10 g/dL, platelet count lower than $100 \times 10^9/L$, and absolute neutrophil count lower than $1.8 \times 10^9/L$. We considered MDS with isolated thrombocytopenia as when the initial neutrophil count was more than $1.8 \times 10^9/L$ and the hemoglobin concentration was more than 100 g/L, but the initial platelet count was less than $100 \times 10^9/L$.^{4,6}

Response criteria were listed in the supplemental Table 1, according to the report of the International Working Group.⁴⁰

Statistical analysis

We summarized patient profiles with frequencies and percentages for categorical variables and medians and interquartile range for continuous variables. We compared the baseline characteristics among groups with a χ^2 test for categorical variables and variance tests (ANOVA) or Kruskal-Wallis H for continuous variables. We measured OS, progression-free survival (PFS), and time to evolution to AML from the time of diagnosis of MDS. We estimated the OS, PFS, and time to evolution to AML with

Kaplan-Meier techniques and used a log-rank test to assess potential differences between subgroups. For univariable and multivariable survival analysis, we used Cox proportional hazards regression to identify the independent prognostic significance of *WT1* and *PRAME* for OS, PFS, and evolution to AML. Only patients with complete information of both *WT1* and *PRAME* transcript levels were included in the multivariable analysis. We constructed Cox models to select variables by enter method, adjusting for clinical and hematological characteristics, with candidate variables having a univariable $P < .10$.

We used receiver operating characteristic (ROC) curves to explore *WT1* and *PRAME* transcript level cutoff values that best differentiated patients with MDS with respect to the higher-risk group. We used the Youden index (sensitivity + specificity - 1) to calculate the optimal thresholds along the receiver operating characteristic curves. We classified a 2-sided $P < .05$ as significant. We carried out statistical analyses using the IBM SPSS software package (version 20.0).

Results

Demographic and clinical features of patients with MDS with thrombocytopenia

We screened 6582 patients with MDS and a total of 1110 patients with MDS with thrombocytopenia (191 isolated thrombocytopenia, 389 bicytopenia, 530 pancytopenia) met the inclusion criteria for

Table 1. Clinical profile of patients with MDS with thrombocytopenia

Feature	Isolated thrombocytopenia (n = 191)	Bicytopenia (n = 389)	Pancytopenia (n = 530)	P
Age at diagnosis, y	56 (37-68)	57 (38.5-66)	53 (38-63)	.056
Sex				.137
Female	95 (49.7)	162 (41.6)	247 (46.6)	
Male	96 (50.3)	227 (58.4)	283 (53.4)	
Cytopenias				
Hb, g/dL	117 (108-131)	94 (69-113)	71 (58-84)	.000
ANC, $\times 10^9/L$	3.0 (2.2-4.7)	1.9 (1.2-3.2)	0.9 (0.5-1.3)	.000
Platelets, $\times 10^9/L$	46 (31-67)	46 (25-70)	35 (18-63)	.000
Severe thrombocytopenia	19 (9.9)	64 (16.5)	152 (28.7)	.000
Moderate thrombocytopenia	87 (45.5)	144 (37.0)	180 (34.0)	
Bone marrow blasts, %	2.5 (1.0-5.0)	4.0 (1.8-8.5)	4.0 (1.2-10.5)	.000
WHO classification				.000
MDS-SLD	26 (13.6)	59 (15.2)	82 (15.5)	
MDS-MLD	9 (4.7)	25 (6.4)	23 (4.3)	
MDS-RS	2 (1.0)	5 (1.3)	6 (1.1)	
5q-	0	0	1 (0.2)	
MDS-EB-1	44 (23.0)	108 (27.8)	103 (19.4)	
MDS-EB-2	19 (9.9)	84 (21.6)	149 (28.1)	
MDS-U	91 (47.6)	108 (27.8)	166 (31.3)	
Cytogenetic risk				
Very good	3 (1.6)	8 (2.1)	7 (1.3)	.037
Good	110 (57.6)	227 (58.4)	291 (54.9)	
Intermediate	63 (33.0)	103 (26.5)	135 (25.5)	
Poor	8 (4.2)	19 (4.9)	38 (7.2)	
Very poor	7 (3.7)	32 (8.2)	59 (11.1)	
MK	13 (6.8)	61 (15.7)	77 (14.5)	.014
IPSS-R				
Very low	25 (13.1)	13 (3.3)	0 (0.0)	.000
Low	85 (44.5)	79 (20.3)	49 (9.2)	
Intermediate	32 (16.8)	144 (37.0)	170 (32.1)	
High	34 (17.8)	91 (23.4)	138 (26.0)	
Very high	15 (7.9)	62 (15.9)	173 (32.6)	
AML evolution	24 (12.6)	61 (15.7)	121 (22.8)	.002

Data are n (%) or median (interquartile range).

5q-, MDS with isolated del(5q); ANC, absolute neutrophil count; Hb, hemoglobin; IPSS-R, Revised International Prognostic Scoring System; MDS-EB, MDS with excess blasts; MDS-MLD, MDS with multilineage dysplasia; MDS-RS, MDS with ring sideroblasts; MDS-SLD, MDS with single-lineage dysplasia; MDS-U, MDS-unclassifiable; MK, monosomal karyotype; WBC, white blood cell.

this study. The demographic details and clinical characteristics were provided in Table 1. In the isolated thrombocytopenia group, the median cell counts at the time of onset were Hb 117 g/dL, ANC $3.0 \times 10^9/L$, and PLT $46 \times 10^9/L$. A total of 118 (61.8%) patients still manifested isolated thrombocytopenia, whereas 45 (23.6%) had progressed to bicytopenia and 28 (14.6%) had progressed to pancytopenia by the time of bone marrow examination diagnosis.

The bone marrow blasts of the isolated thrombocytopenia group were fewer in comparison with the bicytopenia and pancytopenia group (median bone marrow blasts, 2.5%, 4.0%,

and 4.0%; $P = .000$). MDS-U and MDS-EB1 comprised the majority of subtypes in the isolated thrombocytopenia group (representing 47.6% and 23.0% of patients), followed by MDS-SLD (13.6%), MDS-EB2 (9.9%), and MDS-MLD (4.7%). The most frequent World Health Organization (WHO) subtype in patients with bicytopenia and pancytopenia was MDS-EB (49.4% of bicytopenia and 47.5% of pancytopenia). Compared with the other 2 groups, patients with isolated thrombocytopenia had better cytogenetics and lower Revised International Prognostic Scoring System (IPSS-R). With the increase of cytopenia lines, the probability of AML evolution increased (12.6% of isolated thrombocytopenia, 15.7% of bicytopenia, and 22.8% of pancytopenia; $P = .002$; Table 1; Figure 2A).




Figure 2. Probability of AML evolution for patients with MDS according to cytopenia lines and WT1/PRAME transcript levels. (A) Probability of AML evolution for all patients with MDS based on cytopenia lines. (B) Probability of AML evolution for patients with MDS based on WT1 transcript levels. (C) Probability of AML evolution for patients with MDS based on PRAME transcript levels.

Median follow-up of the whole patient cohort was 36 months. OS and PFS for the isolated thrombocytopenia group were longer than the bicytopenia and pancytopenia groups ($P = .005$ for OS

and $P = .008$ for PFS; Figure 3A,D), demonstrating the favorable prognosis.

WT1 and PRAME gene transcript levels

The median *WT1* and *PRAME* expression levels in bone marrow samples from patients with MDS with thrombocytopenia were 2.10% (interquartile range, 0.42%-8.30%) and 1.00% (interquartile range, 0.20%-19.65%). In total, 73.1% of patients with MDS with thrombocytopenia had higher *WT1* transcript levels over the normal range, and 67.3% of patients had higher *PRAME* transcript levels over the normal range, as we previously described.³⁴ *WT1* and *PRAME* were highly expressed in patients with MDS with thrombocytopenia (supplemental Table 2; supplemental Figure 1). The transcript levels of *WT1* and *PRAME* gradually grew with the increase in risk stratification (supplemental Figure 2).

Receiver operating characteristic curve analysis revealed that *WT1* and *PRAME* transcript levels could significantly differentiate patients with respect to higher risk (IPSS-R high risk and very high risk; area under the curve, 0.670 [$P = .000$] for *WT1*; area under the curve, 0.687 [$P = .000$] for *PRAME*). The Youden index indicated that 4.9% and 0.3% were the optimal thresholds for *WT1* and *PRAME*. Therefore, *WT1* transcript levels higher than 4.9% and *PRAME* transcript levels higher than 0.3% were regarded as overexpression in our study (supplemental Figure 3).

The *WT1* overexpression was identified in 27%, 33%, and 39% of patients (difference between isolated thrombocytopenia group and pancytopenia group, $P = .006$), and *PRAME* overexpression was identified in 58%, 66%, and 69% of patients of the isolated thrombocytopenia, bicytopenia, and pancytopenia group, respectively ($P = .061$). The frequency of *PRAME* overexpression was similar among 3 groups.

Clinical associations with WT1 and PRAME

To further correlate the role of *WT1* and *PRAME* transcript levels with prognosis, clinicohematological characteristics were analyzed. For each gene, we compared the clinical features of patients who had *WT1* and *PRAME* overexpression with those of patients who did not. Patients with overexpression of *WT1* and *PRAME* were more likely to have an elevated blast percentage, worse cytogenetics, and higher IPSS-R risk compared with those who had not (Figure 4A,C-D). However, no differences were observed with respect to the severity of thrombocytopenia for both *WT1* and *PRAME* (supplemental Figure 4).

AML evolution association with WT1 and PRAME

We also analyzed the correlation of *WT1* and *PRAME* transcript levels with the risk for AML evolution. Evolution to AML was seen in 206 patients (18.6%), including 9 patients with very low or low-risk disease, 30 patients with intermediate-risk disease, and 167 patients with high-risk or very high risk disease.

Of the 206 patients with evolution to AML, 9 had an unknown *WT1* expression level, 102 had an unknown *PRAME* expression level because of the absence of gene detection, 126 (64.0%) had *WT1* overexpression, and 91 (87.5%) had *PRAME* overexpression. Neither *WT1* nor *PRAME* overexpression was detectable in the other 9 patients who progressed to AML. The combination of *PRAME* and *WT1* could cover more patients for predicting AML evolution.

For patients with MDS, the presence of *WT1* overexpression was associated with a 34.6% risk for AML evolution compared with




Figure 3. OS and PFS of patients with MDS according to cytopenia lines and *WT1*/*PRAME* transcript levels. OS (A) and PFS (D) of patients with MDS based on cytopenia lines. OS (B) and PFS (E) of patients with MDS based on the presence and absence of *WT1* high expression. OS (C) and PFS (F) of patients with MDS based on the presence and absence of *PRAME* high expression.

a 10.5% risk without *WT1* overexpression ($P = .000$). Likewise, the presence of *PRAME* overexpression was also related to a 26.1% risk for AML evolution compared with a 7.2% risk without *PRAME* overexpression ($P = .000$). MDS with *WT1* or *PRAME* overexpression more likely evolved to AML (Figure 2B-C). Multivariate COX analyses showed that *WT1* and *PRAME* overexpression were independent predictors of AML evolution (Table 2).

The prognostic significance of *WT1* and *PRAME* transcript levels

First, we examined the hazard ratio (HR) of death and progression associated with *WT1* and *PRAME*. In this univariate analysis, *WT1* and *PRAME* overexpression were both associated with shorter OS and PFS (Tables 3 and 4; Figure 3). Some patient and disease clinical features are known to influence survival for MDS, including age, sex, cytopenia lines, IPSS-R, and treatment options. We then examined how these variables were associated with OS and PFS in our cohort. Patients who were ≥ 60 years of age, male, higher risk, and in group B and C and who received chemotherapy or HMA had worse OS. Likewise, patients who were ≥ 60 years of age, male, higher risk, and in group B and C had worse PFS. There were no statistically differences in OS for patients stratified by thrombocytopenia (supplemental Figure 5C).

WT1 and *PRAME* overexpression often occur together in the same patient, and both are associated with IPSS-R. To determine the relative

contribution of gene expression level to survival, we generated a multivariable Cox regression model using an enter procedure incorporating age, sex, IPSS-R, cytopenia lines, treatment, and gene transcript levels. As expected, the IPSS-R was strongly associated with OS and PFS. *WT1* and *PRAME* overexpression also emerged as independent predictors of OS and PFS (Table 3-4). This analysis indicated that evaluation of the expression levels of *WT1* and *PRAME* would add additional prognostic value and be complementary to the IPSS-R.

WT1 and *PRAME* overexpression occurred in largely overlapping groups of patients, and approximately 30% had an overexpression in at least 1 of these 2 genes. In addition, patients with MDS with overexpression of both *WT1* and *PRAME* had significantly lower OS and PFS compared with those of the patients with either *WT1* or *PRAME* overexpression ($P = .013$ for OS and $P = .001$ for PFS; Figure 5B,E). Together, the 2 genes identified a population of patients with substantially worse OS and PFS.

The survival analysis for integration of gene expression analysis into the IPSS-R is shown in Figure 5. According to *WT1* and *PRAME* transcript levels, patients with MDS with IPSS-R low risk were classified into 2 prognostic risk groups, a low-favorable group and a low-adverse group, and they had divergent outcomes. The characteristics were shown in supplemental Table 3. MDS-U and MDS-SLD comprised the majority of subtypes in both groups. No statistical difference was observed in the WHO classification, cytogenetic risk, or other clinical variables. The patients in the




Figure 4. Proportions of patients with WT1 and PRAME transcript levels, according to blast percentage, WHO classification, cytogenetics, and IPSS-R. Data are shown for the percentage of blasts in bone marrow aspirate (A), WHO classification (B), cytogenetics (C), and IPSS-R (D) at the time of MDS diagnosis. Overexpression of WT1 and PRAME were associated with elevated blast percentage (defined as $\geq 5\%$; $P = .000$ and $P = .000$ for WT1 and PRAME, respectively) (A), worse cytogenetics ($P = .000$ and $P = .000$ for WT1 and PRAME, respectively) (C), and IPSS-R higher risk ($P = .000$ and $P = .000$ for WT1 and PRAME, respectively) (D).

low-adverse group had an OS similar to that of patients in the intermediate-risk group (Figure 5C,F; supplemental Figure 6). The 2 molecular biomarkers could reclassify patients in the IPSS-R low-adverse group to the intermediate-risk subgroup. These patients may benefit from earlier initiation of disease-modifying therapy.

Response to treatment of patients with WT1/PRAME overexpression

A total of 174 patients with WT1/PRAME overexpression initiated intensive treatment as a result of higher-risk disease at

diagnosis or evidence of disease progression, including 124 patients in the allo-HSCT group and 50 patients in the chemotherapy or HMA group. Data on the patient- and treatment-related characteristics of both treatment groups were listed in supplemental Table 4 and 5.

The OS and PFS of patients with MDS in the HSCT group were better than those in the chemotherapy and HMA group ($P = .000$ for OS and $P = .000$ for PFS; Figure 6). In univariate and multivariate analyses (COX regression), undergoing HSCT was an independent factor associated with better OS (Table 3).

Table 2. Univariable and multivariable analyses (COX regression) identifying independent risk factors for AML evolution of patients with MDS with thrombocytopenia

Variable	AML evolution			
	Univariate Cox		Multivariate Cox	
	HR (95% CI)	P	HR (95% CI)	P
Age (≥60 y vs <60 y)	0.949 (0.714-1.263)	.721		
Sex (male vs female)	1.610 (1.209-2.144)	.001	1.350 (.893-2.042)	.155
IPSS-R (higher risk vs lower risk)	2.386 (2.047-2.780)	.000	4.692 (2.668-8.252)	.000
<i>WT1</i> (≥4.9% vs <4.9%)	3.635 (2.716-4.865)	.000	1.663 (1.099-2.516)	.016
<i>PRAME</i> (≥0.3% vs <0.3%)	3.955 (2.211-7.074)	.000	1.879 (1.019-3.468)	.044

Discussion

To identify the disease features and explore the prognostic value of *WT1* and *PRAME* in patients with MDS with thrombocytopenia, the transcript levels of *WT1* and *PRAME* were detected from a large set of patients with MDS with thrombocytopenia at diagnosis. Our findings suggest that *WT1/PRAME* overexpression at diagnosis is associated with shorter OS and PFS, and high probability of AML evolution. The combination of *WT1/PRAME* transcript analysis with risk group may improve the prognostication precision in clinical practice.

The diagnosis of MDS is usually initially considered by cytopenia. Thrombocytopenia is a common and serious, life-threatening bleeding complications leading to significant mortality. The prevalence of isolated thrombocytopenia as a first manifestation at the onset of MDS was previously reported to be 1% to 12%.⁸ Approximately 30% of these patients were originally presumed to have immune thrombocytopenia.⁸ Patients with MDS with isolated thrombocytopenia are a challenge to clinicians because of the difficulty of diagnosis and lack of well-established treatment strategy. Observations from some prior studies have implied that

patients with MDS with isolated thrombocytopenia have a relatively favorable prognosis.⁷ One recent study came to the opposite conclusion.⁸ However, the data were obtained from cases or smaller sample descriptive reports.^{5,7,8} Here, our data demonstrate that patients with MDS with isolated thrombocytopenia have a favorable prognosis, concordant with the majority of studies. The overexpression of *WT1* or *PRAME* transcript levels are detected in more than half of our patients with thrombocytopenia. Therefore, monitoring *WT1* and *PRAME* transcript levels has the potential to improve the diagnostic accuracy of MDS with thrombocytopenia. In addition, Gonzalez-Porras et al found that severe thrombocytopenia had an adverse prognostic clinical effect in low-risk patients with MDS.⁴¹ The assessment of severe thrombocytopenia was important to risk stratification and to further refining clinical decision making. Our results revealed similar negative effects of *WT1* and *PRAME* overexpression for patients with MDS in a low-risk group. This suggested an underlying association between *WT1/PRAME* and severe thrombocytopenia in low-risk MDS.

MDS are clinically heterogeneous, premalignant disorders that range from conditions with a near-normal life expectancy to situations that are close to AML.⁴² The fact that MDS have highly variable clinical process makes risk stratification important in clinical decision making. Current scoring systems consider the percentage of bone marrow blasts, cytogenetic findings, classification, and cytopenias to predict the prognosis. However, there is poor interobserver concordance and reproducibility in some morphologic evaluation variables, and approximately half of patients lack informative cytogenetic abnormalities, resulting in silent progression of MDS to AML in different age groups.⁴³ In addition, patients with identical chromosomal abnormalities are often clinically heterogeneous.^{9,44} Further study is required to develop more robust prognostic factors.

Recently, many studies have focused on the opportunity afforded by *WT1* and *PRAME* in most patients with AML to provide

Table 3. Univariable and multivariable analyses (COX regression) identifying independent risk factors for OS of patients with MDS with thrombocytopenia


Variable	OS			
	Univariate Cox		Multivariate Cox	
	HR (95% CI)	P	HR (95% CI)	P
Age (≥60 y vs <60 y)*	2.708 (2.066-3.551)	.000	1.458 (0.960-2.214)	.077
Sex (male vs female)	1.575 (1.190-2.085)	.001	1.297 (0.863-1.951)	.211
IPSS-R (higher risk vs lower risk)	2.940 (2.142-4.035)	.000	2.867 (1.715-4.793)	.000
Group (C&B vs A)	1.957 (1.280-2.992)	.002	1.846 (1.003-3.398)	.049
<i>WT1</i> (≥4.9% vs <4.9%)	1.782 (1.348-2.357)	.000	1.522 (1.011-2.292)	.044
<i>PRAME</i> (≥0.3% vs <0.3%)	3.104 (1.852-5.203)	.000	1.855 (1.040-3.308)	.036
Treatment (HSCT vs chemotherapy and HMA)	0.505 (0.346-0.735)	.000	0.355 (0.193-0.653)	.001

Group A, isolated thrombocytopenia group; Group B, bicytopenia group; Group C, pancytopenia group; HSCT, hematopoietic stem cell transplantation; OS, overall survival; PFS, progression-free survival; *PRAME*, preferentially expressed antigen of melanoma; *WT1*, Wilms tumor 1.

*Age was included in the analysis as a categorical variable on the basis of a best-split algorithm showing a statistical difference in OS between patients younger than 60 years and those aged 60 years or older (supplemental Figure 5A-B,D-E).

Table 4. Univariable and multivariable analyses (COX regression) identifying independent risk factors for PFS of patients with MDS with thrombocytopenia

Variable	PFS			
	Univariate Cox		Multivariate Cox	
	HR (95% CI)	P	HR (95% CI)	P
Age (≥ 60 y vs < 60 y)*	2.420 (1.847-3.171)	.000	1.152 (0.801-1.655)	.446
Sex (male vs female)	1.726 (1.302-2.288)	.000	1.388 (0.987-1.952)	.060
IPSS-R (higher risk vs lower risk)	3.467 (2.518-4.773)	.000	3.045 (1.950-4.755)	.000
Group (A and B vs C)	1.937 (1.257-2.985)	.003	1.126 (0.700-1.812)	.624
<i>WT1</i> ($\geq 4.9\%$ vs $< 4.9\%$)	2.147 (1.622-2.842)	.000	1.521 (1.077-2.146)	.017
<i>PRAME</i> ($\geq 0.3\%$ vs $< 0.3\%$)	3.342 (1.993-5.604)	.000	1.801 (1.114-2.914)	.016
Treatment (HSCT vs Chemotherapy and HMA)	0.710 (0.488-1.034)	.074	0.799 (0.514-1.241)	.318



*Age was included in the analysis as a categorical variable on the basis of a best-split algorithm showing a statistical difference in OS between patients younger than 60 years and those aged 60 years or older (supplemental Figure 5A-B,D-E).

a immunogenic antigens candidate for immunotherapy and to advocate *WT1* and *PRAME* as universal markers for minimal residual disease assessment.⁴⁵

WT1 gene encodes a transcription factor; its target genes are important for cellular growth and survival, including growth factors, extracellular matrix, and other transcription factors.^{13,46} It was first identified for its role in the pathogenesis of Wilms tumor.¹³ The *WT1* gene, located at chromosome 11p13, encodes 10 exons and generates a 3-kb mRNA.¹³ Some studies showed that the degree of *WT1* expression was highly associated with the type of MDS, as it was apparently higher in RAEB and s-AML, in contrast to RA.^{29,47} The bone marrow blasts and risk for AML evolution increased in patients with *WT1* overexpression.^{23,29} In addition, a study showed the correlation between the degree of *WT1* transcript levels and the IPSS score.⁴⁸ The results of these studies implied that abnormal *WT1* expression levels could be associated with high risk and may reflect the disease progression. However, these studies lack long-term follow-up observation.


PRAME was first revealed as a tumor-associated antigen via analysis of the specificity of tumor-reactive T-cell clones from the patient with melanoma.⁴⁸ The human *PRAME* gene is located at chromosome 22 (22q11.22) and generates an approximately 12-kb mRNA.⁴⁹ Overexpression of *PRAME* transcripts was initially associated with an advanced tumor stage and poor prognosis in several solid tumors, such as breast cancer, lung cancer, head and neck cancers, and ovarian cancer. In contrast, preliminary reports in AML indicated that patients with *PRAME* overexpression usually had a favorable response to chemotherapy and better survival.^{24,25,50} Information about *PRAME* expression and its prognostic value in MDS is very limited. Our previous study based on small numbers of patients found that *PRAME* was commonly overexpressed in MDS, and the transcript levels of *WT1* in the high-risk group were higher than in the low-risk group.²³ The clinical effect of *PRAME* on the prognosis of patients with MDS with thrombocytopenia has not, however, been evaluated yet. Based on a large cohort of patients with MDS with

thrombocytopenia, our data identify that *WT1* and *PRAME* transcript levels are independent prognostic factors, as their overexpression is correlated with poor OS and PFS, and a higher probability of AML evolution. Simultaneous detection of *WT1* and *PRAME* expression covers more patients with MDS and allows a greater distinction between patients with MDS with significantly different outcomes.

However, the biological explanation of why *WT1* and *PRAME* overexpression is associated with worse prognosis in MDS remains unclear. Recent studies have indicated that *WT1* and *PRAME* are key to regulating hematopoiesis and apoptosis. *WT1* overexpression indices increased apoptosis through transcriptional regulation of the Bcl-2 family member Bcl-2, BAK, and A1.⁵¹⁻⁵⁵ In addition, *WT1* overexpression competes with the differentiation-inducing signal mediated by G-CSFR and constitutively activated Stat3, resulting in the blocking of differentiation,^{56,57} which may be implicated in dysregulated hematopoiesis in MDS. *PRAME* was reported to be a repressor of RA signaling, capable of inhibiting RA-induced hematopoietic differentiation, cell cycle arrest, and apoptosis.¹⁷ Forced *PRAME* expression in normal hematopoietic progenitors inhibited myeloid differentiation.⁵⁸ More important, both *WT1* and *PRAME* are related to the MDS clone. *WT1* and *PRAME* overexpression may be driver factors leading to clonal expansion and disease progression in MDS. Li and colleagues found that *WT1* was expressed predominantly in MDS clonal cells, rather than in nonclonal cells.²⁷ Similarly, another study revealed that elevated *WT1* expression may be related to the presence of preleukemic MDS clones with poor prognostic chromosomal rearrangements.²¹ Our previous work suggested that *PRAME* was also overexpressed in the relative mature cells of clonal MDS, which suggested that their overexpression in MDS was related to malignant clonal characteristics.²³ Collectively, both *WT1* and *PRAME* overexpression may contribute to deregulated apoptosis and ineffective hematopoiesis and are associated with the MDS clone, which constitute the biological basis of incorporating both *WT1* and *PRAME* in the IPSS-R.

Figure 5. OS and PFS of patients with MDS according to IPSS-R and *WT1* and *PRAME* transcript levels.

Patients with MDS were grouped according to favorable (low *WT1* or low *PRAME*) or adverse (high *WT1* or high *PRAME*) expression profiles. Therefore, the proposed score was drawn up by assigning a value of 0 for each gene expressing a favorable RNA level and a value of 1 for each gene with an adverse expression profile. OS (A) and PFS (D) of patients with MDS based on IPSS-R. OS (B) and PFS (E) of patients with MDS based on *WT1/PRAME* score. It distinguished 3 subsets of patients with different OS ($P < .001$) and PFS ($P < .001$). The OS and PFS for patients with both *WT1* and *PRAME* overexpression were significantly lower than that those for patients with either *WT1* or *PRAME* overexpression ($P = .013$ for OS; $P = .001$ for PFS, respectively). OS (C) and PFS (F) of patients with MDS in the next-highest IPSS risk group is included for the purpose of comparison, P values were calculated between low-risk patients with *WT1/PRAME* overexpression and intermediate patients.



Therefore, it is an attractive strategy to apply *WT1*- and *PRAME*-specific immunotherapy to patients with MDS with thrombocytopenia. In addition, *WT1* and *PRAME* transcript levels contribute to distinguishing 2 prognostic subgroups within low-risk patients with MDS. The identification of patients with poor prognosis among those assigned as low risk is important because patients classified as lower risk are not often considered to receive intensive therapy such as bone marrow transplantation and antileukemic chemotherapy. And in Europe, HMA are not licensed to use for patients with MDS in lower risk.⁵⁹ Our study supports the need for monitoring *WT1/PRAME* transcript levels and helping guide decision-making toward treatments that aim to alter the natural history of the disease and induce prolonged survival for the low-risk patients with poor prognosis.

As with any survey study, there are certain weaknesses in our study. The first is the retrospective nature of our work. The data for *WT1* and *PRAME* transcript levels were collected retrospectively in our study, which caused incomplete information. Second, the biological mechanism of why overexpression of *WT1* and *PRAME* is associated with a poor prognosis is unclear. Therefore, we are acutely aware that more prospective validations are needed. Regardless of the limitations, clinical and hematological information is sufficient in the majority of our patients, and samples from bone marrow aspiration were homogeneously collected before treatment. Furthermore, the technology of *WT1* and *PRAME* detection has been well established from our large-scale clinical practice. These allow confident predictions to be made. Overall, our findings serve




Figure 6. OS and PFS of patients with MDS with thrombocytopenia according to treatment. OS (A) and PFS (B) for the 174 patients with MDS, respectively.

as a proof of concept that the integration of *WT1* and *PRAME* significantly increase the ability to capture prognostic information in patients with MDS at diagnosis and may provide a basis for improving clinical decision making.

In conclusion, for patients with MDS with thrombocytopenia, our findings suggest that *WT1/PRAME* overexpression at diagnosis predicts poor survival and AML evolution. The integration of *WT1/PRAME* transcript analysis into risk groups may upstage some low-risk patients and better risk-stratify the patients. Interestingly, the prognosis of patients with MDS with isolated thrombocytopenia is favorable. For patients with higher *WT1/PRAME* transcript levels, allo-HSCT may be superior to chemotherapy and HMA.

Acknowledgments

The corresponding author had full access to all of the data in the study and had final responsibility for the decision to submit for publication.

This work was supported by National Natural Science Foundation of China (No. 81730004 and No. 81670116), Beijing Natural Science Foundation (No. H2018206423 and No. 7171013), Beijing Municipal Science and Technology Commission (No. Z171100001017084), the National Key Research and Development Program of China (No. 2017YFA0105500, No. 2017YFA0105503),

and the Foundation for Innovative Research Groups of the National Natural Science Foundation of China (No.81621001).

The funders of the study had no role in study design, data collection, data analysis, data interpretation, or writing of the report.

Authorship

Contribution: Q.-S.H. and X.-H.Z. were responsible for data collection and management, statistical analysis and interpretation, literature research, and manuscript writing; J.-Z.W., Q.-Z.Z., Q.J., H.J., and J.L. were responsible for data management and statistical analysis; H.-X.L. assisted in statistical analysis; Y.L., J.-B.W., L.S., H.-Y.Z., Z.-L.L., and S.-J.G. were responsible for data collection and management; Y.-Z.Q. and Y.-R.L. performed the gene mutation studies; and B.H., Y.-Y.L., L.-P.X., and X.-J.H. planned, designed, and coordinated the study over the entire period and wrote the manuscript.

Conflict-of-interest disclosures: The authors declare no competing financial interests.

ORCID profile: X.-H.Z., 0000-0003-0245-6792.

Correspondence: Xiao-Hui Zhang, Peking University People's Hospital, No. 11 Xizhimen South St, Xicheng District, Beijing 100044, China; e-mail: zhangxh100@sina.com.

References

- Fenaux P, Haase D, Sanz GF, Santini V, Buske C, Group EGW; ESMO Guidelines Working Group. Myelodysplastic syndromes: ESMO Clinical Practice Guidelines for diagnosis, treatment and follow-up. *Ann Oncol*. 2014;25(Suppl 3):iii57-iii69.
- Malcovati L, Hellström-Lindberg E, Bowen D, et al; European Leukemia Net. Diagnosis and treatment of primary myelodysplastic syndromes in adults: recommendations from the European LeukemiaNet. *Blood*. 2013;122(17):2943-2964.
- Killick SB, Carter C, Culligan D, et al; British Committee for Standards in Haematology. Guidelines for the diagnosis and management of adult myelodysplastic syndromes. *Br J Haematol*. 2014;164(4):503-525.
- Kantarjian H, Giles F, List A, et al. The incidence and impact of thrombocytopenia in myelodysplastic syndromes. *Cancer*. 2007;109(9):1705-1714.
- Kuroda J, Kimura S, Kobayashi Y, Wada K, Uoshima N, Yoshikawa T. Unusual myelodysplastic syndrome with the initial presentation mimicking idiopathic thrombocytopenic purpura. *Acta Haematol*. 2002;108(3):139-143.
- Li W, Morrone K, Kambhampati S, Will B, Steidl U, Verma A. Thrombocytopenia in MDS: epidemiology, mechanisms, clinical consequences and novel therapeutic strategies. *Leukemia*. 2016;30(3):536-544.
- Sashida G, Takaku TI, Shoji N, et al. Clinico-hematologic features of myelodysplastic syndrome presenting as isolated thrombocytopenia: an entity with a relatively favorable prognosis. *Leuk Lymphoma*. 2003;44(4):653-658.

8. Waisbren J, Dinner S, Altman J, et al. Disease characteristics and prognosis of myelodysplastic syndrome presenting with isolated thrombocytopenia. *Int J Hematol.* 2017;105(1):44-51.
9. Haase D, Germing U, Schanz J, et al. New insights into the prognostic impact of the karyotype in MDS and correlation with subtypes: evidence from a core dataset of 2124 patients. *Blood.* 2007;110(13):4385-4395.
10. Jaiswal S, Fontanillas P, Flannick J, et al. Age-related clonal hematopoiesis associated with adverse outcomes. *N Engl J Med.* 2014;371(26):2488-2498.
11. Dao T, Pankov D, Scott A, et al. Therapeutic bispecific T-cell engager antibody targeting the intracellular oncoprotein WT1. *Nat Biotechnol.* 2015;33(10):1079-1086.
12. Gezgin G, Luk SJ, Cao J, et al. PRAME as a potential target for immunotherapy in metastatic uveal melanoma. *JAMA Ophthalmol.* 2017;135(6):541-549.
13. Huff V. Wilms' tumours: about tumour suppressor genes, an oncogene and a chameleon gene. *Nat Rev Cancer.* 2011;11(2):111-121.
14. Ellisen LW, Carlesso N, Cheng T, Scadden DT, Haber DA. The Wilms tumor suppressor WT1 directs stage-specific quiescence and differentiation of human hematopoietic progenitor cells. *EMBO J.* 2001;20(8):1897-1909.
15. Nishida S, Hosen N, Shirakata T, et al. AML1-ETO rapidly induces acute myeloblastic leukemia in cooperation with the Wilms tumor gene, WT1. *Blood.* 2006;107(8):3303-3312.
16. Ikeda H, Lethé B, Lehmann F, et al. Characterization of an antigen that is recognized on a melanoma showing partial HLA loss by CTL expressing an NK inhibitory receptor. *Immunity.* 1997;6(2):199-208.
17. Epping MT, Wang L, Edel MJ, Carlée L, Hernandez M, Bernards R. The human tumor antigen PRAME is a dominant repressor of retinoic acid receptor signaling. *Cell.* 2005;122(6):835-847.
18. Collins SJ. The role of retinoids and retinoic acid receptors in normal hematopoiesis. *Leukemia.* 2002;16(10):1896-1905.
19. Drumea K, Yang ZF, Rosmarin A. Retinoic acid signaling in myelopoiesis. *Curr Opin Hematol.* 2008;15(1):37-41.
20. Collins SJ. Retinoic acid receptors, hematopoiesis and leukemogenesis. *Curr Opin Hematol.* 2008;15(4):346-351.
21. Nagasaki J, Aoyama Y, Hino M, et al. Wilms tumor 1 (WT1) mRNA expression level at diagnosis is a significant prognostic marker in elderly patients with myelodysplastic syndrome. *Acta Haematol.* 2017;137(1):32-39.
22. Nomdedéu JF, Hoyos M, Carricondo M, et al; CETLAM Group. Bone marrow WT1 levels at diagnosis, post-induction and post-intensification in adult de novo AML. *Leukemia.* 2013;27(11):2157-2164.
23. Qin YZ, Zhu HH, Liu YR, et al. PRAME and WT1 transcripts constitute a good molecular marker combination for monitoring minimal residual disease in myelodysplastic syndromes. *Leuk Lymphoma.* 2013;54(7):1442-1449.
24. Santamaria C, Chillón MC, García-Sanz R, et al. The relevance of preferentially expressed antigen of melanoma (PRAME) as a marker of disease activity and prognosis in acute promyelocytic leukemia. *Haematologica.* 2008;93(12):1797-1805.
25. Zhang YH, Lu AD, Yang L, et al. PRAME overexpression predicted good outcome in pediatric B-cell acute lymphoblastic leukemia patients receiving chemotherapy. *Leuk Res.* 2017;52:43-49.
26. Zhao XS, Jin S, Zhu HH, et al. Wilms' tumor gene 1 expression: an independent acute leukemia prognostic indicator following allogeneic hematopoietic SCT. *Bone Marrow Transplant.* 2012;47(4):499-507.
27. Li X, Wu L, Ying S, Chang C, Pu Q. Wilms' tumor gene (WT1) is predominantly expressed in clonal hematopoietic cells in myelodysplastic syndromes. *Leuk Lymphoma.* 2007;48(3):601-604.
28. Van Dijk JP, Knops GH, Van De Loch LT, et al. Abnormal WT1 expression in the CD34-negative compartment in myelodysplastic bone marrow. *Br J Haematol.* 2002;118(4):1027-1033.
29. Ueda Y, Mizutani C, Nannya Y, et al. Clinical evaluation of WT1 mRNA expression levels in peripheral blood and bone marrow in patients with myelodysplastic syndromes. *Leuk Lymphoma.* 2013;54(7):1450-1458.
30. Qin YZ, Wang Y, Zhu HH, et al. Low WT1 transcript levels at diagnosis predicted poor outcomes of acute myeloid leukemia patients with t(8;21) who received chemotherapy or allogeneic hematopoietic stem cell transplantation. *Chin J Cancer.* 2016;35(1):46.
31. Zhao XS, Yan CH, Liu DH, et al. Combined use of WT1 and flow cytometry monitoring can promote sensitivity of predicting relapse after allogeneic HSCT without affecting specificity. *Ann Hematol.* 2013;92(8):1111-1119.
32. Arber DA, Orazi A, Hasserjian R, et al. The 2016 revision to the World Health Organization classification of myeloid neoplasms and acute leukemia. *Blood.* 2016;127(20):2391-2405.
33. Valent P, Horny HP, Bennett JM, et al. Definitions and standards in the diagnosis and treatment of the myelodysplastic syndromes: Consensus statements and report from a working conference. *Leuk Res.* 2007;31(6):727-736.
34. Qin Y, Zhu H, Jiang B, et al. Expression patterns of WT1 and PRAME in acute myeloid leukemia patients and their usefulness for monitoring minimal residual disease. *Leuk Res.* 2009;33(3):384-390.
35. Mo XD, Zhang XH, Xu LP, et al. Haploidentical hematopoietic stem cell transplantation for myelodysplastic syndrome. *Biol Blood Marrow Transplant.* 2017;23(12):2143-2150.
36. Wang Y, Liu DH, Liu KY, et al. Long-term follow-up of haploidentical hematopoietic stem cell transplantation without in vitro T cell depletion for the treatment of leukemia: nine years of experience at a single center. *Cancer.* 2013;119(5):978-985.
37. Wang Y, Wang HX, Lai YR, et al. Haploidentical transplant for myelodysplastic syndrome: registry-based comparison with identical sibling transplant. *Leukemia.* 2016;30(10):2055-2063.

38. Huang XJ, Liu DH, Liu KY, et al. Treatment of acute leukemia with unmanipulated HLA-mismatched/haploidentical blood and bone marrow transplantation. *Biol Blood Marrow Transplant*. 2009;15(2):257-265.
39. Greenberg PL, Tuechler H, Schanz J, et al. Revised international prognostic scoring system for myelodysplastic syndromes. *Blood*. 2012;120(12):2454-2465.
40. Cheson BD, Greenberg PL, Bennett JM, et al. Clinical application and proposal for modification of the International Working Group (IWG) response criteria in myelodysplasia. *Blood*. 2006;108(2):419-425.
41. Gonzalez-Porras JR, Cordoba I, Such E, et al; Spanish Myelodysplastic Syndrome Registry. Prognostic impact of severe thrombocytopenia in low-risk myelodysplastic syndrome. *Cancer*. 2011;117(24):5529-5537.
42. Della Porta MG, Galli A, Bacigalupo A, et al. Clinical effects of driver somatic mutations on the outcomes of patients with myelodysplastic syndromes treated with allogeneic hematopoietic stem-cell transplantation. *J Clin Oncol*. 2016;34(30):3627-3637.
43. Ganguly BB, Kadam NN. Mutations of myelodysplastic syndromes (MDS): An update. *Mutat Res Rev Mutat Res*. 2016;769:47-62.
44. Bejar R, Levine R, Ebert BL. Unraveling the molecular pathophysiology of myelodysplastic syndromes. *J Clin Oncol*. 2011;29(5):504-515.
45. Cillon D, Renneville A, Hermitte F, et al. Real-time quantitative polymerase chain reaction detection of minimal residual disease by standardized WT1 assay to enhance risk stratification in acute myeloid leukemia: a European LeukemiaNet study. *J Clin Oncol*. 2009;27(31):5195-5201.
46. Yang L, Han Y, Suarez Saiz F, Minden MD. A tumor suppressor and oncogene: the WT1 story [published correction appears in *Leukemia*. 2007;21(7):1603]. *Leukemia*. 2007;21(5):868-876.
47. Patmasiriwat P, Fraizer G, Kantarjian H, Saunders GF. WT1 and GATA1 expression in myelodysplastic syndrome and acute leukemia. *Leukemia*. 1999;13(6):891-900.
48. Cillon D, Gottardi E, Messa F, et al; Piedmont Study Group on Myelodysplastic Syndromes. Significant correlation between the degree of WT1 expression and the International Prognostic Scoring System Score in patients with myelodysplastic syndromes. *J Clin Oncol*. 2003;21(10):1988-1995.
49. Wadelin F, Fulton J, McEwan PA, Spriggs KA, Emsley J, Heery DM. Leucine-rich repeat protein PRAME: expression, potential functions and clinical implications for leukaemia. *Mol Cancer*. 2010;9(1):226.
50. Greiner J, Schmitt M, Li L, et al. Expression of tumor-associated antigens in acute myeloid leukemia: Implications for specific immunotherapeutic approaches. *Blood*. 2006;108(13):4109-4117.
51. Mayo MW, Wang CY, Drouin SS, et al. WT1 modulates apoptosis by transcriptionally upregulating the bcl-2 proto-oncogene. *EMBO J*. 1999;18(14):3990-4003.
52. Hewitt SM, Hamada S, McDonnell TJ, Rauscher FJ III, Saunders GF. Regulation of the proto-oncogenes bcl-2 and c-myc by the Wilms' tumor suppressor gene WT1. *Cancer Res*. 1995;55(22):5386-5389.
53. Ito K, Oji Y, Tatsumi N, et al. Antiapoptotic function of 17AA(+)/WT1 (Wilms' tumor gene) isoforms on the intrinsic apoptosis pathway. *Oncogene*. 2006;25(30):4217-4229.
54. Morrison DJ, English MA, Licht JD. WT1 induces apoptosis through transcriptional regulation of the proapoptotic Bcl-2 family member Bak. *Cancer Res*. 2005;65(18):8174-8182.
55. Simpson LA, Burwell EA, Thompson KA, Shahnaz S, Chen AR, Loeb DM. The antiapoptotic gene A1/BFL1 is a WT1 target gene that mediates granulocytic differentiation and resistance to chemotherapy. *Blood*. 2006;107(12):4695-4702.
56. Svedberg H, Richter J, Gullberg U. Forced expression of the Wilms tumor 1 (WT1) gene inhibits proliferation of human hematopoietic CD34(+) progenitor cells. *Leukemia*. 2001;15(12):1914-1922.
57. Inoue K, Tamaki H, Ogawa H, et al. Wilms' tumor gene (WT1) competes with differentiation-inducing signal in hematopoietic progenitor cells. *Blood*. 1998;91(8):2969-2976.
58. Oehler VG, Guthrie KA, Cummings CL, et al. The preferentially expressed antigen in melanoma (PRAME) inhibits myeloid differentiation in normal hematopoietic and leukemic progenitor cells. *Blood*. 2009;114(15):3299-3308.
59. Valcárcel D, Sanz G, Ortega M, et al; Grupo Español de Síndromes Mielodisplásicos (GESMD). Use of newer prognostic indices for patients with myelodysplastic syndromes in the low and intermediate-1 risk categories: a population-based study. *Lancet Haematol*. 2015;2(6):e260-e266.