



Published in final edited form as:

*J Rheumatol.* 2019 December ; 46(12): 1589–1596. doi:10.3899/jrheum.190009.

## Association of Childhood Abuse with Incident Systemic Lupus Erythematosus in Adulthood in a Longitudinal Cohort of Women

Candace H. Feldman, MD, ScD<sup>1</sup>, Susan Malspeis, MS<sup>1</sup>, Cianna Leatherwood, MD, MPH<sup>1</sup>, Laura Kubzansky, PhD<sup>2</sup>, Karen H. Costenbader, MD, MPH<sup>1,\*</sup>, Andrea L. Roberts, PhD<sup>2,\*</sup>

<sup>1</sup>Division of Rheumatology, Immunology and Allergy, Department of Medicine, Brigham and Women's Hospital and Harvard Medical School, Boston, MA

<sup>2</sup>Harvard T.H. Chan School of Public Health, Boston, MA

### Abstract

**Objective:** Exposure to severe stressors may alter immune function and augment inflammation and cytokine release, increasing autoimmune disease risk. We examined if childhood abuse was associated with heightened risk of incident systemic lupus erythematosus (SLE).

**Methods:** Data were from the Nurses' Health Study II, a cohort of U.S. female nurses enrolled in 1989, followed with biennial questionnaires. We measured childhood physical and emotional abuse with the Physical and Emotional Abuse Subscale of the Childhood Trauma Questionnaire and sexual abuse with the Sexual Maltreatment Scale of the Parent-Child Conflict Tactics Scale, both administered in 2001. We identified incident SLE (>4 American College of Rheumatology 1997 classification criteria) through 2015. We used multivariable Cox regression models to evaluate the association between childhood abuse and SLE accounting for potential confounders (e.g. parental education, occupation, home ownership) and mediators (e.g. depression, posttraumatic stress disorder (PTSD)).

**Results:** Among 67,516 women, there were 94 incident SLE cases. In adjusted models, exposure to the highest vs. lowest physical and emotional abuse was associated with 2.57 times greater risk of SLE (95% CI 1.30–5.12). We found that 17% ( $p < 0.0001$ ) of SLE risk associated with abuse could be explained by depression and 23% ( $p < 0.0001$ ) by PTSD. We did not observe a statistically significant association with sexual abuse (HR 0.84, 95% CI 0.40–1.77, highest vs. lowest exposure).

**Conclusion:** We observed significantly increased risk of SLE among women who experienced childhood physical and emotional abuse compared with women who had not. Exposure to childhood adversity may contribute to SLE development.

### Key Terms:

Systemic lupus erythematosus; epidemiology; risk factors

---

**Corresponding author:** Candace H. Feldman, MD, ScD, 60 Fenwood Road, Office 6016P, Boston, MA 02115, cfeldman@bwh.harvard.edu, Phone: 617-525-1035, Fax: 617-264-3019.

\*Authors contributed equally

**Disclosures:** The authors have no relevant conflicts of interest.

## Introduction

Systemic lupus erythematosus (SLE) is a prototypical autoimmune disease with the potential for multisystem organ involvement and significant morbidity. Exposures to stress-related disorders, including posttraumatic stress disorder (PTSD), have been associated with increased risk of subsequent autoimmune diseases.<sup>1, 2</sup> In a longitudinal cohort of U.S. female nurses, women with high PTSD symptoms had nearly three times higher risk of incident SLE compared to women unexposed to trauma.<sup>3</sup> Hypothesized mechanisms include stress-driven alterations to the immune system via inflammatory cytokines and dysregulation of the hypothalamic-pituitary-adrenal axis and the autonomic nervous system, increasing susceptibility to autoimmune diseases and increased disease-related damage.<sup>4-6</sup>

Adverse experiences during childhood, a particularly sensitive time period, have significant effects on long-term neuropsychiatric and physical functioning and health outcomes later in life.<sup>7-9</sup> Exposure to adverse childhood environments and experiences, including socioeconomic disadvantage and maltreatment, has been associated with higher levels of inflammatory markers and cytokines, including C-reactive protein and interleukin-6, as well as to alterations in cell-mediated immune function both later in childhood and in adulthood.<sup>9-11</sup> Childhood adversity has been previously linked to increased risk of chronic diseases and mental illness.<sup>9, 12-14</sup> One study demonstrated that children exposed to two or more adverse childhood experiences were at 2-fold risk for hospitalization with a rheumatic disease compared with unexposed children.<sup>15</sup> Understanding the effects of this quite prevalent exposure on SLE incidence could inform interventions aimed to modify risk among vulnerable individuals during potentially sensitive developmental periods. We therefore investigated this relationship in a large, longitudinal U.S.-based cohort and hypothesized that exposure to childhood maltreatment would be associated with increased risk of developing SLE.

## Methods

### Population

We utilized data from the Nurses' Health Study II (NHSII), a longitudinal cohort of 116,429 female nurses, age 25–44 years at baseline in 1989 from fourteen U.S. states and followed with biennial questionnaires. In 2001, a questionnaire that assessed child abuse was sent to all NHSII participants who returned the most recent biennial questionnaire, with 75% response rate (68,376/91,279). We excluded participants with self-reported connective tissue disease at baseline, <4 1997 American College of Rheumatology (ACR) revised classification criteria for SLE<sup>16</sup>, or diagnosis prior to 1989 (n=672). We also excluded individuals missing child abuse data. This study was approved by the Partners Healthcare Institutional Review Board (#2015P001458); return of the questionnaires implied consent.

### SLE Case Ascertainment

We identified incident SLE cases from June 1991- June 2015. Women were asked to report new physician diagnoses of SLE on biennial questionnaires. Women indicating a new diagnosis were asked to complete the Connective Tissue Disease Screening Questionnaire<sup>17</sup>

and to consent to release of their medical records. Medical records of all nurses who indicated SLE symptoms were independently reviewed by two board-certified rheumatologists. Cases were determined based on >4 1997 ACR criteria.<sup>16</sup>

### Childhood Abuse Exposure Ascertainment

We assessed exposure to physical and emotional abuse using the Physical and Emotional Abuse Subscale of the Childhood Trauma Questionnaire (CTQ).<sup>18</sup> This scale has been shown to have good internal consistency (Cronbach's alpha=0.94), and test-retest reliability (intraclass correlation=0.82).<sup>18</sup> Participants were asked whether as a child (<11 years): "1) people in my family hit me so hard that it left me with bruises and marks, 2) the punishments I received seemed cruel, 3) I was punished with a belt, a board, a cord, or some other hard object, 4) someone in my family yelled and screamed at me, 5) people in my family said hurtful or insulting things to me." We excluded 188 individuals with incomplete CTQ data. Item responses (0: never, 1: rarely, 2: sometimes, 3: often or 4: very often true) were summed, the mean was determined, and then divided into quartiles to assess a dose-response relationship.<sup>19</sup> We also examined the CTQ score as a continuous variable.

We assessed sexual abuse using the two-item Sexual Maltreatment Scale of the Parent-Child Conflict Tactics Scale.<sup>20</sup> These questions were: "Were you ever touched in a sexual way by an adult or an older child or were you forced to touch an adult or an older child in a sexual way when you did not want to?" and "Did an adult or an older child ever force you or attempt to force you into any sexual activity by threatening you, holding you down or hurting you in some way when you did not want to." Respondents were asked to indicate if this happened never, once, or more than once during childhood (up to age 11 years) and as a teenager (age 11–17 years). In keeping with prior work, 0 points was equivalent to no abuse, 1–2 points mild, 3–4 points moderate and 5+ severe sexual abuse; responses were summed across questions.<sup>19</sup> These categories were examined separately and then moderate and severe categories were combined due to small sample sizes. We excluded 224 individuals with incomplete responses.

We conducted a secondary analysis examining physical assault using questions from the Conflict Tactics Scale (CTS)<sup>20</sup>. We included this as a secondary analysis of physical abuse because the full CTS was not queried, and because our primary measure, the CTQ, captured both physical and emotional abuse. The CTS categorizes physical assault by type: "1) spank you for discipline, 2) push grab or shove you, 3) kick, bite or punch you, 4) hit you with something that hurt your body, 5) choke or burn you, 6) physically attack you in some other way," and by frequency (never, once, a few times, more than a few times), during childhood (<11 years) or during adolescence (11–17 years). Responses were summed and divided into none, mild, moderate and severe. Moderate and severe categories were combined to maximize power and because the rate ratio estimates were nearly identical for the two categories. We combined childhood and adolescent abuse because the correlation was high ( $R^2=0.80$ ,  $p<0.001$ ).

## Covariate Ascertainment

Covariates were ascertained from biennial surveys, 1989–2013. We calculated age based on birthdate, and categorized race as white and nonwhite. We examined parental occupation, home ownership and education at the time of birth or infancy as different measures of childhood socioeconomic status, which may be a prior common cause of both childhood abuse and SLE risk. We categorized parental occupation for the parent with the highest occupation level as farmers, laborers, blue and lower white-collar workers (service, craftsmen, sales, clerical, military) and upper white-collar workers (managerial, professional).<sup>21, 22</sup> A separate variable indicated whether the mother worked outside of the home. We dichotomized parental home ownership at the time of birth or infancy (yes/no) and categorized the highest level of education attained by either parent as <high school, some college, and college graduate and above. We included self-reported birthweight (<5.5lbs, 5.5-<10lbs (ref), 10lbs), which has been validated in NHSII and shown to be strongly correlated with birth records.<sup>23</sup> Higher birthweight has been associated with increased SLE risk and we hypothesized a relationship with child abuse as well.<sup>24</sup> We measured childhood obesity, given prior work suggesting it is associated with SLE risk and with childhood maltreatment<sup>25, 26</sup>, using a childhood somatogram score; participants chose a diagram that represented their body type at age 5 (very thin (1) to extremely obese (9), grouped as 1–4 (reference) and 5.<sup>27</sup> We also assessed age at menarche (age 10 versus >10) as this has also been associated with SLE risk and with childhood abuse.<sup>28, 29</sup>

We assessed additional covariates measured during adulthood and updated biennially that may be associated both with prior experiences of childhood abuse and with subsequent SLE risk. Smoking status was categorized as never, past and current and body mass index (BMI) as 18.5-<25, 25-<30 and 30+. We also included measures of depression and of PTSD as both are associated with prior child abuse exposure, and with increased SLE risk.<sup>3, 30–32</sup> We assessed depression (yes/no) as self-reported clinician diagnosis (queried biennially from 2003–2011), self-reported use of antidepressants (queried biennially beginning in 1993) or score <60 on the Mental Health Inventory (MHI-5, measured in 1993, 1997 and 2001).<sup>33</sup> We assessed history of trauma or PTSD in a subsample of respondents who responded in 2008 to the Brief Trauma Questionnaire<sup>34</sup> and the 7-item Short Screening Scale for DSM-4 PTSD<sup>35</sup> categorized as no trauma, trauma and no PTSD, subclinical PTSD (1–3 symptoms) or probable PTSD (4–7 symptoms).<sup>3</sup> We assessed zip code median household income (in 1989), as well as alcohol consumption (none, 0–5g/day and 5g/day), updated biennially.<sup>36</sup>

## Statistical Analyses

We examined the distribution of covariates by physical and emotional abuse (CTQ) and sexual abuse exposure categories. We use Cox proportional hazards regression models to calculate the risk (hazard ratio, HR) of incident SLE associated with physical and emotional abuse, with the lowest quartile of exposure, as the reference. We also considered the CTQ score as a continuous variable to examine SLE risk in association with one standard deviation increase in CTQ score. We first fit models that were age and race-adjusted and then adjusted for childhood socioeconomic factors (parental occupation, education, home ownership), as well as childhood factors (birthweight, somatogram score at age 5, age at menarche), which may confound the association between abuse and SLE risk. We fit the

same models with sexual abuse and physical assault measured by the CTS as the independent variables. We tested the proportional hazards assumption for our primary exposures and did not find violations.

Adult BMI, alcohol consumption, area-level median household income, smoking, depression and PTSD may lie on the causal pathway between abuse exposure and SLE incidence. We therefore used the SAS “mediate” macro to estimate the proportion of the effect of childhood physical and emotional abuse on SLE risk that may be explained by each of these potential mediators in separate age- and race-adjusted, and childhood/parental-factor-adjusted models.<sup>37</sup> To assess mediation, we used Cox models as described above, additionally adjusted for these pathway variables. In sensitivity analyses, we also conducted a prospective analysis that included only SLE cases occurring after women returned the 2001 violence questionnaire, to minimize the possibility of recall bias.

## Results

Of the 67,516 women with CTQ data, 70% with 26 years of follow-up data, 36,224 (54%) reported moderate or high levels of exposure to physical or emotional abuse, and among 67,480 women with sexual abuse data, 6,732 (9.9%) reported moderate or high levels of sexual abuse (Table 1). A slightly higher percentage of non-white versus white women reported abuse. Individuals who were older at the time of report described modestly higher levels of childhood abuse. We found a higher prevalence of abuse among women with parents who had less education, were laborers, blue or lower white-collar workers, and who did not own homes. We also observed higher levels of physical and emotional abuse among women with lower birthweight and who were 10 years old at menarche. In adulthood, women who experienced childhood abuse had higher mean BMIs and were more likely to have smoked compared with women who did not. Depression and PTSD reported during adulthood were positively associated with abuse.

We confirmed 94 cases of SLE (Table 2). Nearly all were ANA positive with mean  $\pm$  SD ACR Criteria of  $5 \pm 1$ . More than half were anti-dsDNA positive, suggesting the potential for more active and severe disease, and the dominant manifestations were arthritis (73%) and hematologic involvement (64%). Of these SLE cases, 36 occurred after 2001 when the child abuse questionnaire was administered.

Considering the relationship between childhood physical and emotional abuse (measured by the CTQ) and SLE risk, we observed a nearly three-times higher risk of SLE comparing the highest level of abuse to no abuse (HR 2.81, 95% CI 1.42–5.56) in age- and race-adjusted models. This was slightly attenuated after adjusting for parental and childhood factors (HR 2.57, 95% CI 1.30–5.12) (Table 3). Each standard deviation increase in CTQ score as a continuous measure was associated with a 28% higher risk of SLE (HR 1.28, 95% CI 1.08–1.53).

With mediation analyses in adjusted models, 5% (95% CI 2.0–11.9,  $p=0.0003$ ) of the effect of the highest level of abuse on SLE risk could be statistically explained by smoking status, and 2.9% (95% CI 1.2–6.7,  $p<0.0001$ ) by BMI. Additionally, we found that 16.7% (95% CI

8–31.7,  $p < 0.0001$ ) of the risk of SLE associated with the highest level of abuse could be explained by depression and 23.3% (95% CI 7.2–54.4,  $p < 0.0001$ ) by PTSD exposure. We included these mediators in our multivariable models adjusted for age, race and childhood/parental factors and found that in models additionally adjusted for smoking and BMI (Table 3a, **Model C**), and depression (Table 3b, **Model D2**), the risk of SLE comparing women who experienced the highest level of abuse vs. the lowest, was modestly attenuated but statistically significant. Adjusting for PTSD (Table 3b, **Model E2**) significantly attenuated the association (HR 1.84, 95% CI 0.83–4.07). Alcohol consumption and zip code median household income were not statistically significant mediators of the association between childhood abuse and SLE risk.

In prospective analyses, the association between the highest degree of childhood physical and emotional abuse (vs. no abuse) and SLE risk remained statistically significant with a larger effect size (HR 3.14, 95% CI 1.03–9.56) in age- and race-adjusted models and was only slightly attenuated after adjusting for childhood and parental factors (HR 3.11, 95% CI 1.01–9.57 (Table 5). In adjusted models, we found a 41% increased risk of SLE for each standard deviation increase in CTQ score (HR 1.41, 95% CI 1.07–1.86).

We separately examined the association between sexual abuse and SLE risk. We did not observe an increased risk of SLE associated with any degree of sexual abuse in either age- and race-adjusted models, or in models additionally adjusted for parental and childhood factors.

### Secondary Analysis

We examined the risk of SLE associated with the alternative measure of physical assault (CTS). The correlation between CTQ and CTS responses was high ( $R^2 = 0.77$ ,  $p < 0.0001$ ). Consistent with our findings with the CTQ, we observed higher SLE risk among women who experienced medium to high levels of physical abuse during childhood or adolescence compared to those who experienced none (HR 1.74, 95% CI 1.11–2.73) (Supplemental Table 1).

### Discussion

In this longitudinal cohort followed for more than 24 years, we observed nearly tripled risk of incident SLE among women who had experienced high levels of childhood physical and emotional abuse compared to women who had not. Adjustment for potential parental and childhood confounders only slightly attenuated this association. In prospective analyses the association persisted, with more than three times higher risk among women exposed to childhood abuse compared to women who were not.

Our findings are in line with prior studies that similarly demonstrated higher risk of autoimmune diseases among individuals exposed to trauma and extreme stress.<sup>1, 3</sup> It is biologically plausible that adversity during childhood plays an important role in the development of autoimmune, inflammatory conditions during adulthood. Children who experienced maltreatment have been shown to have elevated markers of inflammation into adulthood.<sup>11, 38</sup> While the precise mechanism is unknown, one model suggests that

individuals who experienced significant early life adversity may have enhanced psychological and physiological stress sensitivity. When this sensitivity is compounded by fewer social and psychological resources to buffer stress, both psychological and physiological dysregulation may lead to immune dysregulation and inflammation.<sup>39</sup> At the biological level, the concept of “embedding” has been proposed whereby childhood stress results in epigenetic changes, in particular in the DNA of immune system cells, which then have heightened inflammatory tendencies.<sup>40</sup> This results in heightened cytokine and chemokine responses to stress and reduced sensitivity to inhibitory hormonal signals.<sup>40</sup> When combined with other exposures, including higher risk for unhealthy life choices and genetic risk factors, the increased inflammation may lower the threshold for the development of autoimmune diseases such as SLE.

Prior studies have demonstrated associations between perinatal and early life exposures and risk of SLE. Childhood farm residence as well as childhood exposure to agricultural pesticides have been associated with increased SLE risk.<sup>41</sup> Both preterm birth and birthweight >10lbs have similarly been linked to increased rates of SLE compared to at-term birth and normal birthweight, respectively.<sup>24, 41</sup> While childhood adversity has not been previously studied in the context of SLE incidence, in a cohort of 166 patients, four or more adverse childhood experiences was associated with more severe lupus-related damage in adulthood.<sup>42</sup> Although we did not have the ability to examine cumulative lifetime exposure to trauma, prior studies suggest the unique impact of adverse childhood exposures on subsequent inflammatory and autoimmune disease risk.<sup>11, 15</sup>

The percentage of women reporting experiences of childhood abuse in NHSII is comparable with other large U.S.-based studies. Among >17,000 health maintenance organization members, 64% reported at least one of eight categories of adverse childhood experiences, including emotional, physical or sexual abuse.<sup>43</sup> We found a significant association between high levels of childhood physical and emotional abuse and risk of SLE that persisted after adjusting for childhood confounders. Adult BMI, smoking and depression were modest mediators, however the association persisted after adjustment for these factors. PTSD explained the largest percentage (23%) of the association between the highest level of abuse and SLE risk. While we do not know the etiology of the women’s PTSD, prior studies demonstrate that childhood abuse increases vulnerability to PTSD in adulthood.<sup>44</sup> In a study also within the NHSII cohort, the highest number of PTSD symptoms was associated with nearly three times the risk of SLE.<sup>3</sup> It is plausible that individuals who do not develop PTSD related to this childhood trauma may not have as strong of a risk of SLE compared to those who do. There are no studies that examine the role of resilience and other protective factors that may mitigate the risk of autoimmune disease among individuals exposed to severe stressors.

We did not find a significant association between sexual abuse and SLE risk. The lack of association may be a result of under-reporting of sexual abuse, which has been shown previously.<sup>45</sup> In our population, 10% of women reported exposure to moderate or high levels of sexual abuse. In a sample with more individuals exposed to higher levels of sexual abuse, an association might have been observed. It is also plausible that there are more established resources in place to detect and mitigate the effects of sexual abuse but emotional abuse

during childhood may remain hidden and have longer-term psychological and physical effects.<sup>46, 47</sup> It may also be possible that sexual abuse does not contribute to SLE risk. A prior study that demonstrated an association between childhood adverse experiences and autoimmune disease used a composite measure and did not separate sexual abuse from physical and emotional abuse.<sup>15</sup> In this cohort, while we observed a strong correlation, as expected, between the CTQ and the CTS, physical and emotional abuse was not strongly correlated with sexual abuse ( $R^2=0.29$ ,  $p<0.0001$ ).

There are limitations to this work. This is a predominately white cohort of female nurse professionals and our findings may not be broadly generalizable (e.g. to men, other racial/ethnic groups, non-professionals). SLE disproportionately affects non-white racial/ethnic groups and lower socioeconomic status individuals and further studies are needed in diverse cohorts to determine whether these findings replicate. Experiences of childhood abuse were self-reported in adulthood and may be subject to recall bias, particularly among adults with chronic illnesses. However, in prospective analyses restricted to individuals who developed SLE after the report of childhood abuse, the association persisted. There may be misclassification of exposure because women may underreport experiences of both physical and sexual abuse.<sup>45, 48</sup> Individuals may not report abuse for a number of potential reasons including embarrassment, symptoms of victimization, a conscious desire to forget, or a lack of trust in the study.<sup>48, 49</sup> While we were able to consider some markers of parental socioeconomic status, and perinatal and childhood exposures, residual or unmeasured confounding is possible. We did not have information regarding other childhood stressors (e.g. food insecurity, housing instability, parental income) which may confound the observed association, or have an additive effect.

Our study also has several strengths. To our knowledge, this is the first longitudinal cohort study to specifically examine the relationship between childhood physical, emotional and sexual abuse and risk of incident SLE. We utilized a large, richly characterized cohort with more than 24 years of follow-up data including information on parental sociodemographic factors, childhood and adulthood exposures. In addition, we confirmed our SLE cases by chart review and all met ACR criteria. We examined potential confounders and mediators, and we restricted our cases to those occurring after report of SLE to minimize recall bias.

In this cohort of U.S.-based women, we observed a significant association between experiences of childhood physical and emotional abuse and incident SLE. Our study adds further evidence that exposure to severe childhood stressors, even after adjusting for parental socioeconomic factors, increases the risk of autoimmune disease. With findings that suggest that experiences of significant childhood abuse are likely more prevalent than previously appreciated, programs are needed to both develop effective prevention strategies, and to mitigate the far-reaching effects among those previously exposed. In addition, this study provides impetus for providers to actively screen patients for experiences of childhood abuse, as well as for onset of depression and PTSD given their association with autoimmune disease risk.



## Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

## Funding:

This study was supported by NIH K23 AR071500 (CH Feldman) and NIH R01 AR057327 and K24 AR066109 (KH Costenbader). We acknowledge the Channing Division of Network Medicine, Department of Medicine, Brigham and Women's Hospital and Harvard Medical School for its management of the Nurses' Health Study II. The NHSII is supported by UM1 CA176726.

## References

1. Song H, Fang F, Tomasson G, Arnberg FK, Mataix-Cols D, Fernandez de la Cruz L, et al. Association of Stress-Related Disorders With Subsequent Autoimmune Disease. *JAMA* 2018;319:2388–2400. [PubMed: 29922828]
2. O'Donovan A, Cohen BE, Seal KH, Bertenthal D, Margaretten M, Nishimi K, et al. Elevated risk for autoimmune disorders in Iraq and Afghanistan veterans with posttraumatic stress disorder. *Biol Psychiatry* 2015;77:365–374. [PubMed: 25104173]
3. Roberts AL, Malspeis S, Kubzansky LD, Feldman CH, Chang SC, Koenen KC, et al. Association of Trauma and Posttraumatic Stress Disorder With Incident Systemic Lupus Erythematosus in a Longitudinal Cohort of Women. *Arthritis Rheumatol* 2017;69:2162–2169. [PubMed: 28929625]
4. Glaser R, Kiecolt-Glaser JK. Stress-induced immune dysfunction: implications for health. *Nat Rev Immunol* 2005;5:243–251. [PubMed: 15738954]
5. Gutierrez MA, Garcia ME, Rodriguez JA, Rivero S, Jacobelli S. Hypothalamic-pituitary-adrenal axis function and prolactin secretion in systemic lupus erythematosus. *Lupus* 1998;7:404–408. [PubMed: 9736324]
6. Pace TW, Heim CM. A short review on the psychoneuroimmunology of posttraumatic stress disorder: from risk factors to medical comorbidities. *Brain Behav Immun* 2011;25:6–13. [PubMed: 20934505]
7. Taylor SE, Way BM, Seeman TE. Early adversity and adult health outcomes. *Dev Psychopathol* 2011;23:939–954. [PubMed: 21756443]
8. Slopen N, Koenen KC, Kubzansky LD. Cumulative adversity in childhood and emergent risk factors for long-term health. *J Pediatr* 2014;164:631–638 e631–632. [PubMed: 24345452]
9. Slopen N, Kubzansky LD, McLaughlin KA, Koenen KC. Childhood adversity and inflammatory processes in youth: a prospective study. *Psychoneuroendocrinology* 2013;38:188–200. [PubMed: 22727478]
10. Taylor SE, Lehman BJ, Kiefe CI, Seeman TE. Relationship of early life stress and psychological functioning to adult C-reactive protein in the coronary artery risk development in young adults study. *Biol Psychiatry* 2006;60:819–824. [PubMed: 16712805]
11. Danese A, Pariante CM, Caspi A, Taylor A, Poulton R. Childhood maltreatment predicts adult inflammation in a life-course study. *Proc Natl Acad Sci U S A* 2007;104:1319–1324. [PubMed: 17229839]
12. Thomas C, Hypponen E, Power C. Obesity and type 2 diabetes risk in midadult life: the role of childhood adversity. *Pediatrics* 2008;121:e1240–1249. [PubMed: 18450866]
13. Von Korf M, Alonso J, Ormel J, Angermeyer M, Bruffaerts R, Fleiz C, et al. Childhood psychosocial stressors and adult onset arthritis: broad spectrum risk factors and allostatic load. *Pain* 2009;143:76–83. [PubMed: 19251363]
14. Dong M, Giles WH, Felitti VJ, Dube SR, Williams JE, Chapman DP, et al. Insights into causal pathways for ischemic heart disease: adverse childhood experiences study. *Circulation* 2004;110:1761–1766. [PubMed: 15381652]
15. Dube SR, Fairweather D, Pearson WS, Felitti VJ, Anda RF, Croft JB. Cumulative childhood stress and autoimmune diseases in adults. *Psychosom Med* 2009;71:243–250. [PubMed: 19188532]

16. Hochberg MC Updating the American College of Rheumatology revised criteria for the classification of systemic lupus erythematosus. *Arthritis Rheum* 1997;40:1725.
17. Karlson EW, Sanchez-Guerrero J, Wright EA, Lew RA, Daltroy LH, Katz JN, et al. A connective tissue disease screening questionnaire for population studies. *Ann Epidemiol* 1995;5:297–302. [PubMed: 8520712]
18. Bernstein DP, Fink L, Handelsman L, Foote J, Lovejoy M, Wenzel K, et al. Initial reliability and validity of a new retrospective measure of child abuse and neglect. *Am J Psychiatry* 1994;151:1132–1136. [PubMed: 8037246]
19. Roberts AL, Lyall K, Rich-Edwards JW, Ascherio A, Weisskopf MG Association of maternal exposure to childhood abuse with elevated risk for autism in offspring. *JAMA Psychiatry* 2013;70:508–515. [PubMed: 23553149]
20. Straus MA, Hamby SL, Finkelhor D, Moore DW, Runyan D Identification of child maltreatment with the Parent-Child Conflict Tactics Scales: development and psychometric data for a national sample of American parents. *Child Abuse Negl* 1998;22:249–270. [PubMed: 9589178]
21. Lidfeldt J, Li TY, Hu FB, Manson JE, Kawachi I A prospective study of childhood and adult socioeconomic status and incidence of type 2 diabetes in women. *Am J Epidemiol* 2007;165:882–889. [PubMed: 17284723]
22. Liberatos P, Link BG, Kelsey JL The measurement of social class in epidemiology. *Epidemiol Rev* 1988;10:87–121. [PubMed: 3066632]
23. Troy LM, Michels KB, Hunter DJ, Spiegelman D, Manson JE, Colditz GA, et al. Self-reported birthweight and history of having been breastfed among younger women: an assessment of validity. *Int J Epidemiol* 1996;25:122–127. [PubMed: 8666479]
24. Simard JF, Karlson EW, Costenbader KH, Hernan MA, Stampfer MJ, Liang MH, et al. Perinatal factors and adult-onset lupus. *Arthritis Rheum* 2008;59:1155–1161. [PubMed: 18668600]
25. Tedeschi SK, Barbhuiya M, Malspeis S, Lu B, Sparks JA, Karlson EW, et al. Obesity and the risk of systemic lupus erythematosus among women in the Nurses' Health Studies. *Semin Arthritis Rheum* 2017.
26. Sacks RM, Takemoto E, Andrea S, Dieckmann NF, Bauer KW, Boone-Heinonen J Childhood Maltreatment and BMI Trajectory: The Mediating Role of Depression. *Am J Prev Med* 2017;53:625–633. [PubMed: 28928037]
27. Must A, Phillips SM, Stunkard AJ, Naumova EN Expert opinion on body mass index percentiles for figure drawings at menarche. *Int J Obes Relat Metab Disord* 2002;26:876–879. [PubMed: 12037661]
28. Costenbader KH, Feskanich D, Stampfer MJ, Karlson EW Reproductive and menopausal factors and risk of systemic lupus erythematosus in women. *Arthritis Rheum* 2007;56:1251–1262. [PubMed: 17393454]
29. Boynton-Jarrett R, Wright RJ, Putnam FW, Lividoti Hibert E, Michels KB, Forman MR, et al. Childhood abuse and age at menarche. *J Adolesc Health* 2013;52:241–247. [PubMed: 23332491]
30. Roberts AL, Kubzansky LD, Malspeis S, Feldman CH, Costenbader KH Association of Depression With Risk of Incident Systemic Lupus Erythematosus in Women Assessed Across 2 Decades. *JAMA Psychiatry* 2018.
31. Springer KW, Sheridan J, Kuo D, Carnes M Long-term physical and mental health consequences of childhood physical abuse: results from a large population-based sample of men and women. *Child Abuse Negl* 2007;31:517–530. [PubMed: 17532465]
32. Wolfe VV, Gentile C, Wolfe D The impact of sexual abuse on children: A PTSD formulation. *Behavioral Therapy* 1989;20:215–228.
33. Berwick DM, Murphy JM, Goldman PA, Ware JE Jr., Barsky AJ, Weinstein MC Performance of a five-item mental health screening test. *Med Care* 1991;29:169–176. [PubMed: 1994148]
34. Schnurr P, Vieilhauer M, Weathers F. The brief trauma questionnaire. In: *NCf PTSD* (ed). White River Junction, VT, 1999.
35. Breslau N, Peterson EL, Kessler RC, Schultz LR Short screening scale for DSM-IV posttraumatic stress disorder. *Am J Psychiatry* 1999;156:908–911. [PubMed: 10360131]

36. Barbaiya M, Lu B, Sparks JA, Malspeis S, Chang SC, Karlson EW, et al. Influence of Alcohol Consumption on the Risk of Systemic Lupus Erythematosus Among Women in the Nurses' Health Study Cohorts. *Arthritis Care Res (Hoboken)* 2017;69:384–392. [PubMed: 27273851]
37. Hertzmark E, Pazaris M, Spiegelman D. The SAS MEDIATE Macro. Harvard T.H. Chan School of Public Health 2018.
38. Pace TW, Mletzko TC, Alagbe O, Musselman DL, Nemeroff CB, Miller AH, et al. Increased stress-induced inflammatory responses in male patients with major depression and increased early life stress. *Am J Psychiatry* 2006;163:1630–1633. [PubMed: 16946190]
39. Fagundes CP, Glaser R, Kiecolt-Glaser JK Stressful early life experiences and immune dysregulation across the lifespan. *Brain Behav Immun* 2013;27:8–12. [PubMed: 22771426]
40. Miller GE, Chen E, Parker KJ Psychological stress in childhood and susceptibility to the chronic diseases of aging: moving toward a model of behavioral and biological mechanisms. *Psychol Bull* 2011;137:959–997. [PubMed: 21787044]
41. Parks CG, D'Aloisio AA, Sandler DP Early Life Factors Associated with Adult-Onset Systemic Lupus Erythematosus in Women. *Front Immunol* 2016;7:103. [PubMed: 27064771]
42. DeQuattro K, Trupin L, Katz P, Lanata C, Yelin E, Criswell LA THU0600 Cumulative adverse childhood experiences are associated with poor outcomes in adults with systemic lupus erythematosus [abstract]. *BMJ* 2017.
43. Dube SR, Anda RF, Felitti VJ, Chapman DP, Williamson DF, Giles WH Childhood abuse, household dysfunction, and the risk of attempted suicide throughout the life span: findings from the Adverse Childhood Experiences Study. *JAMA* 2001;286:3089–3096. [PubMed: 11754674]
44. McLaughlin KA, Conron KJ, Koenen KC, Gilman SE Childhood adversity, adult stressful life events, and risk of past-year psychiatric disorder: a test of the stress sensitization hypothesis in a population-based sample of adults. *Psychol Med* 2010;40:1647–1658. [PubMed: 20018126]
45. Widom CS, Morris S Accuracy of Adult Recollections of Childhood Victimization: Part 2. Childhood Sexual Abuse. *Psychological Assessment* 1997;9:34–46.
46. Spertus IL, Yehuda R, Wong CM, Halligan S, Seremetis SV Childhood emotional abuse and neglect as predictors of psychological and physical symptoms in women presenting to a primary care practice. *Child Abuse Negl* 2003;27:1247–1258. [PubMed: 14637300]
47. Trickett PK, Mennen FE, Kim K, Sang J Emotional abuse in a sample of multiply maltreated, urban young adolescents: issues of definition and identification. *Child Abuse Negl* 2009;33:27–35. [PubMed: 19178945]
48. Widom CS, Shepard RL Accuracy of Adult Recollections of Childhood Victimization: Part 1. Childhood Physical Abuse. *Psychological Assessment* 1996;8:412–421.
49. Della Femina D, Yeager CA, Lewis DO Child abuse: adolescent records vs. adult recall. *Child Abuse Negl* 1990;14:227–231. [PubMed: 2340430]

**Table 1.** Age-standardized characteristics in 1989 of Nurses' Health Study II Participants in Current Analysis according to Child Trauma Questionnaire (CTQ) response categories (N=67,516) and sexual abuse categories (N=67,480)

Characteristic	CTQ Response Category N=67,516				Sexual Abuse Response Category N=67,480		
	None/Very Low N=14,703	Low N=16,589	Moderate N=20,153	High N=16,071	None N=44,841	Low N=15,907	Moderate/High N=6,732
Age in 1989 - mean ± SD	34.8 ± 4.8	34.5 ± 4.7	34.4 ± 4.6	34.8 ± 4.5	34.5 ± 4.7	34.8 ± 4.6	34.8 ± 4.6
Non-white race - %	4.6	4.9	6.4	8.3	5.3	7.2	8.7
Parental education - %							
high school	43.3	44.9	47.0	49.8	45.2	48.0	50.0
Some college	32.1	32.8	32.4	33.6	32.7	32.4	33.2
College and above	24.6	22.4	20.6	16.6	22.1	19.5	16.8
Parental occupation - %							
Mother works in home	62.9	62.5	61.8	58.7	62.8	60.4	55.5
Farmer	7.1	6.1	5.5	4.3	5.6	5.9	6.0
Laborer	7.8	8.3	9.3	11.1	8.7	9.8	10.8
Blue/lower white collar	46.4	49.3	50.9	52.8	49.4	51.2	51.1
White collar	27.6	25.3	22.9	18.3	25.0	21.4	17.6
Parental home ownership - %	52.3	48.9	47.9	44.0	49.2	47.0	44.6
Birthweight - %							
<5.5 lbs	6.3	6.1	6.6	7.7	80.6	79.3	76.4
5.5-9.9 lbs	81.2	81.5	79.8	77.1	6.6	6.6	7.5
10 lbs	1.1	1.0	1.0	1.1	1.0	1.1	1.2
Age 5 Somatogram 5* - %	7.8	8.2	8.6	10.0	8.4	8.6	10.8
Menarche at age 10 years - %	7.0	7.2	7.6	8.6	7.1	7.8	10.8
Alcohol intake - %							
None	38.2	36.5	37.0	38.6	37.0	37.3	40.9
0-5g/day	42.2	42.7	42.7	41.2	42.9	41.6	39.0
5g/day	19.6	20.9	20.4	20.3	20.1	21.1	20.0
Zip code median household income in 1989 - median \$ (IQR)	56,907 (45,516, 71,151)	56,890 (45,417, 71,795)	56,935 (45,545, 71,743)	56,450 (45,162, 71,082)	58,085 (46,434, 72,520)	56,479 (45,250, 71,250)	55,135 (44,150, 69,389)

Characteristic	CTQ Response Category N=67,516				Sexual Abuse Response Category N=67,480		
	None/ Very Low N=14,703	Low N=16,589	Moderate N=20,153	High N=16,071	None N=44,841	Low N=15,907	Moderate/High N=6,732
Smoking status- %							
Never	71.3	68.6	65.7	59.8	68.6	62.8	58.0
Past	19.0	20.1	22.1	25.8	20.4	23.9	26.3
Current	9.8	11.3	12.2	14.5	11.1	13.3	15.8
BMI- %							
18.5-<25	70.5	69.1	67.7	64.0	69.4	66.1	61.4
25-<30	16.5	17.5	17.9	19.0	16.9	19.3	20.0
30+	8.6	9.5	10.6	13.2	9.6	11.1	15.4
Depression <sup>#</sup> - %							
None	76.2	72.5	66.6	56.0	70.9	64.9	53.4
Depression	23.8	27.5	33.4	44.0	29.1	35.1	46.6
Trauma and PTSD <sup>‡</sup> - %							
No trauma, no PTSD	41.1	35.1	28.7	14.7	36.5	19.9	6.4
Trauma, no PTSD	37.8	39.8	40.5	37.4	38.8	42.5	31.8
PTSD low score	15.1	17.0	19.4	23.9	16.2	22.8	28.1
PTSD moderate + high score	6.0	8.2	11.5	24.1	8.4	14.8	33.7

Values except for age are age-standardized to the age distribution of the study population

\* Age 5 somatogram score represents the diagram that the participant chose that best represents her body type ranging from 1 (very thin) to 9 (very obese)

# Measured by self-reported physician diagnosis, Mental Health Inventory-5 score <60, or use of antidepressant medication for subset of subjects with data available (N=67,469 for CTQ and N=66,904 for sexual abuse)

‡ PTSD available for subset of subjects (N=50,093 for CTQ and N=50,082 for sexual abuse)

**Table 2.**

Characteristics of incident SLE cases in NHS II cohort with 4 ACR classification criteria for SLE (N=94)

Characteristics	
ACR Criteria- mean $\pm$ SD	5.0 $\pm$ 1.0
Age at SLE diagnosis- mean $\pm$ SD	44.9 $\pm$ 8.7
ANA positive- N (%)	93 (98.9)
Anti-dsDNA positive- N (%)	49 (52.1)
Anti-Ro or Anti-La positive- N (%)	17 (18.1)
Anti-Sm positive- N (%)	6 (6.4)
Anti-RNP positive- N (%)	6 (6.4)
Arthritis- N (%)	69 (73.4)
Hematologic involvement- N (%)	60 (63.8)
Renal involvement- N (%)	11 (11.7)
Seen by an ACR member rheumatologist- N (%)	82 (87.2)

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Association of childhood physical and emotional abuse as measured by the CTQ with risk of incident SLE in the NHS II cohort (N=67,516)

**Table 3a.**

CTQ Response	SLE Cases/person-years	Model A HR (95% CI)	Model B HR (95% CI)	Model C HR (95% CI)
None/Very low	11/361,724	<i>Ref.</i>	<i>Ref.</i>	<i>Ref.</i>
Low	24/407,066	1.92 (0.94–3.92)	1.89 (0.92–3.87)	1.85 (0.90–3.79)
Moderate	25/493,200	1.66 (0.81–3.38)	1.58 (0.78–3.23)	1.52 (0.74–3.09)
High	34/386,815	<b>2.81 (1.42–5.56)</b>	<b>2.57 (1.30–5.12)</b>	<b>2.38 (1.19–4.74)</b>
p for trend		<b>p=0.004</b>	<b>p=0.01</b>	<b>p=0.03</b>

Model A: Age and race-adjusted

Model B: Age, race and parental and childhood factor (maternal and paternal occupation, parental education, parental home ownership, birthweight, age at menarche, somatogram score at age 5) adjusted

Model C: Age, race, parental and childhood factor, and adult BMI and smoking adjusted

**Table 3b.**

Association of childhood physical and emotional abuse as measured by the CTQ with risk of incident SLE in the NHS II cohort for subsets with depression (N=67,469) and PTSD data (N=50,093)

CTQ Response	SLE Cases/person-years	Model D1 HR (95% CI)	Model D2 HR (95% CI)	SLE Cases/person years	Model E1 HR (95% CI)	Model E2 HR (95% CI)
None/Very low	8/306,527	<i>Ref.</i>	<i>Ref.</i>	9/271,790	<i>Ref.</i>	<i>Ref.</i>
Low	17/344,728	1.86 (0.80-4.31)	1.77 (0.76-4.11)	23/308,528	<b>2.25 (1.04-4.88)</b>	2.12 (0.97-4.61)
Moderate	21/417,376	1.85 (0.82-4.19)	1.68 (0.74-3.81)	19/370,845	1.46 (0.66-3.25)	1.33 (0.60-2.96)
High	32/325,999	<b>3.31 (1.51-7.22)</b>	<b>2.71 (1.23-5.95)</b>	24/291,448	<b>2.22 (1.02-4.82)</b>	1.84 (0.83-4.07)
p for trend		<b>p=0.001</b>	<b>p=0.01</b>		p=0.18	p=0.44

Model D1: Age, race, parental and childhood factor (maternal and paternal occupation, parental education, parental home ownership, birthweight, age at menarche, somatogram score at age 5) adjusted for subset with depression data available

Model D2: Age, race, parental and childhood factor and adult history of depression adjusted for subset with depression data available

Model E1: Age, race, parental and childhood factor adjusted for subset with PTSD data available

Model E2: Age, race, paternal and childhood factor and adult history of PTSD adjusted for subset with PTSD data available



**Table 4.** Association of childhood sexual abuse with risk of incident SLE in the NHS II cohort (N=67,480)

Sexual Abuse Response Categories	SLE Cases/person-years	Age and race-adjusted model HR (95% CI)	Parental and childhood-factor adjusted model* HR (95% CI)
None	61/1,098,429	<i>Ref.</i>	<i>Ref.</i>
Low	25/388,919	1.14 (0.71–1.81)	1.13 (0.71–1.80)
Moderate/High	8/160,650	0.87 (0.42–1.83)	0.84 (0.40–1.77)
p for trend		p=0.81	p=0.68

\* Model adjusted for age, race, maternal and paternal occupation, parental education, parental home ownership, birthweight, age at menarche, somatogram score at age 5

Association of childhood physical and emotional abuse as measured by the CTQ with risk of incident SLE occurring after assessment of CTQ (2001) in the NHS II cohort (N=67,516)

**Table 5.**

CTQ Response	SLE Cases/person-years	Age and race-adjusted model HR (95% CI)	Parental and childhood-factor adjusted model* HR (95% CI)
None/Very low	4/190,651	<i>Ref.</i>	<i>Ref.</i>
Low	5/213,847	1.09 (0.29–4.07)	1.12 (0.30–4.20)
Moderate	13/258,615	2.38 (0.77–7.33)	2.33 (0.75–7.17)
High	14/200,449	<b>3.14 (1.03–9.56)</b>	<b>3.11 (1.01–9.57)</b>
p for trend		<b>p=0.01</b>	<b>p=0.01</b>

\* Model adjusted for age, race, maternal and paternal occupation, parental education, parental home ownership, birthweight, age at menarche, somatogram score at age 5