

Fatal *Sarcocystis cruzi*–induced eosinophilic myocarditis in a heifer in Uruguay

Journal of Veterinary Diagnostic Investigation
2019, Vol. 31(4) 656–660
© 2019 The Author(s)
Article reuse guidelines:
sagepub.com/journals-permissions
DOI: 10.1177/1040638719856651
jvdi.sagepub.com

Virginia Aráoz, Caroline da Silva Silveira, Gastón Moré,^{ID} Georget Banchemo,
Franklin Riet-Correa, Federico Giannitti¹ ^{ID}

Abstract. *Sarcocystis* spp. are causative agents of bovine eosinophilic myositis and/or myocarditis, which are chronic subclinical myopathies that are occasionally responsible for condemnation at slaughterhouses. *Sarcocystis cruzi* is a protozoan parasite of worldwide distribution transmitted by canids, most commonly associated with subclinical infection in cattle. Although *S. cruzi* infections can rarely lead to fatal systemic disease, fatal cardiac cases with confirmation of the etiologic diagnosis have not been reported, to our knowledge. We describe herein an unusual case of *S. cruzi*–induced fatal bovine eosinophilic myocarditis. A 22-mo-old, Holstein–Hereford heifer, in a group of 110 cattle on pasture, manifested growth retardation and died in February 2017. Autopsy revealed myriad yellow-green 1–3-mm coalescing foci, surrounded by fibrosis, affecting ~75% of the ventricular myocardium. Pulmonary edema, ascites, and hydrothorax were consistent with chronic congestive heart failure. Histology revealed severe eosinophilic, granulomatous, necrotizing myocarditis, with multinucleate giant cells, fibrosis, and mineralization. Numerous thin-walled protozoan cysts resembling *Sarcocystis* spp. were present in the necrotic foci and within the sarcoplasm of adjacent cardiomyocytes. PCR and sequencing of the 18S rRNA gene revealed 99.9–100% homology with *S. cruzi*. Sarcocystosis can be a rare cause of fatal myocarditis in cattle.

Key words: cattle; eosinophilic myocarditis; fatality; *Sarcocystis cruzi*; sarcocystosis.

Sarcocystis spp. are Apicomplexa protozoa of worldwide distribution that infect a wide range of vertebrates,⁴ typically without significant clinical disease. Species differ in life cycle, host specificity, morphology, and pathogenicity. *Sarcocystis* spp. have a life cycle with 2 obligate hosts: an intermediate host in which asexual reproduction occurs with tissue cyst formation, and a carnivore–omnivore definitive host, in which enteric sexual reproduction takes place with the release of sporocysts in feces.⁴

The main *Sarcocystis* spp. that parasitize cattle are *S. cruzi* (syn. *S. bovicanis*), *S. hirsuta*, and *S. hominis*, whose principal definitive hosts are dogs, cats, and humans, respectively. *Sarcocystis rommeli* (formerly misnamed *S. sinensis*) has also been described infecting cattle; however, its definitive host remains unknown.⁵ The species *S. bovifelis* was proposed instead of *S. rommeli* based on cyst morphology and the possible role of felids as definitive hosts.^{4,7} Morphologically, *S. cruzi* has thin-walled cysts (<1 μm); cysts of *S. hirsuta*, *S. rommeli*, and *S. hominis* have thicker walls (2–7 μm).


Generally, canids are definitive hosts for the *Sarcocystis* species that are the most virulent for intermediate hosts. The severity of clinical signs and lesions depends mainly on the number of ingested sporocysts and the immunologic status of the host.⁴ Eosinophilic myositis and myocarditis associated with *Sarcocystis* spp. infections in cattle are usually incidental postmortem findings but can be the reason for condemnation at slaughterhouses. Moreover, histologic

examination of cardiac and skeletal muscles in healthy ruminants from many countries reveals a high frequency of sarcocysts, up to 100%, without associated inflammation.^{13,15,18} Despite high prevalence, these parasites rarely cause clinical disease,^{4,6} thus sarcocystosis has been considered to be a subclinical myopathy (i.e. presence of muscle cysts with or without associated inflammation, but without detectable clinical signs).¹⁹ Herein we describe an unusual case of severe eosinophilic myocarditis with fatal outcome induced by *S. cruzi* infection in a heifer.

Our case occurred in February 2017, at a mixed dairy–beef farm in Colonia, Uruguay. The herd was composed of 110, 8–12-mo-old crossbred heifers and steers, and one 22-mo-old heifer. Cattle grazed on pastures in a 75-hectare area, with water supply from natural watercourses. The affected animal was a 22-mo-old Holstein–Hereford heifer, with a history of

Instituto Nacional de Investigación Agropecuaria (INIA), La Estanzuela, Colonia, Uruguay (Aráoz, Silveira, Banchemo, Riet-Correa, Giannitti); Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), Buenos Aires, Argentina (Moré); Laboratorio de Inmunoparasitología, Facultad de Ciencias Veterinarias, Universidad Nacional de La Plata, La Plata, Buenos Aires, Argentina (Moré).

¹Corresponding author: Federico Giannitti, Instituto Nacional de Investigación Agropecuaria (INIA), Ruta 50 km 11, La Estanzuela, Colonia, 70006, Uruguay. fgiannitti@inia.org.uy



Figures 1–4. Eosinophilic granulomatous myocarditis in a heifer. **Figure 1.** Myriad 1–3-mm, irregular, yellow-green coalescing foci disseminated throughout the myocardium are visible from the epicardial surface. **Figure 2.** Transmural section of the left ventricular free wall with granulomas and white streaks throughout the myocardium. **Figure 3.** A myocardial granuloma composed of a well-demarcated central area of necrosis surrounded by dense interstitial mixed inflammatory cell infiltrates and fibrosis. H&E. **Figure 4.** Abundant eosinophils, fewer lymphocytes, and macrophages infiltrate the myocardial interstitium, which is also expanded by fibrosis. Adjacent cardiomyocytes contain intact intrasarcoplasmic, thin-walled, basophilic *Sarcocystis cruzi* cysts. H&E.

growth retardation, whose clinical condition suddenly declined, and died spontaneously shortly thereafter.

Autopsy revealed myriad, irregular, yellow-green, 1–3-mm coalescing foci with variably distinct borders, and white striations disseminated throughout the myocardium (Figs. 1, 2). Lesions predominantly effaced the ventricular free walls and interventricular septum, and to a lesser extent the atrial myocardium, overall involving ~75% of the heart. In addition, there was moderate diffuse enhancement of the lobular pattern in the liver and diffuse pulmonary congestion and edema, with marked expansion of the interlobular septa, stable froth within the bronchial and tracheal lumens, mild ascites, and hydrothorax. These lesions were compatible with chronic congestive heart failure. No other gross lesions were identified.

Samples of heart, lung, brain, kidney, spleen, lymph node, skeletal muscles, tongue, liver, urinary bladder, and intestine were immersion-fixed in neutral-buffered formalin for 48 h, embedded in paraffin, sectioned at 4–5 μm , and stained with hematoxylin and eosin (H&E) for histologic and micromorphometric evaluation under an optical microscope (Axio-Scope.A1, Carl-Zeiss, Darmstadt, Germany) coupled with a digital camera (AxioCam-512, Carl-Zeiss) and ZEN software (Carl-Zeiss). Fresh heart was collected at autopsy and stored at -20°C .

Histology revealed severe, chronic, disseminated eosinophilic, granulomatous necrotizing myocarditis, with myocardial fibrosis, multinucleate giant cells, and mineralization. The granulomas had a well-demarcated necrotic center surrounded by dense interstitial mixed inflammatory cell infiltrates and fibrosis. The inflammatory infiltrates were characterized by abundant eosinophils, fewer lymphocytes, epithelioid macrophages, and occasional multinucleate giant cells that expanded the myocardial interstitium. Numerous preserved intra- and extra-lesional, ovoid, protozoan cysts containing uniformly sized elongated zoites were observed within the sarcoplasm of cardiomyocytes in, or adjacent to, areas of severe interstitial inflammation. Degenerate sarcocysts were occasionally present within the center of the necrotic foci and surrounded by a dense infiltrate of eosinophils, macrophages, and lymphocytes (Figs. 3, 4, Supplementary Figs. 1–4). Micromorphometric evaluation of 100 preserved cysts revealed that, on average, the major and minor diameters were 46 μm (20–205 μm) and 34 μm (15–73 μm). The cyst wall thickness was $\leq 1 \mu\text{m}$ in all examined cysts, regardless of their diameter. In the lungs, there was diffuse congestion of the alveolar capillaries, with edema and fibrin in the interstitium and alveolar spaces, and microhemorrhages and/or erythrocyte diapedesis into the alveolar

spaces, which occasionally contained uni- or bi-nucleate alveolar macrophages. Diffuse peri-acinar hepatocellular degeneration and necrosis were found in the liver. The pulmonary and hepatic lesions were attributed to chronic heart failure and terminal ischemia, respectively. Additionally, there was focal mild eosinophilic glossitis with very infrequent extralesional intrasarcoplasmic preserved sarcocysts. No significant findings were noted in other tissues; no sarcocysts were detected in skeletal muscle sections other than tongue.

DNA extracted from 2 frozen samples of heart (QIAamp cadaver pathogen mini kit; Qiagen, Hilden, Germany) was processed by real-time PCR with specific probes for *S. cruzi*, *S. hirsuta*, *S. hominis*, and *S. sinensis/S. rommeli*.¹⁴ A positive result was obtained only with the *S. cruzi*-specific probe. Additionally, a fragment of the *Sarcocystis* spp. 18S rRNA gene was amplified by PCR in both DNA samples. Amplification products were purified and sequenced.¹⁷ Sequences were aligned and analyzed (R9 software; Geneious, Auckland, NZ). Consensus sequences were compared with sequences available in GenBank by BLASTn analysis. Sequences of 844 and 843 bp were obtained and registered in GenBank (MK275239–MK275240), and both shared 100–99.9% identity with other sequences of *S. cruzi* (KT901167, KC209738, AB682780, JX679467, among others). *Sarcocystis levinei* (KU247922), a species identified in buffalo, was the next-highest match at 99.4% homology. Conventional PCRs for *Toxoplasma gondii* and *Neospora caninum* were negative.

The extensive and severe cardiac lesions, along with the pulmonary and hepatic lesions, were highly suggestive of chronic heart failure as the cause of death in this heifer. The numerous cysts compatible with *Sarcocystis* spp. within the heart, and the degenerate cysts within the necrotic centers of the myocardial granulomas, in conjunction with the molecular identification of *S. cruzi* and the absence of detection of any other causative agent, strongly argues that *S. cruzi* was the cause of the myocarditis.

Other conditions characterized by eosinophilic granulomatous myocarditis, such as helminthic and fungal infections, were ruled out by the absence of intralesional helminthic and fungal elements on H&E and Gomori methenamine silver stains. Syndromes associated with multisystemic eosinophilic granulomas described in cattle exposed to some toxicants, including plants such as *Vicia villosa* (hairy vetch),¹⁶ were ruled out by the pathology examination, which revealed that severe inflammation was restricted to the heart, and by the lack of history of exposure to these toxicants.

S. cruzi infection in cattle is usually subclinical, although acute and chronic clinical disease have been described in natural and experimental cases.⁴ Acute sarcocystosis, initially called “Dalmeny disease,” was first reported in Canada in 1961.² The clinicopathologic presentation is variable and depends on the dose of sporocysts ingested. Although experimental administration of $\leq 1,000$ sporocysts causes no clinical

signs, doses of ≥ 5 million sporocysts are invariably fatal. Cattle fed 200,000 sporocysts develop clinical disease, some die of acute sarcocystosis, whereas others survive the acute stage and show growth retardation (chronic sarcocystosis).⁴ Acute sarcocystosis, starting at ~ 4 wk postinfection, is characterized clinically by persistent fever, anorexia, weight loss, hair loss, anemia, weakness, muscle twitching, prostration, drooling, neurologic signs, and death; pregnant cows may abort, and lactating cows have reduced milk production.⁴ The histopathologic hallmark of acute sarcocystosis is the presence of numerous *S. cruzi* schizonts in vascular endothelial cells of many organs, along with vasculitis, necrosis, and/or inflammatory infiltrates in muscles, brain, placenta, and other tissues.^{1,4,12} Thus, acute sarcocystosis results from systemic vascular infection with early or immature *S. cruzi* asexual stages.

Unlike the acute presentation, chronic sarcocystosis is characterized clinically by growth retardation, and histopathologically by the presence of mature *S. cruzi* cysts containing bradyzoites within the sarcoplasm of striated myocytes that develop ~ 8 wk postinfection, without immature stages detectable in the vascular endothelium.⁴ Most cases of chronic sarcocystosis are not associated with significant lesions in infected muscles, other than the typical intrasarcoplasmic cysts.^{13,15} However, a small proportion of cattle chronically infected with *S. cruzi* develop eosinophilic myositis or myocarditis with bradyzoite-containing cysts.⁹

Even in cattle that develop considerable lesions, eosinophilic myositis or myocarditis are not apparent clinically. In most cases, the lesions are incidental postmortem findings at slaughterhouses.^{4,9} Typically, affected muscles have characteristic green discoloration detectable grossly and are trimmed from the carcass at meat inspection. Although fatal cases of eosinophilic myocarditis in cattle have been reported, to our knowledge, fatal cases attributable to chronic eosinophilic myocarditis with definitive identification of *S. cruzi* have not been documented.

To date, there is enough evidence, including experimental reproduction of the lesions, to causally associate bovine eosinophilic, granulomatous myositis or myocarditis with various *Sarcocystis* species, including the zoonotic *S. hominis*.^{3,4,18–20} However, the pathogenic mechanisms leading to lesion development are not understood completely. Hypersensitivity reactions are thought to play a role in cattle with *S. cruzi*-associated eosinophilic myocarditis detected at slaughter based on increased *S. cruzi*-specific IgE serum levels in affected animals, in contrast to cattle infected with *S. cruzi* but without lesions of eosinophilic myocarditis.⁹ Additionally, lesions have been reproduced experimentally by intramuscular injection of adjuvanted antigen of *Sarcocystis* obtained from cases of bovine eosinophilic myocarditis at slaughter,¹⁸ which also suggests that the lesion represents an immunologic reaction against *Sarcocystis* spp. antigen.

Cattle with eosinophilic myositis or myocarditis may be genetically predisposed to produce high levels of *Sarcocys-*

tis-specific IgE in response to bradyzoite antigen,⁹ as occurs in individuals with a genetic predisposition to type I hypersensitivity reactions to foreign antigens. This genetic predisposition hypothesis could explain the usual low morbidity of eosinophilic myositis or myocarditis, despite the high *Sarcocystis* spp. prevalence. The manifestation of clinical sarcocystosis with granulomatous and eosinophilic myositis induced by *Sarcocystis fayeri* in 2 genetically related horses by the same stallion also suggests a possible genetic basis for lesion and disease development.¹⁰ Additional theories for potential pathogenesis include infection by more pathogenic strains or genotypes of *Sarcocystis* spp., infective dose, breed, and the immune status of the intermediate hosts,⁴ or increased sarcolemmal fragility.

Although massive *Sarcocystis* spp. infections can lead to acute fatal disease, chronic fatal cases are seldom diagnosed. A fatal case of eosinophilic myocarditis that resulted in sudden death was reported in an 18-mo-old Hereford heifer in Iowa¹¹; however, no association with *Sarcocystis* infection was made. A presumptive case of chronic *Sarcocystis* spp.-induced eosinophilic myocarditis with fatal outcome was reported in a 2-y-old Hereford heifer in Uruguay.⁶ The etiologic diagnosis was based on morphologic evaluation of the protozoan, although the species involved was not confirmed. Interestingly, these cases, including our case, occurred sporadically in Hereford or Hereford-cross heifers 1.5–2 y old, which indirectly favors the hypothesis of a possible genetic or breed predisposition for eosinophilic myocarditis or myositis.

Although the source of infection was not determined in our case, domestic dogs and wild foxes are common in the area. The observation of cysts in myocytes, without schizonts recognized in blood vessels, suggested that the infection had taken place at least 8 wk before death.⁴ The nature of the lesions, particularly the mature interstitial myocardial fibrosis, was also consistent with a protracted course. The lesions involving the heart and the morphology of the thin-walled cysts strongly suggested *S. cruzi* as the species involved.^{4,20} This agrees with other studies in which sarcocysts were analyzed in meat samples from Uruguay, with *S. cruzi* the only species found.^{8,15} *S. cruzi* has been demonstrated to be the species with the highest prevalence (>90%) in several countries, including Argentina,^{13,14} noting that the myocardium is the most frequently affected muscle. Meanwhile, the prevalence of *S. hirsuta* and *S. hominis* is variable in different parts of the world, and these species locate predominantly in the esophagus and other muscles, rather than the myocardium.¹³ Molecular studies identifying *Sarcocystis* spp. infecting Uruguayan cattle are limited.¹⁵

Because several *Sarcocystis* spp. can be associated with eosinophilic myositis, including a zoonotic species, the need has been raised for species confirmation through molecular methods and/or transmission electron microscopy.¹⁹ In our case, *S. cruzi* was the only species identified by PCR and sequencing. *S. cruzi* should be considered an unusual cause of sporadic fatal myocarditis in cattle.

Acknowledgments

We thank Yisell Perdomo from INIA for technical assistance and Dr. Bradd Barr for critically reviewing this manuscript.

Declaration of conflicting interests

The authors declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.


Funding


This work was funded by grant PL-015 N-15156 from INIA.

Supplementary material

Supplementary material for this article is available online.

ORCID iDs

Gastón Moré  <https://orcid.org/0000-0002-0475-717X>

Federico Giannitti  <https://orcid.org/0000-0001-8799-6848>

References

1. Carrigan MJ. An outbreak of sarcocystosis in dairy cattle. *Aust Vet J* 1986;63:22–24.
2. Corner AH, et al. Dalmeny disease. An infection of cattle presumed to be caused by an unidentified protozoan. *Can Vet J* 1963;4:252–264.
3. Dubey JP. Clinical sarcocystosis in calves fed *Sarcocystis hirsuta* sporocysts from cats. *Vet Pathol* 1983;20:90–98.
4. Dubey JP, et al. *Sarcocystosis of Animals and Humans*. 2nd ed. Boca Raton, FL: CRC Press, 2015.
5. Dubey JP, et al. *Sarcocystis rommeli*, n. sp. (Apicomplexa: Sarcocystidae) from cattle (*Bos taurus*) and its differentiation from *Sarcocystis hominis*. *J Eukaryot Microbiol* 2016;63:62–68.
6. Dutra F. Miocarditis sarcocística en vaquillona [Sarcocystic myocarditis in a heifer]. Treinta y Tres, Uruguay: División de Laboratorios Veterinarios, Ministerio de Ganadería Agricultura y Pesca, 2014. *Archivo Veterinario del Este* 22–23. Spanish.
7. Gjerde B. The resurrection of a species: *Sarcocystis bovifelis* Heydorn et al., 1975 is distinct from the current *Sarcocystis hirsuta* in cattle and morphologically indistinguishable from *Sarcocystis sinensis* in water buffaloes. *Parasitol Res* 2016;115:1–21.
8. Gjerde B. Molecular characterization of *Sarcocystis bovifelis*, *Sarcocystis bovini* n. sp., *Sarcocystis hirsuta* and *Sarcocystis cruzi* from cattle (*Bos taurus*) and *Sarcocystis sinensis* from water buffaloes (*Bubalus bubalis*). *Parasitol Res* 2016;115:1473–1492.
9. Granstrom DE, et al. Type-I hypersensitivity as a component of eosinophilic myositis (muscular sarcocystosis) in cattle. *Am J Vet Res* 1989;50:571–574.
10. Herd HR, et al. *Sarcocystis fayeri*-induced granulomatous and eosinophilic myositis in 2 related horses. *Vet Pathol* 2015;52:1191–1194.
11. Jaspers R. Bovine eosinophilic myocarditis. *Iowa State Univ Vet* 1962–3;25(2):article 7.
12. Landsverk T. An outbreak of sarcocystosis in a cattle herd. *Acta Vet Scand* 1979;20:238–244.

13. Moré G, et al. Prevalence of *Sarcocystis* spp. in Argentinean cattle. *Vet Parasitol* 2011;177:162–165.
14. Moré G, et al. Development of a multiplex real time PCR to differentiate *Sarcocystis* spp. affecting cattle. *Vet Parasitol* 2013;197:85–94.
15. Pritt B, et al. Detection of *Sarcocystis* parasites in retail beef: A regional survey combining histological and genetic detection methods. *J Food Protec* 2008;71:2144–2147.
16. Robinson WF, Robinson NA. Cardiovascular system. In: Maxie MG, ed. *Jubb, Kennedy, and Palmer's Pathology of Domestic Animals*. 6th ed. Vol. 3. St. Louis, MO: Elsevier, 2016:36.
17. Scioscia NP, et al. Pampas fox (*Lycalopex gymnocercus*) new intermediate host of *Sarcocystis svanaei* (Apicomplexa: *Sarcocystidae*). *Parasitol Int* 2017;66:214–218.
18. Vangeel L, et al. Intramuscular inoculation of cattle with *Sarcocystis* antigen results in focal eosinophilic myositis. *Vet Parasitol* 2012;183:224–230.
19. Vangeel L, et al. Different *Sarcocystis* spp. are present in bovine eosinophilic myositis. *Vet Parasitol* 2013;197:543–548.
20. Wouda W, et al. Eosinophilic myositis due to *Sarcocystis hominis* in a beef cow. *J Comp Pathol* 2006;135:249–235.