



Published in final edited form as:

J Am Chem Soc. 2017 February 08; 139(5): 1774–1777. doi:10.1021/jacs.6b12307.

Rhodium-Catalyzed Hydrofunctionalization: Enantioselective Coupling of Indolines and 1,3-Dienes


Xiao-Hui Yang, Vy M. Dong*

Department of Chemistry, University of California, Irvine, 4403 Natural Sciences I, Irvine, California, 92697 USA.

Abstract

We communicate a strategy for the hydrofunctionalization of 1,3-dienes *via* Rh-hydride catalysis. Conjugated dienes are coupled to nucleophiles to demonstrate the feasibility of novel C–C, C–O, C–S, and C–N bond forming processes. In the presence of a chiral JoSPOphos ligand, hydroamination generates chiral allylic amines with high regio- and enantioselectivity. Tuning both the pK_a and steric properties of an acid-additive is critical for enantiocontrol.

Graphical abstract



Conjugated dienes are raw materials for polymerization and attractive building blocks for medicinal and natural product synthesis.^{1,2} Inventing enantioselective strategies for functionalizing dienes is an important challenge that has inspired Pd-catalyzed hydroamination,³ Co-catalyzed hydrovinylation,⁴ and Ru-catalyzed hydrohydroxyalkylation.⁵ To expand the power of hydrofunctionalization, we envisioned using Rh-hydride catalysis to couple 1,3dienes and nucleophiles (Figure 1).⁶ Iridium and rhodium-hydrides have been used to transform allenes and/or alkynes into *electrophilic* metal- π -allyl intermediates, which undergo nucleophilic attack to form branched products (Figure 1,A).⁷ Building on this strategy, our laboratory achieved the first enantioselective C–N and C–C bond forming reactions with alkynes.⁸ In analogy to allenes and alkynes, we reasoned that conjugate dienes could be transformed *via* Rh- π -allyl intermediates to produce the corresponding 1,2- and/or 1,4-addition products (Figure 1,B). Herein, we report the generality of Rh-hydride catalysis for enabling atom-economic C–C, C–O, C–N, and C–S bond forming reactions

*Corresponding Author dongv@uci.edu.

ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website.

Experimental procedures and spectral data for all new compounds (PDF)

The authors declare no competing financial interests.

with dienes. In addition, we demonstrate the first hydroamination of unsymmetric dienes with excellent regio- and enantiocontrol.⁹

The mechanism for our proposed hydrofunctionalization of dienes is depicted in Figure 2. It would be critical to identify the right combination of a rhodium catalyst and Brønsted acid (HX) to generate a Rh-hydride catalyst **A** capable of diene insertion. On the basis of steric preference, we reasoned that this Rh-hydride insertion would favor the terminal double bond to form Rh- π -allyl intermediate **B**. From this intermediate, there are two likely pathways for nucleophilic addition. One pathway (path a) involves anion exchange of **B** with nucleophiles to provide complex **C**. Reductive elimination of **C** releases 1,2-addition product **D** or 1,4-addition product **E**. Alternatively, the nucleophile can attack *via* an S_N2' or S_N2 type step (path b) to generate product **D** or **E**.

With this hypothesis in mind, we chose to investigate (*E*)-1-(4-methoxyphenyl)-1,3-butadiene as the model substrate. We investigated a range of achiral bidentate phosphine ligands and acid additives. A summary of our most relevant findings is in Table 1. The 1,3-diene can be cross-coupled with various nucleophiles; promising yields were observed for C–C bond (51% yield), C–N bond (85% yield), C–O bond (37% yield), and C–S bond (61% yield) formations. In all cases, excellent regioselectivity for the 1,2-Markovnikov product was observed (>20:1 *rr*). Using acids with a wide p*K*_a range (2.3 to 6.8) was necessary to encompass nucleophiles such as diketones,¹⁰ indolines,^{8a} alcohols,¹¹ and thiophenols.¹² It appears that weaker nucleophiles performed better in the presence of stronger acid co-catalysts. With this promising reactivity established, we chose to optimize hydroamination due to the value of enantiopure amines in drug discovery.¹³

While there has been interest in enantioselective hydroamination of 1,3-dienes,¹⁴ most studies have focused on intramolecular variants.¹⁵ Prior to our study, Hartwig demonstrated the only example of an enantioselective, intermolecular hydroamination of 1,3-dienes; this study focused on cyclohexadiene.³ In contrast, enantioselective hydroamination of unsymmetric dienes had yet to be achieved. In considering this challenge, we chose indoline **1a** and (*E*)-1-phenyl-1,3-butadiene **2a** as model substrates (Table 2, A). Given our previous work on alkyne hydroamination,^{8a} our initial studies focused on using [Rh(COD)Cl]₂ as the precatalyst and *m*-xylylic acid **A1** as the additive. Ligand (*S,S*)-BDPP **L1**, which gave high enantioselectivity for alkyne hydroamination, promoted diene hydroamination to form allylic amine **3aa** in only 21% *ee* (17% yield, 12:1 *rr*). From an extensive ligand evaluation, we found the Josiphos ligand family **L2–L4** gave excellent regioselectivity (>20:1 *rr*) but only modest reactivity and enantiocontrol. In comparison, the related JoSPOphos ligand **L5**¹⁶ afforded **3aa** in 84% yield, 82% *ee* and >20:1 *rr*.

With ligand **L5** in hand, we next investigated the effect of the acid additive (Table 2, B). The structure and acidity of the carboxylic acid had a marked effect on both yields and enantioselectivities. As highlighted in Table 2, we found that the optimal acid additive was **A4** (80% yield, 90% *ee*, >20:1 *rr*). In comparison, acids bearing less α -substitution, such as 2-phenylacetic acid **A2** and 2,2-diphenylacetic acid **A3**, provided lower *ee*'s, 52% and 78%, respectively. On the other hand, bulky acids that were less acidic, such as **A5** and **A6**, gave diminished yields (40% and 38%, respectively). These results show the significance of

tuning both the pK_a and steric properties of the acid-additive in Rh-hydride catalysis. We anticipated that the carboxylate would undergo exchange more effectively with a methoxide versus chloride ligand.¹⁷ Thus, in an effort to further improve selectivity, we switched from using $[\text{Rh}(\text{COD})\text{Cl}]_2$ to $[\text{Rh}(\text{COD})\text{OMe}]_2$ and observed the desired product in 81% yield and 96% *ee*.

With this protocol, we explored the hydroamination of 1,3-diene **2a** with various indolines **1** (Table 3). The substituent on the indoline had a negligible impact on the yield, regio- and enantioselectivity. In all cases, we observed excellent regioselectivity (>20:1 *rr*) for the Markovnikov 1,2-addition isomers in preference to the *anti*-Markovnikov 1,2-addition or 1,4-addition isomers. This type of regioselectivity was observed by Meek, although *via* π -acidic Rh-catalysis.^{9h} In addition, we obtained the allylic amines with high enantioselectivities (>94% *ee*). A slight decrease in enantioselectivity was observed with indolines containing electron-donating substituents at the 5-position (**3ah**, 81% yield, 88% *ee*) compared with other substituents. Together, these results represent the first intermolecular hydroamination of unsymmetric 1,3-dienes with high regio- and enantiocontrol.⁹

Next, we studied the hydroamination of various 1,3-dienes **2** with indoline **1a** (Table 4). Products bearing both electron-donating and electron-withdrawing groups on the phenyl ring were observed in high regio- and enantioselectivities. Due to steric effects, the coupling of indoline with ortho-substituted substrates **2j** and **2k** gave a slight decrease in yields (73% and 69%, respectively). This protocol tolerated a heterocycle substituted 1,3-dienes such as **2l** (R = 2-furyl) and **2m** (R = 2-thienyl) and afforded the corresponding allylic amines (*R*)-**3la** (78% yield, 95% *ee*) and (*R*)-**3ma** (73% yield, 92% *ee*). In addition, alkyl-substituted 1,3-diene **2n** provided the product **3na** in 68% yield, however the regioselectivity (6:1 *rr*) and enantioselectivity (76% *ee*) were lower compared to aryl substituted 1,3-dienes. For 1,3-pentadiene **2o**, the desired product **3oa** was obtained in 73% yield with 75% *ee*. We derivatized indoline **3oa** *via* oxidation with DDQ to generate the corresponding indole **5**,¹⁸ whose absolute configuration has been reported.¹⁹

Hydrofunctionalization represents a promising way to transform dienes into enantiopure allylic motifs. Krische established the use of Ru- and Ir-hydrides to generate *nucleophilic* π -allyl species.^{5, 20} As a complementary approach, we developed Rh-hydride catalysts that transform dienes into *electrophilic* π -allyl species. By tuning the Rh-hydride's structure and acidity, we achieved an enantioselective hydroamination for indolines and terminal dienes.²¹ Future studies will focus on understanding the origin of 1,2-addition *vs.* 1,4-addition.

Moreover, we plan to design catalysts to expand scope and variants for enantioselective and regioselective C–C, C–N, C–O, and C–S bond formations.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

ACKNOWLEDGMENT

Funding provided by UC Irvine and the National Institutes of Health (GM105938). This manuscript is dedicated to Prof. Qi-Lin Zhou on the occasion of his 60th birthday.

REFERENCES

- (1) (a). For reviews on polymerization of 1,3-dienes, see: Thiele SKH; Wilson DR J. *Macromol. Sci. Part C: Polym. Rev* 2003, C43, 581.(b)Friebe L; Nuyken O; Obrecht W. *Adv. Polym. Sci* 2006, 204, 1.(c)Ricci G; Sommazzi A; Masi F; Ricci M; Boglia A; Leone G. *Coord. Chem. Rev* 2010, 254, 661.(d)Zhang ZC; Cui DM; Wang BL; Liu B; Yang Y. *Struct. Bonding (Berlin)* 2010, 137, 49.(e)Valente A; Mortreux A; Visseaux M; Zinck P. *Chem. Rev* 2013, 113, 3836. [PubMed: 23391234] (f)Takeuchi D. *Stereoselective Polymerization of Conjugated Dienes in Encyclopedia of Polymer Science and Technology*; Wiley: New York, 2013; pp 1–25
- (2) (a). For selected reviews on 1,3-dienes as building blocks, see: Reymond S; Cossy J. *Chem. Rev* 2008, 108, 5359. [PubMed: 18942879] (b)Zhu Y; Cornwall RG; Du H; Zhao B; Shi Y. *Acc. Chem. Res* 2014, 47, 3665. [PubMed: 25402963] (c)Chen JR; Hu X-Q; Lu L-Q; Xiao W-J. *Chem. Rev* 2015, 115, 5301. [PubMed: 25992465] (d)Heravi MM; Ahmadi T; Ghavidel M; Heidari B; Hamidi H. *RSC Adv.* 2015, 5, 101999.(e)Büschleb M; Dorich S; Hanessian S; Tao D; Schenthal KB; Overman LE. *Angew. Chem. Int. Ed* 2016, 55, 4156.
- (3). Löber O; Kawatsura M; Hartwig JF J. *Am. Chem. Soc* 2001, 123, 4366. [PubMed: 11457216]
- (4) (a). Sharma RK; RajanBabu TV J. *Am. Chem. Soc* 2010, 132, 3295. [PubMed: 20163120] (b)Page JP; RajanBabu TV J. *Am. Chem. Soc* 2012, 134, 6556. [PubMed: 22452442] (c)Timsina YN; Sharma RK; RajanBabu TV. *Chem. Sci* 2015, 6, 3994. [PubMed: 26430505]
- (5) (a). Zbieg JR; Yamaguchi E; McInturff EL; Krische MJ. *Science* 2012, 336, 324. [PubMed: 22442385] (b)Park BY; Montgomery TP; Garza VJ; Krische MJ. *J. Am. Chem. Soc* 2013, 135, 16320. [PubMed: 24156560] (c)Nguyen KD; Herkommer D; Krische MJ. *J. Am. Chem. Soc* 2016, 138, 14210. [PubMed: 27762549]
- (6). Hydrofunctionalization; Ananikov VP; Tanaka M, Eds.; In *Topics in Organometallic Chemistry*; Springer: Berlin, Heidelberg, 2014; Vol. 343. For selected examples on difunctionalization of 1,3-dienes, see:(a)Liao L; Jana R; Urkalan KB; Sigman MS. *J. Am. Chem. Soc* 2011, 133, 5784. [PubMed: 21449562] (b)McCammant MS; Liao L; Sigman MS. *J. Am. Chem. Soc* 2013, 135, 4167. [PubMed: 23473452] (c)McCammant MS; Sigman MS. *Chem. Sci* 2015, 6, 1355. [PubMed: 25705367] (d)Liu Y; Xie Y; Wang H; Huang HJ. *Am. Chem. Soc* 2016, 138, 4314. For selected examples on hydrofunctionalization of 1,3-dienes without enantioselectivity, see:(a)Liao L; Sigman MS. *J. Am. Chem. Soc* 2010, 132, 10209. [PubMed: 20608751] (b)Chen Q-A; Kim DK; Dong VM. *J. Am. Chem. Soc* 2014, 136, 3772. [PubMed: 24588202] (c)Saini V; O'Dair M; Sigman MS. *J. Am. Chem. Soc* 2015, 137, 608.(d)McNeill E; Ritter T. *Acc. Chem. Res* 2015, 48, 2330. [PubMed: 26214092]
- (7) (a). For Ir-hydride catalyzed hydrocarboxylation of allene, see: Kim IS; Krische MJ. *Org. Lett* 2008, 10, 513. [PubMed: 18181640] (b)For Rh-hydride catalyzed hydrofunctionalization of allenes and/or alkynes, see: Koschker P; Breit B. *Acc. Chem. Res* 2016, 49, 1524 and references therein. [PubMed: 27455048]
- (8) (a). For Rh-hydride catalyzed C–N bond formation, see: Chen Q-A; Chen Z; Dong VM. *J. Am. Chem. Soc* 2015, 137, 8392. [PubMed: 26107923] (b)For Rh-hydride catalyzed C–C bond formation, see: Cruz FA; Dong VM. *J. Am. Chem. Soc* 2016, DOI: 10.1021/jacs.6b10680.
- (9) (a). For enantioselective hydroamination of symmetric 1,3-dienes, see: ref 3. For intermolecular hydroamination of 1,3-dienes without enantioselectivity, see: Minami T; Okamoto H; Ikeda S; Tanaka R; Ozawa F; Yoshifuji M. *Angew. Chem. Int. Ed* 2001, 40, 4501.(b)Pawlas J; Nakao Y; Kawatsura M; Hartwig JF. *J. Am. Chem. Soc* 2002, 124, 3669. [PubMed: 11929257] (c)Brouwer C; He C. *Angew. Chem. Int. Ed* 2006, 45, 1744.(d)Qin H; Yamagiwa N; Matsunaga S; Shibasaki MJ. *Am. Chem. Soc* 2006, 128, 1611.(e)Johns AM; Liu Z; Hartwig JF. *Angew. Chem. Int. Ed* 2007, 46, 7259.(f)Giner X; Nájera C. *Org. Lett* 2008, 10, 2919. [PubMed: 18553970] (g)Banerjee D; Junge K; Beller M. *Angew. Chem. Int. Ed* 2014, 53, 1630.(h)Goldfogel MJ; Roberts CC;

- Meek SJ *J. Am. Chem. Soc.* 2014, 136, 6227. [PubMed: 24742315] (i) Banerjee D; Junge K; Beller M. *Org. Chem. Front* 2014, 1, 368.
- (10) (a). For C–C bond formation using 1,3-dicarbonyl compounds as nucleophiles, see: Cruz FA; Chen Z; Kurtoic SI; Dong VM *Chem. Comm* 2016, 52, 5836. [PubMed: 27043656] (b) Li C; Grugel CP; Breit B. *Chem. Commun* 2016, 52, 5840. (c) Beck TM; Breit B. *Org. Lett* 2016, 18, 124. [PubMed: 26683497]
- (11). For C–O bond formation using alcohols as nucleophiles, see: Liu Z; Breit B. *Angew. Chem. Int. Ed* 2015, 55, 8440.
- (12). For C–S bond formation using thiols as nucleophiles, see: Pritzius AB; Breit B. *Angew. Chem. Int. Ed* 2015, 54, 3121.
- (13) (a). For recent reviews, see: Michael JP *Nat. Prod. Rep* 2008, 25, 139. [PubMed: 18250900] (b) Nugent TC *Chiral Amine Synthesis: Methods, Developments and Applications*; Wiley-VCH: Weinheim, Germany, 2010. (c) Stereoselective Formation of Amines; Li W; Zhang X, Eds.; In *Topics in Current Chemistry*; Springer: Berlin, Heidelberg, 2014; Vol. 343. (d) Rouden J; Lasne M-C; Blanchet J; Baudoux J. *Chem. Rev* 2014, 114, 712. [PubMed: 24117010] (e) Grangea RL; Clizbe EA; Evans PA *Synthesis* 2016, 48, 2911.
- (14) (a). For recent reviews see: Hong S; Marks TJ *Acc. Chem. Res* 2004, 37, 673. [PubMed: 15379583] (b) Aillaud I; Collin J; Hannedouche J; Schulz E. *Dalton Trans.* 2007, 5105. [PubMed: 17985016] (c) Müller TE; Hultsch KC; Yus M; Foubelo F; Tada M. *Chem. Rev* 2008, 108, 3795. [PubMed: 18729420] (d) Bernoud E; Lepori C; Mellah M; Schulz E; Hannedouche J. *Catal. Sci. Technol* 2015, 5, 2017. (e) Huang L; Arndt M; Gooßen K; Heydt H; Gooßen LJ *Chem. Rev* 2015, 115, 2596. [PubMed: 25721762]
- (15) (a). Examples of intramolecular asymmetric hydroamination of 1,3-dienes, see: Hong S; Kawaoka AM; Marks TJ *J. Am. Chem. Soc.* 2003, 125, 15878. [PubMed: 14677980] (b) Zhang Z; Bender CF; Widenhofer RA *Org. Lett* 2007, 9, 2887. [PubMed: 17595096] (c) Deschamp J; Olier C; Schulz E; Guillot R; Hannedouche J; Collin J. *Adv. Synth. Catal* 2010, 352, 2171. (d) Shapiro ND; Rauniyar V; Hamilton GL; Wu J; Toste FD *Nature*, 2011, 470, 245. [PubMed: 21307938] (e) Kanno O; Kuriyama W; Wang ZJ; Toste FD *Angew. Chem. Int. Ed* 2011, 50, 9919. (f) Dion I; Beauchemin AM *Angew. Chem. Int. Ed* 2011, 50, 8233. (g) Deschamp J; Collin J; Hannedouche J; Schulz E. *Eur. J. Org. Chem* 2011, 3329. (h) Yang Z; Xia C; Liu D; Liu Y; Sugiya M; Imamoto T; Zhang W. *Org. Biomol. Chem* 2015, 13, 2694. [PubMed: 25588159]
- (16) (a). For the applications of JoSPOphos with Rh, see: Landert H; Spindler F; Wyss A; Blaser H-U; Pugin B; Ribourduille Y; Gschwend B; Ramalingam B; Pfaltz A. *Angew. Chem. Int. Ed* 2010, 49, 6873. (b) Wu X; Chen Z; Bai YB; Dong VM *J. Am. Chem. Soc.* 2016, 138, 12013. [PubMed: 27584734]
- (17) (a). Murphy SK; Bruch A; Dong VM *Chem. Sci* 2015, 6, 174. [PubMed: 25580215] (b) Murphy SK; Park J-W; Cruz FA; Dong VM *Science*, 2015, 347, 56. [PubMed: 25554782]
- (18). [See Supporting Information.]
- (19). Zhao Q; Zhuo C-X; You S-L *RSC Adv.* 2014, 4, 10875.
- (20) (a). Chen T-Y; Tsutsumi R; Montgomery TP; Volchkov I; Krische MJ *J. Am. Chem. Soc.* 2015, 137, 1798. [PubMed: 25642996] (b) Oda S; Franke J; Krische MJ *Chem. Sci* 2016, 7, 136. [PubMed: 29861974]
- (21). Under this protocol, amines that differ in basicity from indolines (e.g., aniline, morpholine, and dibenzylamine) showed no reactivity. Internal dienes, such as (1E,3E)-1,4-diphenyl-1,3-butadiene and (1E,3E)-1-phenyl-1,3-pentadiene, also showed no reactivity, presumably due to increased steric bulk.



A: Hydrofunctionalization of alkynes and allenes**B: Hydrofunctionalization of dienes**

Figure 1.
Rh-hydride catalyzed hydrofunctionalization of unsaturated compounds






Figure 2.
Reaction design for the hydrofunctionalization of 1,3-dienes

Table 1.

Constructing different C–X bonds *via* hydrofunctionalization of 1,3-diene^a

^aReaction conditions: NuH (0.2 mmol), 1,3-diene (0.3 mmol), [Rh(COD)Cl]₂ (2 mol%), ligand (4 mol%), acid (50 mol%), DCE (0.4 mL), 24 h. Isolated yields.

Table 2.

Ligand and acid effects on asymmetric hydroamination of 1,3-diene^a

^aReaction conditions: **1a** (0.1 mmol), **2a** (0.15 mmol), [Rh(COD)Cl]₂ (2 mol%), ligand (4 mol%), acid (50 mol%), DCE (0.2 mL), 60 °C, 24 h. Yields determined by ¹H NMR using an internal standard. Regioselectivity determined by ¹H NMR analysis of the unpurified reaction mixture. Enantioselectivity determined by chiral SFC.

^bUsing [Rh(COD)OMe]₂ instead of [Rh(COD)Cl]₂.


Author Manuscript

Author Manuscript

Author Manuscript


Author Manuscript

Table 3.

Hydroamination using various indolines^a

^aReaction conditions: **1** (0.2 mmol), **2a** (0.3 mmol), [Rh(COD)OMe]₂ (2 mol%), JoSPOphos (4 mol%), Ph₃CCOOH (50 mol%), DCE (0.4 mL), 60 °C, 24 h. Isolated yields. Regioselectivity determined by ¹H NMR analysis of the unpurified reaction mixture. Enantioselectivity determined by chiral SFC.

Table 4.

Hydroamination of various 1,3-dienes^a

^aReaction conditions: **1a** (0.2 mmol), **2** (0.3 mmol), $[\text{Rh}(\text{COD})\text{OMe}]_2$ (2 mol%), JoSPOphos (4 mol%), Ph_3CCOOH (50 mol%), DCE (0.4 mL), 60 °C, 24 h. Regioselectivity determined by ^1H NMR analysis of the unpurified reaction mixture. Isolated yields. Enantioselectivity determined by chiral SFC.