

## REVIEW ARTICLE

# Extracellular vesicles in urologic malignancies— Implementations for future cancer care

Zhangsong Wu<sup>1,2,3</sup>  | Zhiqiang Zhang<sup>2,3</sup> | Wuchao Xia<sup>3,4</sup> | Jiajia Cai<sup>3,4</sup> | Yuqing Li<sup>2,3</sup> |  
Song Wu<sup>1,2,3,4,5</sup>

<sup>1</sup>Medical College, Shenzhen University, Shenzhen, China

<sup>2</sup>Department of Urological Surgery, The Third Affiliated Hospital of Shenzhen University, Shenzhen University, Shenzhen, China

<sup>3</sup>Shenzhen Following Precision Medical Institute, The Third Affiliated Hospital of Shenzhen University, Shenzhen University, Shenzhen, China

<sup>4</sup>Medical College, Anhui University of Science and Technology, Huainan, China

<sup>5</sup>Department of Urological Surgery, The First Affiliated Hospital of Guangzhou Medical University, Guangzhou Medical University, Guangzhou, China

## Correspondence

Song Wu, Department of Urological Surgery, The Third Affiliated Hospital of Shenzhen University, Shenzhen University, Shenzhen 518000, China.

Email: doctor\_wusong@126.com

## Funding information

This work was financially supported by a scholarship from The National Key Research and Development Program of China (No. 2017YTA0105900).

## Abstract

Extracellular vesicles (EVs), a heterogeneous group of vesicles differing in size and shape, cargo content and function, are membrane-bound and nano-sized vesicles that could be released by nearly all variations of cells. EVs have gained considerable attention in the past decades for their functions in modulating intercellular signaling and roles as potential pools for the novel diagnostic and prognostic biomarkers, as well as therapeutic targets in several cancers including urological neoplasms. In general, human and animal cells both can release distinct types of EVs, including exosomes, microvesicles, oncosomes and large oncosomes, and apoptotic bodies, while the content of EVs can be divided into proteins, lipids and nucleic acids. However, the lack of standard methods for isolation and detection platforms rein the widespread usage in clinical applications warranted furthermore investigations in the development of reliable, specific and sensitive isolation techniques. Whether and how the EVs work has become pertinent issues. With the aid of high-throughput proteomics or genomics methods, a fully understanding of contents contained in EVs from urogenital tumours, beyond all doubt, will improve our ability to identify the complex genomic alterations in the process of cancer and, in turn, contribute to detect potential therapeutic target and then provide personalization strategy for patient.

## 1 | INTRODUCTION

Urogenital carcinomas are mainly leading to morbidity and mortality worldwide.<sup>1</sup> Although therapeutic strategies (surgery, biotherapy and chemotherapy) for patients suffering from urinary cancers have improved, the curative and monitoring efficacy for these cancers still remains poor, as existing tests (general examinations and biopsies) are not a sufficiently sensitive or specific and heterogeneous peculiarity of these malignancies.<sup>2</sup> Moreover, the deep location of urogenital cancers makes them hard to access to be diagnosed at early stages. Crosstalk between cells and their microenvironment

is a fundamental principle under the normal and pathological condition.<sup>3</sup> EVs, small membrane-bound vesicles, serve as important players of bidirectional communication,<sup>4,5</sup> which released from almost eukaryotic and prokaryotic cells with transmitting complicate messages from donor cells towards anchored cells, and have been discovered in various types of body fluids including urine, blood and bile.<sup>6</sup> According to the ways of production and secretion, EVs are produced by inward budding of intercellular endosomes, which are defined as exosomes (40-1200 nm), EVs straightly shed by budding from the cell membrane, which are almost recognized as large oncosomes or microvesicles, with 1-10 µm or 50-1500 nm, and apoptotic

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2019 The Authors. *Cell Proliferation* Published by John Wiley & Sons Ltd.

vesicles are released during cell undergoing apoptosis ranged from 50 to 2000 nm, respectively.<sup>7-9</sup>

Extracellular vesicles serve as an appealing source for the development of biomarkers as their membrane-bound structure to protect against exogenous proteases and RNases.<sup>10,11</sup> The biological function of EVs is performed by cytosolic lipids, proteins, DNA, mRNA, miRNA, lincRNA and other non-coding RNAs, as well as cell membrane.<sup>12</sup> In addition, cancer cell releases more EVs than that normal one does.<sup>13</sup> Herein, we introduce EVs briefly and provide a comprehensive overview of their biophysical properties, roles and applications in the most common urologic neoplasms, including kidney, prostate and bladder, and discuss potential clinical applications in the future.

## 2 | EV CLASSES, BIOGENESIS AND CONTENTS

Our current understanding of EVs indicates that at least four heterogeneous types of EVs have been identified based on their mechanism of formation and distinguished size: microvesicles, exosomes, oncosomes or large oncosomes, and apoptotic bodies (Table 1). In general, the formation of exosomes and microvesicles is two completely different approaches, but they function similarly. Oncosomes or large oncosomes resemble the way of microvesicles via membrane

budding. Apoptotic bodies specifically arise resulting in indiscriminate membrane bubbling during apoptosis (Figure 1).


Exosomes generally form an early endosome by the endocytosis and internalization of cell-surface receptors into membrane-bound vacuoles in the first step,<sup>14</sup> which then matures to generate a late endosome within undergoing several changes, such as the limiting membrane of the late endosome then buds inward and pinches off as the result of the formation of intraluminal vesicles (ILVs), also known as multivesicular bodies, and ILVs traffick to and fuse with the plasma membrane leading in releasing exosomes eventually.<sup>15</sup> ESCRT-0-III plays significant roles in driving exosome formation<sup>16,17</sup>; in addition, multivesicular bodies could intermediate in the lysosomal degradation pathway.<sup>18</sup> However, the mechanisms related to the fusion of multivesicular bodies with the cellular membrane are uncovered, which may be regulated by several factors including lipid ceramide and Rab GTPase (including Rab5 and Rab7) proteins and ESCRT.<sup>19-21</sup> Numerous literatures have indicated that several biomarkers expressed in the exosome differentially compared with another types of EVs, including heat shock proteins (eg HSP60, 70 and 90), tetraspanins (eg CD9, CD63 and CD81), membrane transporters, fusion proteins, ALG-2-interacting protein X (Alix) and tumour susceptibility gene 101 protein (TSG101).<sup>20,22</sup>

In contrast, microvesicles with nano-sized with 100-1500 nm are straightforwardly shed from the cellular membrane responding to stimuli or physiological conditions.<sup>23</sup> It is believed that ADP ribosylation

**TABLE 1** Details of different extracellular vesicles

	Exosomes	Microvesicles	Oncosomes	Large oncosomes	Apoptotic bodies
Size	40-120 nm	50-1500 nm	100-500 nm	1-10 $\mu$ m	50-2000 nm
Intracellular origin	Endosomes	Plasma membrane	Plasma membrane	Plasma membrane	Plasma membrane
Electron microscopy	Round shape	Irregular shape	Irregular shape	Amoeboid phenotype	Heterogenous
Release	Endolysosomal pathway, internal budding, exocytosis	Membrane budding	Membrane budding	Membrane budding	Generated as a result of apoptotic disintegration, resulting vesicles become part of the extracellular milieu
Marker proteins	Membrane-associated proteins: tetraspanin (CD9, CD63, CD81, CD82). Endosomal sorting complex required for Transport-associate protein: Tsg101, ALIX. Cytoplasmic proteins: Hsp70, Hsp90. Membrane transport and fusion proteins: Rab GTPases, annexins	Integrins, selectins, CD40 ligand	Integrins, selectins, Membrane-associated proteins	Integrins, selectins, VDAC $\frac{1}{2}$ SLC25A3/5/6 ITGA5/6	Histones, C3b, Annexin V, Caspase 3
Contents	Proteins, lipids, mRNA, miRNA and cytosol	Proteins, lipids, mRNA, miRNA and cytosol	Proteins, lipids, mRNA, miRNA and cytosol	Proteins, lipids, mRNA, miRNA and cytosol	Proteins, lipids, DNA, rRNA, organelles and cytosol

Abbreviations: C3b, complement component 3; HSP, heat shock proteins; ITGA5/6, human integrin alpha 5/6; miRNA, microRNA; SLC25A3/5/6, solute carrier family 25 member3/5/6; VDACS, Voltage-dependent anion channels.



**FIGURE 1** Release and uptake mechanisms of extracellular vesicles. Extracellular vesicles can be classed as exosomes, microvesicles and apoptotic bodies, based on the mechanism by which they are released from cells and differentiated based on their size and content. MVs, and oncosomes or large oncosomes are directly shed or bud from the plasma membrane. Apoptotic bodies are released from the cell undergoing programmed cell death. Exosomes are formed by inward budding of multivesicular bodies

factor 6 (ARF6) can mediate the freeing of protease-loaded vesicles from the cellular membrane due to the crosstalk with Rho signalling pathways.<sup>24-26</sup> Moreover, microvesicles are specifically produced in the cellular membrane regions that are linked to be enriched in cholesterol, ceramide and lipid rafts.<sup>27</sup> TSG101 is also known to interact with accessory proteins Alix and arresting domain-containing protein-1 (ARRDC1) during releasing microvesicles, illustrating that microvesicles sharing some same characteristics with exosomal biogenesis.<sup>28</sup> As described for microvesicles, oncosomes and large oncosomes are generated by plasma membrane budding, with amoeboid-like phenotype. Notably, oncosomes and large oncosomes specially derived from cancerous cells are indicated to play vital roles in malignancies invasion. The term "oncosome" is firstly referred that the EVs with size range from 100 to 500 nm. Subsequently, large non-apoptotic EVs were detected in prostate tumours with their unusual size in 1-10  $\mu\text{m}$ , called large oncosomes. Currently, some studies demonstrate that oncosomes and large oncosomes are definitely different variety of EVs respecting to their size, cargo contents and target effects, and additional studies are therefore of the essence to clarify differences between oncosomes and large oncosomes.

Apoptotic bodies, imposing an effect on the cellular response by transmitting their substance towards receptor cells, are generated during undergoing programmed cell death with 500-4000 nm, and their content contains fragmented cytoplasmic organelles as well as destructive nuclei.<sup>29,30</sup>

### 3 | ISOLATION TECHNIQUES OF EVS

No remarkable consensus is in existence of the best approach for isolation, qualitative and quantitative analysis of EVs. There are listing several methods for the isolation of EVs (Figure 2) and demonstrate the available disadvantages and advantages as well (Table 2).<sup>31</sup>


### 4 | CENTRIFUGATION

In recent, differential centrifugation is the most common technique in responding to isolating EVs, and this approach is consisting of three main centrifugation processes: low speed to eliminate a main portion of the cells, then intermediate speed to subside cell debris and aggregate biopolymers and the other structures with density higher than that of EVs and finally high speed to pellet extracellular vesicles. The advent of density gradient ultracentrifugation increases the efficiency of particle separation according to their buoyant density. However, an important disadvantage of differential centrifugation cannot thoroughly separate protein and other non-exosomal particles from EVs, limiting its efficacy and usage in clinical studies for diagnosis, to large extent, the advent of density gradient ultracentrifugation that reverses the poor separation efficiency due to their buoyant density, and it is frequently used for EV isolation though with a considerable loss of EVs.<sup>31,32</sup>

## 5 | FILTRATION

Due to the micropores or nanopores, EVs can also be isolated by numerous protocols (eg ultrafiltration and hydrostatic dialysis). Ultrafiltration is currently available method used for EV isolation,

which represents an efficient alternative to ultracentrifugation. It involves the use of membrane filters with a narrow range of pore size distribution to deplete the proteins with molecular weight over 100 kDa, cell debris and floating cells. However, the size of filters is especially suitable for cell culture media and urine samples, which



**FIGURE 2** The common methods to isolate EVs. Straight brackets: isolated EVs; yellow: soluble components; and blue: buffer. A, In differential centrifugation, separation is based on sedimentation velocity, largely depended by size; B, in density gradient centrifugation, separation is relied on buoyant and density; C, size exclusion chromatography uses a porous matrix (dotted circles) that separates on size; D, in ultrafiltration, separation is based on size; E, in immunocapture assays, EVs are captured based on the presence of specific EVs surface molecules. F, in precipitation, EVs isolation via adding some water-excluding polymers to sample to force the precipitation of small EVs out. G, In microfluidic device, EVs isolation via combining several methods such as immunoaffinity and filtration systems. Copyright 2017, University of Helsinki, Frank AW Coumans<sup>31</sup>

**TABLE 2** Summary of EVs isolation techniques

Methods	Isolation method	Isolation Principle	Advantages	Limitations
Centrifugation	Differential centrifugation	Sedimentation velocity	<ol style="list-style-type: none"> <li>1. Broad application</li> <li>2. Standardization</li> <li>3. Ease of use</li> <li>4. Reproducibility</li> </ol>	<ol style="list-style-type: none"> <li>1. High equipment costs, cumbersome long run times and low portability.</li> <li>2. Time-consuming.</li> <li>3. Recovery based on sedimentation efficiency. No absolute separation of vesicle subpopulations.</li> <li>4. Risk of contamination and formation of protein aggregates</li> </ol>
	Density gradient centrifugation	Buoyant density	<ol style="list-style-type: none"> <li>1. Lower contamination risks because proteins partition into different density layers than EVs</li> <li>2. Large sample capacity and yield</li> <li>3. Good purity and preserved morphology of isolated EVs</li> <li>4. Unlike sucrose, iodixanol forms iso-osmotic solutions at all densities, thus better preserving the MV size</li> </ol>	<ol style="list-style-type: none"> <li>1. Requires expensive ultracentrifuge</li> <li>2. Time-consuming</li> <li>3. Sucrose toxicity might limit downstream functional studies</li> <li>4. No absolute separation of vesicle subpopulations owing to overlapping density</li> </ol>
Filtration	Ultrafiltration	Size	<ol style="list-style-type: none"> <li>1. Easy to use</li> <li>2. Quick technique</li> <li>3. Reproducible</li> </ol>	<ol style="list-style-type: none"> <li>1. Small sample volume limitations,</li> <li>2. Protein contamination</li> <li>3. Loss of yield owing to trapping in filter pores</li> </ol>
	Chromatography	Size/charge	<ol style="list-style-type: none"> <li>1. Increases purity and integrity</li> <li>2. Suitable for isolation from complex biofluids</li> </ol>	<ol style="list-style-type: none"> <li>1. Requires specialized equipment</li> <li>2. Small sample volume limitations</li> <li>3. Time-consuming</li> </ol>
Immunoaffinity	Immunological separation	Presence of specific EVs surface molecules	<ol style="list-style-type: none"> <li>1. Isolation of all or specific subtypes of EVs</li> <li>2. Higher purity of EVs than with UC</li> <li>3. Possible quantification and characterization of EV protein</li> </ol>	<ol style="list-style-type: none"> <li>1. Requires prior knowledge of vesicle characteristics</li> <li>2. Requires specific antibody</li> <li>3. Not suitable for large sample volumes</li> <li>4. Captured vesicles might not retain functionality after elution</li> </ol>
Precipitation based	Polymeric precipitation	PEG precipitation	<ol style="list-style-type: none"> <li>1. High speed</li> <li>2. Simple procedure</li> <li>3. High yield</li> </ol>	<ol style="list-style-type: none"> <li>1. Low purity caused by contamination</li> <li>2. Low specificity</li> </ol>
	Protein organic solvent precipitation	The ion-pairing effect	<ol style="list-style-type: none"> <li>1. Technique overcomes the disadvantages of co-precipitation of proteins</li> <li>2. MV denaturation found with highspeed UC</li> <li>3. Isolation efficiency higher than that of UC</li> </ol>	<ol style="list-style-type: none"> <li>1. Co-precipitation of other non-EV contaminants (proteins, lipoproteins and polymeric materials)</li> <li>2. Long run times, tedious sample preparation and lengthy pre- and post-clean-up</li> </ol>
Microfluidics based	Microfluidics	Presence of specific molecules, Physical properties such as size, Microfluidic filtration	<ol style="list-style-type: none"> <li>1. Increases throughput and allow multiplexing</li> <li>2. Reduced cost, sample size and processing time</li> </ol>	<ol style="list-style-type: none"> <li>1. Lack of standardization and clinically applicable methods</li> <li>2. Microfluidic devices can damage MVs due to shear stress</li> <li>3. There are scalability and validation issues in clinical practice</li> <li>4. Drawback of capturing only specific EV populations with IC-based methods, and low recovery with sieving approaches</li> </ol>

Abbreviations: PEG, polyethylene glycol.

limit their applications to clinical routine tests. Hydrostatic filtration dialysis (HFD), another approach developed for isolating EVs from urine samples, shows its advantages on the removal of ultracentrifugation via multiple steps and the possibility of isolating EVs from highly diluted solutions. Gel filtration, also called size exclusion chromatography, is one of the methods to collect EVs based on the basis of the size differences, and the disadvantages of this method are its low yield and rather expensive chromatographic sorbent.<sup>33,34</sup>

## 6 | PRECIPITATION METHODS

The current research provides numerous protocols for EV isolation via adding some water-excluding polymers to sample, and subsequently force the precipitation of small EVs out, its time-saving and easy usage make it suitable for clinical use, though show unavoidable contamination of the isolated EVs with proteins, protein complexes, lipoproteins and nucleoproteins, as well as viral and other particles.<sup>35,36</sup> There are several polymer kits available that have already applied for purifying the EVs, such as hydrophilic polymers, protamine, sodium acetate and proteins with organic solvent (PROSPR) with along their disadvantages and advantages demonstrating in Table 2.

## 7 | IMMUNOAFFINITY ISOLATION METHODS

Immunoaffinity isolation is another approach to isolate the EVs with increasing purity, owing to selectively exploiting the presence of specific molecules in the small EV surface<sup>37</sup>; for example, the lipids, proteins and polysaccharides are common substances that exposed on the surface of EVs, as a result, showing potency in being ligands for manifold molecules. Generally, there are five main methods for the isolation of EVs based on immunoaffinity, including antibodies to EV receptors,<sup>38</sup> phosphatidylserine-binding proteins,<sup>39</sup> heparin-modified sorbents<sup>40</sup> and binding of heat shock proteins,<sup>34</sup> as well as lectins.<sup>41</sup> Although along with evident advantages of the EV purified isolation, the expensive costs, and the insufficient efficiency of isolation, and difficulties encountering in the process of isolation the large volumes of EVs, which substantially limits the applicability of immunoaffinity isolation methods.

## 8 | MICROFLUIDIC DEVICES

Microfluidic devices are composed of a network of microchannels with different sizes, which have been implicated for EV isolation from cell culture and various tissue fluids on the basis of the immunoaffinity principle, as well as systems. However, some issues are yet to be removed; for instance, the inputted sample shows great possibility to block channels and the efficacy of isolation of EVs is extremely slow, consequently, decreasing their diagnostic potential,<sup>38</sup>

and overcome some of the challenges involving in EVs detection, such as the problem of the small size and lacking in distinct biomarkers, which contributing to get a comprehensive understanding function of their contents (eg protein, RNA and lipid).

Several qualitative and quantitative analysis techniques are currently available (Table 3). For instance, transmission electron microscopy (TEM) could be combined with immunogold staining to represent structural details and delineate the subpopulations of EVs.<sup>42</sup> A study indicates that cryo-electron microscopy might be more suitable for depicting the morphology of EVs as its no fixation or staining.<sup>43</sup> The size, morphology and intactness of EVs also could be determined by scanning electron microscopy (SEM) and atomic force microscopy (AFM).<sup>44</sup> Measuring the size and number distribution of single EVs can be made by dynamic light scattering (DLS) and nanoparticle tracking analysis (NTA).<sup>45</sup> Both the conventional flow cytometry and novel fluorescence-based flow cytometry could be promising tools to qualitatively and quantitatively analyse the EVs.<sup>46</sup> Western blot, enzyme-linked immunosorbent assay (ELISA) and EVs arrays are used to present purity and enrichment. Micronuclear magnetic resonance [ $\mu$ NMR] system and a photosensitizer-bead detection system (ExoScreen) are other sensitive qualitative and quantitative approaches.<sup>47-49</sup>

As mentioned above (Table 2), each type of isolation approach has intrinsic advantages and restrictions with respect to cost-effectivity, complexity, purity, yield and functionality of the EVs. In urological neoplasms similarly, one of the current challenges is how to develop the sensitive detection platforms and robust and remarkable isolation techniques, with promising potency in the identification of EVs and their subpopulations to trigger reliable prognosis and precise prediction of treatment response, or to provide novel neoplasm grading and staging via analysing of easier accessible and minimal-invasive body fluids. Therefore, many standards should be established to develop such interesting EVs isolation and capture tools. Initially, it is highlighted that the optimal specimen pools should be determined, such as give priority to the sample safety and accessibility, target EVs quantity, and simple and convenient manoeuvrability, especially urine for urogenital tumours; second, the preservation conditions such as temperature, time and additives should be standardized because the quantity and variation of EVs could diminish differential preservation conditions, consequentially giving rise to the deflection of results but also difficulty in continued supervision of cancer development; and third, promising isolation and detection techniques that satisfy the clinical application in a hospital setting should be established.

## 9 | GENERAL FUNCTIONS OF EVS IN MALIGNANCIES

Bioactive molecules of EVs secreted by both cancer cells and tumour-associated cells provide the essential signals for favouring tumour growth via remodelling the architectures in tumour microenvironments and forming pre-metastatic niches. Different mechanisms of EVs-mediated tumour proliferation and progression will be discussed in the following sections (Figure 3).

**TABLE 3** Techniques for extracellular vesicle detection and characterization

Methods	Size detection range/detection limit	Size distribution	Concentration	Marker detection
Quantitative methods				
DLS	1 nm-6 $\mu$ m	+	-	-
qNano	70 nm-10 $\mu$ m	+	+	-
Qualitative methods				
Western blot and ELISA	NA	-	-	+
Extracellular vesicle array	NA	-	-	+
TEM	<1 nm	+	-	+
SEM	~1 nm	+	-	+
Cryo-EM	<1 nm	+	-	+
AFM	<1 nm	+	-	-
Quantitative and qualitative methods				
NTA	50 nm-1 $\mu$ m	+	+	+
Conventional flow cytometry	$\geq$ 300 nm	-	+	+
	<300 nm	-	-	+
TRPS	70 nm-10 $\mu$ m	+	+	-
Fluorescence high-resolution flow cytometry	~100 nm	-	+	+
$\mu$ NMR system	50-150 nm	-	+	+
nPLEX assay	NA	-	+	+
ExoScreen	NA	-	+	+

Note: "+" indicates variable can be measured, "-" indicates it cannot.

Abbreviations: AFM, atomic force microscopy; Cryo-EM, Cryo-electron microscopy; DLS, dynamic light scattering; nPLEX, nanoplasmonic exosome; NTA, nanoparticle tracking analysis; SEM, scanning electron microscopy; TEM, transmission electron microscopy; TRPS, Tunable resistive pulse sensing;  $\mu$ NMR, micronuclear magnetic resonance.

## 10 | PROMOTION OF ANGIOGENESIS

Tumour progression is a dynamic and multistep process requiring continuous nutrient and oxygen supplied by sufficient blood conducts, while also serving to remove waste materials. The advent of cancer stem cells (CSCs) has provided a novel mechanism for the development and progression of the tumour via differentiating into endothelial cells to contribute to the angiogenesis.<sup>50,51</sup> In addition, a research indicates, for example, that miRNAs, secreted from exosomes, regulate transcription, proliferation, metabolic processing and mRNAs encode vascular endothelial growth factor (VEGF), fibroblast growth factor (FGF), angiopoietin1, Ephrin A3, matrix metalloproteinase-2 (MMP-2), and MMP-9 and growth factors in contrast to CD105-negative CSCs.<sup>19</sup>


## 11 | EPITHELIAL-TO-MESENCHYMAL TRANSITION

Epithelial-to-mesenchymal transition (EMT) is a developmentally vital reversible process of which fully differentiated cells lose their epithelial features (eg E-cadherin,  $\beta$ -catenin and plakoglobin), acquiring a migratory mesenchymal phenotype (eg N-cadherin and vimentin). EMT also contributes to the metastatic potential of tumours.<sup>52</sup> Exosome mediates that growth in migration and invasion by the way

of EMT has been observed in many other studies.<sup>53-55</sup> Urothelial cells exhibit the EMT after exposing to tumour-derived exosomes.<sup>56</sup>

## 12 | FORMATION OF PRE-METASTATIC NICHES

Primary tumours can release some biological factors that migrate to preferred metastatic regions and dynamically remodel these sites before spreading to a distant organ, which means that form predetermined metastatic microenvironments, also referred to as pre-metastatic niches.<sup>57</sup> In general, exosomes display the characteristics of organ tropism, and the process of the construction of pre-metastatic niche involves with initial tumour-derived exosomes releasing into the circulation system and then escaping from the vascular beds to migrate to distant secondary organ.<sup>58,59</sup> During the process, the crucial initial step is how vascular leaking exosomes can target organ tissues; nowadays, it is induced by complicated processes involved combination of stromal cells and released cancer cell-derived exosomes resulting in reprogramming of these cells<sup>60-63</sup> and activation of several vital signalling pathways,<sup>64,65</sup> which alter the local chemokine repertoire of the tumour microenvironment (TME) and remodel the components of the extracellular matrix (ECM) in turns.<sup>61,62,66</sup> Moreover, the vast researches indicate that there are exerting cooperations between bone



**FIGURE 3** Physiological processes influenced by EVs. Extracellular vesicles are involved in most physiological processes that are associated with intercellular communication, and the content of extracellular vesicles, including mRNAs, microRNAs (miRNAs), lipids and proteins, is depicted

marrow-derived cells and exosomes in the primary phase of pre-metastatic niche that can stimulate the mobilization of these cells into the circulatory system circulation and disseminate to distant sites, sequentially generate a local pro-inflammatory focus with pro-tumorigenic immunosuppression.<sup>60-62</sup>

### 13 | MODULATION OF THE TUMOUR MICROENVIRONMENT

Accumulating evidence support tumour progression is the consequence of communication between tumour cells and cells within the tumour microenvironment via paracrine or autocrine, such as adipocytes, fibroblasts, immune cells and cells of the vascular. It has been shown that tumour cells have a higher propensity to secrete larger quantity of exosome<sup>67,68</sup>; for instance, cancer-associated fibroblasts (CAFs) support tumour cells in proliferation, delaying senescence and resisting against drugs, which induced by exosome-secreted miR-9 and the human telomerase reverse transcriptase (hTERT).<sup>69,70</sup> Activation of signalling pathways by exosomes is one of the way to modulate the tumour microenvironment, such as irritation of the TGF- $\beta$ /Smad pathway by transferring TGF- $\beta$  to human umbilical cord-derived mesenchymal cells (hucMSCs), subsequently differentiating into CAFs.<sup>71</sup> Moreover, recent research shows that transferring TGF- $\beta$  also contributes fibroblasts converse to myofibroblasts, which could secrete insulin-like growth factor 1 (IGF-1), activin A and VEGF to induce tumour progression.<sup>72</sup> The irritation of another signalling pathway by exosomes in the bidirectional crosstalk between cancer cells and normal stromal cells, such as nuclear factor kappa B (NF- $\kappa$ B) and epidermal growth factor receptor (EGFR) signalling, also plays vital roles in the proliferation and migration of tumour. Endothelial cells show

enhanced cell motility and tube formation ability after re-educated by tumour-derived exosomes<sup>73</sup>; moreover, RNA secreted from EVs develops hepatocyte growth factor synthesis through the activation of ERK1/2 and AKT signalling pathways.<sup>74</sup> It is widely believed that tumour-derived EVs impose significant effects in mediating communication between immune and cancer cells of renal cell cancer (RCC),<sup>75</sup> such as immune evasion of tumours.<sup>76,77</sup> MiR-222-3p induces polarization of tumour-associated macrophages by the activation of SOCS3/STAT3 pathway to facilitate tumourigenesis and cancer progression.<sup>78</sup> Additionally, Rab27a supports exosome could modify the tumour microenvironment via advancing recruitment and differentiation of bone marrow-derived neutrophils to cancer cells.<sup>79</sup> Furthermore, a few studies suggest that tumour-derived EVs facilitate cancer progression by attenuating immune and more specific EVs could diminish the cytotoxicity of natural killer cells and T cell in immunoreaction.<sup>80-82</sup> Tumour-derived EVs also could influence the cancer cells themselves via autocrine to irritate the invasion and migration, and reduce adhesion abilities as well via enhancing MMP-9 or chemokine receptor type 4 (CXCR4).<sup>74</sup>

### 14 | MANAGEMENT OF UROLOGIC MALIGNANCIES

In the past several decades, although with the increasing development of renewable treatments for urological tumours, for instance chemotherapies and molecular targeted therapies and renowned immunotherapies, the prognosis remains poor. Recently, sufficient researches on EVs in urinary tumours provide a deeper understanding of biogenesis and pathogenesis and might be offered underlying therapeutic targets in urologic cancers. Herein, we review the



current published research on EV for commonly urogenital carcinomas including bladder cancer (BCa), kidney cancer (RCC) and prostate cancer (PCa).

## 15 | KIDNEY CANCER (RENAL CELL CANCER)

Renal cell cancers (RCCs) represent 2%-3% of all cancers.<sup>83,84</sup> Many kidney cancers remain asymptomatic until the late disease stages with 50% of patients are detected incidentally by non-invasive imaging investigating, and approximately 30% of patients with metastasis in the primary time of diagnosis.<sup>85,86</sup> The utility of urinary EVs could be recognized as potential diagnostic and prognostic markers in RCCs (Table 4).<sup>87</sup> Proteomic analysis of urinary EVs differs from the patients and healthy groups with showing an RCC-specific signature of the effectiveness of proteins.<sup>58,88</sup> Another proteins, for instance, podocalyxin (PODXL), WNT signalling pathway inhibitor 4 (DKK4), ceruloplasmin, carbonic anhydrase IX (CAIX) and MMP-9, were validated by using immunoblotting method.<sup>89</sup> In addition, among RCC patients, the mRNA levels of glutathione S-transferase alpha 1 (GSTA1), CCAAT enhancer binding protein alpha (CEBPA) and PCBD1 in EVs decrease compared with healthy groups.<sup>90</sup> Moreover, a report demonstrates that the lipids within urinary EV show difference between RCC patients and healthy groups.<sup>91</sup> Based on microRNA expression screening, the cluster of miRNAs including miR-449a, miR-126-3p, miR-486-5p and miR-34b-5p could differentiate clear cell RCCs from benign subjects<sup>92</sup> and could be recognized as potential diagnostic

and prognostic markers.<sup>58,93,94</sup> Serum-derived EVs have also been recognized as prominently diagnostic and prognostic tools for clear cell RCCs.<sup>93,94</sup> Recently, azurocidin as a permeabilizer for vascular endothelial cells has been isolated from serum-free medium within incubating clear cell RCC tissues samples.<sup>95</sup>

Renal cell cancers generate EVs that could educate endothelial<sup>95</sup> and immune cells,<sup>96</sup> with promoting angiogenesis and immunosuppressive activity, respectively. As for immune systems, increasing evidence shows that tumour-derived EVs result in immune evasion of tumours partly via the activation of caspase pathway to trigger apoptosis in T lymphocytes.<sup>97</sup> In addition, both EVs-derived antigens and Hsp70 can inhibit the immunoreaction though induction tumour growth factors of and pro-inflammatory cytokines similarly.<sup>98,99</sup> Moreover, EVs also can be applied for cancer immunotherapy, mainly due to promoting cytotoxic effects and proliferation of T cells via releasing interferons.<sup>100</sup> Furthermore, RCCs also could secrete EVs to interact with endothelial cells to promote lymphopoiesis and angiogenesis and thereby metastasis. A study shows that EVs derived from RCC cell line (eg 786-O) increase the expression of vascular endothelial growth factor in human umbilical vein endothelial cells (HUVECs) with resulting in tubular formation of HUVECs,<sup>97</sup> involving the downregulation of hepatocyte cell adhesion molecule by upregulating phosphorylated AKT in RCCs.<sup>101</sup> What is more, CD105<sup>+</sup> stem cells of RCC release EVs that promote and trigger the formation of a pre-metastatic niche by upregulating MMP2 and VEGF<sup>58,102</sup>; although sunitinib is a first-line targeted regimen for metastatic RCCs,<sup>103</sup> its efficacy of biotherapy for the long time is controversial in respect to drug resistance, which has

**TABLE 4** Candidate biomarkers for kidney cancer derived from EVs

Source	Methodologies	End point	Type of marker	Markers	Reference
Urine	Ultracentrifugation	Diagnosis	mRNA	GSTA1, CEBPA and PCBD1	De Palma et al <sup>89</sup>
Urine	Density gradient ultracentrifugation	Diagnosis	Proteins	MMP-9, PODXL, DKK4, CAIX and ceruloplasmin	Raimondo et al <sup>88</sup>
Cancer stem cells	Ultracentrifugation, Flow cytometry immunohistochemistry	Diagnosis	Proteins	VEGF, FGF, angiopoietin 1, Ephrin-A3, MMP-2, MMP-9	Grange et al <sup>57</sup>
RCC cells	Centrifugation Filtration Flow cytometry Western blot ELISA	Diagnosis	Proteins	Fas ligand, Bcl2-L-4	Yang et al <sup>87</sup>
Viable human tissue	Ultracentrifugation Mass spectrometry Western blot	Diagnosis	Proteins	Azurocidin 1	Jingushi et al <sup>93</sup>
Serum	Immunoaffinity magnetic beads	Diagnosis	miRNA	miR-210 and miR-1233	Zhang et al <sup>92</sup>
Serum	Total Exosome isolation kit	Prognosis	miRNA	miR-224	Fujii et al <sup>93</sup>
Cancer stem cells	Ultracentrifugation Microarray analysis qRT-PCR	Diagnosis	miRNA	miR-200c, miR-92, miR-141, miR-19b, miR-29a, miR-29c, miR-650, miR-151	Grange et al <sup>94</sup>
Urine	Centrifugation Urine exosome RNA isolation kit	Diagnosis	miRNA	miR-126-3p, miR-449a, miR-34b-5p, miR-486-5p	Butz et al <sup>91</sup>
Urine	Ultracentrifugation Mass spectrometry	Diagnosis	Lipids	Lysophosphatidylethanolamine metabolite	Del Boccio et al <sup>90</sup>

been confirmed that regulated by lncARSR by promoting MET and AXL expression in RCC cells.<sup>104,105</sup> Therefore, all those support that EV-based targets display a promising potency for the development of novel cancer therapies.

EVs also can be served as a vaccine for RCCs. A report reveals that RCCs derived from EVs could increase immunogenicity by proliferating T cells and releasing interferons subsequently<sup>100</sup>; the effectiveness of vaccines when dendritic cells (DCs) load with EVs is higher rather than whole tumour lysate.<sup>98</sup>

## 16 | THE BLADDER CANCER

Bladder cancer (BCa) is the seventh most commonly diagnosed cancer in the male population worldwide, and the diagnosis for BCa is usually on the basis of cytology, urinalysis and cystoscopy. Cytology is a highly specific test, but low in the sensitivity for the diagnosis of BCa.<sup>106</sup> Cystoscopy is the gold standard to diagnose the BCa, while this method is expensive and invasive, even for flexible cystoscopy, and the risk of developing urinary infections is up to 10%<sup>107</sup>; non-invasive and reliable biomarkers are therefore required in the future. Given that, urine is an excellently suitable fluid for biomarkers discovery in BCa. The biomarkers (mainly including proteins, mRNA, lncRNA and miRNA) within EVs isolated from BCa were investigated by different research groups, which could be promising molecules to identify the BCa and predict the progression of the BCa (Table 5). Based on proteomic analysis of urinary EVs, several studies have been identified cargoes of possible biomarkers for BC patient, but not in healthy volunteers.<sup>108,109</sup> Among them, research shows that the levels of tumour-associated calcium-signal transducer 2 (TACSTD2) are correlative with BCa, compared with high-grade BCa. The author identified seven proteins differentially expressed in the low-risk group (Table 5).<sup>109</sup> Proteome profiling of urinary exosomes indicates H2B1K and alpha 1-antitrypsin as prognostic and diagnostic biomarkers for urothelial bladder cancer, which could be verified in immunohistochemistry (IHC).<sup>110</sup> Additionally, HEXB, S100A4 and SND1 significantly identified in EV derived from the MIBC cell line also are upregulating in urinary EV from MIBC patients when vs to normal groups.<sup>111</sup> There are other proteins could be recognized as potential diagnostic and prognostic markers for BCa.<sup>112-115</sup> Using a whole transcriptome array, a study reports the potential application of mRNAs in urinary EVs for diagnoses, such as LASS2 and GALT1 involving progression with only in BCa patients, and ARHGEF39 and FOXO3 while only expressing in healthy controls to suppress the tumour.<sup>116</sup> In addition, based on the microarray analysis of miRNA, great studies pay their attention on the roles of diagnostic and prognostic.<sup>117-121</sup> Interestingly, research shows that several microRNAs from urinary EVs significantly upregulate in BCa, but not in plasma from same patients, which suggests that different biofluids may harbour different molecules.<sup>122</sup>

Furthermore, to unravel the roles of lncRNA in BCa, recent researches show that lncRNAs, such as HOX-AS-2, HOTAIR, ANRIL and lnc-RoR, are enriched in BCs cancer cell line EVs as well as

urinary EVs from high-grade BCa patients, demonstrating that lncRNAs have potential as biomarkers for BCa.<sup>123</sup> Additionally, recent reports demonstrate that the CD63<sup>+</sup> urinary EVs could be a biomarker for the detection of BCa.<sup>114,124</sup> EVs derived from BCa cell line also influence local regions microenvironments or distant cells by transferring their content, as result of facilitating proliferation, angiogenesis, invasion, and migration and the inhibition of apoptosis<sup>56,112,125</sup>; for example, EGF-like repeat and discoidin I-like domain-containing protein 3 (EDIL-3) from invasive BCa cell lines could stimulate the migration and angiogenesis of urothelial and endothelial cells.<sup>112</sup> Periostin, another factor from invasive BCa cell lines, could contribute low-grade BCa cancer cells to gain the aggressiveness within via activating ERK oncogenic pathway.<sup>115</sup> Urothelial cells exposed to EVs from cancer cell lines or patient specimen show the phenomena of EMT.<sup>56</sup> Similarly, lncRNA-UCA1, which may play an important role in causing intratumoural hypoxia, could irritate tumour progression via the EMT as well.<sup>126</sup> However, exosomes also can discard tumour-suppressive miRNAs contributed to BCa progression, such as miR-23b and miR-921,<sup>121</sup> all those could provide underlying targets for the future therapies for the BCa. Recently, a report demonstrates the effectiveness of EVs as a vector to carry siRNAs in BCa; therefore, in the future, EVs have the potential functions to stably deliver substantial therapeutic cargoes including miRNAs and siRNAs, to anchor organs with the development of tissue engineering technology.<sup>127</sup>

## 17 | PROSTATE CANCER

Prostate cancer (PCa) is the second most commonly diagnosed cancer in men, accounting for 15% of all cancers diagnosed.<sup>128</sup> Although PSA testing contributes to identify and manage PCa in the early phase, it still has some limitations, for example, the specificity of discrimination of benign prostate diseases, such as acute prostatitis and benign hyperplasia.<sup>129</sup> Thus, the more specific and ideal substrate (eg urine, prostatic plasmas and blood samples) for PCa are urgently developed rather than invasive prostate biopsies.<sup>130</sup> Some studies have presented the usefulness of urinary EVs as diagnostic factors (Table 6).<sup>131-134</sup> EV-involved transmembrane proteins CD63 and CD9 are sufficient in urine from PCa.<sup>135</sup> Integrin  $\alpha$ 3,  $\delta$ -Catenin, integrin  $\beta$ 1 and FABP5 proteins are identified in urinary EVs of PC patients with the significantly increased levels of PCa patients during the process of the investigation of the proteomic cargo of urinary PC-derived EV.<sup>136-139</sup> Moreover, on the basis of the mass spectrometry proteome analysis, thousands of proteins encapsulated on and within vesicles are identified as biomarker candidates from urinary EVs or cell lines of PCa,<sup>139-146</sup> whereas the value as biomarkers is still controversial, and several types of research regenerate the previous biomarkers by the using targeted proteomics and immuno-assays.<sup>134</sup> Based on a proximity ligation assay, PC-derived EVs in blood has also shown to contain proteins specific to PCa, such as phosphatase and tensin homolog gene (PTEN), survivin and other factors with decreased level compared to benign prostatic hyperplasia or health

**TABLE 5** Candidate biomarkers for bladder cancer derived from EVs

Source	Methodologies	End point	Type of marker	Markers	Reference
BCC/urine	Ultracentrifugation Flow cytometry In-gel digestion Mass spectrometry	Diagnosis	Proteins	$\beta$ 1 and $\alpha$ 6 integrins, CD36, CD44, CD73, CD10, MUC1, basigin, 5T4	Welton et al <sup>107</sup>
Urine	Ultracentrifugation	Diagnosis	Proteins	APOA1, CD5L, FGA, FGB, FGG, HPR, HP	Chen et al <sup>108</sup>
Urine	Differential ultracentrifugation	Diagnosis Prognosis	Proteins	Alpha-1 antitrypsin, histone H2B1K	Lin et al <sup>109</sup>
Urine	Differential ultracentrifugation	Diagnosis	Proteins	HEXB, S100A4, SND1	Silvers et al <sup>110</sup>
BCC/urine	Sucrose/D2O cushion Ultracentrifugation	Diagnosis	Proteins	EDIL-3	Beckham et al <sup>111</sup>
Urine	Ultracentrifugation In-gel digestion Mass spectrometry	Diagnosis	Proteins	Resistin, GTPase NRas, MUC4, EPS8L1, EPS8L2, EHD4, G3BP, RAI3, GSA	Smalley et al <sup>112</sup>
Urine	Centrifugation Filtration Integrated double-filtration Microfluidic device	Prognosis	Proteins	CD63 + EV signal intensity	Liang et al <sup>113</sup>
Urine	Ultracentrifugation	Prognosis	Proteins	Periostin	Silvers et al <sup>114</sup>
Urine	Ultracentrifugation NanoSight microarray PCR	Diagnosis	mRNA	LASS2, GALNT1	Perez et al <sup>115</sup>
Urine	Differential ultracentrifugation	Diagnosis	miRNA	miR-21-5p	Matsuzaki et al <sup>116</sup>
Urine	Differential ultracentrifugation Filtration	Diagnosis Prognosis	miRNA proteins	miR-375, miR-146a, apoB	Andreu et al <sup>117</sup>
Urine	Differential centrifugation Total exosome isolation kit	Prognosis	miRNA	miR-141-3p, miR-200a-3p, miR-205-5p	Baumgart et al <sup>118</sup>
Urine	Differential ultracentrifugation	Prognosis	miRNA	miR-940	Long et al <sup>119</sup>
Urine	Nanostring miRNA assays Droplet digital PCR	Diagnosis	miRNA	miR-205, miR-200c-3p, miR-29b-3p; miR-921, miR-23b	Ostenfeld et al <sup>120</sup> Berrondo et al <sup>122</sup>
Urine	Centrifugation Exosome RNA isolation kit	Diagnosis	miRNA	miR-4454, miR-21, miR-720	Armstrong et al <sup>121</sup>
Urine	Differential ultracentrifugation	Diagnosis Prognosis	mRNA; lncRNA	HOTAIR, HOX-AS-2, MALAT1, SOX2, OCT4, HYMA1, LINC00477, LOC 100506688, OTX2-AS1	Berrondo et al <sup>122</sup>

subjects.<sup>133,147-149</sup> PC-based EVs show great value in comprehensively mapping nucleic acid changes in PCa via urine- or blood-based liquid biopsies, and it has been known that the EVs is important pool resource for circulating-free DNA (cfDNA)<sup>150</sup>; as such, some factors demonstrated good clinical usefulness and diagnostic value in predicting for high-grade PC, including TP53 mutations, ERG and PCA3, can be detected in the EVs.<sup>150</sup> ExoDx Prostate IntelliScore urine exosome assay has been developed as a non-invasive detective tool to distinguish the high-grade PCa from low-grade groups and benign diseases at initial biopsy.<sup>151</sup> Several mRNAs could be

recognized as promising diagnostic and prognostic tools for the PCa<sup>152-156</sup>; for instance, the transcripts of CDH3 from EVs were significantly decreased compared with benign hyperplasia<sup>156</sup>; contrarily, the mRNA level of PTEN gene can only be detected in the patient of PC,<sup>147</sup> and nevertheless, both of them are expected to be powerful for the diagnosis and monitoring of PCa.<sup>152</sup> RT-PCR, microarray and RNA sequencing technologies have focused on the non-coding RNA content within EVs so far; for example, in the urinary sample, the levels of lncRNA-p21, a suppressor of p53 signalling, contribute to detecting PC from benign disease.<sup>157</sup> Next-generation

**TABLE 6** Candidate biomarkers for prostate cancer derived from EVs

Source	Methodologies	End point	Type of marker	Markers	Reference
Plasma	Ultracentrifugation Western blot ELISA	Diagnosis	Proteins	Survivin	Khan et al <sup>132</sup>
Urine	Differential ultracentrifugation Filtration	Diagnosis Prognosis	Proteins	TGM4, ADSV, PPAP, PSA, CD63, SPHM, GLPK5 TMEM256, flotillin 2, Rab3B, PARK7, LAMTOR1 TM256, LAMTOR1, ADIRF TMEM256, flotillin 2, Rab3B, PARK7, LAMTOR1	Wang et al <sup>133</sup> Sequeiros et al <sup>145</sup>
Urine	Differential ultracentrifugation	Diagnosis Prognosis	Proteins	$\delta$ -catenin Integrin $\alpha$ 3, Integrin $\beta$ 1 FABP5	Liu et al <sup>135</sup> Lu et al <sup>136</sup> Fujita et al <sup>138</sup>
Tissue Urine	Differential ultracentrifugation Filtration	Diagnosis Prognosis	Proteins	CD63, ANXA1-3, FASN, FOLH1, GDF15, MDR1, XPO1, TGM4, TIMP1, SFN, TMEM256, LAMTOR1, ADIRF, ITGA3, and ITGB1	Bijnsdorp et al <sup>137</sup>
Tissue	Ultracentrifugation Gel filtration Chromatography, 2D-PAGE Mass spectrometry	Diagnosis Prognosis	Proteins	ANXA1, ANXA3, ANXA5, DDAH1	Ronquist et al <sup>140</sup>
Cell lines	Ultracentrifugation Mass spectrometry Bead immuno-isolation Western blot	Diagnosis Prognosis	Proteins	CDCP1, CD151, CD147	Sandvig et al <sup>141</sup>
Cell lines Urine	Ultracentrifugation immunoprecipitation Western blot Electron microscopy	Diagnosis Prognosis	Proteins	ANXA2, CLSTN1, FASN, FLNC, FOLH1, GDF15ACPP, LTF, DDP4, TGM4, MME, PSA, SEMG1, AZGP1, ANPEP, G3BP, PSMA, TMPRSS2, FASN, LGALS3, PSCA, KLK2, KLK11, TIMP1 PDCD6IP, XPO-1, ENO1	Duijvesz et al <sup>142</sup> Utleg et al <sup>143</sup> Principe et al <sup>144</sup>
Plasma	Ultracentrifugation Western blot immunofluorescence	Diagnosis	Proteins	PTEN	Gabriel et al <sup>134</sup>
Serum	Differential centrifugation	Predictive Monitoring	Proteins	ABCB1, ABCB4, PABPC4	Kato et al <sup>147</sup>
Urine Plasma	Differential ultracentrifugation Filtration Chromatography	Diagnosis	Proteins	Afamin, cardiotrophin-1, CDON, endoplasmic re- ticulum aminopeptidase 1, FGF19, IL17RC, NAMPT, IL1RAPL2, CD226, IGFBP2, CCL16, TNFSF18, IGFBP5; Aromatic-L-amino-acid decarboxylase	Welton et al <sup>148</sup>
Urine	Centrifugation Filtration Ultrafiltration	Diagnosis Prognosis	mRNA	PCA3, TMPRSS2-ERG AGR2, SV-G, AGR2 SV-H CDH3	Neeb et al <sup>151</sup> Donovan et al <sup>152</sup> Hendriks et al <sup>153</sup> Motamedinia et al <sup>154</sup> Royo et al <sup>155</sup>

(Continues)

TABLE 6 (Continued)

Source	Methodologies	End point	Type of marker	Markers	Reference
Plasma/ Serum/Urine	ExoMIR extraction Filtration qRT-PCR	Diagnosis	miRNA	miR-107, miR-130b, miR-181a-2, miR141, miR-301a, miR-326, miR-331-3p, miR-375, miR-432, miR-574-3p, miR-22110, miR-625 miR-1290, miR-375, miR-574-3p, miR-141-5p, and miR-21-5p miRNA-21, let-7c, miR-196a-5p, miR- 501-3p, miR-19b, miR-145	Samsonov et al <sup>41</sup> Fojt et al <sup>130</sup> Bryant et al <sup>158</sup> Bryzgunova et al <sup>159</sup> Rodriguez et al <sup>131</sup> Wani et al <sup>160</sup> Xu et al <sup>161</sup>
Urine	Differential centrifugation Urine exosome RNA isolation kit	Diagnosis	lincRNA	lincRNA-p21	Işin et al <sup>156</sup>
Urine	Differential ultracentrifugation Filtration	Diagnosis	Lipids	Lactosylceramide, phosphatidylserine, phosphatidylg- lycerol, diacylglycerol, triacylglycerol	Skotland et al <sup>164</sup> Yang et al <sup>165</sup>
Urine	Differential ultracentrifugation Filtration	Diagnosis	Metabolites	Adenosine, glucuronate, isobutyryl-L-carnitine, D- ribose 5-phosphate	Puhka et al <sup>166</sup>
Urine	Differential ultracentrifugation	Diagnosis Prognosis	Glycomic	N-linked glycans	Nyalwidhe et al <sup>167</sup>

sequencing reveals the potential values for miRNA served as diagnostic and prognostic biomarkers for PCa within serum or plasma EVs,<sup>41,131,132,158-162</sup> such as miR-141 and miR-375 in serum, have been correlated with metastatic PCa.<sup>159,163</sup> Another study indicates that exosomal miR-1290 and miR-375 could be as prognostic markers in castration-resistant prostate cancer (CRPC).<sup>164</sup> In recent, several research works demonstrate that the lipids including diacylglycerol and triacylglycerol are differentially enriched in PCa rather than healthy groups.<sup>165,166</sup> Glycomic and metabolomic profiling of urinary EV reveal several small molecule metabolites could be novel biomarkers to predict the development of PCa, for example levels of N-linked glycans, glucuronate, adenosine, D-ribose-5-phosphate and isobutyryl-L-carnitine.<sup>167,168</sup>

The intercellular crosstalk through EVs could stimulate tumour progression. Several proteins presenting on and in EVs from PCa cell lines are recognized as significant mediators for the biological communication between cancer cells and tumour microenvironment or surrounding cell, including cytokine CX3CL1, MMPs and transforming growth factor B, play significant roles in the proliferation and differentiation of fibroblasts.<sup>169,170</sup> In addition, integrins ITGA3 and ITGB1 can affect invasion and migration of normal prostate epithelial cells.<sup>138</sup> Several studies suggest that complicate intercellular interactions between cancer cells, osteoclasts and osteoblasts contribute to bone metastasis.<sup>171,172</sup> It is the first protein that has been reported in the EVs originated from human hormone-refractory PCa cells to facilitate mouse pre-osteoblast differentiation.<sup>171</sup> Another molecular miR-141-3p is involved with the osteoblastic metastasis of prostate cancer via reducing the expression of Deleted in Liver Cancer 1 (DLC1) and activating the p38MAPK and OPG/RANKL pathway in osteoblasts<sup>173</sup>; however, another research fixes their attention on the role of EVs derived from PCa cell line in the osteoclast for the bone metastasis promoting osteoclast proliferation and identifies that tumour cell-derived EVs play important roles in impairing the osteoclast formation and differentiation through the underlying mechanism is unknown yet.<sup>172</sup> Moreover, the effect of biological crosstalk between PCa cell and immune cells though EVs causes the induction immune suppression via down-regulating the NKG2D cytotoxicity receptor and diminishing the IL-2 response.<sup>174,175</sup>

PCa-derived EVs also involve drug resistance. ABCB1, ABCB4, PABPC4 and SH3GL1 are much more sufficient in EVs from docetaxel-resistant prostate cancer cell lines and potentially higher in serum EVs in men with docetaxel-resistant PCa.<sup>148</sup> AR-V7 is correlated with resistance to enzalutamide and abiraterone in metastatic CRPC patients, which could be a biomarker to predict the CRPC.<sup>176,177</sup> Finally, EVs-derived PCs can be used as vaccine vesicles that present prostate-associated antigens such as PSA and PAP on their membrane to exert an anti-tumour immune response.<sup>98,178,179</sup>

## 18 | DISCUSSION

With the advent of novel concepts involving EVs in many physiologic as well as pathologic conditions, the field of EV research develops

much excitement in the urologic malignancies. Unlike conventional biopsies of that only consist of a small amount of tumour solid masses with ignoring heterogeneity, EVs, as liquid biopsies, could capture and obtain overall tumour heterogeneity owing to directly releasing from all cells in the cancer tissue and its microenvironment. Beyond a doubt, specific bioactive contents contained in circulating EVs have great promise as reliable surrogates of urological cancers; therefore, their molecular cargoes including nucleic acids, protein and lipid composition, as well as their numbers, are representing as diagnostic, prognostic biomarkers for urinary tract diseases, and immense promising for therapeutic advancements. To conclude, we have a faith for the implement of EVs in urological cancers diagnosis and therapeutics owing to their enormous potencies in several aspects, as described below (Figure 4).<sup>19</sup>


## 19 | ROLES IN DIAGNOSTIC AND PREDICTIVE BIOMARKERS

As more is understood about the fundamentals of EVs biology and roles involved in tumorigenesis and therapy resistance, EVs-based analytical methods are increasingly interesting targets for clinical application. EVs are directly released from heterogeneous tumorous and reflect a snapshot of the current state of the neoplasm; therefore, EVs have great potential as remarkable, specific and sensitive biomarkers of oncogenesis, treatment response and therapy resistance. In urogenital cancers, it is thought that increased exosomes are produced by more advanced cancers, and it thus has been suggested that total circulating exosome burden may serve as indicators for disease surveillance. Exosomal contents can also identify disease or predict treatment response, such as several proteins (eg PD-1, PD-L1) or some nucleic acids (eg miRNA) with the roles as diagnostic biomarkers for cancer, indicators for therapeutics, worse still, research to date strongly indicate EVs involve treatment irresponsiveness. Such phenomenon was observed in various malignancies also including urogenital cancers; for example, docetaxel-sensitive cell lines of prostate carcinoma undesirably acquire drug resistance again when co-culture with EVs derived from the drug-resistant cells.<sup>180</sup> To date, increased pumping agents out of tumorous cells or omics alternations induced by the cargo of EVs are two most common opinions, but the underlying mechanisms are still entirely unknown. Further investigations that tailored clinical studies are now warranted to determine how best to prevent this occurring, in the interest of patients and also for economic benefit. Additionally, endothelial cell-derived EVs can reflect transient cellular stress conditions and could be useful as predictors of anti-angiogenic therapy effectiveness and cancer cell status. However, issues with interpretation of studies and reproducibility have arisen due to the deficit of standard isolation and characterization, and nomenclature employed, and as the result of the publication of the Minimal Information for Studies of Extracellular Vesicles 2014 (MISEV 2014) with emphasizing that build the set of biophysical, biochemical and functional standard that help in detecting particular biological cargo or functions in

extracellular vesicles.<sup>181</sup> Until 2018, the updated MISEV guidelines were published, and it continued to standardize the experimental parameters for EV isolation and characterization to provide more reproducible and robust outcomes.<sup>182</sup> Continued efforts to systematically catalogue the protein, nucleic acid and lipid constituents of EVs isolated from richly annotated specimens could ultimately attribute to rapidly evolve and expand the development of selective and sensitive capture platforms directed towards specific EVs.

## 20 | ROLES IN THERAPEUTICS

Directly or indirectly, EVs derived from tumour or TME can influence urogenital neoplasm via intercellular crosstalk or modification of TME, respectively. As discussed above, cargoes of tumour-derived EVs attribute to cancer development. Thus, the blockage of exosome production, secretion and ablation of specific active exosomal contents, as well as exosome-mediated cell-cell communication between cancer and TME, have been proposed as alternative therapeutic strategies. Importantly, it is essential to note that EVs show promising potency for immunotherapy. As we all known, immunotherapy has revolutionized cancer therapy, especially the advent of immune checkpoint blockades (ie PD-1/PD-L1). Recently, it has been demonstrated that cancer-derived exosomes transfer functional PD-L1 and inhibit immune responses,<sup>183</sup> while suppression of exosomal PD-L1 induces systemic anti-tumour immunity and memory in urological carcinomas,<sup>184</sup> and clinical trials have already been initiated to explore their safety and efficacy in humans. Interestingly, EVs-based vaccines can serve as new candidates that have shown their potential as novel cancer intervention in some clinical trials, indicating that rely on their role as tumour antigens and facilitate an anti-tumour immunity in turns. The popular EVs-based vaccines mainly derive from dendritic cell (Dex immunotherapy), but the clinical efficacy is not ideal to date; further research will be required to reassess clinical applications with taking the defects in current prospective designs into considerations such as lack of preselection criteria and small sample size. EVs, on the other hand, have garnered much attention as several characteristics of an optimal delivery system. First of all, the nanometric-sized EVs confer the effective assimilation and intracellular trafficking for recipient cells. Second, the EV bioactive molecules are protected from degradation in the extracellular milieu and circulation due to lipid bilayer-membrane structure of EVs.<sup>185</sup> Third, autologous EVs show lower immunogenicity and toxicity than other conventional drug-delivery platforms.<sup>186</sup> Furthermore, EVs possessing specific surface proteins (eg integrins) could bear intrinsic targeting properties that are able to interact with target cells or organs.<sup>187</sup> Although EVs have many advantages as described above, this enormous promise therapeutic delivery tool requires further study for clinical applications. These include the identification of the optimal EV donor cell type, large-scale EVs isolation, preservation of EV structural integrity during drug loading, scalable manufacture and storage. A further challenge remains improving methods to shift in vivo biodistribution



**FIGURE 4** Future implements for EVs in urological cancer. EVs impact the multistep process of cancer; therefore, EVs should be a novel treatment strategy by inhibiting intercellular crosstalk. EVs could serve as promising diagnostic and prognostic biomarkers to dynamically trace the changes in cancer due to their high specificity and sensitivity. In addition, EVs have the potential functions to stably deliver substantial therapeutic cargoes like miRNAs and siRNAs with stability, few side effects and organ specificity. Furthermore, several studies have reported the potential of EVs derived from dendritic cells used as vaccine vesicles. Copyright 2018, The Jikei University School of Medicine, Fumihiko Urabe<sup>19</sup>

of EVs from non-specific sites towards accumulation in desired tissues. Although considerable efforts by engineering EVs to present cell type-specific ligands have been made in guaranteeing rich accumulation in target tissues, one of the major obstacles remains low delivery efficacy. The elucidation to these questions will enhance rationality and reliability to irritate the utility of EV-involving molecular cargoes as cancer diagnostics in the clinical practices.

## 21 | CONCLUSION

In recent, EVs have gained rocketing interest in the field of urological tumour research owing to their multifaceted role in the development and treatment of cancer, and their perspective as a weapon to the armoury for cancer treatment. Since EVs play pivotal effects on intercellular interactions in variable biological fluids, their numbers, protein, nucleic acids, lipid and signalling/epigenetic regulators components could be transfer to recipient cells and subsequently affect the pathologic process of the receptor cells, eventually result in abnormal proliferation, EMT, angiogenesis and metastasis by regulation of the TME to cause drug resistance, and preparation of pre-metastatic niches, to enhance

dissemination of cancer cells and cause relapse after a prolonged period of dormancy; thereby, targeting this communication will offer a novel therapeutic strategy for urologic cancer eradication. Apart from that, EVs could serve as a non-invasive liquid biopsy and have been emerged as new potential diagnostic/prognostic biomarkers, as well as playing provoking roles in predicting anti-cancer drug responses. In additions, we could use EVs as cancer vaccines or as drug delivery modules with promising therapeutic applications, unless more researches are required for clinical applications. However, the research in EVs is encountered with urgent challenges, including the standardization of approaches for the isolation, quantification and analysis of EVs from complicated tissues sample (mainly from cancer line cell medium or low numbers of patient urine samples). Great efforts have been made to precisely determine EV particles and, nevertheless, still have a great way to standardize EVs enumeration in some particular specimens, such as blood. Moreover, a further challenge is what EVs, their contents or their ratio should best be quantified as robust biomarkers in the surveillance of urological diseases staging is yet unknown; thus, in the future studies, we need pay more attention to develop stereospecific antibodies to map the topography of EVs. Furthermore, another big problem in the field is their half-life in human samples

are yet unexplored. With the increasing knowledge of their roles and development of the next-generation sequencing, mass spectrometry-based metabolomics and proteomics, we are enthusiastically sure that EVs will contribute to play clinical applications for urological cancer treatment and management in the near future.

## ACKNOWLEDGEMENTS

We gratefully acknowledged the help from staffs at the Department of Urology of the Third Affiliated Hospital of Shenzhen University for the data assistance.

## CONFLICT OF INTEREST

The authors declare that they have no competing interests.

## AUTHOR CONTRIBUTIONS

Z.W., S.W. and Z.Z. designed the review and made a retrieval strategy; Z.W. and Y.L. drafted the review text; W.X. and J.C. drafted the tables and figures; and both authors contributed to revision and finalization of the manuscript.

## DATA ACCESSIBILITY

Research data are not shared.

## CONSENT FOR PUBLICATION

The patient has given his consent for his case report to be published.

## ORCID

Zhangsong Wu  <https://orcid.org/0000-0002-7432-5997>

## REFERENCES

1. Siegel L, Miller KD, Jemal A. Cancer statistics, 2018. *CA Cancer J Clin*. 2018;68:7-30.
2. Dy GW, Gore JL, Forouzanfar MH, Naghavi M, Fitzmaurice C. Global burden of urologic cancers, 1990–2013. *Eur Urol*. 2017;71(3):437-446.
3. Pitt JM, Kroemer G, Zitvogel L. Extracellular vesicles: masters of intercellular communication and potential clinical interventions. *J Clin Invest*. 2016;126(4):1139.
4. Colombo M, Raposo G, Théry C. Biogenesis, secretion, and intercellular interactions of exosomes and other extracellular vesicles. *Annu Rev Cell Dev Biol*. 2014;30(1):255-289.
5. Xu R, Greening DW, Zhu H-J, Takahashi N, Simpson RJ. Extracellular vesicle isolation and characterization: toward clinical application. *J Clin Invest*. 2016;126(4):1152-1162.
6. de Pablos Torró LM, Retana Moreira L, Osuna A. Extracellular vesicles in Chagas disease: a new passenger for an old disease. *Front Microbiol*. 2018;9:1190.
7. Li W, Wang H, Zhao Z, et al. Emerging nanotechnologies for liquid biopsy: the detection of circulating tumor cells and extracellular vesicles. *Adv Mater*. 2018;e1805344.
8. Shao H, Im H, Castro CM, Breakefield X, Weissleder R, Lee H. New technologies for analysis of extracellular vesicles. *Chem Rev*. 2018;118(4):1917-1950.
9. Nguyen H, Simpson RJ, Salamonsen LA, Greening DW. Extracellular vesicles in the intrauterine environment: challenges and potential functions. *Biol Reprod*. 2016;95(5):109.
10. Vickers KC, Palmisano BT, Shoucri BM, Shamburek RD, Remaley AT. Corrigendum: microRNAs are transported in plasma and delivered to recipient cells by high-density lipoproteins. *Nat Cell Biol*. 2014;17(1):423-433.
11. Nawaz M, Camussi G, Valadi H, et al. The emerging role of extracellular vesicles as biomarkers for urogenital cancers. *Nature Reviews Urology*. 2014;11(12):688-701.
12. Zaborowski MP, Balaj L, Breakefield XO, Lai CP. Extracellular vesicles: composition, biological relevance, and methods of study. *Bioscience*. 2015;65(8):783-797.
13. Latifkar A, Cerione RA, Antonyak MA. Probing the mechanisms of extracellular vesicle biogenesis and function in cancer. *Biochem Soc Trans*. 2018;46(5):1137-1146.
14. Cocucci E, Meldolesi J. Ectosomes and exosomes: shedding the confusion between extracellular vesicles. *Trends Cell Biol*. 2015;25(6):364-372.
15. Yanez-Mo M, Siljander PR, Andreu Z, et al. Biological properties of extracellular vesicles and their physiological functions. *J Extracell Vesicles*. 2015;4:27066.
16. Hurley JH. ESCRTs are everywhere. *EMBO J*. 2015;34(19):2398-2407.
17. Tai Y-L, Chen K-C, Hsieh J-T, Shen T-L. Exosomes in cancer development and clinical applications. *Cancer Sci*. 2018;109(8):2364-2374.
18. Zhao G-N, Zhang P, Gong J, et al. Tmbim1 is a multivesicular body regulator that protects against non-alcoholic fatty liver disease in mice and monkeys by targeting the lysosomal degradation of Tlr4. *Nat Med*. 2017;23(6):742-752.
19. Urabe F, Kosaka N, Kimura T, Egawa S, Ochiya T. Extracellular vesicles: toward a clinical application in urological cancer treatment. *Int J Urol*. 2018;25(6):533-543.
20. Xu R, Rai A, Chen M, Suwakulsiri W, Greening DW, Simpson RJ. Extracellular vesicles in cancer – implications for future improvements in cancer care. *Nat Rev Clin Oncol*. 2018;15(10):617-638.
21. Ostrowski M, Carmo NB, Krumeich S, et al. Rab27a and Rab27b control different steps of the exosome secretion pathway. *Nat Cell Biol*. 2010;12(1):19-30.
22. Junker K, Heinzelmann J, Beckham C, Ochiya T, Jenster G. Extracellular vesicles and their role in urologic malignancies. *Eur Urol*. 2016;70(2):323-331.
23. Wilhelm EN, Mourot L, Rakobowchuk M. Exercise-derived microvesicles: a review of the literature. *Sports Med*. 2018;48(9):2025-2039.
24. Li R, Peng C, Zhang X, Wu Y, Pan S, Xiao Y. Roles of Arf6 in cancer cell invasion, metastasis and proliferation. *Life Sci*. 2017;182:80-84.
25. Yoo JH, Shi DS, Grossmann AH, et al. ARF6 is an actionable node that orchestrates oncogenic GNAQ signaling in uveal melanoma. *Cancer Cell*. 2016;29(6):889-904.
26. Li B, Antonyak MA, Zhang J, Cerione RA. RhoA triggers a specific signaling pathway that generates transforming microvesicles in cancer cells. *Oncogene*. 2012;3(4):219-224.
27. Becker A, Thakur BK, Weiss JM, Kim HS, Peinado H, Lyden D. Extracellular vesicles in cancer: cell-to-cell mediators of metastasis. *Cancer Cell*. 2016;30(6):836-848.
28. Nabhan JF, Hu R, Oh RS, Cohen SN, Lu Q. Formation and release of arrestin domain-containing protein 1-mediated microvesicles (ARMVs) at plasma membrane by recruitment of TSG101 protein. *Proc Natl Acad Sci USA*. 2012;109(11):4146-4151.
29. Bergsmedh A, Szeles A, Henriksson M, et al. Horizontal transfer of oncogenes by uptake of apoptotic bodies. *Proc Natl Acad Sci USA*. 2001;98(11):6407-6411.



30. Zhu Z, Zhang D, Lee H, et al. Macrophage-derived apoptotic bodies promote the proliferation of the recipient cells via shuttling microRNA-221/222. *J Leukoc Biol.* 2017;101(6):1349-1359.
31. Coumans F, Brisson AR, Buzas EI, et al. Methodological guidelines to study extracellular vesicles. *Circ Res.* 2017;120(10):1632-1648.
32. Merchant ML, Rood IM, Deegens J, Klein JB. Isolation and characterization of urinary extracellular vesicles: implications for biomarker discovery. *Nat Rev Nephrol.* 2017;13(12):731-749.
33. Woo H-K, Sunkara V, Park J, et al. Exodisc for rapid, size-selective, and efficient isolation and analysis of nanoscale extracellular vesicles from biological samples. *ACS Nano.* 2017;11(2):1360-1370.
34. Konoshenko MY, Lekchnov EA, Vlassov AV, Laktionov PP. Isolation of extracellular vesicles: general methodologies and latest trends. *Biomed Res Int.* 2018;2018:8545347.
35. Van Deun J, Mestdagh P, Sormunen R, et al. The impact of disparate isolation methods for extracellular vesicles on downstream RNA profiling. *J Extracell Vesicles.* 2014;3(1):24858.
36. Lobb RJ, Becker M, Wen Wen S, et al. Optimized exosome isolation protocol for cell culture supernatant and human plasma. *J Extracell Vesicles.* 2015;4:27031.
37. Salih M, Fenton RA, Knipscheer J, et al. An immunoassay for urinary extracellular vesicles. *Am J Physiol Renal Physiol.* 2016;310(8):F796-F801.
38. Ingato D, Lee JU, Sim SJ, Kwon YJ. Good things come in small packages: overcoming challenges to harness extracellular vesicles for therapeutic delivery. *J Control Release.* 2016;241:174-185.
39. Nakai W, Yoshida T, Diez D, et al. A novel affinity-based method for the isolation of highly purified extracellular vesicles. *Sci Rep.* 2016;6:33935.
40. Balaj L, Atai NA, Chen W, et al. Heparin affinity purification of extracellular vesicles. *Sci Rep.* 2015;5:10266.
41. Samsonov R, Shtam T, Burdakov V, et al. Lectin-induced agglutination method of urinary exosomes isolation followed by mi-RNA analysis: application for prostate cancer diagnostic. *Prostate.* 2016;76(1):68-79.
42. Jung MK, Mun JY. Sample preparation and imaging of exosomes by transmission electron microscopy. *J Vis Exp.* 2018;131:e56482.
43. Brisson AR, Tan S, Linares R, Gounou C, Arraud N. Extracellular vesicles from activated platelets: a semiquantitative cryo-electron microscopy and immuno-gold labeling study. *Platelets.* 2017;28(3):263-271.
44. Chuo S-Y, Chien J-Y, Lai C-K. Imaging extracellular vesicles: current and emerging methods. *J Biomed Sci.* 2018;25(1):91.
45. Szatanek R, Baj-Krzyworzeka M, Zimoch J, Lekka M, Siedlar M, Baran J. The methods of choice for extracellular vesicles (EVs) characterization. *Int J Mol Sci.* 2017;18(6):1153.
46. Libregts S, Arkesteijn G, Németh A, Nolte-t Hoen E, Wauben M. Flow cytometric analysis of extracellular vesicle subsets in plasma: impact of swarm by particles of non-interest. *J Thromb Haemost.* 2018;16(7):1423-1436.
47. Im H, Shao H, Park YI, et al. Label-free detection and molecular profiling of exosomes with a nano-plasmonic sensor. *Nat Biotechnol.* 2014;32(5):490-495.
48. Shao H, Chung J, Balaj L, et al. 443 Protein typing of circulating microvesicles allows real-time monitoring of glioblastoma therapy. *Nat Med.* 2012;48(12):137-138.
49. Yoshioka Y, Kosaka N, Konishi Y, et al. Ultra-sensitive liquid biopsy of circulating extracellular vesicles using ExoScreen. *Nat Commun.* 2014;5:3591.
50. Ricci-Vitiani L, Pallini R, Biffoni M, et al. Tumour vascularization via endothelial differentiation of glioblastoma stem-like cells. *Nature.* 2010;468(7325):824-828.
51. Audia A, Conroy S, Glass R, Bhat K. The impact of the tumor microenvironment on the properties of glioma stem-like cells. *Front Oncol.* 2017;7:143.
52. Singh M, Yelle N, Venugopal C, Singh SK. EMT: Mechanisms and therapeutic implications. *Pharmacol Ther.* 2018;182:80-94.
53. Aga M, Bentz GL, Raffa S, et al. Exosomal HIF1 $\alpha$  supports invasive potential of nasopharyngeal carcinoma-associated LMP1-positive exosomes. *Oncogene.* 2014;33(37):4613-4622.
54. Gopal SK, Greening DW, Rai A, et al. Extracellular vesicles: their role in cancer biology and epithelial-mesenchymal transition. *Biochem J.* 2017;474(1):21-45.
55. Greening DW, Gopal SK, Mathias RA, et al. Emerging roles of exosomes during epithelial-mesenchymal transition and cancer progression. *Semin Cell Dev Biol.* 2015;40:60-71.
56. Franzen CA, Blackwell RH, Todorovic V, et al. Urothelial cells undergo epithelial-to-mesenchymal transition after exposure to muscle invasive bladder cancer exosomes. *Oncogenesis.* 2015;4:e163.
57. Peinado H, Zhang H, Matei IR, et al. Pre-metastatic niches: organ-specific homes for metastases. *Nat Rev Cancer.* 2017;17(5):302.
58. Grange C, Tapparo M, Collino F, et al. Microvesicles released from human renal cancer stem cells stimulate angiogenesis and formation of lung premetastatic niche. *Cancer Res.* 2011;71(15):5346-5356.
59. Smyth T, Kullberg M, Malik N, Smith-Jones P, Graner MW, Anchordoquy TJ. Biodistribution and delivery efficiency of unmodified tumor-derived exosomes. *J Control Release.* 2015;199:145-155.
60. Peinado H, Alečković M, Lavotshkin S, et al. Melanoma exosomes educate bone marrow progenitor cells toward a pro-metastatic phenotype through MET. *Nat Med.* 2012;18(6):883-891.
61. Liu Y, Gu Y, Han Y, et al. Tumor exosomal RNAs promote lung pre-metastatic niche formation by activating alveolar epithelial TLR3 to recruit neutrophils. *Cancer Cell.* 2016;30(2):243-256.
62. Costa-Silva B, Aiello NM, Ocean AJ, et al. Pancreatic cancer exosomes initiate pre-metastatic niche formation in the liver. *Nat Cell Biol.* 2015;17(6):816-826.
63. Fong MY, Zhou W, Liu L, et al. Breast-cancer-secreted miR-122 reprograms glucose metabolism in premetastatic niche to promote metastasis. *Nat Cell Biol.* 2015;17(2):183-194.
64. Choi D, Lee TH, Spinelli C, Chennakrishnaiah S, D'Asti E, Rak J. Extracellular vesicle communication pathways as regulatory targets of oncogenic transformation. *Semin Cell Dev Biol.* 2017;67:11-22.
65. Nabet BY, Qiu YU, Shabason JE, et al. Exosome RNA unshielding couples stromal activation to pattern recognition receptor signaling in cancer. *Cell.* 2017;170(2):352-366.e313.
66. Hood JL, San RS, Wickline SA. Exosomes released by melanoma cells prepare sentinel lymph nodes for tumor metastasis. *Cancer Res.* 2011;71(11):3792.
67. Ruivo CF, Adem B, Silva M, Melo SA. The biology of cancer exosomes: insights and new perspectives. *Cancer Res.* 2017;77(23):6480-6488.
68. Sung BH, Weaver AM. Exosome secretion promotes chemotaxis of cancer cells. *Cell Adh Migr.* 2017;11(2):187-195.
69. Baroni S, Romero-Cordoba S, Plantamura I, et al. Exosome-mediated delivery of miR-9 induces cancer-associated fibroblast-like properties in human breast fibroblasts. *Cell Death Dis.* 2016;7(7):e2312.
70. Mishra VK, Subramaniam M, Kari V, et al. Kruppel-like transcription factor KLF10 suppresses TGF $\beta$ -induced epithelial-to-mesenchymal transition via a negative feedback mechanism. *Cancer Res.* 2017;77(9):2387-2400.
71. Gu J, Qian H, Shen LI, et al. Gastric cancer exosomes trigger differentiation of umbilical cord derived mesenchymal stem cells to carcinoma-associated fibroblasts through TGF- $\beta$ /Smad pathway. *PLoS ONE.* 2012;7(12):e52465.
72. Corrado C, Saieva L, Raimondo S, Santoro A, De Leo G, Alessandro R. Chronic myelogenous leukaemia exosomes modulate bone

- marrow microenvironment through activation of epidermal growth factor receptor. *J Cell Mol Med.* 2016;20(10):1829-1839.
73. Todorova D, Simoncini S, Lacroix R, Sabatier F, Dignat-George F. Extracellular vesicles in angiogenesis. *Circ Res.* 2017;120(10):1658-1673.
  74. Du T, Ju G, Wu S, et al. Microvesicles derived from human Wharton's jelly mesenchymal stem cells promote human renal cancer cell growth and aggressiveness through induction of hepatocyte growth factor. *PLoS ONE.* 2014;9(5):e96836.
  75. Berlato C, Khan MN, Schioppa T, et al. A CCR7 antagonist reverses the tumor-promoting microenvironment of renal cancer. *J Clin Invest.* 2017;127(3):801-813.
  76. Katlinski KV, Gui J, Katlinskaya YV, et al. Inactivation of interferon receptor promotes the establishment of immune privileged tumor microenvironment. *Cancer Cell.* 2017;31(2):194-207.
  77. Saha D, Martuza RL, Rabkin SD. Macrophage polarization contributes to glioblastoma eradication by combination immunovirotherapy and immune checkpoint blockade. *Cancer Cell.* 2017;32(2):253-267.e255.
  78. Li X, Wang X. The emerging roles and therapeutic potential of exosomes in epithelial ovarian cancer. *Mol Cancer.* 2017;16(1):92.
  79. Guo D, Lui G, Lai SL, et al. RAB27A promotes melanoma cell invasion and metastasis via regulation of pro-invasive exosomes. *Int J Cancer.* 2019;144:3070-3085.
  80. Seo N, Shirakura Y, Tahara Y, et al. Activated CD8(+) T cell extracellular vesicles prevent tumour progression by targeting of lesional mesenchymal cells. *Nat Commun.* 2018;9(1):435.
  81. Sun Z, Yang S, Zhou Q, et al. Emerging role of exosome-derived long non-coding RNAs in tumor microenvironment. *Mol Cancer.* 2018;17(1):82.
  82. Zhou R, Chen KK, Zhang J, et al. The decade of exosomal long RNA species: an emerging cancer antagonist. *Mol Cancer.* 2018;17(1):75.
  83. Torre LA, Bray F, Siegel RL, Ferlay J, Lortet-Tieulent J, Jemal A. Global cancer statistics, 2012. *CA Cancer J Clin.* 2015;65(2):69-90.
  84. Barata PC, Rini BI. Treatment of renal cell carcinoma: current status and future directions. *CA Cancer J Clin.* 2017;67(6):507-524.
  85. Novara G, Ficarra V, Antonelli A, et al. Validation of the 2009 TNM version in a large multi-institutional cohort of patients treated for renal cell carcinoma: are further improvements needed? *Eur Urol.* 2010;58(4):588-595.
  86. Fisher R, Gore M, Larkin J. Current and future systemic treatments for renal cell carcinoma. *Semin Cancer Biol.* 2013;23(1):38-45.
  87. Schubert M, Junker K, Heinzlmann J. Prognostic and predictive miRNA biomarkers in bladder, kidney and prostate cancer: Where do we stand in biomarker development? *J Cancer Res Clin Oncol.* 2016;142(8):1673-1695.
  88. Yang L, Wu X, Wang D, Luo C, Chen L. Renal carcinoma cell-derived exosomes induce human immortalized line of Jurkat T lymphocyte apoptosis in vitro. *Urol Int.* 2013;91(3):363-369.
  89. Raimondo F, Morosi L, Corbetta S, et al. Differential protein profiling of renal cell carcinoma urinary exosomes. *Mol Biosyst.* 2013;9(6):1220-1233.
  90. De Palma G, Sallustio F, Curci C, et al. The three-gene signature in urinary extracellular vesicles from patients with clear cell renal cell carcinoma. *J Cancer.* 2016;7(14):1960-1967.
  91. Del BP, Raimondo F, Pieragostino D, et al. A hyphenated microLC-Q-TOF-MS platform for exosomal lipidomics investigations: application to RCC urinary exosomes. *Electrophoresis.* 2012;33(4):689-696.
  92. Butz H, Nofech-Mozes R, Ding Q, et al. Exosomal microRNAs are diagnostic biomarkers and can mediate cell-cell communication in renal cell carcinoma. *Eur Urol Focus.* 2016;2(2):210-218.
  93. Zhang W, Ni M, Su Y, et al. MicroRNAs in serum exosomes as potential biomarkers in clear-cell renal cell carcinoma. *Eur Urol Focus.* 2018;4:412-419.
  94. Fujii N, Hirata H, Ueno K, et al. Extracellular miR-224 as a prognostic marker for clear cell renal cell carcinoma. *Oncotarget.* 2017;8(66):109877-109888.
  95. Jinguishi K, Uemura M, Ohnishi N, et al. Extracellular vesicles isolated from human renal cell carcinoma tissues disrupt vascular endothelial cell morphology via azurocidin. *Int J Cancer.* 2018;142(3):607-617.
  96. Milella M, Felici A. Biology of metastatic renal cell carcinoma. *J Cancer.* 2011;2:369-373.
  97. Zhang L, Wu X, Luo C, et al. The 786-0 renal cancer cell-derived exosomes promote angiogenesis by downregulating the expression of hepatocyte cell adhesion molecule. *Mol Med Report.* 2013;8(1):272-276.
  98. Gu X, Erb U, Büchler MW, Zöller M. Improved vaccine efficacy of tumor exosome compared to tumor lysate loaded dendritic cells in mice. *Int J Cancer.* 2015;136(4):E74-E84.
  99. Diao J, Yang X, Song X, et al. Exosomal Hsp70 mediates immunosuppressive activity of the myeloid-derived suppressor cells via phosphorylation of Stat3. *Med Oncol.* 2015;32(2):453.
  100. Zhang Y, Luo CL, He BC, Zhang JM, Cheng G, Wu XH. Exosomes derived from IL-12-anchored renal cancer cells increase induction of specific antitumor response in vitro: a novel vaccine for renal cell carcinoma. *Int J Oncol.* 2010;36(1):133-140.
  101. Jiang XL, Zhang Y, Tan B, Luo CL, Wu XH. Renal tumor-derived exosomes inhibit hepaCAM expression of renal carcinoma cells in a p-AKT-dependent manner. *Neoplasma.* 2014;61(4):416.
  102. Bussolati B, Bruno S, Grange C, Ferrando U, Camussi G. Identification of a tumor-initiating stem cell population in human renal carcinomas. *FASEB J.* 2008;22(10):3696-3705.
  103. Posadas EM, Limvorasak S, Figlin RA. Targeted therapies for renal cell carcinoma. *Nat Rev Nephrol.* 2017;13(8):496-511.
  104. Molina AM, Lin X, Korytowsky B, et al. Sunitinib objective response in metastatic renal cell carcinoma: analysis of 1059 patients treated on clinical trials. *Eur J Cancer.* 2014;50(2):351-358.
  105. Qu LE, Ding J, Chen C, et al. Exosome-transmitted lncARSR promotes sunitinib resistance in renal cancer by acting as a competing endogenous RNA. *Cancer Cell.* 2016;29(5):653-668.
  106. Babjuk M, Böhle A, Burger M, et al. EAU Guidelines on non-muscle-invasive urothelial carcinoma of the bladder: update 2016. *Eur Urol.* 2017;71(3):447-461.
  107. Almallah YZ, Rennie CD, Stone J, Lancashire M. Urinary tract infection and patient satisfaction after flexible cystoscopy and urodynamic evaluation. *Urology.* 2000;56(1):37-39.
  108. Welton JL, Khanna S, Giles PJ, et al. Proteomics analysis of bladder cancer exosomes. *Mol Cell Proteomics.* 2010;9(6):1324-1338.
  109. Chen C-L, Lai Y-F, Tang P, et al. Comparative and targeted proteomic analyses of urinary microparticles from bladder cancer and hernia patients. *J Proteome Res.* 2012;11(12):5611.
  110. Lin S-Y, Chang C-H, Wu H-C, et al. Proteome profiling of urinary exosomes identifies alpha 1-antitrypsin and H2B1K as diagnostic and prognostic biomarkers for urothelial carcinoma. *Sci Rep.* 2016;6:34446.
  111. Silvers CR, Miyamoto H, Messing EM, Netto GJ, Lee Y-F. Characterization of urinary extracellular vesicle proteins in muscle-invasive bladder cancer. *Oncotarget.* 2017;8(53):91199-91208.
  112. Beckham CJ, Olsen J, Yin P-N, et al. Bladder cancer exosomes contain EDIL-3/Del1 and facilitate cancer progression. *J Urol.* 2014;192(2):583-592.
  113. Smalley DM, Sheman NE, Nelson K, Theodorescu D. Isolation and identification of potential urinary microparticle biomarkers of bladder cancer. *J Proteome Res.* 2008;7(5):2088-2096.
  114. Liang K, Liu F, Fan J, et al. Nanoplasmonic quantification of tumour-derived extracellular vesicles in plasma microsomes for diagnosis and treatment monitoring. *Nat Biomed Eng.* 2017;1(4):21-21.

115. Silvers CR, Liu Y-R, Wu C-H, Miyamoto H, Messing EM, Lee Y-F. Identification of extracellular vesicle-borne periostin as a feature of muscle-invasive bladder cancer. *Oncotarget*. 2016;7(17):23335-23345.
116. Perez A, Loizaga A, Arceo R, et al. A pilot study on the potential of RNA-associated to urinary vesicles as a suitable non-invasive source for diagnostic purposes in bladder cancer. *Cancers (Basel)*. 2014;6(1):179-192.
117. Matsuzaki K, Fujita K, Jingushi K, et al. MiR-21-5p in urinary extracellular vesicles is a novel biomarker of urothelial carcinoma. *Oncotarget*. 2017;8(15):24668-24678.
118. Andreu Z, Otta Oshiro R, Redruello A, et al. Extracellular vesicles as a source for non-invasive biomarkers in bladder cancer progression. *Eur J Pharm Sci*. 2017;98:70-79.
119. Baumgart S, Hölter S, Ohlmann C-H, et al. Exosomes of invasive urothelial carcinoma cells are characterized by a specific miRNA expression signature. *Oncotarget*. 2017;8(35):58278.
120. Long JD, Sullivan TB, Humphrey J, et al. A non-invasive miRNA based assay to detect bladder cancer in cell-free urine. *Am J Transl Res*. 2015;7(11):2500-2509.
121. Ostenfeld MS, Jeppesen DK, Laurberg JR, et al. Cellular disposal of miR23b by RAB27-dependent exosome release is linked to acquisition of metastatic properties. *Cancer Res*. 2014;74(20):5758-5771.
122. Armstrong DA, Green BB, Seigne JD, Schned AR, Marsit CJ. MicroRNA molecular profiling from matched tumor and bio-fluids in bladder cancer. *Mol Cancer*. 2015;14(1):194.
123. Berrondo C, Flax J, Kucherov V, et al. Expression of the long non-coding RNA HOTAIR correlates with disease progression in bladder cancer and is contained in bladder cancer patient urinary exosomes. *PLoS ONE*. 2016;11(1):e0147236.
124. Liang L-G, Kong M-Q, Zhou S, et al. An integrated double-filtration microfluidic device for isolation, enrichment and quantification of urinary extracellular vesicles for detection of bladder cancer. *Sci Rep*. 2017;7:46224.
125. Yang L, Wu X-H, Wang D, Luo C-L, Chen L-X. Bladder cancer cell-derived exosomes inhibit tumor cell apoptosis and induce cell proliferation in vitro. *Mol Med Report*. 2013;8(4):1272-1278.
126. Xue M, Chen W, Xiang AN, et al. Hypoxic exosomes facilitate bladder tumor growth and development through transferring long non-coding RNA-UCA1. *Mol Cancer*. 2017;16(1):143.
127. Greco KA, Franzen CA, Foreman KE, Flanigan RC, Kuo PC, Gupta GN. PLK-1 silencing in bladder cancer by siRNA delivered with exosomes. *Urology*. 2016;91(241):e241-247.
128. Ferlay J, Soerjomataram I, Dikshit R, et al. Cancer incidence and mortality worldwide: sources, methods and major patterns in GLOBOCAN 2012. *Int J Cancer*. 2015;136(5):E359-386.
129. Rao AR, Motiwala HG, Karim OM, et al. The discovery of prostate-specific antigen. *BJU Int*. 2008;101(1):5-10.
130. Crawford ED, Abrahamsson P-A. PSA-based screening for prostate cancer: how does it compare with other cancer screening tests? *Eur Urol*. 2008;54(2):262-273.
131. Foj L, Ferrer F, Serra M, et al. Exosomal and non-exosomal urinary miRNAs in prostate cancer detection and prognosis. *Prostate*. 2017;77(6):573-583.
132. Rodríguez M, Bajo-Santos C, Hessvik NP, et al. Identification of non-invasive miRNAs biomarkers for prostate cancer by deep sequencing analysis of urinary exosomes. *Mol Cancer*. 2017;16(1):156.
133. Khan S, Jutzy J, Valenzuela M, et al. Plasma-derived exosomal survivin, a plausible biomarker for early detection of prostate cancer. *PLoS ONE*. 2012;7(10):e46737.
134. Wang L, Skotland T, Berge V, Sandvig K, Llorente A. Exosomal proteins as prostate cancer biomarkers in urine: from mass spectrometry discovery to immunoassay-based validation. *Eur J Pharm Sci*. 2016;98:80.
135. Duijvesz D, Versluis C, van der Fels C, et al. Immuno-based detection of extracellular vesicles in urine as diagnostic marker for prostate cancer. *Int J Cancer*. 2015;137(12):2869-2878.
136. Liu C, Guo J, Tian F, et al. Field-free isolation of exosomes from extracellular vesicles by microfluidic viscoelastic flows. *ACS Nano*. 2017;11(7):6968-6976.
137. Lu Q, Zhang J, Allison R, et al. Identification of extracellular delta-catenin accumulation for prostate cancer detection. *Prostate*. 2009;69(4):411-418.
138. Bijnsdorp IV, Geldof AA, Lavaei M, Piersma SR, van Moorselaar R, Jimenez CR. Exosomal ITGA3 interferes with non-cancerous prostate cell functions and is increased in urine exosomes of metastatic prostate cancer patients. *J Extracell Vesicles*. 2013;2:22097.
139. Fujita K, Kume H, Matsuzaki K, et al. Proteomic analysis of urinary extracellular vesicles from high Gleason score prostate cancer. *Sci Rep*. 2017;7:42961.
140. Øverbye A, Skotland T, Koehler CJ, et al. Identification of prostate cancer biomarkers in urinary exosomes. *Oncotarget*. 2015;6(30):30357-30376.
141. Ronquist KG, Ronquist G, Larsson A, Carlsson L. Proteomic analysis of prostate cancer metastasis-derived prostasomes. *Anticancer Res*. 2010;30(2):285-290.
142. Sandvig K, Llorente A. Proteomic analysis of microvesicles released by the human prostate cancer cell line PC-3. *Mol Cell Proteomics*. 2012;11(7):M111.012914.
143. Duijvesz D, Burnum-Johnson KE, Gritsenko MA, et al. Proteomic profiling of exosomes leads to the identification of novel biomarkers for prostate cancer. *PLoS ONE*. 2013;8(12):e82589.
144. Utleg AG, Yi EC, Xie T, et al. Proteomic analysis of human prostasomes. *Prostate*. 2010;56(2):150-161.
145. Simona Principe E, Kim Y, Sinha A, et al. In-depth proteomic analyses of exosomes isolated from expressed prostatic secretions in urine. *Proteomics*. 2013;13(10-11):1667-1671.
146. Sequeiros T, Rigau M, Chiva C, et al. Targeted proteomics in urinary extracellular vesicles identifies biomarkers for diagnosis and prognosis of prostate cancer. *Oncotarget*. 2016;8(3):4960.
147. Gabriel K, Ingram A, Austin R, et al. Regulation of the tumor suppressor PTEN through exosomes: a diagnostic potential for prostate cancer. *PLoS ONE*. 2013;8(7):e70047.
148. Kato T, Mizutani K, Kameyama K, et al. Serum exosomal P-glycoprotein is a potential marker to diagnose docetaxel resistance and select a taxoid for patients with prostate cancer. *Urol Oncol*. 2015;33(9):385.e315-320.
149. Welton JL, Brennan P, Gurney M, et al. Proteomics analysis of vesicles isolated from plasma and urine of prostate cancer patients using a multiplex, aptamer-based protein array. *J Extracell Vesicles*. 2016;5(1):31209.
150. Lázaro-Ibáñez E, Sanz-García A, Visakorpi T, et al. Different gDNA content in the subpopulations of prostate cancer extracellular vesicles: apoptotic bodies, microvesicles, and exosomes. *Prostate*. 2014;74(14):1379-1390.
151. McKiernan J, Donovan MJ, O'Neill V, et al. A novel urine exosome gene expression assay to predict high-grade prostate cancer at initial biopsy. *JAMA Oncol*. 2016;2(7):882-889.
152. Neeb A, Hefele S, Bormann S, et al. Splice variant transcripts of the anterior gradient 2 gene as a marker of prostate cancer. *Oncotarget*. 2014;5(18):8681-8689.
153. Donovan MJ, Noerholm M, Bentink S, et al. A molecular signature of PCA3 and ERG exosomal RNA from non-DRE urine is predictive of initial prostate biopsy result. *Prostate Cancer Prostatic Dis*. 2015;18(4):370-375.
154. Hendriks RJ, Dijkstra S, Jannink SA, et al. Comparative analysis of prostate cancer specific biomarkers PCA3 and ERG in whole urine, urinary sediments and exosomes. *Clin Chem Lab Med*. 2016;54(3):483-492.
155. Motamedinia P, Scott AN, Bate KL, et al. Urine exosomes for non-invasive assessment of gene expression and mutations of prostate cancer. *PLoS ONE*. 2016;11(5):e0154507.

156. Royo F, Zuñiga-Garcia P, Torrano V, et al. Transcriptomic profiling of urine extracellular vesicles reveals alterations of CDH3 in prostate cancer. *Oncotarget*. 2016;7(6):6835-6846.
157. Işın M, Uysaler E, Özgür E, et al. Exosomal lncRNA-p21 levels may help to distinguish prostate cancer from benign disease. *Front Genet*. 2015;6:168.
158. Koppers-Lalic D, Hackenberg M, deMenezes R, et al. Noninvasive prostate cancer detection by measuring miRNA variants (isomiRs) in urine extracellular vesicles. *Oncotarget*. 2016;7(16):22566-22578.
159. Bryant RJ, Pawlowski T, Catto J, et al. Changes in circulating microRNA levels associated with prostate cancer. *Br J Cancer*. 2012;106(4):768-774.
160. Bryzgunova OE, Zaripov MM, Skvortsova TE, et al. Comparative study of extracellular vesicles from the urine of healthy individuals and prostate cancer patients. *PLoS ONE*. 2016;11(6):e0157566.
161. Wani S, Kaul D, Mavuduru RS, Kakkar N, Bhatia A. Urinary-exosomal miR-2909: a novel pathognomonic trait of prostate cancer severity. *J Biotechnol*. 2017;259:135-139.
162. Xu Y, Qin S, An T, Tang Y, Huang Y, Zheng L. MiR-145 detection in urinary extracellular vesicles increase diagnostic efficiency of prostate cancer based on hydrostatic filtration dialysis method. *Prostate*. 2017;77(10):1167-1175.
163. Hao X-K, Li Z, Ma Y-Y, et al. Exosomal microRNA-141 is upregulated in the serum of prostate cancer patients. *Onco Targets Ther*. 2016;9(1):139-148.
164. Huang X, Yuan T, Liang M, et al. Exosomal miR-1290 and miR-375 as prognostic markers in castration-resistant prostate cancer. *Eur Urol*. 2015;67(1):33-41.
165. Skotland T, Ekroos K, Kauhanen D, et al. Molecular lipid species in urinary exosomes as potential prostate cancer biomarkers. *Eur J Cancer*. 2017;70:122-132.
166. Yang JS, Lee JC, Byeon SK, Rha KH, Moon MH. Size dependent lipidomic analysis of urinary exosomes from patients with prostate cancer by flow field-flow fractionation and nanoflow liquid chromatography-tandem mass spectrometry. *Anal Chem*. 2017;89(4):2488-2496.
167. Puhka M, Takatalo M, Nordberg M-E, et al. Metabolomic profiling of extracellular vesicles and alternative normalization methods reveal enriched metabolites and strategies to study prostate cancer-related changes. *Theranostics*. 2017;7(16):3824-3841.
168. Nyalwidhe JO, Betesh LR, Powers TW, et al. Increased bisecting N-acetylglucosamine and decreased branched chain glycans of N-linked glycoproteins in expressed prostatic secretions associated with prostate cancer progression. *Proteomics Clin Appl*. 2013;7(9-10):677-689.
169. Webber JP, Spary LK, Sanders AJ, et al. Differentiation of tumour-promoting stromal myofibroblasts by cancer exosomes. *Oncogene*. 2015;34(3):290-302.
170. Castellana D, Zobairi F, Martinez MC, et al. Membrane microvesicles as actors in the establishment of a favorable prostatic tumoral niche: a role for activated fibroblasts and CX3CL1-CX3CR176 axis. *Cancer Res*. 2009;69(3):785-793.
171. Itoh T, Ito Y, Ohtsuki Y, et al. Microvesicles released from hormone-refractory prostate cancer cells facilitate mouse pre-osteoblast differentiation. *J Mol Histol*. 2012;43(5):509-515.
172. Karlsson T, Lundholm M, Widmark A, Persson E. Tumor cell-derived exosomes from the prostate cancer cell line TRAMP-C1 impair osteoclast formation and differentiation. *PLoS ONE*. 2016;11(11):e0166284.
173. Ye Y, Li S-L, Ma Y-Y, et al. Exosomal miR-141-3p regulates osteoblast activity to promote the osteoblastic metastasis of prostate cancer. *Oncotarget*. 2017;8(55):94834-94849.
174. Lundholm M, Schröder M, Nagaeva O, et al. Prostate tumor-derived exosomes down-regulate NKG2D expression on natural killer cells and CD8+ T cells: mechanism of immune evasion. *PLoS ONE*. 2014;9(9):e108925.
175. Clayton A, Mitchell JP, Court J, Mason MD, Tabi Z. Human tumor-derived exosomes selectively impair lymphocyte responses to interleukin-2. *Cancer Res*. 2007;67(15):7458-7466.
176. Seitz AK, Thoene S, Bietenbeck A, et al. AR-V7 in peripheral whole blood of patients with castration-resistant prostate cancer: association with treatment-specific outcome under abiraterone and enzalutamide. *Eur Urol*. 2017;72(5):828.
177. Del Re M, Biasco E, Crucitta S, et al. The detection of androgen receptor splice variant 7 in plasma-derived exosomal RNA strongly predicts resistance to hormonal therapy in metastatic prostate cancer patients. *Eur Urol*. 2017;71(5):680-687.
178. Seifalian A. The application of exosomes as a nanoscale cancer vaccine. *Int J Nanomed*. 2010;5(1):889-900.
179. Rountree RB, Mandl SJ, Nachtwey JM, et al. Exosome targeting of tumor antigens expressed by cancer vaccines can improve antigen immunogenicity and therapeutic efficacy. *Cancer Res*. 2011;71(15):5235-5244.
180. Corcoran C, Rani S, O'Brien K, et al. Docetaxel-resistance in prostate cancer: evaluating associated phenotypic changes and potential for resistance transfer via exosomes. *PLoS ONE*. 2012;7(12):e50999.
181. Lötvall J, Hill AF, Hochberg F, et al. Minimal experimental requirements for definition of extracellular vesicles and their functions: a position statement from the International Society for Extracellular Vesicles. *J Extracell Vesicles*. 2014;3:26913.
182. Thery C, Witwer KW, Aikawa E, et al. Minimal information for studies of extracellular vesicles 2018 (MISEV2018): a position statement of the International Society for Extracellular Vesicles and update of the MISEV2014 guidelines. *J Extracell Vesicles*. 2018;7(1):1535750.
183. Chen G, Huang AC, Zhang W, et al. Exosomal PD-L1 contributes to immunosuppression and is associated with anti-PD-1 response. *Nature*. 2018;560(7718):382-386.
184. Poggio M, Hu T, Pai C-C, et al. Suppression of exosomal PD-L1 induces systemic anti-tumor immunity and memory. *Cell*. 2019;177(2):414-427.e413.
185. Ha D, Yang N, Nadihe V. Exosomes as therapeutic drug carriers and delivery vehicles across biological membranes: current perspectives and future challenges. *Acta pharmaceutica Sinica B*. 2016;6(4):287-296.
186. Vader P, Mol EA, Pasterkamp G, Schiffelers RM. Extracellular vesicles for drug delivery. *Adv Drug Deliv Rev*. 2016;106(Pt A):148-156.
187. Willms E, Cabañas C, Mäger I, Wood M, Vader P. Extracellular vesicle heterogeneity: subpopulations, isolation techniques, and diverse functions in cancer progression. *Front Immunol*. 2018;9:738.

**How to cite this article:** Wu Z, Zhang Z, Xia W, Cai J, Li Y, Wu S. Extracellular vesicles in urologic malignancies—Implementations for future cancer care. *Cell Prolif*. 2019;52:e12659. <https://doi.org/10.1111/cpr.12659>