

# Processing of False Belief Passages During Natural Story Comprehension: An fMRI Study


Katerina D. Kandylaki,<sup>1,2,\*</sup> Arne Nagels,<sup>1</sup> Sarah Tune,<sup>3</sup> Richard Wiese,<sup>2</sup>  
Ina Bornkessel-Schlesewsky,<sup>4</sup> and Tilo Kircher<sup>1</sup>

<sup>1</sup>Department of Psychiatry and Psychotherapy, Philipps-University Marburg,  
Rudolf-Bultmann-Str. 8, Marburg 35039, Germany

<sup>2</sup>Department of Germanic Linguistics, Philipps-University Marburg, Deutschhausstr. 3,  
Marburg 35032, Germany

<sup>3</sup>Department of Neurology, University of California, Irvine, Biological Sciences III, Irvine,  
California, USA

<sup>4</sup>Cognitive Neuroscience Laboratory, School of Psychology, Social Work and Social Policy,  
University of South Australia, Box 2471, Adelaide, South Australia 5001, Australia



**Abstract:** The neural correlates of theory of mind (ToM) are typically studied using paradigms which require participants to draw explicit, task-related inferences (e.g., in the false belief task). In a natural setup, such as listening to stories, false belief mentalizing occurs incidentally as part of narrative processing. In our experiment, participants listened to auditorily presented stories with false belief passages (implicit false belief processing) and immediately after each story answered comprehension questions (explicit false belief processing), while neural responses were measured with functional magnetic resonance imaging (fMRI). All stories included (among other situations) one false belief condition and one closely matched control condition. For the implicit ToM processing, we modeled the hemodynamic response during the false belief passages in the story and compared it to the hemodynamic response during the closely matched control passages. For implicit mentalizing, we found activation in typical ToM processing regions, that is the angular gyrus (AG), superior medial frontal gyrus (SmFG), precuneus (PCUN), middle temporal gyrus (MTG) as well as in the inferior frontal gyrus (IFG) bilaterally. For explicit ToM, we only found AG activation. The conjunction analysis highlighted the left AG and MTG as well as the bilateral IFG as overlapping ToM processing regions for both implicit and explicit modes. Implicit ToM processing during listening to false belief passages, recruits the left SmFG and bilateral PCUN in addition to the “mentalizing network” known from explicit processing tasks. *Hum Brain Mapp* 36:4231–4246, 2015. © 2015 Wiley Periodicals, Inc.

**Key words:** theory of mind; false beliefs; fMRI; auditory stories



Additional Supporting Information may be found in the online version of this article.

Contract grant sponsor: Funded by Philipps-University Marburg and by the state of Hesse under the LOEWE research focus “Exploring Fundamental Linguistic Categories”.

\*Correspondence to: Katerina D. Kandylaki; Rudolf-Bultmann-Str. 8, 35039 Marburg, Germany. E-mail: kandylak@med.uni-marburg.de

Received for publication 11 February 2015; Revised 12 June 2015; Accepted 6 July 2015.

DOI: 10.1002/hbm.22907

Published online 10 September 2015 in Wiley Online Library (wileyonlinelibrary.com).

## INTRODUCTION

Theory of Mind (ToM) is the cognitive capacity to attribute mental states to self and others [Goldman et al., 2012; Premack and Woodruff, 1978]. This capacity, which is also referred to as mentalizing [e.g., Frith and Frith, 1999, 2010], is used incidentally in a broad range of naturally occurring social situations. Consider, for example, a situation in which two friends are talking and a third joins the discussion after 2 min. The first two will be aware that the third will have no knowledge of their preceding conversation: they have a “Theory of Mind” [Premack and Woodruff, 1978] that allows them to take on the third person’s perspective. Although the term ToM may appear relatively straightforward from the perspective of this initial definition, the rich facet of social situations where ToM is used remains to be systematically categorized [Schaafsma et al., 2015]. ToM is thought to involve a variety of sub- and super-processes [Schaafsma et al., 2015] and can be deconstructed based on criteria such as implicit versus explicit and cognitive versus affective [e.g., Schlawffke et al., 2014]. In addition to the deconstruction of the concept of ToM, and to quantify ToM as precisely as possible, a reconstruction of ToM components from basic building blocks is needed: for example, face recognition and gaze processing are essential for completing the reading the mind in the eyes test, which is a measure of mentalizing ability [Baron-Cohen et al., 2001]. fMRI research on specific ToM tasks, such as false belief processing, can inform this reconstruction with brain maps capturing instances of ToM, for details see Schaafsma et al. [2015]. In spite of these multifaceted aspects of ToM, several meta-analyses of functional neuroimaging studies connected to ToM processing [Mar, 2011; Northoff et al., 2006; Schurz et al., 2014; Spreng, Mar, and Kim, 2009; Van Overwalle, 2009] suggest that a number of regions appear to be involved in ToM regardless of the specific task. These include the bilateral temporo-parietal junction (TPJ), which corresponds to the inferior portion of the angular and supra-marginal gyri (AG, SMG) as well the posterior superior temporal sulcus, superior parts of the frontal gyrus (SFG), medial prefrontal cortex (mPFC), and precuneus (PCUN).

Van Overwalle and coworkers first investigated another aspect of spontaneous ToM processing in trait attributions [Kestemont et al., 2013; Ma et al., 2011, 2012] and causal attributions [Kestemont et al., 2015]; in these studies, spontaneous and intentional ToM processing was manipulated between participants via the experimental instructions. In another study, *implicit* mentalizing was elicited via visually presented images in the contrast of false versus true belief processing [Sommer et al., 2007]. According to a recent meta-analysis [Schurz et al., 2014], trait attribution and false belief processing show very similar activation patterns. This seems meaningful conceptually because causal attributions refer to beliefs about a temporary event just like false beliefs. However, stimulus presentation was very similar in many of the studies for trait attribution

and false belief processing that were included in the meta-analysis: all stimuli were visual and included a minimal amount of contextual information.

Here, by contrast, we present the first study to use auditory story stimuli in conjunction with a rich situational context, within which the false belief situations were embedded. We assume that ToM processing in such embedded false belief passages is *implicit* because it occurs during listening to a story with the simple goal of keeping track of the narrative plot. If the story involves situations that require the attribution of mental states to others, mentalizing would be a prerequisite for successful understanding of the plot. This notion is similar to van der Wel et al.’s. [2014] automatic belief tracking, although they measured behavior in the movement trajectory of a cursor while the subject was giving their response, after having watched a short movie which involved belief tracking. By contrast, a situation or task in which a participant was asked to actively answer questions or make judgements about a person’s mental state can be described as *explicit* ToM with regard to belief processing. In this case, the inferences drawn regarding another person’s mental state are clearly task-induced.

### False Belief Processing in Story Comprehension

Story-based approaches as opposed to highly controlled experimental setups provide the opportunity of testing false belief processing in the context of a semantically rich, cohesive story.

The first story-based neuroimaging study on ToM was reported by Fletcher et al. [1995]. The authors presented short stories each involving one false belief or control situation and examined the contrast between false belief versus physical stories versus unlinked sentences. Participants were instructed to read the stories and answer one question immediately after each story. They were, however, informed about the type of the story, that is, in the case of false belief stories, that they should pay attention to people’s beliefs and intentions. Also, the authors modeled false belief processing during the reading of false belief and control passages and during the answering of a related question together. For the critical contrast between ToM and physical stories, this study observed activation in left medial frontal regions and in the anterior and posterior cingulate cortex (aCC, pCC).

The majority of subsequent story-based neuroimaging investigation of ToM processing used the original false belief vignettes from the Fletcher et al. [1995] positron emission tomography (PET) study or translations thereof and compared them to different control conditions. All of them presented the stories visually and modeled participants’ reading process. For example, Vogeley et al. [2001] modified this material to cross ToM and self versus other processing. Story, question, and silent answer were all modeled together in a block design; thus, there was no

differentiation between *implicit* and *explicit* false belief processing. For the main effect of ToM, Vogeley et al. found differences in (among others): the right aCC, right SFG, and left lateral prefrontal cortex.

Using fMRI, Saxe and Kanwisher [2003] contrasted false belief and mechanical inference stories in two reading experiments and found ToM-related activation in the TPJ both for reading a false belief story (Experiment 1) and for reading the story and performing the question answering task (Experiment 2). This region has shown robust activation across a number of story-based studies on ToM processing [Aichhorn et al., 2009; Gallagher et al., 2000; Kobayashi et al., 2006; Lee et al., 2011; Saxe and Kanwisher, 2003; Spengler et al., 2009].

Aichhorn et al. [2009] tested the contrast of false belief versus photograph during story processing and were the first to explicitly distinguish two different modes in ToM processing: Time point 1: *Story*, when the participants read the story (again, *implicit* false belief processing in reading) and Time point 2: *Question*, when the participants read and answered the question about the story (*explicit* false belief processing in reading). Their results showed activation in the middle and superior temporal gyri (MTG, STG), superior and inferior frontal gyri (SFG, IFG) and in the TPJ for false belief compared with false photograph conditions at both time points. Also, PCUN activation was found only for the task but not for the story.

In summary, the story-based paradigms which investigated *explicit* false belief processing have identified a network comprising the following regions: mPFC, SFG, IFG, TPJ, MTG, and PCUN. However, all previous story-based studies used a block design in which one story and one answer formed one condition, either the false belief or the control condition. Most of these studies modeled *implicit* and *explicit* ToM together in the same block [Fletcher et al., 1995; Saxe and Kanwisher, 2003; Vogeley et al., 2001], with the notable exception of Aichhorn et al. [2009], who modeled them separately (but in reading comprehension and in short stories). One previous fMRI study did use auditory linguistic stimuli to contrast causality-related inferences with ToM-related inferences [Ferstl and Cramon, 2002]. However, this experiment only used minimal context and did not distinguish between ToM-related inferences drawn during listening and task performance.

The design details of previous studies (mostly visual stimulation with minimal context and an explicit task) may have contributed to some of the inconsistencies about the involvement of the TPJ in false belief processing that are apparent in the literature [Callejas et al. 2011]. More specifically, most of the above mentioned studies did not differentiate between components such as text processing or memory maintenance. The current study, by contrast, extended the work of Ferstl and Cramon [2002] using linguistically rich narratives and by embedding false belief passages into these larger contexts. This allowed us to model the processing of false belief passages in language

comprehension (*implicit* ToM) separately from overall story processing and decision-making during question answering. In comparison to previous designs, this manipulation should render results less sensitive, for example, to individual processing speed (as in reading paradigms). In addition, we aimed to capture ToM-related processes when reading and answering a question about false beliefs. To this end, we modeled reading of both question and answer options during which the participants manipulated the story information to perform the task (*explicit* ToM).

Ferstl and Cramon [2002] raised the question of the relationship between coherence processes in narrative comprehension and ToM processing. For both logical (coherence) and person-related (ToM) inferences, the results mainly shared frontomedian cortex (FMC) activation. The authors connect this activation to a domain general function of the FMC, “the initiation and maintenance of nonautomatic cognitive processes” [Ferstl and Cramon, 2002, p. 1,610]. This view accords well with a recent review on ToM, which argues that ToM-related processes should be decomposed into smaller subprocesses blocks [Schaafsma et al. 2015]. From this perspective, we could assume that both language and ToM are high-level functions that comprise a number of basic subprocesses [for a recent neurobiological perspective that advocates the decomposition of language into basic submechanisms, see Bornkessel-Schlesewsky et al., 2015]. Some of the subprocesses might overlap: for example, one basic subprocess of ToM [according to Schaafsma et al., 2015] may be the understanding of causality, which is also a basic subprocess in the semantics of language comprehension [Kuperberg et al. 2011]. It may be the case that, when these subprocesses overlap, brain activations in the respective networks are enhanced. This might be one possibility of how linguistic and social processes like ToM interact in the brain, although this interaction remains to be studied more systematically.

### The Present Study

The present fMRI study tested *implicit* and *explicit* false belief processing by means of a novel paradigm: (a) we presented stories (approximately 2 min in length) with false belief and control situations embedded among a range of other scenarios; (b) these stories were presented auditorily; and (c) this design allowed us to model the hemodynamic response to false belief and control events based on their onset and duration within the two-minute-long story. We chose to have participants listen to the stories (instead of reading them) to increase naturalness in the setup, as reading is a culturally recent innovation that is less than 6,000 years old (see also [Dehaene and Cohen, 2011]’s “neuronal recycling hypothesis”). To also examine *explicit* mentalizing, we presented participants with two questions subsequent to each story, targeting both false

belief as well as control story content. The questions and answers were presented visually, to provide participants with various information modalities, which would keep them alert throughout the whole experiment.

Based on previous research on story-based false belief processing, we expected to observe activation for the contrast of false belief ToM versus control passages during story listening (*implicit* false belief processing) in the mentalizing network: mPFC, SFG, IFG, MTG, PCUN, and TPJ. Based on the Aichhorn et al. [2009] findings, we expected *explicit* false belief processing activation to manifest itself mainly in the right TPJ and in the bilateral SFG as part of the mPFC (see Schurz et al. [2014] for the role of the mPFC in mentalizing).

## METHODS

### Participants

Twenty-two monolingual native speakers of German participated in the study, all right-handed (Edinburgh Inventory of Handedness) (age mean = 24.3 years,  $sd = 2.1$  years, male  $N = 6$ ), recruited from postings at the University of Marburg. We had to exclude data from two participants due to movement artefacts, resulting in a total of 20 datasets that entered the final analysis for the current study. The study was approved by the ethics committee of the Faculty of Medicine of the Philipps-University Marburg. All participants gave written informed consent before participating in the study and were paid 30 euros for participation.

### Stimuli

For testing *implicit* ToM processing we created 20 stories with a length of 2 min ( $\pm 10$  s; mean and standard deviation of story length 306 (13) words, 23 (4) sentences). All stories included one false belief condition (im-TOM) and one control condition (im-NONTOM). The 40 (im-TOM and im-NONTOM) situations included a variety of social interactions: 12 every day situations for example, *cooking, playing, driving*, 12 scenarios related to hobbies such as *hiking, visiting an art exhibition, sports*, 10 work situations in for example, *shop, office, conference*, 4 school situations (*school trip, chess competition*) and 2 university situations (*department party*). The common pattern of all situations was that there were two persons involved and one of them had a false belief. To comprehend these passages, the participants needed to use their ToM. The control passages were matched in length to the false belief passages and were part of a physical chain of events, in which one event led to the next one. The important difference between false belief and control passages was the existence of “different minds”, the situation in which two story participants have different beliefs. In the analysis of the results, the event chain pas-

sages which did not require mentalizing are referred to as control or NONTOM passages.

We developed two versions of each story as follows: version A included im-TOM at the end of the first minute of the story and im-NONTOM (control passage) at the end of the second minute of the story. In version B, the manipulation was reversed: im-NONTOM (control passage) at the end of the first minute and im-TOM at the end of the second minute of the story. This design resulted in a total of 40 stories ( $20 \times 2$  minimal pairs). To avoid one participant hearing two versions of the same story (with a similar plot but alternating critical passages), we split the 40 stories into two lists of 20 stories each. One participant heard one of the two lists in an individually randomized order: for example Participant 1 would hear list 1 which contained story 1A, 2B, 3B, 4A, and so on, and Participant 2 would hear list 2 which contained story 1B, 2A, 3A, 4B, and so on. Examples of the critical passages from two versions of the same story (1A and 1B) were as follows (translated from the German original):

#### Story 1A

- Within the first half: “[...] but his wife was so busy taking pictures of the idyllic landscape, so she didn’t realise, that her husband ate all the sandwiches. When later they arrived at the summit of Brocken, she also wanted to eat a sandwich, but found only drinks in her bag. *She thought that maybe she had forgotten the food in the car.*<sub>im-TOM</sub> [...]”
- Within the second half: “[...] The man took the camera from his backpack and gave it to the hiker. The hiker put in a lot of effort and *took a whole series of pictures from all different perspectives.*<sub>im-NONTOM</sub> The couple thanked him and went on walking. [...]”

#### Story 1B

- Within the first half: “[...] A few minutes later she came back to her husband and he had already eaten three sandwiches. At the summit of the Brocken the woman got also hungry and she found the last sandwich. *With delight she ate it and drank a few sips of apple juice.*<sub>im-NONTOM</sub> [...]”
- Within the second half: “[...] The man gave the camera to the hiker who accidentally packed it up in his backpack after photographing the couple. Then he went on hiking. *The couple was looking for the camera without success.*<sub>im-TOM</sub> until the hiker came back and apologised many times. [...]”

Stimuli were spoken by a professionally trained female speaker of German at a normal speech rate. We recorded the stimuli in a sound proof electroencephalogram (EEG) laboratory cabin with a sampling rate of 44.1 kHz and a 16bit (mono) sample size. For sampling, we used the sound recording and analysis software Amadeus Pro

**TABLE I. Question and answer details for the contrast ex-TOM versus ex-NONTOM**

Metric	All	ex-TOM		ex-NONTOM
	Mean (standard deviation)			
Question length in words	8.9 (3)	9.5 (3.2)	$P = 0.06$	8.3 (2.8)
Number of clauses of the question	1.5 (0.5)	1.6 (0.5)	$P = 0.11$	1.4 (0.5)
Answer length in words	3.7 (1.9)	3.9 (2.2)	$P = 0.08$	3.5 (1.5)
	Percentage (Number)			
Location content of question	58.8% (47/80)	62.5% (25/40)	$P = 0.49$	55% (22/40)

The inferential statistics represent model comparison of two models: (a) the null model, in which only the random factor of story is included and (b) the main effect of ToM, in which the type of the question (ex-TOM vs. ex-NONTOM) as well as the random factor of story is included.

(version1.5.3, HairerSoft) and an Electret microphone (Beyerdynamic MC930C). Two example stories are available in the Supporting Information.

We pretested the stories prior to the imaging study to validate their quality. In an online questionnaire, we gathered ratings from 177 participants. The questionnaire was distributed through a students' mailing list. Participants who did not fulfil the language criteria (monolingually raised German native speakers) were excluded from the final analysis. The participants were asked to judge comprehensibility ("How comprehensible was this passage?") and naturalness ("How natural was this passage?") of the auditory stimuli. Ratings were collected on a 4-point scale from 1 (very unnatural/incomprehensible) to 4 (extremely natural/comprehensible). The use of earphones was highly recommended in the instructions of the questionnaire. An analysis using linear mixed effects models using R statistical software [Team, 2014] and the lme4 package [Bates et al., 2014] with fixed factor of condition and random effects (only intercepts due to convergence problems) of story and subject showed that ratings for im-TOM and im-NONTOM passages did not differ significantly: Comprehensibility means (standard deviations): im-TOM 3.62 (0.62) versus im-NONTOM 3.64 (0.57),  $P = 0.67$ , Naturalness means (sds): im-TOM 3.23 (0.79) versus im-NONTOM 3.22 (0.77),  $P = 0.87$ .

In the scanner, the stories were presented auditorily, while the subject was looking at a fixation point in the center of a computer display. After each story, two questions and two possible answers for each question were presented visually. The questions referred to the im-TOM and im-NONTOM part of the story, thus creating the *explicit* conditions ex-TOM and ex-NONTOM. For example, the questions and answers for the above-mentioned example passages were:

#### Story 1A

- ex-TOM: Where did the woman think that the sandwiches were? Answers: a. In the car versus b. In her husband's stomach

- ex-NONTOM: Who had the camera when there was a series of pictures taken? Answers: a. The hiker versus b. The old man


#### Story 1B

- ex-NONTOM: How many sandwiches were left for the woman, after the man had finished eating? Answers: a. One versus b. Three
- ex-TOM: Where was the camera in the opinion of the couple, after the hiker took pictures of them? Answers: a. In the backpack of the couple versus b. In the backpack of the hiker

To rule out possible alternative explanations of the results for the ex-TOM versus ex-NONTOM contrast, questions and answers were analyzed according to the metrics in Table I. The following metrics were analyzed using linear mixed effects models (package lme4) in R: (1) question length in words, (2) number of clauses of question (as a measurement of syntactic complexity), (3) answer length in words, and (4) question type, whether it asked about location or not. In the inferential statistics, we used likelihood ratio tests to compare: (a) the null model, in which only the random factor of story is included and (b) the main effect of ToM, in which the type of the question (ex-TOM vs. ex-NONTOM) as well as the random factor of story is included. For question and answer length, the main effect of ToM model showed a marginally significant improvement over the null model ( $P = 0.0643$  and  $P = 0.08183$ , respectively). For the criteria of number of clauses and question type (location versus non-location) there was no significant difference between the null and the main effect of ToM models ( $P = 0.1137$  and  $P = 0.495$ , respectively).

#### Imaging Procedure and Behavioral Data Acquisition

Prior to the scanning procedure, a training session outside the scanner was performed. Participants listened to



**Figure 1.**

One example trial. The trial began with a fixation cross, then a fixation point was shown during the whole story (approximately 2 min). Next, the question screen appeared for 5 s and after this the answer screen (which still included the question) was shown for maximally 3 s. After that, the screens of second question and answer appeared with the same durations as the first question and answer screens.

two stories and answered two questions subsequent to each story. The practice stimuli were not part of the experimental stimuli. In the scanner, the participants listened to 20 stories and answered 40 questions (two after each story). Participants heard the stories through MRI compatible earphones. Sound quality and loudness was optimized in the scanner before starting the experiment. The order of the stories was assigned randomly and was different for each participant, to avoid sequence effects. The stories were divided into four blocks of five stories each. After each block, the participant had a break of 45 s. During the break, the participants saw the visual message "Short break!" in the middle of the screen, while the scanner was still running.

One story trial consisted of the following events: first a fixation cross was shown in the middle of the screen for 500 ms before the story started. The cross was then

replaced by a fixation point and at the same time the story started. The duration of the story was approximately 2 min. After the story there was a jitter between 1.5 and 4.5 s (duration assigned randomly), after which the first question was presented visually. The question was presented all at once, centered and toward the top of the screen for 5 s. After that, the possible answers appeared toward the bottom of the screen, clearly separated from each other; each answer began with an index letter (a) always on the left, and (b) always on the right side of the screen below the question (see Fig. 1 for a graphical representation of the question and answer screens). The possible answers stayed on the screen until participants made their decision; however, they disappeared if participants took longer than 3 s to respond (duration pretested to ensure a natural pace of the experiment). Participants gave their answers by pressing the left or right button on a

button box, which was fixed to their left leg, with their left middle or index finger accordingly. The left hand was chosen as a response hand to minimize left hemispheric artefacts which could overlap with linguistic processing [Callan et al. 2004]. The position of the correct answer was counterbalanced. Presentation of stimuli was time-jittered between story and questions and also between first and second question. All visual stimuli (cross, fixation point, questions, and answers) were presented in dark gray on light gray background. Figure 1 shows a graphical representation of an example trial. The procedure was implemented and presented with the software package Presentation (Neuro-behavioral Systems, San Francisco, CA).

### Behavioral Data Analyses

For the behavioral data analyses, we used R statistical software [Team, 2014] and the lme4 package [Bates et al., 2014]. For both responses and reaction times (RTs), we calculated models with fixed factors of condition and question order and random factors story and subject. The first question always referred to the first manipulation in the story (irrespective of the condition) and the second question always tested the information of the second manipulation of the story (also irrespective of the condition). We used logistic regression (because both dependent and independent variables were categorical) in combination with the maximal random effects structure [random slopes and intercept per condition and answer order for story and subject; see Barr et al., 2013] for the response analyses (R function glmer). Due to convergence problems in the models with the maximal random effects structure in the RTs, we included in the models (calculated with the R function lmer) the most complex random effects structure that reached convergence (random slopes and intercept per condition for story and subject). To assess the effects of the different factors on the response times and responses, we used a forward model selection procedure within which we used likelihood ratio tests to compare a base model including only an intercept with successively more complex models (function anova in R).

### fMRI Data Acquisition

During the MR-session, a series of echo-planar-images was gathered to record the time course of the subjects' brain activity. Measurements were performed on a 3 Tesla MRI system (Trio, A Tim System 3T, Siemens, Erlangen, Germany) with a 12 channel head matrix receive coil. Functional images were acquired using a T2\* weighted single shot echo planar imaging (EPI) sequence: parallel imaging factor of 2 (GRAPPA), TE = 25 ms, TR = 1,450 ms, flip angle 90°, slice thickness 4.0 mm and 0.6 mm gap, matrix 64 × 64, field of view = 224 × 224 mm, in-plane resolution 3.5 × 3.5 mm<sup>2</sup>, bandwidth 2,232 Hz/pixel, EPI factor of 64, and an echo spacing of 0.53 ms. Transversal

slices oriented to the AC–PC line were gathered in ascending order.

The initial five images were removed from the analyses to avoid saturation and stabilization effects. Head movements of the participants were minimized using foam paddings.

A whole head T1 weighted dataset was acquired with a 3d MPRage sequence (parallel imaging factor of 2 (GRAPPA), TE = 2.26 ms, TR = 1,900 ms, flip angle 9°, 1 mm isometric resolution, 176 sagittal slices, 256 × 256 matrix).

### fMRI Data Analyses

All analyses for the fMRI data were calculated in SPM8 (Wellcome Trust Centre for Neuroimaging), implemented in MATLAB (Mathworks, Sherborn, MA).

A slice time correction (to the 15th slice) was performed first. Then, images were realigned to the first image to correct for head movement artefacts. We normalized the volumes into standard stereotaxic anatomical Montreal neurological institute (MNI) space using the transformation matrix calculated from the first EPI scan of each subject and the EPI template. On the normalized data (resliced voxel size 2 mm<sup>3</sup>), we applied an 8 mm full-width-at-half-maximum Gaussian smoothing kernel to compensate for intersubject anatomical variation.

For the single-subject analysis, the design matrix for each subject was created individually, based on the log files from the fMRI-session, because each participant heard the stories in a different order. We modeled im-TOM and im-NONTOM conditions in seconds (mean duration of event = 4,326 ms, standard deviation = 1,525 ms). As critical events we modeled one sentence from each passage: for im-TOM it was the sentence in which the protagonist had a false belief and for im-NONTOM it was a length-matched sentence from the control passage. The events of the previous examples were as follows:

#### Story IA

- *She thought that maybe she had forgotten the food in the car.*<sub>im-TOM</sub>
- *took a whole series of pictures from all different perspectives.*<sub>im-NONTOM</sub>

#### Story IB

- *With delight she ate it and drank a few sips of apple juice.*<sub>im-NONTOM</sub>
- *The couple was looking for the camera without success.*<sub>im-TOM</sub>

For ex-TOM and ex-NONTOM conditions, we modeled the question (5 s) and answer (RTs) trial together as critical events (mean duration: 7.426 s, sd: 1.030 s); these events did not involve the motor response. The two trials for which

**TABLE II. Mean percentage of correct, incorrect, and missed responses per condition**

Answer	ex-TOM	ex-NONTOM
Correct	92	88
Incorrect	7	11
Not answered	1	2

there was no response were not modeled. As factors of no interest we modeled separately: the rest of the stories (excluding the im-TOM and im-NONTOM parts), the button presses (motor responses) and the jitters before each question and story. Our baseline consisted of the three 45 s pauses between blocks. To remove movement artefacts for each individual session the realignment parameters were entered as multiple regressors in the first-level analysis.

On the group-level analysis, we modeled two  $T$ -contrasts between the following first-level conditions: (a) im-TOM versus im-NONTOM and (b) ex-TOM versus ex-NONTOM. Brain activations were plotted on the anatomical MRIcron (<http://www.mccauslandcenter.sc.edu/mricro/mricron/>) high resolution template (the Colin brain). We used the cluster extent thresholding algorithm by [Slotnick et al., 2003], which implements a FWE correction using a Monte Carlo simulation approach, to correct for multiple comparisons. We set the desired correction threshold for multiple comparisons to  $P < 0.05$  and the assumed voxel type I error to  $P < 0.001$ ; after 10,000 iterations our cluster extend threshold was estimated at 48 voxels. For all fMRI results reported for *implicit* (im-TOM vs. im-NONTOM) and *explicit* (ex-TOM vs. ex-NONTOM) contrasts (and the reverse contrasts), we used a whole brain analysis and used an individual voxel threshold of  $P < 0.001$  with a cluster extend threshold of 48 voxels.

### Contrasts of Interest

In addition to the contrasts im-TOM versus im-NONTOM and ex-TOM versus ex-NONTOM, which test *implicit* and *explicit* ToM separately, we were interested in the common regions activated for both contrasts. Therefore, we performed a conjunction analysis using statistical parametric maps (SPMs) of the minimum  $T$ -statistic over the previous contrasts (im-TOM vs. im-NONTOM and ex-TOM vs. ex-NONTOM). Inference was based on  $P$ -values adjusted for the search volume using random field theory [for details on the exact procedure see Friston et al., 2005]. The SPM8 algorithm for conjunction (testing the conjunction null hypothesis as recommended in Nichols et al. [2005]) assumes that the  $P$ -value of the conjunction is the square root of the  $P$ -value of the involved contrasts. We set the  $P$ -value for the conjunction to 0.05, thereby implicitly thresholding each individual contrast at  $P < 0.0025$ . We further corrected the conjunction results for multiple comparisons by setting a cluster threshold on 120 voxels, as


estimated by the Slotnick et al. [2003] algorithm after 10,000 iterations (assumed voxel type I error to  $P < 0.05$ ).

## RESULTS

Subjects achieved a mean of 90% (sd = 5.61) correctness in the answers. The mean percentages of correct, incorrect, and missed responses per condition are presented in Table II. In a logistic mixed effects models analysis (see “Behavioural data analyses” for details), we found no significant main effect of condition (ex-TOM vs. ex-NONTOM) for the responses (correct, incorrect, and missed response). A comparison of the logistic mixed effects models (using the function `anova` in R) showed no significant improvement of model fit ( $P$ 's > 0.7) of the main effects and interaction models in comparison to the null model (which included only an intercept in addition to the maximal random effects structure). None of the single main effects models (only main effect of condition or only main effect of question order) improved model fit compared with the null model ( $P$ 's > 0.5).


We also analyzed the RTs with mixed effects models (for the results see Fig. 2). We found no significant improvement of the model fit ( $P = 0.26$ ) when comparing the main effects model of condition and question order to the null model (which included only the intercept and random effects). None of the single main effects of condition or question order improved the model in comparison to both the null model and the main effects of condition and question order models ( $P$ 's > 0.1).

In the fMRI analyses, we found significant activation for im-TOM versus im-NONTOM in the bilateral AG, left MTG, right middle temporal pole, bilateral PCUN, left

**Figure 2.**

RTs for the first and second question after each story for the ex-TOM and ex-NONTOM conditions. Abbreviations: cor, correct response; inc, incorrect response. [Color figure can be viewed in the online issue, which is available at [wileyonlinelibrary.com](http://wileyonlinelibrary.com).]





**Figure 3.**

The contrast im-TOM versus im-NONTOM is depicted in red and ex-TOM versus ex-NONTOM in yellow, superimposed on a high resolution brain template (the Colin brain) of the MRIcron software ( $P < 0.001$ , cluster extend threshold of 48 voxels, Monte Carlo corrected). Abbreviations: AG, angular gyrus;

SmFG, superior medial frontal gyrus; mPFC, medial prefrontal cortex; MTG, middle temporal gyrus; MFG, middle frontal gyrus; CE, cerebellum; PCUN, precuneus. Coordinates: a) MNI:  $-56, -58, 32$ , b) MNI:  $-60, -24, -10$ , c) MNI:  $46, 22, 38$ , d) MNI:  $-8, -48, 44$ .

cerebellum (CE) (crus 2, VIII), right CE (IX), as well as bilateral MFG and IFG. For ex-TOM versus ex-NONTOM, we found significant activation in the left AG. For the reverse implicit contrast im-NONTOM versus im-TOM, we found activation in the right MFG and left posterior central gyrus ( $P < 0.001$ , cluster extend threshold of 48 voxels). For the reverse explicit contrast ex-NONTOM versus ex-TOM, we found activation in the right hippocampus (also  $P < 0.001$ , cluster extend threshold of 48 voxels).

An overview of the results for the contrasts of interest with coordinates (MNI),  $T$ -scores and cluster sizes is presented in Table III (see Fig. 3 for the localization of the results on the brain template).

Table IV shows the suprathreshold clusters activated for the conjunction. We found suprathreshold activation for the conjunction in the left AG, MTG, MFG, bilateral IFG as well as in the CE (crus 1) (for the localization of the effects see Fig. 4).

**TABLE III. False belief activation peaks with their local maxima coordinates for the contrasts im-TOM versus im-NONTOM and ex-TOM versus ex-NONTOM ( $P < 0.001$ , cluster extend threshold 48 voxels, Monte Carlo corrected)**

Contrast	Anatomical region	H	MNI coordinates	$T$	Cluster size
im-TOM versus im-NONTOM	Angular gyrus (AG)	L	-58 -64 30	7.64	2,121
	Cerebellum (CE) - Crus 2	L	-24 -82 -34	7.19	2,853
	Angular gyrus (AG)	R	56 -60 30	7.1	2,812
	Precuneus (PCUN)	L	-8 -48 44	6.23	860
	Middle frontal gyrus (MFG)	L	-22 52 24	5.89	1,620
	Superior medial frontal gyrus (SmFG) - mPFC	L	-4 48 38	5.68	1,620
	Middle frontal gyrus (MFG)	L	-40 12 46	5.28	806
	Middle temporal pole (MTP)	R	50 8 -28	5.03	257
	Middle frontal gyrus (MFG)	R	24 24 44	4.97	236
	Middle temporal gyrus (MTG)	L	-60 -24 -10	4.71	410
	Precuneus (PCUN)	R	12 -50 40	4.4	860
	Superior frontal gyrus (SFG)	R	26 58 18	4.4	88
	Middle frontal gyrus (MFG)	R	46 22 38	4.3	217
	Superior frontal gyrus (SFG)	L	-32 52 0	4.3	52
	Inferior frontal gyrus (IFG) - Pars triangularis	L	-56 26 6	4.11	297
	Inferior frontal gyrus (IFG) - Pars orbitalis	L	-50 28 -6	4	297
	ex-TOM versus ex-NONTOM	Angular gyrus (AG)	L	-56 -58 32	3.76

Coordinates (x, y, z) are listed in MNI atlas space (H, hemisphere; mPFC, medial prefrontal cortex).

The changes of the BOLD signal, as reflected in the contrast estimates (first principal component of the signal) in the peak voxel, in the left AG across all contrasts of interest revealed more activation for the im-TOM and ex-TOM in comparison to the im-NONTOM and ex-NONTOM condition, respectively (see Fig. 5). In the barplots of the contrast estimates for the implicit contrast (see Figs. 5 and 6, under the red label), it is the differences between im-TOM and im-NONTOM that are responsible for the *implicit* ToM activation. In the ex-TOM versus ex-NONTOM plots (see Fig. 5, under the yellow label) the differences between ex-TOM and ex-NONTOM are driving the suprathreshold activation. Please note that in both implicit and explicit contrasts (ex-TOM vs. ex-NONTOM under the yellow label and im-TOM vs. im-NONTOM under the red label) we have plotted all conditions of interest for the sake of com-

pleteness, even though only two are relevant for the activation of the contrast. For the conjunction contrast (see Figs. 5 and 6 under the green label), all four conditions contribute to the common activation of implicit and explicit contrasts by showing the same tendencies in the differences between TOM and NONTOM conditions: a left lateralized network comprising the AG, MTG, MFG, and IFG showed stronger increases in BOLD signal for im-TOM and ex-TOM in comparison to im-NONTOM and ex-NONTOM.


## DISCUSSION

In this study, participants listened to two-minute long stories with embedded *implicit* false belief and control passages. After each story, participants answered one false

**TABLE IV. Activation peaks with their local maxima coordinates for the conjunction of the contrasts im-TOM versus im-NONTOM and ex-TOM versus ex-NONTOM (threshold  $P < 0.05$ , cluster extend threshold 120 voxels, Monte Carlo corrected)**

Contrast	Anatomical region	H	MNI Coordinates	$T$	Cluster size
Conjunction	Angular gyrus (AG)	L	-56 -58 32	3.76	1,363
	Middle frontal gyrus (MFG)	L	-40 20 48	2.61	487
	Middle temporal gyrus (MTG)	L	-52 -36 -14	2.42	484
	Inferior frontal gyrus (IFG) - Pars triangularis	R	54 30 28	2.37	135
	Cerebellum (Crus 1)	R	28 -80 -32	2.33	293
	Inferior frontal gyrus (IFG)	L	-42 28 -12	2.31	139

Coordinates (x, y, z) are listed in MNI atlas space (H, hemisphere).



**Figure 4.**

The activations of the conjunction of the contrasts im-TOM versus im-NONTOM and ex-TOM versus ex-NONTOM are shown in green (conjunction voxel threshold of  $P < 0.05$  and cluster extend threshold of 120 voxels, Monte Carlo corrected). Abbreviations:

AG, angular gyrus; MTG, middle temporal gyrus; MFG, middle frontal gyrus; IFG, inferior frontal gyrus. Coordinates: a) MNI:  $-56, -58, 32$ , b) MNI:  $-52, -46, -14$ , c) MNI:  $-42, 28, -12$ .

belief and one control question (*explicit* false belief processing). Our main finding was that processing false beliefs *incidentally* in a rich and natural narrative context recruits ToM processing regions (AG, MTG, mPFC in the SmFG, PCUN)—known from the *explicit* ToM literature. For

*explicit* false belief processing, we could replicate previous results in our left AG findings. The conjunction analysis revealed a left lateralized network of the AG, MTG, MFG, and IFG as the common pattern activated during both *implicit* and *explicit* false belief processing.




Figure 5.

Bar plots of BOLD signal changes (contrast estimates) at the peak voxel of the left (LH) angular gyrus (AG) for implicit and explicit false belief processing as well as their conjunction. Threshold for the implicit and explicit contrasts:  $P < 0.001$  and cluster extend threshold of 48 voxels (Monte Carlo corrected), for the conjunction  $P < 0.05$  and cluster extend threshold of 120

voxels (Monte Carlo corrected). The error bars represent 90% confidence intervals. Please note that the peak voxel of the LH AG is the same for the explicit contrast and the conjunction, therefore, the barplots are also the same. [Color figure can be viewed in the online issue, which is available at [wileyonlinelibrary.com](http://wileyonlinelibrary.com).]

For *implicit* false belief processing (im-TOM vs. im-NONTOM), we found activation in the AG bilaterally. The AG is an anatomical subdivision of the posterior inferior parietal lobule, which is considered part of the so-called TPJ [Mar, 2011; Seghier, 2013]. TPJ activation has been previously found in several story-based ToM paradigms [Aichhorn et al., 2009; Fletcher et al., 1995; Gallagher et al., 2000; Kobayashi et al., 2006; Lee et al., 2011; Mitchell, 2008; Saxe and Kanwisher, 2003; Vogeley et al., 2001]. Especially the right TPJ is assumed to be heavily involved in mentalizing processes [Perner et al., 2006; Saxe and Kanwisher, 2003] throughout many different situations: answering an open question about visually presented and cartoon stories [Gallagher et al., 2000], answering in a multiple choice question after visually presented stories [Spengler et al., 2009], second-order false belief tasks [Kobayashi et al., 2006], false belief stories and desires also presented visually [Saxe and Kanwisher, 2003], reading TOM and SELF stories and covertly answering questions [Vogelely et al., 2001]. All of these previous studies did not investigate or model *implicit* false belief processing separately, although it was also measured during stimulation. The only study from the literature that attempted to capture *implicit* false belief processing was Aichhorn et al. [2009], in which the reading of false belief stories was modeled separately to answering the question in two different first level analyses. Although another instance of automatic ToM processing has been tested in trait judgements [Ma et al., 2012, 2011], the current study was the first to test *implicit* ToM in false belief processing (a) in listening and (b) embedded in a rich and natural linguistic context. As we found TPJ activation for *implicit* false belief processing in this novel setup, we conclude that the TPJ may not only be active in decision-based setups but that it may also be

recruited incidentally in implicit false belief tasks (such as mentalizing during narrative processing).

Our contribution adds to the existing literature on social cognition as part of language processing [Ferstl and Cramon, 2002; Ferstl et al., 2008; Mason and Just, 2009] by revealing ToM processing regions when people listen to stories including false belief situations. Our finding of SmFG (mPFC) in the implicit contrast and in the conjunction in particular is in line with Ferstl and Cramon [2002]’s findings of FMC activation for narrative coherence based on social cues. However, a more systematic study of the interplay between social and linguistic cues in narrative processing is needed to draw conclusions about how the two domains interact on a neurobiological level.

The TPJ is one of the regions of a ToM processing *network*. This network includes (among others) the following regions: mPFC, IFG, MFG, MTG, CE, and PCUN [Aichhorn et al., 2009; Gallagher et al., 2000; Kobayashi et al., 2006; Lee et al., 2011; Saxe and Kanwisher, 2003; Spengler et al., 2009]. In this study, we provided further support for this ToM network especially during *implicit* false belief processing when listening to narrative stories. Our findings specifically provide evidence for the claim that ToM processing regions are recruited spontaneously when false beliefs are embedded in natural context.

Parts of the “classical” ToM processing network have been claimed to be strongly involved in the default mode network (DMN). Especially the AG and the PCUN have repeatedly shown reliable activation during resting state fMRI experiments [Shehzad et al., 2009; Seghier, 2013; Utevsky et al. 2014]. In connection to our findings and given our experimental setup, our results strengthen the claims for these areas (AG, PCUN) to be involved in (automatic) belief tracking, as explained by van der Wel et al.




Figure 6.

Bar plots of BOLD signal changes (contrast estimates) at the peak voxel of each cluster for implicit false belief processing and the conjunction of implicit and explicit contrasts. Abbreviations: RH, right hemisphere; LH, left hemisphere; AG, angular gyrus; SmFG, superior medial frontal gyrus; mPFC, medial prefrontal cortex; MTG, middle temporal gyrus; MTP, middle temporal pole; MFG, middle frontal gyrus; PCUN, precuneus. Please note that we have placed the plots for precuneus, AG and IFG next to each other for illustration reasons and with absolutely no

intention of denoting that they are overlapping regions. For explicit and implicit contrasts: individual voxel threshold of  $P < 0.001$  and cluster extend threshold of 48 voxels (Monte Carlo corrected). For the conjunction: individual voxel threshold of  $P < 0.05$  and cluster extend threshold of 120 voxels (Monte Carlo corrected). The error bars represent 90% confidence intervals. [Color figure can be viewed in the online issue, which is available at [wileyonlinelibrary.com](http://wileyonlinelibrary.com).]

[2014]. A similar concept to the DMN is the human “reorienting” system [Corbetta et al., 2008], which distinguishes between a dorsal and a ventral attention system. Even though these systems are formulated in connection to visual cognition we would like to attempt a connection to auditory processing. In this view, the AG and the PCUN belong to the dorsal attention system and support top-down attentional control. Conversely, the ventral attention network includes the MFG, the ventral frontal cortex, and the inferior parts of the TPJ (posterior STG) and is responsible for bottom-up reorientation. This framework could offer an alternative interpretation for our results in terms of attentional reorienting [for example as in Rothmayr et al., 2011].

Our *explicit* false belief contrast showed suprathreshold activation only in the left AG. As there has been only one study which aimed to disentangle implicit and explicit false belief processing [Aichhorn et al., 2009] we can attempt a comparison of our explicit contrast results with their findings at *Time Point 2: Question*. Their main contrast was false belief (FB) versus false photograph (PH). As this contrast was used to define the regions of interest (ROIs) for the remaining contrasts, a fact that highlights the importance of this contrast in comparison to the remaining contrasts of interest, we can compare it to our explicit contrast (ex-TOM vs. ex-NONTOM). Their results also showed activation in the left AG, as in our manipulation, but they included additional areas such as the anterior

MTG, temporal pole, IFG, PCUN, and mPFC. These were precisely the areas that showed signal changes for our implicit contrast, while they did not survive the threshold for the explicit contrast. However, the conjunction pointed to the same clusters (left lateralized AG, MTG, IFG, MFG, CE) as commonly activated areas for implicit and explicit contrasts. This suggested that the explicit contrast mask extended to the other ToM processing areas but the effect sizes were not strong enough to survive the  $P < 0.001$  threshold. Moreover, methodological details of design and modeling might also explain the differences in the findings of our explicit contrast to the FB versus PH contrast of Aichhorn et al. [2009]. The design differed in four respects: the presentation modality (listening vs. reading stories); the length of the stories and, therefore, the amount of provided context (23 sentences vs. 2 sentences); the first level analysis (modeled critical events in one first level analysis vs. modeled critical events in two different first level analyses); the second level analysis (whole-brain vs. combined whole-brain and ROI analysis). Finally, due to the absence of jitters (else, the presence of a constant ISI of 2 s) between the modeled events in Aichhorn et al. [2009] it is unclear how reliably the contributions of each conditions to the overlapping BOLD response could have been estimated. In contrast to this, in our study of narrative stories, the context of the story created a natural jitter between the events of the implicit contrast. Also, for the explicit contrast we optimized our design for efficient modeling of the hemodynamic response by introducing ISIs (jitters) of random duration (1.5–4.5 s) between story and question as well as between the first and second question.

Despite the single cluster activation for the explicit contrast, the pattern of left lateralized activation revealed by the conjunction of the two contrasts of interest is in line with the ToM processing network known from the imaging literature. This left lateralization might be connected to the nature of the stimuli, which comprised of false belief situations presented auditorily and embedded in rich linguistic context.

We have to acknowledge two minor limitations in our paradigm. First, it might be the case that the questions (*explicit* ToM) pointed participants toward what was tested in the study. However, these questions comprised only half of the total number of asked questions. There was always a control question in addition to the false belief question after each story and the order of the two questions was counterbalanced across the whole experiment. Moreover, the formulations of the false belief questions were very variable, so that we did not repeat the words “think,” “opinion,” and so forth, too often during the experiment. In addition, the stories include a wide variety of situations of which the experimental manipulations constitute not even half of the total duration of one story. Thus, participants can be keeping track of more things at different times during the story: for example, they might have followed the path that the hikers took (in the hiking story mentioned in the Stimuli section).

The second limitation is related to the order of the presented conditions; in our design the *implicit* condition always preceded the *explicit* condition, which means that the two conditions were not totally independent in time. This might have caused more common activation than if these two conditions were measured completely independently. However, other experimental solutions, such as asking the false belief questions after the fMRI scan session, would also be suboptimal, as remembering the plots of 20 stories would be very demanding and lead to a high number of incorrect answers. We chose to ask two questions after each story not only to test false belief processing in *explicit* mode, but also to make sure that the participants were alert and paying attention to the stories. As two of our participants almost fell asleep repeatedly during scanning (these datasets were excluded from the analyses), we have to accept that it might be difficult to lay in the MRI scanner for 1 h listening to stories without falling asleep, despite the noisy environment of the scanning procedure and the task.

## CONCLUSIONS

In this study, we showed that the “classical” ToM processing network (e.g., TPJ, mPFC, MTG, PCUN) is activated during *implicit* false belief processing, while listening to short stories. We were the first to reliably disentangle *implicit* and *explicit* ToM processing by modeling short false-belief sentences within longer stories and by separating them further from the explicit false belief task. Our study, therefore, offers insights to the neural underpinnings of auditory language processing and social competence, integral parts of human nature long before the use of written communication.

## ACKNOWLEDGMENTS

The authors would like to thank Karen Henrich and Marie-Josephine Rocholl for their help in recording the stories as well as Jens Sommer, Mechthild Wallnig, and Rita Werner for their help in data collection.

## REFERENCES

- Aichhorn M, Perner J, Weiss B, Kronbichler M, Staffen W, Ladurner G (2009): Temporo-parietal junction activity in theory-of-mind tasks: Falseness, beliefs, or attention. *J Cogn Neurosci* 21:1179–1192. doi:10.1162/jocn.2009.21082
- Baron-Cohen S, Wheelwright S, Hill J, Raste Y, Plumb I (2001): The “Reading the Mind in the Eyes” test revised version: A study with normal adults, and adults with Asperger syndrome or high-functioning autism. *J Child Psychol Psychiatry* 42:241–251.
- Barr DJ, Levy R, Scheepers C, Tily HJ (2013): Random effects structure for confirmatory hypothesis testing: Keep it maximal. *J Mem Lang* 68:255–278.
- Bates D, Mächler M, Bolker B, Walker S (2014): Fitting linear mixed-effects models using lme4. *arXiv preprint arXiv:1406.5823*.

- Bornkessel-Schlesewsky I, Schlesewsky M, Small SL, Rauschecker JP (2015): Neurobiological roots of language in primate audition: common computational properties. *Trends Cogn Sci* 19: 142–150.
- Callan DE, Jones JA, Callan AM, Akahane-Yamada R (2004): Phonetic perceptual identification by native-and second-language speakers differentially activates brain regions involved with acoustic phonetic processing and those involved with articulatory-auditory/orosensory internal models. *NeuroImage* 22: 1182–1194.
- Callejas A, Shulman GL, Corbetta M (2011): False Belief vs. False Photographs: A Test of Theory of Mind or Working Memory? *Front Psychol* 2:316.
- Corbetta M, Patel G, Shulman GL (2008): The reorienting system of the human brain: From environment to theory of mind. *Neuron* 58:306–324. doi:10.1016/j.neuron.2008.04.017
- Dehaene S, Cohen L (2011): The unique role of the visual word form area in reading. *Trends Cogn Sci* 15:254–262. doi:10.1016/j.tics.2011.04.003
- Ferstl EC, von Cramon DY (2002): What does the frontomedian cortex contribute to language processing: Coherence or theory of mind? *NeuroImage* 17:1599–1612. doi:10.1006/nimg.2002.1247
- Ferstl EC, Neumann J, Bogler C, von Cramon DY (2008): The extended language network: A meta-analysis of neuroimaging studies on text comprehension. *Hum Brain Mapp* 29:581–593. doi:10.1002/hbm.20422
- Fletcher PC, Happé F, Frith U, Baker SC, Dolan RJ, Frackowiak RS, Frith CD (1995): Other minds in the brain: A functional imaging study of “theory of mind” in story comprehension. *Cognition* 57:109–128. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/8556839>
- Friston KJ, Penny WD, Glaser DE (2005): Conjunction revisited. *NeuroImage* 25:661–667. doi:10.1016/j.neuroimage.2005.01.013
- Frith C, Frith U (1999): Interacting minds: A biological basis. *Science* 286:1692–1695. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10576727> <http://www.sciencemag.org/content/286/5445/1692.short>
- Frith U, Frith C (2010): The social brain: Allowing humans to boldly go where no other species has been. *Philos Trans R Soc Lond B Biol Sci* 365:165–176. doi:10.1098/rstb.2009.0160
- Gallagher HL, Happé F, Brunswick N, Fletcher PC, Frith U, Frith CD (2000): Reading the mind in cartoons and stories: An fMRI study of ‘theory of mind’ in verbal and nonverbal tasks. *Neuropsychologia* 38:11–21. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10617288>
- Goldman AI (2012): Theory of mind. *The Oxford handbook of philosophy of cognitive science*, 402–424.
- Kestemont J, Vandekerckhove M, Ma N, Van Hoek N, Van Overwalle F (2013): Situation and person attributions under spontaneous and intentional instructions: An fMRI study. *Soc Cogn Affect Neurosci* 8:481–493. doi:10.1093/scan/nss022
- Kestemont J, Ma N, Baetens K, Clément N, Van Overwalle F, Vandekerckhove M (2015): Neural correlates of attributing causes to the self, another person and the situation. *Soc Cogn Affect Neurosci* 10:114–121.
- Kobayashi C, Glover GH, Temple E (2006): Cultural and linguistic influence on neural bases of “Theory of Mind”: An fMRI study with Japanese bilinguals. *Brain Lang* 98:210–220. doi:10.1016/j.bandl.2006.04.013
- Kuperberg GR, Paczynski M, Ditman T (2011): Establishing causal coherence across sentences: An ERP study. *J Cogn Neurosci* 23:1230–1246.
- Lee J, Quintana J, Nori P, Green MF (2011): Theory of mind in schizophrenia: Exploring neural mechanisms of belief attribution. *Soc Neurosci* 6:569–581.
- Ma N, Vandekerckhove M, Van Overwalle F, Seurinck R, Fias W (2011): Spontaneous and intentional trait inferences recruit a common mentalising network to a different degree: Spontaneous inferences activate only its core areas. *Soc Neurosci* 6:123–138. doi:10.1080/17470919.2010.485884
- Ma N, Vandekerckhove M, Baetens K, Overwalle FV, Seurinck R, Fias W (2012): Inconsistencies in spontaneous and intentional trait inferences. *Soc Cogn Affect Neurosci* 7:937–950. doi:10.1093/scan/nsr064
- Mar RA (2011): The neural bases of social cognition and story comprehension. *Ann Rev Psychol* 62:103–134. doi:10.1146/annurev-psych-120709-145406
- Mason RA, Just MA (2009): The role of the theory-of-mind cortical network in the comprehension of narratives. *Linguist Lang Compass*, 3:157–174. doi:10.1111/j.1749-818X.2008.00122.x
- Mitchell JP (2008): Activity in right temporo-parietal junction is not selective for theory-of-mind. *Cereb Cortex (New York, N.Y.: 1991)*, 18:262–271. doi:10.1093/cercor/bhm051
- Nichols T, Brett M, Andersson J, Wager T, Poline JB (2005): Valid conjunction inference with the minimum statistic. *NeuroImage* 25:653–660. doi:10.1016/j.neuroimage.2004.12.005
- Northoff G, Heinzel A, Greck M, de Bermpohl F, Dobrowolny H, Panksepp J (2006): Self-referential processing in our brain: A meta-analysis of imaging studies on the self. *NeuroImage* 31: 440–457. doi:10.1016/j.neuroimage.2005.12.002
- Perner J, Aichhorn M, Kronbichler M, Staffen W, Ladurner G (2006): Thinking of mental and other representations: The roles of left and right temporo-parietal junction. *Soc Neurosci* 1:245–258. doi:10.1080/17470910600989896
- Premack D, Woodruff G (1978): Does the chimpanzee have a theory of mind? *Behav Brain Sci* 1:515–526.
- Rothmayr C, Sodian B, Hajak G, Döhnell K, Meinhardt J, Sommer M (2011): Common and distinct neural networks for false-belief reasoning and inhibitory control. *Neuroimage* 56:1705–1713.
- Saxe R, Kanwisher N (2003): People thinking about thinking people. The role of the temporo-parietal junction in “theory of mind”. *NeuroImage* 19:1835–1842. doi:10.1016/S1053-8119(03)00230-1
- Schaafsma SM, Pfaff DW, Spunt RP, Adolphs R (2015): Deconstructing and reconstructing theory of mind. *Trends Cogn Sci* 19:65–72.
- Schlaffke L, Lissek S, Lenz M, Juckel G, Schultz T, Tegenthoff M, Tobias Schmidt-Wilcke, Brüne, M. (2014). Shared and non-shared neural networks of cognitive and affective theory-of-mind: A neuroimaging study using cartoon picture stories. *Hum Brain Mapp* 36:29–39. doi:10.1002/hbm.22610
- Schurz M, Radua J, Aichhorn M, Richlan F, Perner J (2014): Fractionating theory of mind: A meta-analysis of functional brain imaging studies. *Neurosci Biobehav Rev* 42C:9–34. doi:10.1016/j.neubiorev.2014.01.009
- Seghier ML (2013): The angular gyrus multiple functions and multiple subdivisions. *Neuroscientist* 19:43–61.
- Shehzad Z, Kelly AC, Reiss PT, Gee DG, Gotimer K, Uddin LQ, Lee SH, Margulies DS, Roy AK, Biswal BB, Petkova E, Castellanos FX, Milham MP (2009): The resting brain: Unconstrained yet reliable. *Cereb Cortex* 19:2209–2229.
- Slotnick SD, Moo LR, Segal JB, Hart J (2003): Distinct prefrontal cortex activity associated with item memory and source

- memory for visual shapes. *Brain Res Cogn Brain Res* 17:75–82. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12763194>
- Sommer M, Döhnel K, Sodian B, Meinhardt J, Thoermer C, Hajak G (2007): Neural correlates of true and false belief reasoning. *Neuroimage* 35:1378–1384.
- Spengler S, Cramon DY, von Brass M (2009): Control of shared representations relies on key processes involved in mental state attribution. *Hum Brain Mapp* 30:3704–3718. doi:10.1002/hbm.20800
- Spreng RN, Mar RA, Kim ASN (2009): The common neural basis of autobiographical memory, prospection, navigation, theory of mind, and the default mode: A quantitative meta-analysis. *J Cogn Neurosci* 21:489–510. doi:10.1162/jocn.2008.21029
- Team RC (2014). R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria, 2012.
- Utevsky AV, Smith DV, Huettel SA (2014): Precuneus is a functional core of the default-mode network. *J Neurosci* 34:932–940.
- Van Overwalle F (2009): Social cognition and the brain: A meta-analysis. *Hum Brain Mapp* 30:829–858. doi:10.1002/hbm.20547
- Vogeley K, Bussfeld P, Newen A, Herrmann S, Happé F, Falkai P, Maier W, Shah NJ, Fink GR, Zilles K (2001): Mind reading: Neural mechanisms of theory of mind and self-perspective. *NeuroImage* 14:170–181. doi:10.1006/nimg.2001.0789
- van der Wel RP, Sebanz N, Knoblich G (2014): Do people automatically track others' beliefs? Evidence from a continuous measure. *Cognition* 130:128–133. doi:10.1016/j.cognition.2013.10.004