



Published in final edited form as:

Drug Alcohol Depend. 2019 November 01; 204: 107512. doi:10.1016/j.drugalcdep.2019.05.032.

Adolescents' Awareness of the Nicotine Strength and E-cigarette Status of JUUL E-cigarettes

Meghan E. Morean, Ph.D^{*,a,b}, Krysten W. Bold, Ph.D^b, Grace Kong, Ph.D^b, Ralitza Gueorguieva^c, Deepa Camenga, M.D^d, Patricia Simon, Ph.D^e, Asti Jackson, Ph.D^b, Dana A. Cavallo, Ph.D^b, Suchitra Krishnan-Sarin, Ph.D^b

^aOberlin College, 120 West Lorain Street, Oberlin, OH, 44074, USA

^bYale University School of Medicine, 34 Park Street, New Haven, CT, 06519, USA

^cYale University School of Public Health, 60 College St, New Haven, CT, 06520, USA

^dYale University School of Medicine, 464 Congress Avenue (Suite 260), New Haven, CT, 06519, USA

^eYale University School of Medicine, 389 Whitney Avenue New Haven, CT, 06511, USA

Abstract

Background: JUUL e-cigarettes are popular among youth. However, it is unknown whether adolescents understand that 5% JUUL pods contain a high nicotine concentration or consider JUULs to be e-cigarettes.

Method: 3170 students from 4 Connecticut high schools completed a school-based survey (May-October 2018). Students reported on lifetime and past-month JUUL use and perceived JUUL nicotine strength (low/medium/high/don't know) when no information about nicotine concentration was provided and, subsequently, when informed JUULs contain 5% nicotine. Students reported whether they believe JUULs are e-cigarettes (no/yes/don't know).

Results: Students were never JUUL users (56.6%), ever users (13.2%), and past-month users (30.2%). When no information was provided, students reported that JUULs contain low (10.5%), medium (26.9%), or high nicotine levels (31.1%); 31.4% did not know. When informed JUULs contain 5% nicotine, students were more likely to believe JUUL's nicotine strength was low

*Corresponding Author: Meghan E. Morean, Department of Psychology, Oberlin College, 120 West Lorain Street, Oberlin, OH, 44074, USA. meghan.morean@gmail.com.

Contributors

Drs. Morean, Bold, Kong, Camenga, Simon, Jackson, Cavallo, and Krishnan-Sarin collaborated on the development of the original study and conceptualized the current research question. Drs. Morean and Gueorguieva completed the statistical analyses. Dr. Morean drafted the original version of the manuscript and all other authors provided substantive feedback on further drafts. All authors approved the submitted manuscript.

Conflict of Interest

No conflicts declared.

Declarations of Interest: None

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

(29.5%) or medium (29.3%) than high (21.3%) and less likely to report not knowing (19.9%). 39% of students believed JUULs are not e-cigarettes or did not know.

Discussion: Most students were unaware of JUUL’s high nicotine concentration, with more believing that JUULs contain low or medium nicotine concentrations when informed JUULs contain 5% nicotine. Thus, youth may misinterpret the nicotine concentration printed on JUUL pod packaging, raising concerns about inadvertent exposure to high nicotine levels and dependence risk. Further, 39% of adolescents believed JUULs are not e-cigarettes or were unsure. Regulatory efforts are needed to establish understandable nicotine concentration labels, require products to be labeled accordingly, and clarify what products constitute e-cigarettes.

Keywords

adolescent; electronic cigarette; e-cigarette; vaping; JUUL; nicotine

1. INTRODUCTION

JUUL e-cigarettes resemble USB flash drives and use disposable pods containing nicotine salt and flavors (e.g., tobacco, mango). JUUL is the most popular American e-cigarette brand (Carver, 2018), and, at the time the study was conducted, its pods uniformly contained among the highest nicotine concentrations commercially available (5% or 59mg/ml; JUUL Labs Inc, 2018). JUUL Labs Inc. (2018) claims that JUULs are a “satisfying alternative to cigarettes” for adult smokers. However, JUULs are popular among youth (Huang et al., 2019; Kavuluru et al., 2019; Willett et al., 2019), with a recent study showing that JUULs were the most popular vaping device used by high school students (Krishnan-Sarin et al., 2019) JUULs’ popularity combined with their high nicotine content raises concerns about exposing youth to the deleterious effects of nicotine on the developing brain and addiction risk (Morean et al., 2018; U.S. Department of Health and Human Services [USDHHS], 2012).

Importantly, youth often are unaware of the nicotine content of e-liquids (Morean et al., 2016), and most youth are unaware that JUULs always contain nicotine (Willett et al., 2019). Beginning in August 2018, the US Food and Drug Administration (FDA) mandated that all e-liquid packaging display a warning (“This product contains nicotine. Nicotine is an addictive chemical”), which may improve youths’ knowledge of whether a product contains any nicotine. However, this warning does not address nicotine strength. Although no reporting standard exists, e-liquid nicotine concentrations typically are presented as milligrams/milliliter or as percent nicotine. At the time of the study, all JUUL pod packages were labeled as “5% strength.” While “5% strength” refers to the fact that pods contain 5% nicotine, the word “nicotine” was absent from the label (although it has since been added). It is important to understand how youth interpret this information, as the labeling of JUUL nicotine pods could lead to misunderstanding. For example, 5% sounds like a small amount, so youth may incorrectly believe that JUUL pods contain a low nicotine concentration.

Further, youth commonly use the term “JUULing” to refer to JUUL use instead of terms more commonly used to refer to e-cigarette use like “vaping,” raising questions about whether youth view JUULs as e-cigarettes (Huang et al., 2019; Kavuluru et al., 2019).

Understanding whether youth consider JUULs to be e-cigarettes is critical to informing youth-focused regulatory and prevention efforts (FDA, 2018). If youth do not think JUULs are e-cigarettes, this may represent an additional pathway through which youth inadvertently expose themselves to vaping-related harms and suggests that products should be labeled clearly as e-cigarettes.

We address these aforementioned issues by examining adolescents' perceptions of JUUL pods' nicotine strength when no information about nicotine concentration explicitly was provided versus when adolescents were informed JUUL pods contain 5% nicotine. We included the no information condition to assess adolescents' baseline knowledge of JUULs' nicotine strength. We showed adolescents a picture of JUUL pod packaging and highlighted that "JUULs contain 5% nicotine" to understand how adolescents perceive the nicotine concentration label. Finally, we evaluated whether adolescents view JUULs as e-cigarettes.

2. METHODS

2.1 Participants

All students from 4 Connecticut high schools who were in attendance on the dates of survey administration (May-October 2018) were invited to participate ($N = 3730$). 85.0% of students ($N = 3170$) completed the anonymous, computerized survey (52.4% female, 60.4% White; $15.87[SD=1.29]$ years old).

2.2 Procedures

The Institutional Review Board of Yale University, the school boards, and the participating schools approved the study. Passive parental permission was obtained prior to survey administration. Three parents refused their child's participation. All participants were informed that participation was voluntary. Completing the survey indicated participants' consent/assent.

2.3 Measures

Participants reported on lifetime and past-30-day JUUL use (Table 1). Participants also reported on perceived nicotine strength of JUUL pods (low, medium, high, I don't know), first when no information about nicotine concentration was provided, and, subsequently, when informed that JUULs contain "5% nicotine." The nicotine strength questions were presented separately, and participants could not use the "back/forward" buttons to view or change previous responses. Finally, participants reported whether they believe JUULs are e-cigarettes (no, yes, I don't know).

2.4 Data Analyses

We categorized students as never JUUL users, ever JUUL users (tried a JUUL but no past-month use), and past-30-day (past-month) JUUL users. We then ran descriptive statistics within the total sample and by JUUL use status. To evaluate whether there were significant differences in perceived nicotine strength when no information was provided versus when 5% nicotine was specified, we ran a multinomial Generalized Estimating Equation (GEE) model to assess the effects of the within-subjects factor "information" (no information vs.

5% content) and the between-subjects factor “JUUL use status” (never, ever, past-month) on perceived nicotine strength (low, medium, high, don’t know). Tests of the interaction of information by JUUL use status and of the main effects were performed. Significant results were followed by tests of effect slices (i.e., assessments of the effect of information for each type of JUUL user) and pairwise comparisons. School (4 levels) was entered as a fixed effect in the model to control for potential differences by school. Significance was set at 0.05. Odds ratios and 95% confidence intervals were calculated to evaluate the magnitude of effects. Finally, we ran a chi-square evaluating whether JUUL use status influenced beliefs that JUULs are e-cigarettes.

3. RESULTS

The sample comprised 56.6% never JUUL users, 13.2% ever users, and 30.2% past-month users (Table 1). When no information about nicotine strength was provided, 37.4% of all students thought JUULs contain a low or medium nicotine strength and 31.4% were unsure. After informing students that JUULs contain 5% nicotine, 58.8% of all students thought JUULs contain a low or medium nicotine strength and 19.9% were unsure. The GEE model produced a statistically significant interaction between information and JUUL use status ($\chi^2(6) = 101.73, p < .001$) and significant main effects for information ($\chi^2(3) = 499.19, p < .001$) and JUUL use ($\chi^2(6) = 164.80, p < .001$). The significant interaction was explained by a larger difference between the no information and information conditions in predicting perceived nicotine strength among never users than ever users ($\chi^2(3) = 29.52, p < .001$) and past month users ($\chi^2(3) = 98.52, p < .001$; Figure 1). However, there were no significant differences in the impact of information on perceived nicotine strength between ever and past month users ($\chi^2(3) = 4.59, p = 0.20$). Never users were more likely to rate nicotine strength as low (OR=7.50, 95% CI: [6.21, 9.06]) or medium (2.06, 95% CI: [1.76, 2.42]) rather than high when informed that JUULs contain 5% nicotine. Ever and past month users also were more likely to rate nicotine strength as lower after being informed that JUULs contain 5% nicotine, but the effects were smaller than for never users (low versus high nicotine: ever users OR=2.97, 95% CI: [2.25, 3.92], past-month users OR=2.22, 95% CI: [1.87, 2.62]; medium versus high nicotine: ever users OR=1.53, 95% CI: [1.17, 2.01], past-month users OR=1.19, 95% CI: [1.03, 1.36]). Past month users also were less likely to respond “don’t know” compared to rating the nicotine strength as high when they were informed that JUULs contain 5% nicotine (OR=0.76, 95% CI: [0.63, 0.91]). Taken together, the findings suggest that most students, including JUUL users, were unaware of JUULs’ high nicotine concentration, with even more students indicating that JUULs contain low or medium nicotine (versus high) when they were informed that JUULs contain 5% nicotine.

In total, 61.1% of adolescents believed that JUULs are e-cigarettes; 15.3% believed that JUULs are not e-cigarettes, and 23.7% did not know whether JUULs are e-cigarettes. Believing JUULs are e-cigarettes differed by JUUL use status ($\chi^2[4] = 94.73, p < .001$). Ever users (66.3%) and past-month users (68.9%) were more likely to think that JUULs are e-cigarettes than never users (55.7%). In addition, ever users (17.4%) and past-month users (14.4%) were less likely to report not knowing if JUULs are e-cigarettes compared to never users (30.0%).

4. DISCUSSION

At the time of the study, JUUL pods contained a uniformly high nicotine concentration (5%). However, when adolescents' baseline knowledge of JUULs nicotine strength was assessed, 37.4% of all adolescents thought JUULs were low or medium nicotine strength (including 59.2% of past-month users) and 31.4% did not know. When adolescents were told "JUULs contain 5% nicotine," all adolescents were more likely to think that JUULs contain low or medium nicotine (compared to high) than when they were given no information about nicotine content. Differences based on JUUL use status suggested that the findings were most pronounced among never JUUL users, which may be expected given their lack of familiarity with the product. However, believing JUULs contain a low level of nicotine may be a risk factor for experimentation/initiation among never users. With regard to JUUL users, the impact of being informed that "JUULs contain 5% nicotine" was not negligible; ever and past-month users were more than twice as likely to believe JUULs contain low nicotine after reading that JUULs contain 5% nicotine. These findings may be due to the fact that 5% sounds like a small amount, although this was not tested in the study. Further research is needed to determine whether labeling nicotine concentrations in milligrams/milliliter, using cigarettes as a reference point (e.g., 1 pod = 20 cigarettes), or developing a novel method of labeling nicotine concentrations increases youths' ability to understand nicotine strength. Irrespective of why youth misunderstand JUULs' nicotine strength, the findings are concerning for several reasons. First, adolescent JUUL users could be inadvertently exposing themselves to high nicotine levels, which is associated with deleterious effects on the developing brain including impaired cognitive development, executive functioning, and impulse control (USDHHS, 2012). Second, adolescents are more susceptible to developing nicotine dependence than adults (USDHHS, 2012), and recent evidence demonstrates that adolescents can experience nicotine dependence via using e-cigarettes (Morean et al., 2018). Thus, adolescents using high nicotine content products like JUUL may be especially vulnerable to addiction.

Furthermore, 39% of all adolescents did not consider JUULs to be e-cigarettes or were unsure, and a sizeable percentage (31%) of JUUL users did not recognize they are using an e-cigarette. It is possible that adolescents simply do not understand what constitutes a product being considered an e-cigarette, and, as a result, also may be unaware that they are using a tobacco product; this idea needs to be explored in future studies. Alternatively, adolescents may consider JUULs to be separate devices entirely, consistent with using the term "JUULing." Given that many adolescents did not realize that JUULs are e-cigarettes, our findings indicate that it is critical to ensure that questions assessing "e-cigarette" use are easily understood by participants (Alexander et al., 2016), and we suggest including popular brand examples in definitions. Alternatively, assessing the use of specific e-cigarette devices including JUUL may yield more accurate use rates, consistent with prior studies (Morean et al., 2019).

The study findings should be interpreted in light of several limitations. First, we informed students that JUULs contain 5% nicotine, which is not exactly what was printed on JUUL packaging at the time of the study (i.e., JUUL packages used the terminology "5% strength"). It is possible that youth may have been even more confused about what "5%

strength” means. Second, we focused on 5% nicotine JUUL pods, but did not assess adolescents’ ability to understand the nicotine strength of alternative brand pods that are compatible with JUUL yet are available in variable nicotine concentrations (e.g., Ziip pods; 1.8%, 3%) or newer 3% nicotine JUUL pods. While additional research is needed, the central finding that youth did not understand what 5% nicotine means likely extends to other concentrations labeled as percent nicotine. Third, the response options for the nicotine strength questions (i.e., low, medium, high) were somewhat vague, did not undergo cognitive testing, and no comparison point was provided (e.g., tobacco cigarettes). While future research is needed to quantify what constitutes low, medium, and high nicotine concentrations, at the time of the study, JUULs’ nicotine concentration was among the highest commercially available, clearly situating JUUL in the “high nicotine” category. Finally, data were obtained from four Connecticut high schools, and replication is needed in nationally representative samples.

In sum, the current study is the first to provide direct evidence that 1) most adolescents are unaware that JUUL pods contain a high nicotine concentration (when they are given no information about JUUL’s nicotine strength), 2) most adolescents do not understand that the nicotine concentration printed on JUUL pod packaging corresponds to a high nicotine level, and 3) nearly 40% of adolescents do not think JUULs are e-cigarettes or do not know. As previously noted, a standardized nicotine warning now is required on all e-liquid packaging. Although this warning informs users that a product contains nicotine, it does not speak to the strength of the product. Our results suggest that labeling nicotine concentrations using percent nicotine may be misleading, especially among youth. Regulatory efforts are needed to develop labels that convey nicotine concentrations in a meaningful way to all consumers and to ensure that products are labeled accordingly. Further, considering the known neurotoxic effects of nicotine on the developing brain, regulators should consider including content on labels that educates youth about the negative consequences of nicotine exposure. In addition, given that nearly 40% of students did not think JUULs are e-cigarettes or were unsure, efforts are needed to determine the best ways to clarify for youth what products constitute e-cigarettes and that e-cigarettes are tobacco products. Health risk communication and prevention efforts targeting youth e-cigarette use should include specific language about JUUL and other pod-based systems, including nicotine levels, to ensure that youth understand that the risks associated with e-cigarette use extend to these products. Alternatively, given the ever evolving landscape of nicotine-delivering products, efforts may be more effective if they simply educate youth about the dangers of inhaling nicotine in any form.

Acknowledgments

We wish to thank all participants who contributed to this work.

Role of Funding Source

Research reported in this publication was supported by the National Institutes of Health/NIDA and FDA Center for Tobacco Products (CTP) (P50DA036151; Yale Tobacco Center of Regulatory Science, TCORS). The content is solely the responsibility of the authors and does not necessarily represent the official views of the funding agencies, including the NIH or the Food and Drug Administration. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

References

- Alexander JP, Coleman BN, Johnson SE, Tessman GK, Tworek C, Dickinson DM, 2016 Smoke and vapor: exploring the terminology landscape among electronic cigarette users. *Tob Regul Sci*, 2(3), 204–213. [PubMed: 27430008]
- Carver R Juul exceeds 60 percent market share for e-cigs. *Winston-Salem Journal*. 5 1, 2018 https://www.journalnow.com/business/juul-exceeds-percent-market-share-for-e-cigs/article_1fd229f1-df7d-5436-ac90-9263c418dc1c.html. Accessed February 12, 2019.
- Huang J, Duan Z, Kwok J, Binns S, Vera LE, Kim Y, Szczypka G, Emery SL, 2019 Vaping versus JUULing: how the extraordinary growth and marketing of JUUL transformed the US retail e-cigarette market. *Tob Control*, 28(2), 146–151. [PubMed: 29853561]
- JUUL Labs Inc. Our Mission, <https://www.juul.com/mission-values>. Accessed February 12, 2019.
- Kavuluru. R, Han S, Hahn EJ, 2019 On the popularity of the USB flash drive-shaped electronic cigarette Juul. *Tob Control*. 28,110–112. doi:10.1136/tobaccocontrol-2018-054259 [PubMed: 29654121]
- Krishnan-Sarin S, Jackson A, Morean M, Kong G, Bold KW, Camenga DR, Cavallo DA, Simon P, Wu R, 2019 E-cigarette devices used by high-school youth. *Drug Alcohol Depend*. 194, 395–400. [PubMed: 30497057]
- Morean ME, Camenga D, Bold K, Kong G, Jackson A, Simon P, Krishnan-Sarin S, 2019 Querying about the use of specific e-cigarette devices may enhance accurate measurement of e-cigarette prevalence rates among high school students. *Nicotine Tob Res*. 2019; doi:10.1093/ntr/nty240.
- Morean ME, Kong G, Cavallo DA, Camenga DR, Krishnan-Sarin S, 2016 Nicotine concentration of e-cigarettes used by adolescents. *Drug Alcohol Depend*. 167, 224–227. [PubMed: 27592270]
- Morean ME, Krishnan-Sarin S, O'Malley SS, 2018 Assessing nicotine dependence in adolescent e-cigarette users: The 4-item Patient-Reported Outcomes Measurement Information System (PROMIS) Nicotine Dependence Item Bank for electronic cigarettes. *Drug Alcohol Depend*. 188, 60–63. [PubMed: 29753155]
- U.S. Department of Health and Human Services, 2012 Preventing tobacco use among youth and young adults: A Report of the Surgeon General. https://www.ncbi.nlm.nih.gov/books/NBK99237/pdf/Bookshelf_NBK99237.pdf Accessed February 12, 2019.
- US Food and Drug Administration, 2018 "Covered" Tobacco Product and Roll-Your-Own/Cigarette Tobacco Labeling and Warning Statement Requirements. <https://www.fda.gov/TobaccoProducts/Labeling/Labeling/ucm524470.htm#nicotine>. Accessed February 12, 2019.
- US Food and Drug Administration, 2018 Statement from FDA Commissioner Scott Gottlieb, M.D., on new enforcement actions and a Youth Tobacco Prevention Plan to stop youth use of, and access to, JUUL and other e-cigarettes [4 24, 2018]. <https://www.fda.gov/newsevents/newsroom/pressannouncements/ucm605432.htm>. Accessed February 12, 2019.
- Willett JG, Bennett M, Hair EC, Xiao H, Greenberg MS, Harvey E, Cantrell J and Vallone D, 2019 Recognition, use and perceptions of JUUL among youth and young adults. *Tob Control*. 28, 115–116. [PubMed: 29669749]
- Ziip Pods. <https://www.vapor4life.com/vape-pods/ziip-pods/>. Accessed February 12, 2019.

Highlights

- JUULs can contain 5% nicotine, which teens may not understand is a high nicotine level
- When no information about nicotine strength was provided, most teens thought JUULs contain low or medium nicotine or did not know
- When told JUULs contain 5% nicotine, more teens thought JUULs contain low or medium nicotine
- 38.9% of teens thought JUULs were not a type of e-cigarette or were unsure




Figure 1. Perceived nicotine strength varies based on information about nicotine content provided (no information vs. informed JUULs contain 5% nicotine) and JUUL use status
Note. * denotes responses that differ significantly from a response of “high nicotine” after adolescents were informed that JUULs contain 5% nicotine.

Table 1. Descriptive statistics for central study variables within the total sample and by JUUL use status

Study Variables	Study Questions and Responses by Study Sample				
	Total Sample (N = 3170)	Never JUUL Users (n = 1795, 6.6%)	Ever JUUL Users (n = 419; 3/2%)	Past-Month JUUL Users (n = 956; 30.2%)	
Birth Sex	<i>At birth, what was your sex?</i>				
Female (%)	52.4	52.1	51.8	53.1	
Age	<i>How old are you?</i>				
Mean years (std. dev.)	15.87 (1.29)	15.69 (1.28)	15.95 (1.25)	16.17 (1.26)	
Race	<i>How would you describe your race?</i>				
White (%)	60.4	55.7	62.5	68.1	
Ever JUUL Use	<i>Have you ever tried a JUUL?</i>				
Yes (%)	43.4	0.0	100.0	100.0	
Past 30 Day JUUL Use	<i>Approximately how many days out of the past 30 days did you use a JUUL?</i>				
Frequency in # Days	--	0.0	0.0	13.59 (11.68)	
Any Past 30 Day Use (%)	30.2	0.0	0.0	100.0	
JUUL Nicotine Strength (No Info)	<i>Which of the following do you think best describes the amount of nicotine in a JUUL?</i>				
Low (%)	10.5	7.3	15.6	14.4	
Medium (%)	26.9	23.6	27.4	33.0	
High (%)	31.1	29.4	28.8	35.4	
I don't know (%)	31.4	39.7	28.1	17.2	
JUUL Nicotine Strength (5% Specified)	<i>The nicotine content of a JUUL is 5%. Which of the following do you think best describes the amount of nicotine in a JUUL?</i>				
Low (%)	29.5	30.3	32.9	26.5	
Medium (%)	29.3	27.4	29.8	32.7	
High (%)	21.3	16.8	20.9	29.9	
I don't know (%)	19.9	25.6	16.3	10.8	
JUUL as a Type of E-cigarette	<i>Do you consider a JUUL to be a type of e-cigarette?</i>				
No (%)	15.3	14.3	16.2	16.6	
Yes (%)	61.1	55.7	66.3	68.9	
I don't know (%)	23.7	30.0	17.4	14.4	