



RESEARCH PAPER

# Facultative crassulacean acid metabolism in a C<sub>3</sub>–C<sub>4</sub> intermediate

Klaus Winter<sup>1,\*</sup>, Rowan F. Sage<sup>2, </sup>, Erika J. Edwards<sup>3</sup>, Aurelio Virgo<sup>1</sup> and Joseph A. M. Holtum<sup>1,4, </sup>

<sup>1</sup> Smithsonian Tropical Research Institute, PO Box 0843-03092, Balboa, Ancón, Republic of Panama

<sup>2</sup> Department of Ecology and Evolutionary Biology, University of Toronto, Toronto, ON, M5S 3B2, Canada

<sup>3</sup> Department of Ecology and Evolutionary Biology, Yale University, New Haven, CT 06511, USA

<sup>4</sup> College of Science and Engineering, James Cook University, Townsville, QLD, 4811, Australia

\* Correspondence: [winterk@si.edu](mailto:winterk@si.edu)

Received 18 October 2018; Editorial decision 12 February 2019; Accepted 14 February 2019

Editor: John Cushman, University of Nevada, USA

## Abstract

The *Portulacaceae* enable the study of the evolutionary relationship between C<sub>4</sub> and crassulacean acid metabolism (CAM) photosynthesis. Shoots of well-watered plants of the C<sub>3</sub>–C<sub>4</sub> intermediate species *Portulaca cryptopetala* Speng. exhibit net uptake of CO<sub>2</sub> solely during the light. CO<sub>2</sub> fixation is primarily via the C<sub>3</sub> pathway as indicated by a strong stimulation of CO<sub>2</sub> uptake when shoots were provided with air containing 2% O<sub>2</sub>. When plants were subjected to water stress, daytime CO<sub>2</sub> uptake was reduced and CAM-type net CO<sub>2</sub> uptake in the dark occurred. This was accompanied by nocturnal accumulation of acid in both leaves and stems, also a defining characteristic of CAM. Following rewatering, net CO<sub>2</sub> uptake in the dark ceased in shoots, as did nocturnal acidification of the leaves and stems. With this unequivocal demonstration of stress-related reversible, i.e. facultative, induction of CAM, *P. cryptopetala* becomes the first C<sub>3</sub>–C<sub>4</sub> intermediate species reported to exhibit CAM. *Portulaca molokiniensis* Hobdy, a C<sub>4</sub> species, also exhibited CAM only when subjected to water stress. Facultative CAM has now been demonstrated in all investigated species of *Portulaca*, which are well sampled from across the phylogeny. This strongly suggests that in *Portulaca*, a lineage in which species engage predominately in C<sub>4</sub> photosynthesis, facultative CAM is ancestral to C<sub>4</sub>. In a broader context, it has now been demonstrated that CAM can co-exist in leaves that exhibit any of the other types of photosynthesis known in terrestrial plants: C<sub>3</sub>, C<sub>4</sub> and C<sub>3</sub>–C<sub>4</sub> intermediate.

**Keywords:** C<sub>4</sub> photosynthesis, crassulacean acid metabolism, *Portulaca cryptopetala*, *Portulaca molokiniensis*, Portulacaceae.

## Introduction

An estimated 10% of terrestrial vascular plants express either crassulacean acid metabolism (CAM) or C<sub>4</sub> photosynthesis (Smith and Winter, 1996; Winter *et al.*, 2015; Sage, 2016). Both photosynthetic pathways have evolved independently over 60 times. CAM is documented in more than 30 angiosperm families, and in one family in each of the cycads, gnetophytes, ferns, and lycophytes (Smith and Winter, 1996), while C<sub>4</sub> is known in 19 families of angiosperms (Sage and Sultman, 2016).

The CAM and C<sub>4</sub> pathways are comparable in many respects (Osmond, 1978; Hatch, 1987). Using a similar complement of enzymes, each pathway concentrates CO<sub>2</sub> in the vicinity of Rubisco thereby reducing the competitive inhibition by molecular oxygen of CO<sub>2</sub> uptake. In both pathways, atmospheric CO<sub>2</sub> initially fixed as HCO<sub>3</sub><sup>-</sup> using oxygen-insensitive phosphoenolpyruvate carboxylase (PEPc) is incorporated into a four-carbon intermediate from which CO<sub>2</sub> is ultimately

Abbreviations: BS, bundle sheath; FM, fresh mass; PEPc, phosphoenolpyruvate carboxylase; PFD, photosynthetic photon flux density.

© The Author(s) 2019. Published by Oxford University Press on behalf of the Society for Experimental Biology.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited. For commercial re-use, please contact [journals.permissions@oup.com](mailto:journals.permissions@oup.com)

liberated in the vicinity of Rubisco. At the site of Rubisco, CO<sub>2</sub> attains concentrations that ensure that the enzyme functions overwhelmingly as a carboxylase.

Despite similarities, CAM and C<sub>4</sub> differ in important aspects. In C<sub>4</sub> plants, all of the processes associated with photosynthetic CO<sub>2</sub> assimilation occur during the light. PEPc and Rubisco are simultaneously active but are separated spatially, usually in two distinct types of cells (for exceptions, see e.g. [Edwards and Voznesenskaya, 2011](#)). The primary carboxylation by PEPc typically occurs in thin-walled mesophyll cells that surround thicker walled bundle-sheath (BS) cells. The four-carbon intermediate is transferred via plasmodesmata to BS cells where CO<sub>2</sub> is liberated and the refixation of the CO<sub>2</sub> by Rubisco takes place. In contrast to C<sub>4</sub> photosynthesis, CAM is essentially a single-cell phenomenon, during which the PEPc- and Rubisco-catalysed carboxylations operate at different times of the day–night cycle, i.e. their activity is separated temporally. During the night, CO<sub>2</sub> is fixed by PEPc and the four-carbon intermediate, malic acid, is stored in large vacuoles. During the following light period, the stomata close, PEPc is inactivated, and CO<sub>2</sub> released from the decarboxylation of malic acid is refixed by Rubisco.

Across the phylogenetic tree of angiosperms ([Ogburn and Edwards, 2010, 2012](#)), CAM and C<sub>4</sub> origins cluster in numerous distinct clades suggesting that certain plant lineages are prone to evolve both pathways ([Edwards and Ogburn, 2012](#)). It has been proposed that the distinct anatomical requirements of the CAM and C<sub>4</sub> pathways, coupled with differences in timing and regulation of their respective biochemical pathways, reduce the likelihood that both co-occur in the same organ. The possibility that parts of the C<sub>4</sub> and CAM cycles take place in the same cell has been suggested to be even less likely ([Sage, 2002](#)). In accordance, the reported instances of co-expression of CAM and C<sub>4</sub> within plants with Kranz anatomy are rare, with the only known cases being observed in *Portulaca* ([Koch and Kennedy, 1980](#); [Guralnick \*et al.\*, 2002](#)). *Portulaca* is the only genus of the Portulacaceae, a family assigned to the order Caryophyllales in which CAM and C<sub>4</sub> have evolved multiple times ([Christin \*et al.\*, 2014](#)). Phylogenetic relationships of families within the Caryophyllales and the currently known distribution of CAM and C<sub>4</sub> photosynthesis among them have been recently featured in [Holtum \*et al.\* \(2018\)](#) (see their [Fig. 3](#)).

Originally, CAM and C<sub>4</sub> were reported for two species of *Portulaca* (*P. oleracea* and *P. grandiflora*), and in each, it appeared that CAM and C<sub>4</sub> were confined to different cell or tissue regions. More recent studies indicate that the co-existence of CAM and C<sub>4</sub> in the same photosynthetic organ is common in *Portulaca* ([Winter and Holtum, 2014, 2017](#); [Holtum \*et al.\*, 2017a](#); [Winter, 2019](#)). CAM has been demonstrated by CO<sub>2</sub> gas exchange and quantification of nocturnal acidification in seven species from four of the six major phylogenetic clades of *Portulaca*. In each case of CAM and C<sub>4</sub> co-expression, the expression of CAM was facultative ([Guralnick \*et al.\*, 2002](#); [D'Andrea \*et al.\*, 2014](#); [Winter and Holtum, 2014, 2017](#); [Holtum \*et al.\*, 2017a](#)). CAM-type gas-exchange patterns and nocturnal acidification were not detected in well-watered plants, but were induced when the plants were subjected to water stress. When stressed plants were rewatered, their physiology returned to the original well-watered pattern. The observation of widespread

CAM in *Portulaca* is supported by the evolutionary history of PEPc genes in *Portulaca* ([Christin \*et al.\*, 2014](#)). The putative gene encoding CAM-specific PEPc was apparently present before the divergence of *Portulaca*, and is similarly used for CAM in relatives of *Portulaca*, whereas PEPcs optimized for C<sub>4</sub> metabolism in *Portulaca* originated from a duplication event of a different paralog, which occurred at the base of *Portulaca*.

The coexistence of C<sub>4</sub> and CAM in leaves of C<sub>4</sub> *Portulaca* species raises interesting questions about the location of both pathways, i.e. whether they occur in different regions of the leaf or whether there is cell sharing. This issue is not yet fully resolved. In *Portulaca oleracea*, CAM-type nocturnal CO<sub>2</sub> fixation presumably takes place in centripetally located large parenchyma cells, yet critical daytime reactions of the CAM cycle may occur in the C<sub>4</sub> bundle-sheath cells ([Lara \*et al.\*, 2003, 2004](#)). By contrast, for *P. grandiflora* separate operation of the C<sub>4</sub> and CAM pathways in different regions of the leaf has been postulated, with C<sub>4</sub> in mesophyll cells associated with the bundle sheath cells and the complete CAM cycle taking place in the centripetal parenchyma cells ([Guralnick and Jackson, 2001](#); [Guralnick \*et al.\*, 2002](#); [Holtum \*et al.\*, 2017a](#)).

Species in five of the six phylogenetic clades of *Portulaca* are thought to use C<sub>4</sub> as the principal pathway of carbon acquisition ([Ocampo \*et al.\*, 2013](#); [Voznesenskaya \*et al.\*, 2017](#)). All species examined exhibit C<sub>4</sub>-type δ<sup>13</sup>C values, Kranz anatomies, enzyme complements, and gas-exchange characteristics. *Portulaca* is not known to contain C<sub>3</sub> species *senso strictu*, but three species in the *Cryptopetala* clade, *P. cryptopetala*, *P. hirsutissima* and *P. mucronata*, have been characterized as C<sub>3</sub>–C<sub>4</sub> intermediates on the basis of C<sub>3</sub>-type δ<sup>13</sup>C values, anatomy, location of glycine decarboxylase, and CO<sub>2</sub> compensation points ([Voznesenskaya \*et al.\*, 2010, 2017](#); [Ocampo \*et al.\*, 2013](#)). It was inferred that the C<sub>3</sub>–C<sub>4</sub> *Cryptopetala* clade evolved from C<sub>4</sub> progenitors and that it represents a reversion from a C<sub>4</sub> state ([Ocampo and Columbus, 2012](#); [Ocampo \*et al.\*, 2013](#)). The reversion hypothesis was questioned by [Christin \*et al.\* \(2014\)](#) who argued, on the basis of the composition of PEPc genes, the distinct leaf anatomy in each major clade, and the diversity of the de-carboxylating enzymes used by the different clades, that C<sub>4</sub> evolved multiple times in parallel. The *Cryptopetala* clade may therefore be a lineage of *Portulaca* with a photosynthetic complement that reflects a pre-C<sub>4</sub> stage.

CAM in the *Cryptopetala* clade would strengthen the argument that CAM represents an ancestral state in *Portulaca*, being present prior to the evolution of C<sub>4</sub> photosynthesis. If so, the relationship between C<sub>3</sub>–C<sub>4</sub> metabolism and CAM remains unclear. In C<sub>3</sub>–C<sub>4</sub> intermediates, CO<sub>2</sub> is concentrated into BS-like compartments via the localization of the photorespiratory enzyme glycine decarboxylase (GDC) in the BS, and the shuttling of photorespiratory glycine into the BS for decarboxylation. This metabolism, termed C<sub>2</sub> photosynthesis, can raise CO<sub>2</sub> concentrations in the BS two to three times above the atmospheric value, but does not greatly alter δ<sup>13</sup>C values from what are present in C<sub>3</sub> species ([Keerberg \*et al.\*, 2014](#); [Sage \*et al.\*, 2014](#)). As proposed for C<sub>4</sub> plants, dual expression of C<sub>2</sub> metabolism and CAM could interfere with the optimal function of each, and hence it could be hypothesized that the two metabolic types are segregated either to different tissues or to different phases of development. Here we use gas exchange and

measurements of titratable acidity to explore whether CAM is present in the annual/biennial *P. cryptopetala*, and in the perennial *P. molokiniensis* (Hobdy, 1987). *Portulaca cryptopetala* is a C<sub>3</sub>–C<sub>4</sub> intermediate and a member of one of the three clades of *Portulaca* in which CAM has not yet been reported. *Portulaca molokiniensis* is a C<sub>4</sub> species that belongs to the C<sub>4</sub> Oleracea clade that is sister to the *Cryptopetala* clade.

## Materials and methods

Seeds of *P. cryptopetala* and *P. molokiniensis* were obtained from the laboratory stock of one of us (RFS). Plants were grown from seed in either 0.5 litre terracotta pots with an upper diameter of 10 cm, or in 1 litre terracotta pots with an upper diameter of 13 cm. Pots contained potting mix (Miracle-Gro Lawn Products, Marysville, OH, USA). Plants were 1–3 months old when studied.

Two laboratory gas-exchange systems were used to measure 24 h patterns of CO<sub>2</sub> gas exchange of plants. Whole shoots were enclosed in either an 11×11×10 cm or an 11×11×16 cm Perspex cuvette. Roots plus pot remained outside the cuvette. The gas-exchange cuvettes were located inside controlled-environment chambers operating under 12 h light (28 °C):12 h dark (22 °C) cycles. Light was provided by LED grow lights (model LL4L-GP300, GrowPro300). Photon flux density at the level of the cuvettes is specified in the corresponding figure legends. Cuvettes were supplied with air containing 400 ppm CO<sub>2</sub> at flow rates of either 1.26 or 2.5 l min<sup>-1</sup>. Net CO<sub>2</sub> exchange was measured in flow-through gas-exchange systems consisting of Walz components (gas mixing units, air pumps, cold traps, dew point mirrors; Walz GmbH, Effeltrich, Germany), LI-6252 CO<sub>2</sub> analyzers (Li-Cor, NE, USA) and CR-1000 data loggers (Campbell Scientific, UT, USA) (Holtum and Winter, 2003). For measurements at 2% O<sub>2</sub>, N<sub>2</sub> flowing at 4.75 l min<sup>-1</sup> was added to ambient air flowing at 0.5 l min<sup>-1</sup>. CO<sub>2</sub> was removed by passing the mixture through soda-lime and then re-added via a mass-flow controller to obtain 400 ppm CO<sub>2</sub> before the gas mixture entered the cuvette. Exposures to air containing 2% O<sub>2</sub> lasted 30–60 min.

Well-watered plants were watered daily to field capacity. Drought treatments were imposed by withholding irrigation until net CO<sub>2</sub> uptake in the light was reduced to close to, at most, 10% of the value for well-watered plants, after which the plants were rewatered daily.

In a separate set of experiments, nine plants of each species were grown in the laboratory under 12 h light–12 h dark cycles. Photosynthetically active photon flux density (PPFD) was 600 μmol m<sup>-2</sup> s<sup>-1</sup> supplied by a LED grow light (300 W Diamond series, Advanced LED Lights, Hiwasse, AR, USA). Temperature was 26 °C during light periods and 24 °C during dark periods. Plants watered daily to field capacity were deprived of water for several days and then rewatered as specified in the corresponding figure legends. Mature leaves were excised at the end of the light and dark periods from each well-watered, drought-stressed and rewatered plants, and then the fresh mass (FM) obtained, and leaf area measured using a LI-3100 area meter (Li-Cor). Samples were then frozen in liquid nitrogen and freeze-dried. After determination of dry mass, samples were boiled in 80 ml of 50% ethanol until the volume had about halved. Water was then added to bring the volume back to 80 ml and the extract was boiled until the volume again decreased by about half. The extracts were brought to the original volume with water, cooled to room temperature, and titrated with 5 mM KOH to pH 6.5.

## Results


Well-watered plants of *P. cryptopetala* exhibited net CO<sub>2</sub> uptake during the day and net CO<sub>2</sub> loss at night (Fig. 1; see also Supplementary Fig. S1 at JXB online). The net rates of CO<sub>2</sub> exchange during the day and night increased as the plants grew. In the experiment of Fig. 1, 3 d after watering ceased

(day 5 of the experiment), net CO<sub>2</sub> exchange began to decline in the light and the dark. The shape of the CO<sub>2</sub> exchange curve in the dark was noticeably more curved, and nocturnal CO<sub>2</sub> exchange approached the CO<sub>2</sub> compensation point. On day 6, net CO<sub>2</sub> uptake was present for the first time at night. Nocturnal uptake peaked during the night of day 7 and remained approximately constant until the night of day 9, the day prior to rewatering. Within 6 h of rewatering the plant on day 10, CO<sub>2</sub> uptake during the light had almost recovered to the rates observed before the imposition of water stress. No nocturnal net CO<sub>2</sub> uptake was present during the following dark periods.


On day 4, when the *P. cryptopetala* plant shown in Fig. 1 was still exhibiting the well-watered pattern of CO<sub>2</sub> uptake in the light and CO<sub>2</sub> loss at night, the transfer of shoots during the light from an air-stream containing 21% O<sub>2</sub> to an air-stream containing 2% O<sub>2</sub> was accompanied by an increase in the rate of net CO<sub>2</sub> uptake of up to 46%. When air containing 21% O<sub>2</sub> was resupplied, the rate of CO<sub>2</sub> uptake reattained the control pre-2% O<sub>2</sub> rate. A total of nine 2% O<sub>2</sub> treatments were performed on three plants and resulted in an increase of CO<sub>2</sub> uptake by 39±7% (mean ±SD, n=3). The range was 31–46%.

When a plant of *P. cryptopetala* was exposed to sequential watering, droughting, and rewatering cycles, the stress-related induction of net CO<sub>2</sub> uptake in the dark was observed during each period of water stress (Fig. 2). The shoot inside the gas-exchange cuvette continued to grow during the experiment as evidenced by the progressive increase in net CO<sub>2</sub> uptake during the light.

Leaves of well-watered *P. cryptopetala* either did not exhibit nocturnal acidification or, if it was present, the end of night/end of day differences in acidity were very low (Fig. 3). Following the imposition of water stress, strong nocturnal



**Fig. 1.** Eleven days of net CO<sub>2</sub> exchange by the shoot (leaves plus stems) of a *Portulaca cryptopetala* plant growing in a pot. Watering was withheld on day 2 (red arrow) and recommenced on day 10 (blue arrow). During the light period of day 4, the shoot was exposed to air containing 2% O<sub>2</sub> (black arrow) for approx. 1 h. Shaded areas represent the 12 h dark periods. PPFD incident to the top of the gas-exchange cuvette was 1000 μmol m<sup>-2</sup> s<sup>-1</sup>. At the end of the experiment, total leaf area was 71.1 cm<sup>2</sup>, and leaf and stem dry masses were 0.26 and 0.095 g, respectively.



**Fig. 2.** Twenty-five days of net CO<sub>2</sub> exchange by the shoot (leaves and stems) of a potted *Portulaca cryptopetala* that was exposed to two wetting and drying cycles. Watering was withheld on days 3 and 16 (red arrows) and recommenced on days 14 and 23 (blue arrows). Shaded areas represent the 12 h dark periods. PFD incident to the top of the gas-exchange cuvette was 1350  $\mu\text{mol m}^{-2} \text{s}^{-1}$ . At the end of the experiment, total leaf area was 152 cm<sup>2</sup> and leaf and stem dry masses were 0.578 and 0.372 g, respectively.

acidification was induced, reaching about 110  $\mu\text{mol H}^+ \text{g}^{-1}$  FM. At the end of the night, the absolute leaf H<sup>+</sup> content was about 25-fold greater than in unstressed plants. Following rewatering, nocturnal leaf acidification was reduced markedly such that the end of the night–end of the day differences in H<sup>+</sup> levels were close to zero. The expression in Fig. 3 of acid levels on fresh mass, dry mass, and leaf area bases enables the calculation of acid concentrations in leaves, permits estimation of the effects of changes in leaf-water content that occur during the droughting process, and facilitates comparison with gas-exchange measurements of CO<sub>2</sub> exchange.


In stems of *P. cryptopetala*, in a manner similar to leaves, marked nocturnal acidification was induced when the plants were subjected to water stress (Fig. 4), with the end of the night acid pool increasing by about 10-fold in comparison to unstressed plants. In contrast to leaves, the stems of rewatered plants continued to exhibit nocturnal acidification, although the levels on a fresh mass basis were only about 14% of those observed in stems of droughted plants.

Well-watered shoots of *P. molokiniensis* exhibited net CO<sub>2</sub> uptake during the light and net CO<sub>2</sub> loss in the dark (Fig. 5). Following the imposition of water stress, a marked decrease in CO<sub>2</sub> uptake was accompanied by the induction of net CO<sub>2</sub> uptake in the dark. Rewatering was followed by a recovery of net CO<sub>2</sub> uptake during the light and a loss of nocturnal net CO<sub>2</sub> uptake. As was observed for *P. cryptopetala*, the exposure of shoots of *P. molokiniensis* to sequential watering, droughting, and rewatering cycles was accompanied by the stress-related

induction of net CO<sub>2</sub> uptake at night during each period of water stress. The continued increase in net CO<sub>2</sub> uptake during the light demonstrated that the shoots of *P. molokiniensis* continued to grow during the experiment. The stress-induced, reversible induction of net dark CO<sub>2</sub> fixation shown in Fig. 5 was fully confirmed in three additional gas-exchange experiments with three different *P. molokiniensis* plants (see Supplementary Figs S2–S4).

In well-watered *P. molokiniensis*, the transfer of shoots during the light from an air-stream containing 21% O<sub>2</sub> to an air-stream containing 2% O<sub>2</sub> was accompanied by an increase in the rate of net CO<sub>2</sub> uptake by 8±3% (mean ±SD, n=3 different plants; total of 10 measurements) (e.g. Fig. 5; Supplementary Fig. S2). The range was 5–14%. As with *P. cryptopetala*, when air containing 21% O<sub>2</sub> was resupplied, the rate of CO<sub>2</sub> uptake reattained the control pre-2% O<sub>2</sub> rate.


In a manner similar to *P. cryptopetala*, nocturnal acidification was either not present or barely detectable in leaves of well-watered *P. molokiniensis* (Fig. 6). Leaf acidity increased at the end of the light and the dark periods when plants were stressed. The increase in acidity at the end of the dark was much greater than at the end of the light period, resulting in substantial net acidification during the night. Nocturnal acidification of similar magnitude in droughted *P. molokiniensis* has been observed previously (L. Guralnik, unpublished data, personal communication). The nocturnal acidification was completely lost following rewatering, although the background [H<sup>+</sup>] remained somewhat greater than acidity levels at the beginning of the experiment.



**Fig. 3.** Titratable acidity in recently fully expanded leaves of *Portulaca cryptopetala* at the end of the 12 h light period (red) and the end of the 12 h dark period (blue) in plants that were well-watered (left-hand column), droughted (middle column; 9 d without irrigation) and droughted and rewatered (right-hand column; 3 d with irrigation). The data are expressed on a fresh mass basis (A), a dry mass basis (B) and a leaf area basis (C). Mean  $\pm$ SD ( $n=5$  leaves; at a given time point each leaf was harvested from a different plant). The numerical values shown above the bars are  $P$  values (one-tailed  $t$ -test). Bold letters indicate that the values at the end of the dark period were significantly greater than those at the end of the day at  $P \leq 0.05$ .

## Discussion

The demonstration of CAM in *P. cryptopetala* and in *P. molokiniensis* adds a new facet to our understanding of the diversity in origins, functioning, expression, and interrelationships of C<sub>3</sub>, C<sub>4</sub>, and CAM photosynthesis. CAM, long known to be co-expressed alongside C<sub>3</sub> photosynthesis in plants with succulent tissues, is now documented in eight C<sub>4</sub> species, all within *Portulaca* (Koch and Kennedy, 1980; Guralnick *et al.*, 2002; Christin *et al.*, 2014; Winter and Holtum, 2014, 2017; Holtum *et al.*, 2017a). With the evidence presented here for CAM in *P. cryptopetala*, we can now conclude that CAM can also co-occur in leaves with C<sub>3</sub>-C<sub>4</sub> photosynthesis. In *Portulaca*, the C<sub>4</sub> and C<sub>3</sub>-C<sub>4</sub> intermediate species that express CAM are dispersed across five of the six clades of *Portulaca* (Fig. 7). CAM




**Fig. 4.** Titratable acidity in stems of *Portulaca cryptopetala* at the end of the 12 h light period (red) and the end of the 12 h dark period (blue) in plants that were well-watered (left-hand column), droughted (middle column; 10 d without irrigation), and droughted and rewatered (right-hand column; 2 d with irrigation). The data are expressed on a fresh mass basis (A) and a dry mass basis (B). Mean  $\pm$ SD ( $n=5$  stems; at a given time point each stem was harvested from a different plant). The numerical values shown above the bars are  $P$  values (one-tailed  $t$ -test). Bold letters indicate that the values at the end of the dark period were significantly greater than those at the end of the day at  $P \leq 0.05$ .

has been detected in species with all of the forms of anatomy described for *Portulaca* (Atriplicoid, Pilosoid, Portulaceloid and C<sub>3</sub>-C<sub>4</sub>) and in both NAD-ME C<sub>4</sub> species (*P. oleracea* and *P. molokiniensis*) and in NADP-ME C<sub>4</sub> species (*P. pilosa*, *P. grandiflora* and *P. umbraticola*) (Voznesenskaya *et al.*, 2010, 2017; Ocampo *et al.*, 2013).

In *P. cryptopetala* and *P. molokiniensis*, as in other *Portulaca* with CAM, the expression of CAM is unmistakably facultative. Compared with rates of C<sub>3</sub> and C<sub>4</sub> photosynthesis in unstressed plants, the magnitudes of CAM-type dark CO<sub>2</sub> uptake and nocturnal acidification are relatively low, but both characters are clearly present in water-stressed plants and are absent, or close to absent, in well-watered plants (Figs 1–6). The observation that CAM can be repeatedly induced or lost following cycles of water supply and water stress in *P. cryptopetala* and *P. molokiniensis* (Figs 2, 5) reveals a tight relationship between the environmental trigger, in this case water stress, and the physiological reaction of the plants, independent of ontogeny (Winter and Holtum, 2007).

At present, in the absence of field studies, we can only speculate as to how a combination of C<sub>4</sub> and CAM traits in a single plant might potentially be of adaptive significance. The most obvious conclusion is that C<sub>4</sub> and C<sub>2</sub> provide a capacity for enhanced productivity and that CAM increases the ability to cope with



**Fig. 5.** Twenty-seven days of net CO<sub>2</sub> exchange by shoots of a potted *Portulaca molokiniensis* that was exposed to two wetting and drying cycles. Watering was withheld on days 5 and 16 (red arrows) and recommenced on days 14 and 23 (blue arrows). During the light periods of days 6, 8, and 15 the shoots were exposed to air containing 2% O<sub>2</sub> (black arrows). Shaded areas represent the 12 h dark periods. PFD incident to the top of the gas-exchange cuvette was 1000  $\mu\text{mol m}^{-2} \text{s}^{-1}$ . At the end of the experiment, total leaf area was 97.5 cm<sup>2</sup>, and leaf and stem dry masses were 0.405 and 0.060 g, respectively.


water stress (Winter and Ziegler, 1992). Particularly in warmer climates, the C<sub>4</sub> component could enable rapid growth and high nitrogen-use efficiency, and CAM could contribute to survival via its ability to reduce carbon and water loss when the supply of water is constrained. The rapid switching from CAM back to C<sub>4</sub> would be expected to enable a prompt response to rainfall events, an ability of relevance to species that are fast-growing, generally annual, weedy ecological opportunists of disturbed sites, e.g. *P. cryptopetala*, *P. grandiflora*, *P. oleracea*, and *P. pilosa*.

While an intermediate CO<sub>2</sub> compensation point and other characteristics (Voznesenskaya *et al.*, 2010, 2017; Ocampo *et al.*, 2013) support the notion that *P. cryptopetala* is not a C<sub>4</sub> species but rather a C<sub>3</sub>–C<sub>4</sub> species, the stimulation of photosynthesis by up to 46% when *P. cryptopetala* was exposed to air containing 2% O<sub>2</sub> (Fig. 1; Supplementary Fig. S1), together with C<sub>3</sub>-type  $\delta^{13}\text{C}$  values, suggests that it is an intermediate in which, at current ambient CO<sub>2</sub> concentrations, uptake of atmospheric CO<sub>2</sub> in the light is catalysed largely by Rubisco. Presumably, this Rubisco signal is contributed to by Rubisco in C<sub>3</sub>–C<sub>4</sub> tissue and CAM tissue.

In contrast to *P. cryptopetala*, the exposure of *P. molokiniensis* to 2% O<sub>2</sub> resulted in up to a 14% stimulation of photosynthesis (Fig. 5; Supplementary Fig. S2), a response more similar to that of C<sub>4</sub> plants. In C<sub>4</sub> plants, photosynthesis is typically unaffected by a transfer from 21 to 2% O<sub>2</sub> but, at current ambient [CO<sub>2</sub>], it is not uncommon for plants to exhibit a small stimulation in photosynthesis as [O<sub>2</sub>] is lowered from 21 to 5–10%

followed by a small inhibition as [O<sub>2</sub>] is further reduced to 2% or lower (Maroco *et al.*, 1997, 1998). The inhibition is thought to be related to a greater requirement for O<sub>2</sub>-dependent ATP generation by C<sub>4</sub> photosynthesis compared with C<sub>3</sub> photosynthesis. The ATP is required to regenerate PEP, the primary substrate of the C<sub>4</sub> cycle. The [O<sub>2</sub>] at which the stimulation-to-inhibition transition occurs is apparently species-specific and may be anywhere between 10 and 2%. The small stimulation in CO<sub>2</sub> uptake in well-watered plants of *P. molokiniensis* following exposure to 2% O<sub>2</sub> is probably not an effect of O<sub>2</sub> on C<sub>4</sub> metabolism; rather it reflects the effect of [O<sub>2</sub>] on reducing photosynthesis in the large-celled chloroplast-containing parenchyma (Kim and Fisher, 1990) in which C<sub>3</sub> photosynthesis presumably occurs in well-watered plants, and in which CAM is induced when the plants are drought-stressed.

Christin *et al.* (2014) suggest that the occurrence of CAM and C<sub>4</sub> in *Portulaca* is the product of a partially shared evolutionary trajectory in which *Portulaca* was ancestrally a C<sub>3</sub>–CAM plant. C<sub>4</sub> photosynthesis subsequently evolved multiple times while a functional CAM cycle was maintained. For enzymes other than PEPc, *Portulaca* co-opted the ancestral CAM genes for C<sub>4</sub> photosynthesis, but the C<sub>4</sub> PEPc genes appear to have arisen via *Portulaca*-specific gene duplication, and were independently optimized in each *Portulaca* clade. It is possible that the *Cryptopetala* clade may represent an ancestral C<sub>3</sub>–CAM state common to all extant *Portulaca* (but see Hancock and Edwards (2014) for challenges to this type of inference).




**Fig. 6.** Titratable acidity in recently fully expanded leaves of *Portulaca molokiniensis* at the end of the 12 h light period (red) and at the end of the 12 h dark period (blue) in plants that were well-watered (left-hand column), droughted (middle column; 12 d without irrigation), and droughted and rewatered (right-hand column; 6 d with irrigation). The data are expressed on a fresh mass basis (A), a dry mass basis (B), and a leaf area basis (C). Mean  $\pm$ SD ( $n=5$  leaves; at a given time point each leaf was harvested from a different plant). The numeric values shown above the bars are  $P$  values (one-tailed  $t$ -test). Bold letters indicate that the values at the end of the dark period were significantly greater than those at the end of the day at  $P \leq 0.05$ .

Nevertheless, the presence of CAM but not full C<sub>4</sub> in *P. cryptopetala* is consistent with the notion that in *Portulaca* CAM is an ancestral state that has persisted despite the subsequent repeated evolution of C<sub>4</sub> photosynthesis.

As is the case for leaves in the C<sub>3</sub>-C<sub>4</sub> *P. cryptopetala*, the stems of *Portulaca* species in general lack Kranz anatomy and the C<sub>4</sub> pathway (Voznesenskaya *et al.*, 2010). Observations of nocturnal acidification in stems as well as leaves of the C<sub>3</sub>-C<sub>4</sub> *P. cryptopetala* (Figs 3, 4) and the C<sub>4</sub> species *P. oleracea* (Koch and Kennedy, 1980) and *P. grandiflora* (Guralnick *et al.*, 2002) may also be taken as evidence in support of the concept of a pre-C<sub>4</sub> presence of CAM in *Portulaca*.

There is little sign that the evolution of C<sub>4</sub> in organs with CAM has systematically enhanced or retarded the CAM phenotype in *Portulaca*. Nocturnal acidification in leaves of the C<sub>3</sub>-C<sub>4</sub> *P. cryptopetala* does not overly differ in magnitude from



**Fig. 7.** Phylogenetic relationships of species within the genus *Portulaca*, based upon the analyses of Ocampo *et al.* (2013) and Moore *et al.* (2018), showing the currently known distribution of C<sub>4</sub> (red) and C<sub>3</sub>-C<sub>4</sub> (blue) photosynthesis among them. The C<sub>4</sub> distribution is from Ocampo *et al.* (2013), and the CAM distribution is from Koch and Kennedy (1980), Guralnick *et al.* (2002), Winter and Holtum (2017), Holtum *et al.* (2017a), Winter (2019), and the current study.

acidification in the C<sub>4</sub>-CAM species in the other three clades known with CAM. Nocturnal acid accumulation in leaves of *P. cryptopetala* of ca. 100  $\mu\text{mol H}^+ \text{g}^{-1} \text{FM}$  (Fig. 3A) is comparable to values reported for *P. oleracea* (Koch and Kennedy, 1980) and *P. grandiflora* (Guralnick *et al.*, 2002), but greater than levels of ca. 75  $\mu\text{mol H}^+ \text{g}^{-1} \text{FM}$  reported for *P. australis*, *P. digyna*, *P. molokiniensis*, and *P. pilosa*, and far in excess of the 8  $\mu\text{mol H}^+ \text{g}^{-1} \text{FM}$  reported for *P. cyclophylla* (Holtum *et al.*, 2017b; Winter and Holtum, 2017). Although water-stressed *P. molokiniensis* (Fig. 6) accumulated less acid at night than did water-stressed *P. cryptopetala*, in terms of the absolute acidity stored in tissues, the acid levels in *P. molokiniensis* were greater. The reason for the difference was that following the imposition of stress, the background levels of acid increased in *P. molokiniensis* but not in *P. cryptopetala*. To further address the question of possible differences in the capacity for nocturnal acid accumulation between different species of *Portulaca*, a rigorous comparison of acid levels from a wide range of species growing under identical conditions is warranted.

Similarities exist between the expression of CAM in *Portulaca* (Portulacaceae) and in the Australian *Calandrinia* (Montiaceae) (Winter and Holtum, 2011; Holtum *et al.*, 2017b; Hancock *et al.*, 2018). Both are located in the sub-order Portulacineae (Carophyllales) where they nest among lineages in which CAM and succulence are common (Moore *et al.*, 2018; Ogburn and Edwards, 2013), and both are mainly composed of small, short-lived, succulent-leaved herbs of open arid to semi-arid sites (Eggle 2004; Nyffeler *et al.*, 2008; Kapitany, 2007). Indeed, in Australia it is not uncommon to see species of *Portulaca* and *Calandrinia* growing alongside each other. Facultative CAM appears widespread in both groups but, although full C<sub>4</sub> is present in *Portulaca*, there is currently no evidence of strong constitutive CAM in either lineage, despite both having diverged from their respective progenitors around 30 Ma ago (Arakaki

*et al.*, 2011; Hancock *et al.*, 2018). In each of the lineages, it is unclear why full CAM has not evolved but facultative CAM has. The answer undoubtedly lies in historical contingencies that are the products of interactions between genetic composition and ecological opportunity over space and time (Edwards and Donoghue, 2013; Christin *et al.*, 2014, 2015).

In the case of the  $C_4$  pathway, detailed analyses of phylogeny, anatomy, genes, and physiological phenotypes in the ~40  $C_3$ – $C_4$  intermediates known from ca. 20 monocot and eudicot genera has markedly assisted conceptualization of the importance of parallel and convergent evolution to the multiple emergence of the  $C_4$  pathway, and of the processes that constrain and enable it (Sage *et al.*, 2011, 2014; Christin *et al.*, 2015). If plants with low-level CAM or facultative CAM are the CAM equivalent of  $C_3$ – $C_4$  intermediates, then many more  $C_3$ –CAM intermediates are known than are  $C_3$ – $C_4$  intermediates (Winter *et al.*, 2015). Presumably, as has been demonstrated for the  $C_4$  pathway intermediates, the  $C_3$ –CAM intermediates contain a subset of the anatomical and biochemical components of the CAM  $CO_2$  pump that improve physiological performance over the  $C_3$  system in the places where the plants are found (Heckmann, 2016). Addressing the core questions of CAM origins and expression will benefit from rigorous comparisons across lineages of genes and traits that have been acquired repeatedly during evolution of CAM.

## Supplementary data

Supplementary data are available at *JXB* online.

Fig. S1. Fourteen days of net  $CO_2$  exchange of *Portulaca cryptopetala* during a wet–dry–wet cycle.

Fig. S2. Eleven days of net  $CO_2$  exchange of *Portulaca molokiniensis* during a wet–dry–wet cycle.

Fig. S3. Sixteen days of net  $CO_2$  exchange of *Portulaca molokiniensis* during a wet–dry–wet cycle.

Fig. S4. Twelve days of net  $CO_2$  exchange of *Portulaca molokiniensis* during a wet–dry–wet cycle.

## Acknowledgements

The research was supported by Australian Research Council Discovery Grant DP160100098, US National Science Foundation grant DEB-1252901, and by the Smithsonian Tropical Research Institute.

## References

- Arakaki M, Christin PA, Nyffeler R, Lendel A, Eggli U, Ogburn RM, Spriggs E, Moore M, Edwards EJ. 2011. Contemporaneous and recent radiations of the world's major succulent plant lineages. *Proceedings of the National Academy of Sciences, USA* **108**, 8379–8384.
- Christin PA, Arakaki M, Osborne CP, *et al.* 2014. Shared origins of a key enzyme during the evolution of  $C_4$  and CAM metabolism. *Journal of Experimental Botany* **65**, 3609–3621.
- Christin PA, Arakaki M, Osborne CP, Edwards EJ. 2015. Genetic enablers underlying the clustered evolutionary origins of  $C_4$  photosynthesis in angiosperms. *Molecular Biology and Evolution* **32**, 846–858.
- D'Andrea RM, Andreo CS, Lara MV. 2014. Deciphering the mechanisms involved in *Portulaca oleracea* ( $C_4$ ) response to drought: metabolic changes including crassulacean acid-like metabolism induction and reversal upon re-watering. *Physiologia Plantarum* **152**, 414–430.
- Edwards EJ, Donoghue MJ. 2013. Is it easy to move and easy to evolve? Evolutionary accessibility and adaptation. *Journal of Experimental Botany* **64**, 4047–4052.
- Edwards EJ, Ogburn RM. 2012. Angiosperm responses to a low  $CO_2$  world: CAM and  $C_4$  photosynthesis as parallel evolutionary trajectories. *International Journal of Plant Science* **173**, 724–733.
- Edwards GE, Vozsenskaya E. 2011.  $C_4$  photosynthesis: Kranz forms and single-cell  $C_4$  in terrestrial plants. In: Raghavendra AS, Sage RF, eds.  $C_4$  photosynthesis and related  $CO_2$  concentrating mechanisms. Dordrecht: Springer, 29–61.
- Eggli U. 2004. *Illustrated handbook of succulent plants: dicotyledons*. Berlin, Heidelberg: Springer Verlag.
- Guralnick LJ, Edwards GE, Ku MSB, Hockema B, Franceschi VR. 2002. Photosynthetic and anatomical characteristics in the  $C_4$ -crassulacean acid metabolism-cycling plant, *Portulaca grandiflora*. *Functional Plant Biology* **29**, 763–773.
- Guralnick LJ, Jackson MD. 2001. The occurrence and phylogenetics of crassulacean acid metabolism in the Portulacaceae. *International Journal of Plant Sciences* **162**, 257–262.
- Hancock L, Edwards EJ. 2014. Phylogeny and the inference of evolutionary trajectories. *Journal of Experimental Botany* **65**, 3491–3498.
- Hancock LP, Obbens F, Moore AJ, Thiele K, de Vos JM, West J, Holtum JAM, Edwards EJ. 2018. Phylogeny, evolution, and biogeographic history of *Calandrinia* (Montiaceae). *American Journal of Botany* **105**, 1021–1034.
- Hatch MD. 1987.  $C_4$  photosynthesis, a unique blend of modified biochemistry, anatomy and ultrastructure. *Biochimica et Biophysica Acta* **895**, 81–106.
- Heckmann D. 2016.  $C_4$  photosynthesis evolution: the conditional Mt. Fuji. *Current Opinion in Plant Biology* **31**, 149–154.
- Hobdy RW. 1987. *Portulaca molokiniensis* (Portulacaceae); a new species from the Hawaiian Islands. *Pacific Science* **41**, 64–67.
- Holtum JAM, Hancock LP, Edwards EJ, Winter K. 2017a. Optional use of CAM photosynthesis in two  $C_4$  species, *Portulaca cyclophylla* and *Portulaca digyna*. *Journal of Plant Physiology* **214**, 91–96.
- Holtum JAM, Hancock LP, Edwards EJ, Winter K. 2017b. Facultative CAM photosynthesis (crassulacean acid metabolism) in four species of *Calandrinia*, ephemeral succulents of arid Australia. *Photosynthesis Research* **134**, 17–25.
- Holtum JAM, Hancock LP, Edwards EJ, Winter K. 2018. Crassulacean acid metabolism in the Basellaceae (Caryophyllales). *Plant Biology* **20**, 409–414.
- Holtum JA, Winter K. 2003. Photosynthetic  $CO_2$  uptake in seedlings of two tropical tree species exposed to oscillating elevated concentrations of  $CO_2$ . *Planta* **218**, 152–158.
- Kapitany A. 2007. *Australian succulent plants*. Boronia, Victoria, Australia: Kapitany Concepts.
- Keerberg O, Pärnik T, Ivanova H, Bassüner B, Bauwe H. 2014.  $C_2$  photosynthesis generates about 3-fold elevated leaf  $CO_2$  levels in the  $C_3$ – $C_4$  intermediate species *Flaveria pubescens*. *Journal of Experimental Botany* **65**, 3649–3656.
- Kim I, Fisher DG. 1990. Structural aspects of the leaves of seven species of *Portulaca* growing in Hawaii. *Canadian Journal of Botany* **68**, 1803–1811.
- Koch K, Kennedy RA. 1980. Characteristics of crassulacean acid metabolism in the succulent  $C_4$  dicot, *Portulaca oleracea* L. *Plant Physiology* **65**, 193–197.
- Lara MV, Disante KB, Podestá FE, Andreo CS, Drincovich MF. 2003. Induction of a Crassulacean acid like metabolism in the  $C_4$  succulent plant, *Portulaca oleracea* L.: physiological and morphological changes are accompanied by specific modifications in phosphoenolpyruvate carboxylase. *Photosynthesis Research* **77**, 241–254.
- Lara MV, Drincovich MF, Andreo CS. 2004. Induction of a crassulacean acid-like metabolism in the  $C_4$  succulent plant, *Portulaca oleracea* L.: study of enzymes involved in carbon fixation and carbohydrate metabolism. *Plant & Cell Physiology* **45**, 618–626.
- Maroco JP, Ku MSB, Edwards GE. 1997. Oxygen sensitivity of  $C_4$  photosynthesis: evidence from gas exchange and fluorescence analysis with different  $C_4$  sub-types. *Plant, Cell & Environment* **20**, 1525–1533.
- Maroco JP, Ku MSB, Lea PJ, Dever LV, Leegood RC, Furbank RT, Edwards GE. 1998. Oxygen requirement and inhibition of  $C_4$  photosynthesis. An analysis of  $C_4$  plants deficient in the  $C_3$  and  $C_4$  cycles. *Plant Physiology* **116**, 823–832.



- Moore AJ, De Vos JM, Hancock LP, Goolsby E, Edwards EJ.** 2018. Targeted enrichment of large gene families for phylogenetic inference: phylogeny and molecular evolution of photosynthesis genes in the Portulacaceae (Caryophyllales). *Systematic Biology* **67**, 367–383.
- Nyffeler R, Egli U, Ogburn M, Edwards EJ.** 2008. Variations on a theme: repeated evolution of succulent life forms in the Portulacaceae (Caryophyllales). *Haseltonia* **14**, 26–36.
- Ocampo G, Columbus JT.** 2012. Molecular phylogenetics, historical biogeography, and chromosome number evolution of *Portulaca* (Portulacaceae). *Molecular Phylogenetics and Evolution* **63**, 97–112.
- Ocampo G, Koteyeva NK, Voznesenskaya EV, Edwards GE, Sage TL, Sage RF, Columbus JT.** 2013. Evolution of leaf anatomy and photosynthetic pathways in Portulacaceae. *American Journal of Botany* **100**, 2388–2402.
- Ogburn RM, Edwards EJ.** 2010. The ecological water-use strategies of succulent plants. *Advances in Botanical Research* **55**, 179–225.
- Ogburn RM, Edwards EJ.** 2012. Quantifying succulence: a rapid, physiologically meaningful metric of plant water storage. *Plant, Cell & Environment* **35**, 1533–1542.
- Ogburn RM, Edwards EJ.** 2013. Repeated origin of three-dimensional leaf venation releases constraints on the evolution of succulence in plants. *Current Biology* **23**, 722–726.
- Osmond CB.** 1978. Crassulacean acid metabolism: a curiosity in context. *Annual Review of Plant Physiology* **29**, 379–414.
- Sage RF.** 2002. Are crassulacean acid metabolism and C<sub>4</sub> photosynthesis incompatible? *Functional Plant Biology* **29**, 775–785.
- Sage RF.** 2016. A portrait of the C<sub>4</sub> photosynthetic family on the 50<sup>th</sup> anniversary of its discovery: species number, evolutionary lineages, and Hall of Fame. *Journal of Experimental Botany* **67**, 4039–4056.
- Sage RF, Christin PA, Edwards EJ.** 2011. The C<sub>4</sub> plant lineages of planet Earth. *Journal of Experimental Botany* **62**, 3155–3169.
- Sage RF, Khoshravesh R, Sage TL.** 2014. From proto-Kranz to C<sub>4</sub> Kranz: building the bridge to C<sub>4</sub> photosynthesis. *Journal of Experimental Botany* **65**, 3341–3356.
- Sage RF, Sultmanis S.** 2016. Why are there no C<sub>4</sub> forests? *Journal of Plant Physiology* **203**, 55–68.
- Smith JAC, Winter K.** 1996. Taxonomic distribution of crassulacean acid metabolism. In: Winter K, Smith JAC, eds. *Crassulacean acid metabolism*. Berlin, Heidelberg: Springer Verlag, 427–436.
- Voznesenskaya EV, Koteyeva NK, Edwards GE, Ocampo G.** 2010. Revealing diversity in structural and biochemical forms of C<sub>4</sub> photosynthesis and a C<sub>3</sub>-C<sub>4</sub> intermediate in genus *Portulaca* L. (Portulacaceae). *Journal of Experimental Botany* **61**, 3647–3662.
- Voznesenskaya EV, Koteyeva NK, Edwards GE, Ocampo G.** 2017. Unique photosynthetic phenotypes in *Portulaca* (Portulacaceae): C<sub>3</sub>-C<sub>4</sub> intermediates and NAD-ME C<sub>4</sub> species with Piloid-type Kranz anatomy. *Journal of Experimental Botany* **68**, 225–239.
- Winter K.** 2019. Ecophysiology of constitutive and facultative CAM photosynthesis. *Journal of Experimental Botany* **70**, 6495–6508.
- Winter K, Holtum JA.** 2007. Environment or development? Lifetime net CO<sub>2</sub> exchange and control of the expression of crassulacean acid metabolism in *Mesembryanthemum crystallinum*. *Plant Physiology* **143**, 98–107.
- Winter K, Holtum JAM.** 2011. Induction and reversal of crassulacean acid metabolism in *Calandrinia polyandra*: effects of soil moisture and nutrients. *Functional Plant Biology* **38**, 576–582.
- Winter K, Holtum JA.** 2014. Facultative crassulacean acid metabolism (CAM) plants: powerful tools for unravelling the functional elements of CAM photosynthesis. *Journal of Experimental Botany* **65**, 3425–3441.
- Winter K, Holtum JAM.** 2017. CO<sub>2</sub>-exchange patterns demonstrate facultative CAM photosynthesis (crassulacean acid metabolism) in four small Australian C<sub>3</sub> and C<sub>4</sub> leaf-succulents. *Australian Journal of Botany* **65**, 103–108.
- Winter K, Holtum JA, Smith JA.** 2015. Crassulacean acid metabolism: a continuous or discrete trait? *New Phytologist* **208**, 73–78.
- Winter K, Ziegler H.** 1992. Induction of crassulacean acid metabolism in *Mesembryanthemum crystallinum* increases reproductive success under conditions of drought and salinity stress. *Oecologia* **92**, 475–479.