



HHS Public Access

Author manuscript

Cell. Author manuscript; available in PMC 2020 October 17.

Published in final edited form as:

Cell. 2019 October 17; 179(3): 787–799.e17. doi:10.1016/j.cell.2019.09.016.

Optical pooled screens in human cells

David Feldman^{1,2,10}, Avtar Singh^{1,10}, Jonathan L. Schmid-Burgk¹, Rebecca J. Carlson^{1,3}, Anja Mezger^{1,4}, Anthony J. Garrity¹, Feng Zhang^{1,5,6,7,8}, Paul C. Blainey^{1,5,9,11,*}

¹Broad Institute of MIT and Harvard, Cambridge, Massachusetts, 02142, USA.

²Department of Physics, MIT, Cambridge, Massachusetts, 02142, USA.

³Department of Health Sciences and Technology, MIT, Cambridge, Massachusetts, 02142, USA.

⁴Department of Medical Biochemistry and Biophysics, Karolinska Institute, Stockholm, Sweden.

⁵Department of Biological Engineering, MIT, Cambridge, Massachusetts, 02142, USA.

⁶McGovern Institute for Brain Research at MIT, Cambridge, Massachusetts, 02142, USA.

⁷Department of Brain and Cognitive Science, MIT, Cambridge, Massachusetts, 02142, USA.

⁸Howard Hughes Medical Institute, MIT, Cambridge, Massachusetts, 02142, USA.

⁹Koch Institute for Integrative Cancer Research, MIT, Cambridge, Massachusetts, 02142, USA.

¹⁰These authors contributed equally to this work.

¹¹Lead contact.

SUMMARY

Genetic screens are critical for the systematic identification of genes underlying cellular phenotypes. Pooling gene perturbations greatly improves scalability, but is not compatible with imaging of complex and dynamic cellular phenotypes. Here, we introduce a pooled approach for optical genetic screens in mammalian cells. We use targeted *in situ* sequencing to demultiplex a library of genetic perturbations following image-based phenotyping. We screened a set of 952 genes across millions of cells for involvement in NF- κ B signaling by imaging the translocation of RelA (p65) to the nucleus. Screening at a single time point across 3 cell lines recovered 15 known

*Correspondence to: pblainey@broadinstitute.org.

AUTHOR CONTRIBUTIONS

D.F. designed the approach with input from all authors. D.F., A.S., A.M., A.J.G., R.J.C. and J.S.B performed experiments. D.F. analyzed data. J.S.B and D.F. designed the NF-KB screen. P.C.B. and F.Z. supervised the research. D.F., A.S., J.S.B, and P.C.B. wrote the manuscript with contributions from all authors.

DATA AND CODE AVAILABILITY

Python scripts, including a complete pipeline from example data to sequencing reads, will be made available, along with plasmid sequences and updated protocols at <https://github.com/blaineylab/OpticalPooledScreens>. RNA-seq data is deposited in the Gene Expression Omnibus (GSE132704). Raw and processed image data is available on request.

DECLARATION OF INTERESTS


J.S.B and A.M. declare no competing interests. A.M is an employee at AstraZeneca. D.F. is an employee at insitro. A.J.G. is an employee of and holds equity in Arbor Biotechnologies, Inc. P.C.B is a consultant to and holds equity in 10X Genomics, General Automation Lab Technologies, Celsius Therapeutics, and Next Gen Diagnostics, LLC. P.C.B is a consultant to insitro. F.Z. is a founder, consultant to and holds equity in Arbor Biotechnologies, Beam Therapeutics, Editas Medicine, Pairwise Plants, and Sherlock Biosciences. The Broad Institute and MIT have filed U.S. patent applications on work described in this manuscript and may seek to license the technology. P.C.B, D.F., R.J.C., A.S. are listed as inventors.

pathway components, while repeating the screen with live-cell imaging revealed a role for Mediator complex subunits in regulating the duration of p65 nuclear retention. These results establish a highly multiplexed approach to image-based screens of spatially and temporally defined phenotypes with pooled libraries.

In Brief

A screening approach that combines high-content imaging with in situ sequencing can identify genes that affect spatially and temporally defined phenotypes like morphology and subcellular localization, expanding the list of scientific questions that can be asked with CRISPR-based tools.

Graphical Abstract



INTRODUCTION

Genetic perturbation screens are powerful tools for establishing causal links between genes and phenotypes in mammalian systems. Pooling genetic perturbations has greatly improved the ability to perform screens of many genetic elements but has been limited to enrichment-based phenotypic assays (Gilbert et al., 2014; Parnas et al., 2015; Shalem et al., 2014; Wang et al., 2014) or single-cell molecular profiling (Adamson et al., 2018; Datlinger et al., 2017; Dixit et al., 2016; Jaitin et al., 2016; Rubin et al., 2019). Conversely, imaging can capture a rich set of phenotypes at scale, including dynamics, but has not been compatible with pooled approaches, limiting the ability to systematically screen for genes that impact cell function.

Pooling genetic perturbations offers several advantages, making it practical to conduct comprehensive, genome-scale screens to find relevant components in a biological process. Pooled libraries can be constructed, introduced into cells, and read out as single samples, dramatically reducing the cost, effort and time needed to perform large-scale screens. Moreover, combining all perturbations into a single sample reduces batch effects and improves statistical power since all perturbed and control cells experience the same conditions. These advantages make pooled screening a preferred method in the biological community.

Historically, pooled screens have relied on enriching populations of cells for a phenotype of interest, followed by next-generation sequencing (NGS) to measure changes in perturbation abundance. Common enrichment-based phenotypes include differential cell fitness (e.g., under drug selection) (Shalem et al., 2014; Wang et al., 2014) and differential fluorescence of a marker (e.g., a genetic reporter or immunostained protein), followed by separation of a target population via fluorescence-activated cell sorting (FACS) (Parnas et al., 2015). Although highly scalable, these approaches are limited to population-level measurement,

often rely on indirect reporters for the biological activity of interest, and necessarily limit phenotypes to one or a few parameters. Recently, pooled perturbations were integrated with single cell RNA-seq (Adamson et al., 2016; Datlinger et al., 2017; Dixit et al., 2016; Jaitin et al., 2016), and subsequently with chromatin accessibility profiling by ATAC-seq and protein detection by mass cytometry (Rubin et al., 2019; Wroblewska et al., 2018) to achieve high-dimensional readout for pooled screens. These molecular profiling screens capture high-dimensional representations of cell state, yielding more information about perturbation effects. However, the relationship between RNA or other molecular profiles and cellular functions is often unknown. Moreover, existing single cell molecular profiling approaches are destructive and thus cannot be used to directly monitor dynamic processes over time in individual cells.

Imaging offers an attractive alternative as it can collect many spatially and temporally resolved parameters from millions of individual cells. A multitude of optical assays have been optimized for genetic screens of protein localization, enzyme activity, metabolic state, molecular/cellular dynamics and cellular morphology in a variety of biological contexts, including mitosis (Moffat et al., 2006; Neumann et al., 2010), endocytosis (Collinet et al., 2010), viral infection (Karlas et al., 2010), differentiation (Chia et al., 2010), metabolism (Guo et al., 2008), DNA damage (Floyd et al., 2013), autophagy (Orvedahl et al., 2011) and synaptogenesis (Linhoff et al., 2009). However, these and other high-content screens required expensive and laborious testing of arrayed perturbations. Image-based screening of pooled perturbations has thus far only been demonstrated with fluorescence *in situ* hybridization (FISH)-based approaches that required high-magnification imaging with limited scalability for mammalian applications (Lawson et al., 2017; Wang et al., 2019).

Here, we report an optical pooled screening method that greatly expands the range of phenotypes amenable to large-scale pooled perturbation screens in mammalian cells. Our approach uses microscopy to determine first the phenotype and then the perturbation identity in each cell. We identify perturbations by targeted *in situ* sequencing (Ke et al., 2013) of either compact perturbations (e.g., CRISPR sgRNAs) or short associated barcodes. *In situ* sequencing uses enzymatic amplification to generate high signal levels, permitting imaging at low magnification, which made it feasible for us to screen millions of cells from a pooled library. We first vetted this approach by screening a synthetic reporter activated by CRISPR-induced mutations to estimate the accuracy of mapping perturbations to phenotypes *in situ*. Next, we performed optical pooled CRISPR loss-of-function screens across three cell lines to study p65 nuclear translocation in response to pro-inflammatory signals, identifying core NF- κ B pathway members and additional factors as hits. We used optical pooled screens with time-lapse imaging to uncover a role for Mediator components as regulators of p65 nuclear retention time and subsequently found that *MED12/MED24* knockouts led to sustained expression of key NF- κ B target genes. The approach demonstrated here paves the way to pooled image-based screens of genome-scale libraries.

RESULTS

A single integrated genetic perturbation can be identified by *in situ* RNA sequencing

We developed an optical barcoding strategy to enable pooled screens of microscopy-based phenotypes (Figure 1A). In our approach, perturbations are identified via targeted *in situ* sequencing of an expressed barcode (Figure 1B). As pooled screens in mammalian cells generally use single-copy lentiviral integration to deliver perturbations, we focused on establishing reliable *in situ* sequencing of barcode transcripts in this format. To optically barcode CRISPR perturbations, we modified an existing lentiviral CRISPR guide RNA (sgRNA) expression vector (LentiGuide-Puro) to express both an sgRNA and a 12-nt barcode, and termed the resulting vector LentiGuide-BC. The barcodes and constant flanking sequences were inserted into the 3' UTR of the Pol II-transcribed antibiotic resistance gene, a highly expressed mRNA suitable for *in situ* detection.

To test *in situ* identification of perturbations, we transduced a LentiGuide-BC library containing 40 sgRNA- barcode pairs into HeLa-TetR-Cas9 cells at low multiplicity of infection (MOI < 0.01). We prepared samples for targeted *in situ* sequencing using a padlock-based approach (Ke et al., 2013) in which variable sequences within an RNA transcript are converted to single-stranded DNA and enzymatically amplified via *in situ* reverse transcription, padlock extension/ligation, and rolling circle amplification (RCA) (Figures 1B, 2A and S1A). Sequencing the amplified DNA with a 4-color sequencing-by-synthesis chemistry over 12 cycles generated high quality reads with excellent uniformity among barcodes (Figure 2B and S1B-G). After image segmentation and base calling analysis, more than 85% of sequence reads mapped exactly to one of the 40 expected barcodes out of the $4^{12} = 16.7$ million possible 12-nt sequences (Figure 2C, STAR Methods).

Perturbation detection *in situ* is compatible with the demands of large screens

Next, we demonstrated that an *in situ* sequencing approach can meet the demands of large pooled screens: (1) measurement across millions of cells quickly; (2) high efficiency and accuracy of perturbation mapping per cell; (3) detection of one or more perturbations delivered independently; and (4) compatibility with a large number of perturbations.

We showed that the *in situ* readout step can process millions of cells within a few days to provide the high coverage (typically 100–1,000 cells/perturbation) needed for high-throughput optical screening. High signal intensity allows accurate sequence data to be obtained at low optical magnification across large fields of view, each containing thousands of cells (Figure 2B, Movie S1, Table S1). We maximized fluorescence signal-to-background by optimizing the barcode amplification protocol, including the post-fixation step that follows reverse transcription and the conditions for padlock extension/ligation (Figure S2A). Using the optimized protocol, *in situ* sequencing spots were readily visible at 10X optical magnification, with at least one exactly mapped read detected in more than 77% of transduced cells (Figures 2B, 2D and 2E).

Next, we constructed two types of large barcoded perturbation libraries using oligo microarray synthesis. In the first approach, we designed a set of 83,314 12-nt barcodes using

criteria that balanced GC content, minimized homopolymer repeats, and maintained a minimum edit distance between barcodes of three (STAR Methods). This allows rejection of reads containing up to two single base insertion/deletion/substitution errors arising from oligo synthesis or *in situ* processing (Figure 2F, STAR Methods) (Buschmann and Bystrykh, 2013). We then used a two-step procedure to clone a library of sgRNAs and associated barcodes into LentiGuide-BC (STAR Methods). In contrast to perturbation barcoding by random pairing (Wang et al., 2019), this approach pairs sgRNAs with specific barcodes *in silico*, ensuring efficient use of available barcodes. We addressed a known problem whereby barcode-sgRNA associations are swapped due to reverse transcription-mediated recombination during lentiviral infection (Adamson et al., 2018; Hill et al., 2018; Sack et al., 2016; Xie et al., 2018) by using a modified lentiviral packaging protocol (Feldman et al., 2018) that reduced the frequency of cells exhibiting swapped barcodes from >28% to <5%.

In the second approach, we used the CROP-seq vector to directly sequence a Pol II-transcribed copy of the sgRNA, rather than relying on auxiliary barcodes (Datlinger et al., 2017). We observed accurate sequencing of sgRNAs in HeLa-TetR-Cas9 cells, but with reduced fluorescent signal intensity compared to LentiGuide-BC. We then systematically tested a set of 84 CROP-seq-targeting padlocks with a range of padlock binding arm sequences, padlock backbone lengths and backbone sequences (STAR Methods, Figure S2B, Table S2) to yield a top-performing CROP-seq padlock with a 4-fold increase in reads per cell and 1.7-fold increase in signal intensity relative to the median padlock.

Moreover, we showed that *in situ* sequencing could be used to read out multiple perturbations within the same cell. To this end, we transduced HeLa-TetR-Cas9 cells with one CROP-seq library carrying a puromycin selection marker, followed by a second CROP-seq library carrying a zeocin selection marker. After serial transduction and antibiotic selection, we performed *in situ* sequencing on both libraries simultaneously. Most cells (81%) contained reads mapping to two sgRNAs (Figure S2C,D). The ability to independently sequence perturbations delivered by separate vectors suggests a straightforward and general route to higher-order combinatorial screens.

Accurate mapping of phenotype to genotype in an optical pooled screen

We next showed that our approach can correctly map genetic perturbations to cell phenotypes *in situ* by performing a reporter imaging screen, in which a lentiviral reporter produces an HA-tagged, nuclear-localized H2B protein after a Cas9-induced +1 frameshift in a target region (Figures 3A and S3A, STAR Methods). Cells expressing the reporter can either be screened *in situ* or by FACS. The reporter is highly specific and sensitive, with a mean *in situ* activation across 5 targeting sgRNAs of $65 \pm 2.7\%$ and background of <0.001 % in the absence of a targeting sgRNA (Figures 3B and S3B). We transduced cells stably expressing the frameshift reporter with a LentiGuide-BC library containing 972 barcodes redundantly encoding 5 targeting and 5 control sgRNAs (average of 97 barcodes per sgRNA) (Table S3). We then induced Cas9 expression, measured reporter activation by immunofluorescence, and determined barcode sequences by *in situ* sequencing.

All barcodes encoding targeting sgRNAs were distinguishable from control sgRNAs by HA⁺ fraction, with a per-cell identification accuracy of 90.6% (based on false positive events in

which HA⁺ cells were assigned control sgRNAs, STAR Methods). Indeed, there were no errors in barcode-to-phenotype assignment even for barcodes represented by very few cells (Figures 3B and S3C, STAR Methods). Screening the same cell library by FACS with NGS readout showed similar enrichment of targeting sgRNAs (Figures 3C and S3D), with consistent representation of most barcodes in both contexts (95% within 5-fold abundance, Figure S1). We achieved comparably robust mapping of CRISPR sgRNAs to the frameshift reporter phenotype with the CROP-seq vector in HeLa cells as well as U2-OS, HCT116, A375, HT1080 and HEK293 cells (Figure 3D), with per-cell identification accuracy ranging from 83–98%. Thus, the combined errors that may arise from oligonucleotide synthesis, library cloning, lentiviral delivery, barcode diffusion during *in situ* processing, barcode readout by *in situ* sequencing, or incorrect assignment of reads to cells during image processing are infrequent enough to permit pooled functional screens.

An optical pooled screen in millions of cells for regulators of NF- κ B activation

After demonstrating the ability to screen genetic knockouts for optical phenotypes, we sought to identify genes required for activation of NF- κ B, a family of transcription factors (p50, p52, p65, RelB and c-Rel) that translocate to the nucleus in response to a host of stimuli (Gewurz et al., 2012; Hayden and Ghosh, 2012; Pahl, 1999). At baseline, NF- κ B dimers are maintained in an inactive state in the cytoplasm by inhibitory I κ B proteins, which mask nuclear localization signals to prevent NF- κ B translocation into the nucleus. Upon stimulation of cell surface receptors, a signal cascade is initiated, leading to downstream activation of the IKK complex. IKK- β then phosphorylates I κ B proteins, triggering their phosphorylation-dependent ubiquitination and subsequent degradation by the proteasome, releasing NF- κ B from inhibition. Free NF- κ B dimers may then translocate to the nucleus and induce the expression of genes promoting cell proliferation, survival, and pro-inflammatory responses. It is known that post-translational modifications play a key role in regulating the baseline state and activation of NF- κ B. In addition to its role in proteolytic degradation, ubiquitin has been shown to function in IKK activation and the activity of many core NF- κ B pathway members is controlled by alterations in ubiquitination state.

We used an established nuclear translocation assay to measure the localization of a p65-mNeonGreen reporter in HeLa cells following stimulation with either IL-1 β or TNF α , cytokines that activate NF- κ B via different pathways (Figure 4A). We screened 3,063 sgRNAs targeting 963 genes using the LentiGuide-BC design. We included all GO-annotated ubiquitin ligase and deubiquitinase enzymes, as well as 425 immune-related genes in the library, hypothesizing that ubiquitin signaling may play as-yet unrecognized roles in NF- κ B activation and relaxation (return to the baseline cellular distribution) (Chen, 2005) (Figure 4A, Table S3). After stimulation with either IL-1 β or TNF α , we imaged p65-mNeonGreen translocation, and then performed *in situ* sequencing to identify sgRNAs. A total of 3,037,909 cells were retained for analysis after filtering cells based on reporter expression, nuclear morphology, and exact barcode mapping (952 out of 963 genes retained for analysis) (Figure S3E). We scored the degree of p65-mNeonGreen translocation in each cell by cross-correlation with a DAPI nuclear stain. We ranked the perturbations by the difference of their translocation score distribution from negative control sgRNAs to identify

gene knockouts that led to defects in response to IL-1 β and/or TNF α (Figures 4B, 4C and S4A, Tables S1 and S3, STAR Methods).

The screen recovered most known NF- κ B regulators and uncovered novel candidates

The hits in our screen included known pathway components annotated by KEGG (Kanehisa and Goto, 2000) for NF- κ B activation by IL-1 β signaling (5/5 genes), TNF α signaling (4/7 genes) and downstream components (5/7 genes), including cytokine-specific receptors, adaptor proteins, and factors that activate the shared regulator MAP3K7 (Figure 4D) (Gewurz et al., 2012). Hits common to both cytokine stimulations included *MAP3K7* and its target, the IKK complex (*CHUK*, *IKBKB*, *IKBKG*), as well as components of the SKP1-CUL1-F-box ubiquitin ligase complex and proteasome subunits, which together promote degradation of the inhibitor *NFKBIA*/ κ B α and nuclear translocation of p65.

We confirmed the results of the pooled screen by individual CRISPR knockouts, with 19 out of 20 top-ranked hits validated (z-score threshold 3.75, Figure 4E and Table S3). Phenotype strength was well correlated between the primary screening and validation ranks (Spearman's $\rho = 0.84$ and 0.73 for IL-1 β and TNF α , respectively), emphasizing the quantitative nature of the primary screen (Figure S4B). The p65-mNeonGreen screen showed high sensitivity, detecting genes known to be involved in NF- κ B activation that show only modest translocation defects in our model system. This set includes genes such as *RBCK1/HOIL1* (LUBAC complex, poly-ubiquitination of *IKBKG/NEMO* and *RIPK1*), *DCUNID1* (neddylation of CUL1) and *COPS5* (COP9 signalosome, regulates neddylation).

Interestingly, the set of screening hits validated by arrayed knockout using multiple individual sgRNAs included known negative regulators of NF- κ B activation, such as *NFKBIA*/ κ B α and *CYLD*. Knockout of these negative regulators may lead to permanent baseline activation of NF- κ B signaling, which in turn causes negative feedback activation (Jäättelä et al., 1996), potentially rendering perturbed cells refractory to induced NF- κ B activation. The screen also identified several candidate regulators, including *BAP1*, *HCFC1*, and *KCTD5*. Among these, *BAP1* has been previously described to deubiquitinate *HCFC1* (Machida et al., 2009), with relevance for controlling metabolism (Bononi et al., 2017), ER-stress signaling (Dai et al., 2017), cell-cycle progression (Misaghi et al., 2009), and viral gene expression (Johnson et al., 1999).

In order to investigate cell line-dependent wiring of the NF- κ B signaling pathway, we repeated the same screen with an antibody against endogenous p65 in HeLa, A549 and HCT116 cells. The cytokine concentration for each cell line was determined by imaging p65 translocation in wild type cells 40 minutes after stimulation and using the lowest concentration that saturated translocation (STAR Methods). This titration was found to improve screen sensitivity, but the remainder of the screening protocol was identical across cell lines. The antibody screens detected many of the KEGG-annotated regulators found in the HeLa p65-mNeonGreen screen (17/19 in HeLa, 18/19 in A549 and 11/19 in HCT116, Table S3). Notably, some annotated regulators were not detected as hits in any of the three cell lines. Among these, *TRAF5* and *BIRC3* have functionally redundant counterparts (*TRAF2* and *BIRC2*) (Mahoney et al., 2008; Tada et al., 2001). Indeed, *TRAF2* scored as a positive regulator in all three cell lines, while *BIRC2* was only detected in HeLa, suggesting

its role may vary among the different cell lines. Meanwhile, knockdown of *TABI* was previously shown not to impact IKK activation (Chen, 2005; Wang et al., 2001), consistent with our finding that *TABI* was not detected as a hit in any cell line. Comparing high-scoring genes across cell lines revealed that the E2 ubiquitin-conjugating enzyme *UBE2N* was strongly involved in IL-1 β -dependent p65 activation in A549 cells, while being largely dispensable in the other cell lines tested (Figure S4C). Similarly, *BIRC2* was detected only in HeLa cells to be involved in TNF α -dependent p65 translocation. The robustness and scalability of the optical pooled screening protocol makes it possible to conduct in-depth analyses of biological pathways across cellular backgrounds.

High-content analysis of morphology distinguishes regulators by function

In addition to the translocation defect measured in knockouts of NF- κ B positive regulators, we observed that certain classes of genes led to similar morphological changes in cells. For example, proteasomal knockouts induced a distinct morphology consisting of rounded cells (low eccentricity) with enlarged nuclei. We hypothesized that loss of genes with related functions might induce similar morphological changes, providing additional dimensions to interpret our screening results, and thus re-scored each cell in the primary HeLa p65-mNeonGreen screen for cell and nuclear morphology and nuclear stain intensity (STAR Methods). Members of the CUL1-RBX1-SKP1-FBXW11 complex and its substrate *NEDD8* showed increased cell and nuclear area, with 4 out of 5 genes grouped together by PCA (Figures S4D-H). *UFD1*, a member of the VCP/p97-UFD1-NPL4 complex that mediates post-ubiquitination degradation of I κ B α , shared a similar morphological profile to *COP55*, which interacts directly with *VCP/p97* as part of the COP9 signalosome (Cayli et al., 2009; Li et al., 2014). Disruption of the chromatin remodeler *INO80*, as well as *NOP53/GLTSCR2*, and *UBA52*, genes with roles in ribosome biogenesis, showed a decrease in cell and nuclear area. The ability to group classify knockouts into functional categories based on morphological changes is a key benefit of image-based screening, which could be further enhanced by staining additional cellular markers to extract more information from each cell (Gustafsdottir et al., 2013).

A live-cell imaging screen identifies *MED12* and *MED24* as regulators of NF- κ B translocation kinetics

Activation of NF- κ B involves a cascade of signaling events and feedback loops whose kinetics determine the dynamic response of the pathway. As optical pooled screens can be readily combined with live-cell imaging, we screened a CROP-seq library of sgRNAs targeting the same 952 genes (5,638 sgRNAs) detected in the initial activation screen for variability in the timing of p65 activation and relaxation (434,505 cells analyzed, Table S3). Cells were stimulated with IL-1 β or TNF α and imaged at 23 minute intervals for 6 hours post-stimulation. Nuclei were tracked using a cell-permeable DNA stain and p65-mNeonGreen nuclear translocation was assessed at each time point. Following live-cell analysis, cells were fixed and the perturbation in each cell was read out by *in situ* sequencing of the sgRNA sequences. The live-cell screen closely reproduced hits from the fixed-cell activation screen described above, with 15 out of the 20 top positive regulators shared when the live cell analysis was restricted to a single matched time point (STAR Methods).

Hierarchical clustering of mean translocation time profiles revealed distinct populations of positive and negative regulators (Figure 5A). To quantify changes in translocation kinetics, we defined a cumulative defect metric for each gene. First, we integrated over time the difference between each cell's translocation score and the mean translocation score of non-targeting controls. Then, for each gene, the distribution of cell-level cumulative defects was tested for statistically significant deviations from the non-targeting control population (STAR Methods, Table S4). By integrating differences in translocation over time, we were able to identify known key negative regulators of the pathway with effects on p65 relaxation that did not score in the initial static screen, including *TNFAIP3* (an NF- κ B target gene that provides negative feedback by deubiquitinating multiple upstream signaling components) (Chen, 2005), *KEAP1* (a ubiquitin ligase involved in degradation of IKK β) (Lee et al., 2009) and *USP7* (a deubiquitinase that slows ubiquitination and proteasomal degradation of NF- κ B) (Colleran et al., 2013).

Next, we performed a pooled validation screen for 61 genes that showed evidence for a kinetic phenotype (10 sgRNAs/gene). We optimized conditions for live-cell imaging and cell tracking by using faster microscope hardware, allowing us to link time-resolved p65 translocation data to perturbation barcodes from 2,595,514 cells (median of 3,611 cells/sgRNA) at 30 minute intervals. At this high level of sampling, individual sgRNAs targeting a given gene showed excellent concordance (Figure 5B and Figure S5). The pooled validation screen confirmed several genes that were initially detected as lower-ranked hits in the activation screen, while adding additional kinetic details. For example, at later time points, *RIPK1* showed an increased translocation defect while *KCTD5* and *HCFC1* were seen to negatively regulate p65.

The validation screen also confirmed *MED12* and *MED24* as previously unknown negative regulators of p65 nuclear translocation (Figures 5B, 5C and S5). Clonal homozygous knockouts of each of these Mediator complex subunits confirmed delayed relaxation relative to wild type cells (Figure 5D). Of note, a mono-allelic *MED24* knockout clone showed a much weaker phenotype, underscoring the high sensitivity of the primary optical screen despite the heterogeneity of alleles generated by Cas9 cleavage (Figure 5D). We performed RNA sequencing of clonal knockout lines after stimulation with either TNF α or IL-1 β . Among highly induced genes (>50-fold increase in wild type cells after 4 hours of stimulation with either cytokine, adjusted p-value < 10⁻³), we identified 6 genes (*CCL2*, *CCL20*, *CXCL1*, *CXCL2*, *CXCL3*, *CXCL8*) that were differentially expressed in either *MED12* or *MED24* clonal knockout cells (>8-fold change versus wild type cells at 4 hours post-stimulation with either cytokine, adjusted p-value < 10⁻³) (Figure 5E, STAR Methods), all of which encode for chemokines that attract immune cells to sites of inflammation. Even more strikingly, pro-IL-1 β , which is a well-established NF- κ B target gene in the context of Toll-like receptor activation (Sims and Smith, 2010), was transcriptionally induced in *MED12* or *MED24* deficient HeLa cells but not in wild type cells upon TNF α stimulation (Figure 5F). These results suggest that, apart from the well-studied function of the Mediator complex in recruiting active RNA polymerase-II to DNA-bound transcription factors (Malik and Roeder, 2005) including to p65 (Guermah et al., 1998), Mediator components may be involved in restricting the transcriptional response of induced genes during pro-inflammatory NF- κ B signaling.

DISCUSSION

Optical pooled screens enable systematic analysis of the genetic components underpinning a wide range of spatially and temporally defined phenotypes, including subcellular localization, live-cell dynamics, and high-content morphological profiling. The *in situ* sequencing framework is compatible with any perturbation that can be identified by a short expressed sequence and readily scales to millions of cells and genome-scale libraries.

We applied pooled CRISPR loss-of-function screening targeting 952 genes in multiple cell lines to identify regulators of p65 translocation, recovering nearly all annotated regulators of TNF α - and IL-1 β -induced NF- κ B activation. Moreover, by performing a pooled live-cell imaging screen, we discovered two components of the Mediator complex as previously unknown negative regulators of pro-inflammatory signaling that we validated in clonal knockout cell lines. Although the Mediator complex has been implicated in p65-mediated gene transcription (Essen et al., 2009), Mediator components have not previously been shown to be involved in negative regulation of either the nuclear translocation or the transcriptional response of NF- κ B. As one of the most potent pyrogens in the human body, IL-1 β is secreted only upon transcriptional (NF- κ B) as well as post-transcriptional (Caspase 1) licensing. Our results indicated that Mediator components play a role in suppressing pro-IL-1 β induction by TNF α signaling, which might provide a mechanism for preventing aberrant feedback activation of immune signaling pathways (Bauernfeind et al., 2016). The ability to screen dynamic responses, such as p65 translocation kinetics, adds a rich temporal dimension to the profiling of gene knockouts, permitting direct measurement of response onset and duration in addition to strength, directionality (positive/negative) and cytokine specificity.

The use of *in situ* sequencing to read out perturbations has several advantages over alternative approaches. Amplification by RCA enables fast *in situ* sequencing at low magnification (10X), greatly increasing throughput. The 12-nt sequences used here can robustly distinguish >80,000 perturbations, sufficient to encode genome-scale perturbation libraries. Existing CRISPR sgRNA libraries can be read out directly (using the CROP-seq vector) while other perturbations (e.g., ORFs, non-coding sequences) can be paired with short barcodes (Melnikov et al., 2012; Yang et al., 2011). By comparison, reported methods for highly multiplexed FISH require higher imaging magnification (60X) and barcodes longer than 200 bp, precluding cost-effective direct oligo array synthesis. Epitope-based protein barcodes are a promising method for enzyme-free decoding of pooled elements, but are currently limited in scale to ~100 barcodes (Wroblewska et al., 2018).

Despite the rich phenotypic information provided by imaging assays, optical phenotyping has been underused for screening applications due to the substantial cost, time and labor required to execute large-scale arrayed perturbation screens. In recent years, there has been wide use of pooled genome-scale CRISPR screens based on enriching cells via fitness or reporter fluorescence in various model systems, but only one large-scale (2,281 sgRNA) CRISPR imaging screen (Groot et al., 2018). Pooling improves data quality and dramatically reduces the cost and labor required, making genome-scale screens with image-based phenotyping accessible to many laboratories.

Optical pooled screening serves as an important complement to single-cell molecular profiling, which also provides high-content data but is not yet able to deliver dynamic or spatially resolved information. The LentiGuide-BC and CROP-seq libraries described in this study both generate mRNA encoding perturbation identity, so the same libraries of perturbed cells can be screened *in situ* to read dynamic and/or spatially resolved information and by complementary molecular profiling methods, such as single cell RNA-seq. This approach could assist in establishing functional relationships between molecular profiles and many cellular phenotypes (e.g., morphology, motility, electrical depolarization, cell-to-cell interactions).

Our approach is broadly applicable across many settings. We demonstrate the identification of multiple perturbations within the same cell, providing a straightforward route to study higher order genetic interactions. The potential to integrate optical screening with high-dimensional morphological profiling (Gustafsdottir et al., 2013) and *in situ* multiplexed gene expression analysis (Chen et al., 2015; Ke et al., 2013; Lee et al., 2014; Lubeck et al., 2014; Wang et al., 2014) raises the prospect of learning phenotypes from high-content data rather than pre-specifying phenotypes of interest. Libraries of endogenous or engineered protein variants could be screened for effects on cell structure or other optically-defined phenotypes. Optical pooled screening in a 2D or 3D co-culture system could be used to analyze non-cell autonomous phenotypes based on physical contact (e.g., formation of adhesion complexes, direct contact signaling, neurotransmission). Existing protocols for *in situ* sequencing in tissue samples (Ke et al., 2013; Wang et al., 2018) highlight the exciting possibility of perturbing cells *in vivo* and measuring the resulting phenotypes within the native spatial context.

STAR ★METHODS

LEAD CONTACT AND MATERIALS AVAILABILITY

Further information and requests for resources and reagents should be directed to and will be fulfilled by the Lead Contact, Paul Blainey (pblainey@broadinstitute.org)

Plasmids generated in this study have been deposited to Addgene (additional details provided in the Key Resources Table).

EXPERIMENTAL MODEL AND SUBJECT DETAILS

Tissue culture

HEK293FT cells (Thermo Fisher Scientific R70007) were cultured in DMEM with sodium pyruvate and GlutaMAX (Life 10569044) supplemented with heat-inactivated fetal bovine serum (Seradigm 97068–085) and 100 U/mL penicillin-streptomycin (Thermo Fisher Scientific 15140163). HeLa cells were cultured in the same media with serum substituted for 10% tetracycline-screened fetal bovine serum (Hyclone SH30070.03T). All other cell lines (A549, HCT116, HT1080, A375) were cultured in the same media with 10% heat-inactivated fetal bovine serum (Sigma F4135).

In preparation for *in situ* analysis, cells were seeded onto glass-bottom plates (6-well: Cellvis P06-1.5H-N, 24-well: Greiner Bio-one 662892, 96-well: Greiner Bio-one 655892) at a density of 50,000 cells/cm² and incubated for 2 days to permit proper cell attachment, spreading, and colony formation.

Selection of inducible HeLa Cas9 clone

Parental HeLa-TetR-Cas9 cells were a gift from Iain Cheeseman. In order to select a clone with optimal induction and Cas9 activity for further experiments, single cells were sorted into a 96-well plate (Sony SH800), clonally expanded, and screened for Cas9 activity after 8 days with and without 1 µg/mL doxycycline induction. Cas9 activity was assessed by transducing each clone with pXPR_011 (Addgene #59702), a reporter vector expressing GFP and an sgRNA targeting GFP (Doench et al., 2014), and using FACS to read out efficiency of protein knockdown. Additionally, gene editing was directly assessed by transducing HeLa-TetR-Cas9 clones with a guide targeting *TFRC* (sgRNA:CTATACGCCACATAACCCCC). Genomic DNA was extracted from both uninduced and induced clones by resuspending in cell lysis buffer (10 mM Tris pH 7.5, 1 mM CaCl₂, 3 mM MgCl₂, 1 mM EDTA, 1% Triton X-100, and 0.2 mg/mL Proteinase K), and heating for 10 minutes at 65° and 15 minutes at 95°. The guide target region was amplified by PCR (P5 primer: CTCTTTCCCTACACGACGCTCTTCCGATCTATGACCTTAGGCTTATTTAACTTAAT C, P7 primer: CTGGAGTTCAGACGTGTGCTCTTCCGATCTGAGTCTCATGCACTGTTTTGC) and sequenced on an Illumina MiniSeq. The best clones showed efficient indel generation (97%) in the presence of doxycycline and minimal cutting (2%) in its absence.

METHOD DETAILS

Lentivirus production

HEK293FT cells were seeded into 15-cm plates or multi-well plates at a density of 100,000 cells/cm². After 20 hours, cells were transfected with pMD2.G (Addgene #12259), psPAX2 (Addgene #12260), and a lentiviral transfer plasmid (2:3:4 ratio by mass) using Lipofectamine 3000 (Thermo Fisher Scientific L3000015). Media was exchanged after 4 hours and supplemented with 2 mM caffeine 20 hours post-transfection to increase viral titer. Viral supernatant was harvested 48 hours after transfection and filtered through 0.45 µm cellulose acetate filters (Corning 431220).

Lentiviral transduction

Cells were transduced by adding viral supernatant supplemented with polybrene (8 µg/mL) and centrifuging at 1000g for 2 hours at 33°. At 5 hours post-infection, media was exchanged. At 24 hours post-infection, cells were passaged into media containing selection antibiotic at the following concentrations: 1 µg/mL puromycin (Thermo Fisher Scientific A1113802), 300 µg/mL hygromycin (Invivogen ant-hg-1), 30 µg/mL blasticidin (Thermo Fisher Scientific A1113903), and 300 µg/mL zeocin (Thermo Fisher Scientific R25001).

For lentiviral transduction of LentiGuide-BC libraries, a carrier plasmid was utilized to minimize recombination between distant genetic elements (e.g., sgRNA and associated barcode). Libraries were packaged following the above protocol, with the library transfer plasmid diluted in integration-deficient carrier vector pR_LG (1:10 mass ratio of library to carrier, Addgene #112895) (Feldman et al., 2018).

Library cell line validation

For library transductions, multiplicity of infection (Table S1) was estimated by counting colonies after sparse plating and antibiotic selection. Genomic DNA was also extracted for NGS validation of library representation.

Next generation sequencing of libraries

Genomic DNA was extracted using an extraction mix as described above. Barcodes and sgRNAs were amplified by PCR from a minimum of 10^6 genomic equivalents per library using NEBNext 2X Master Mix (initial denaturation for 5 minutes at 98°C, followed by 28 cycles of annealing for 10 seconds at 65°C, extension for 25 seconds at 72°C, and denaturation for 20 seconds at 98°C).

Library design and cloning

A set of 12-nt barcodes was designed by selecting 83,314 barcodes from the set of 16.7 million possible 12-nt sequences by filtering for GC content between 25% and 75%, no more than 4 consecutive repeated bases, and minimum substitution and insertion/deletion edit distance (Levenshtein distance) of 3 between any pair of barcodes. Ensuring a minimum edit distance is useful for detecting and correcting errors, which arise mainly from DNA synthesis and *in situ* reads with low signal-to-background ratios. The E-CRISP web tool was used to select sgRNA sequences targeting genes of interest (Heigwer et al., 2014). Barcode-sgRNA pairs were randomly assigned and co-synthesized on a 125-nt 90K oligo array (CustomArray/Genscript) (Table S3). Individual libraries were amplified from the oligo pool by dial-out PCR (Schwartz et al., 2012) and cloned into LentiGuide-BC or LentiGuide-BC-CMV (the latter contains the CMV promoter instead of the EF1a promoter) via two steps of Golden Gate assembly using BsmBI restriction sites. Then, the sgRNA scaffold sequence and desired resistance cassette were inserted using BbsI restriction sites. Libraries were transformed in electrocompetent cells (Lucigen Endura) and grown in liquid culture for 18 hours at 30°C before extracting plasmid DNA. The sgRNA-barcode association was validated by Sanger sequencing individual colonies from the final library.

Padlock-based RNA detection

Preparation of targeted RNA amplicons for *in situ* sequencing was adapted from published protocols with modifications to improve molecular detection efficiency and amplification yield (Ke et al., 2013; Larsson et al., 2010). Cells were seeded onto glass-bottom dishes two days prior to *in situ* processing. For D458 cells, glass bottom plates were coated with poly-L-lysine (0.01% w/v in water, Sigma Aldrich) for 30 minutes and washed with PBS prior to seeding. Cells were fixed with 4% paraformaldehyde (Electron Microscopy Sciences 15714) for 30 minutes, washed with PBS, and permeabilized with 70% ethanol for 30 minutes.

Permeabilization solution was carefully exchanged with PBS-T wash buffer (PBS + 0.05% Tween-20) to minimize sample dehydration. Reverse transcription mix (1× RevertAid RT buffer, 250 μM dNTPs, 0.2 mg/mL BSA, 1 μM RT primer, 0.8 U/μL Ribolock RNase inhibitor, and 4.8 U/μL RevertAid H minus reverse transcriptase) was added to the sample and incubated for 16 hours at 37°C. Following reverse transcription, cells were washed 5 times with PBS-T and post-fixed with 3% paraformaldehyde and 0.1% glutaraldehyde for 30 minutes at room temperature, then washed with PBS-T 5 times. After this step, cells expressing p65-mNeonGreen were imaged. Samples were thoroughly washed again with PBS-T, incubated in a padlock probe and extension-ligation reaction mix (1× Ampligase buffer, 0.4 U/μL RNase H, 0.2 mg/mL BSA, 100 nM padlock probe, 0.02 U/μL TaqIT polymerase, 0.5 U/μL Ampligase and 50 nM dNTPs) for 5 minutes at 37°C and 90 minutes at 45°C, and then washed 2 times with PBS-T. Circularized padlocks were amplified with rolling circle amplification mix (1× Phi29 buffer, 250 μM dNTPs, 0.2 mg/mL BSA, 5% glycerol, and 1 U/μL Phi29 DNA polymerase) at 300°C for either 3 hours or overnight.

***In situ* sequencing**

Rolling circle amplicons were prepared for sequencing by hybridizing a mix containing sequencing primer oSBS_LentiGuide-BC or oSBS_CROP-seq (1 μM primer in 2X SSC + 10% formamide) for 30 minutes at room temperature. Barcodes were read out using sequencing-by-synthesis reagents from the Illumina MiSeq 500 cycle Nano kit (Illumina MS-103–1003). First, samples were washed with incorporation buffer (Nano kit PR2) and incubated for 3 minutes in incorporation mix (Nano kit reagent 1) at 60°C. Samples were then thoroughly washed with PR2 at 60°C (6 washes for 3 minutes each) and placed in 200 ng/mL DAPI in 2x SSC for fluorescence imaging. Following each imaging cycle, dye terminators were removed by incubation for 6 minutes in Illumina cleavage mix (Nano kit reagent 4) at 60°C, and samples were thoroughly washed with PR2.

Padlock optimization

Improved padlock sequences were found for the CROP-seq and CROP-seq-v2 (contains a modified sgRNA scaffold for increased CRISPR efficiency) vectors by pooled testing of a set of 84 padlocks (per vector) for *in situ* RNA detection efficiency and RCA amplification yield. Sequences varied in the length of the cDNA binding region (melting temperature from 45°C, 49°C, or 54°C), and in overall length (72, 77, 83, or 89 nt for CROP-seq and 59, 69, 79, or 90 nt for CROP-seq-v2). The padlock backbone sequence is not functionally constrained; here, backbone sequences were randomly generated with G nucleotides excluded to reduce secondary structure. The backbone sequences were then filtered to have C content of 50±5% and no homopolymers exceeding 3 nt. Padlock sequences with the most positive minimum free energies were selected from these filtered sequences using mfold's zipfold utility (Zuker, 2003). For compatibility with pooled testing, the backbone sequences for each vector were required to begin with a unique 5-mer barcode. Padlocks were individually synthesized (Integrated DNA Technologies) and pooled prior to phosphorylation with T4 PNK (New England Biolabs) according to the manufacturer's instructions. Padlock-based RNA detection and *in situ* sequencing of the padlock backbones were performed as previously described in HeLa-TetR-Cas9 cells, using a sequencing primer that targeted the 5-mer padlock barcode. After identifying padlocks by 5 cycles of *in situ*

sequencing, the padlocks were hybridized with a fluorescently labeled detection oligo to give a barcode-independent estimate of relative RCA yield. Relative RNA detection efficiency (fraction of sequencing reads from a given padlock) was used as a proxy for absolute RNA detection efficiency.

Fluorescence microscopy

All *in situ* sequencing images were acquired using a Ti-E Eclipse inverted epifluorescence microscope (Nikon) with automated XYZ stage control and hardware autofocus. An LED light engine (Lumencor Sola SE FISH II) was used for fluorescence illumination and all hardware was controlled using Micromanager software (Edelstein et al., 2010). *In situ* sequencing cycles were imaged using a 10X 0.45 NA CFI Plan Apo Lambda objective (Nikon) with the following filters (Semrock) and exposure times for each base: G (excitation 534/20 nm, emission 572/28 nm, dichroic 552 nm, 200 ms); T (excitation 575/25 nm, emission 615/24 nm, dichroic 596 nm, 200 ms); A (excitation 635/18 nm, emission 680/42 nm, dichroic 652 nm, 200 ms); C (excitation 661/20 nm, emission 732/68 nm, dichroic 695 nm, 800 ms).

Frameshift reporter screen

HeLa-TetR-Cas9 cells were stably transduced at MOI > 2 with pL_FR_Hygro and selected with hygromycin for 7 days to generate the HeLa-TetR-Cas9-FR cell line. Cells transduced with the pL_FR_Hygro lentiviral vector express an open reading frame consisting of a 50-nt frameshift reporter target sequence, followed by an H2B histone coding sequence with C-terminus HA epitope tag (+1 frameshift), followed by a second H2B sequence with C-terminus myc tag (+0 frameshift) and hygro antibiotic resistance cassette (+0 frameshift). The H2B-HA, H2B-myc, and hygromycin resistance sequences are preceded by self-cleaving 2A peptides in the same reading frame. Before generation of Cas9-mediated indel mutations, cells express the coding sequences with +0 frameshift. Subsequent activation of the reporter by co-expression of Cas9 and a targeting sgRNA leads to mutations in the target sequence, which may alter the downstream reading frame. A frameshift of +1 leads to expression of the H2B-HA protein, which can be visualized by immunofluorescence and detected by microscopy or flow cytometry. Integration of multiple copies of reporter per cell increases the likelihood of generating a +1 frameshift in at least one copy.

HeLa-TetR-Cas9-FR cells were used to screen targeting and control sgRNAs. A barcoded sgRNA perturbation library with 972 barcodes, each encoding one of 5 targeting or 5 control sgRNAs, was synthesized and cloned into LentiGuide-BC-CMV. This library was transduced into HeLa-TetR-Cas9-FR cells at MOI < 0.05 in three replicates, which were independently cultured and screened. Following 4 days of puromycin selection, cells were collected to validate library representation by NGS. Cas9 expression was induced by supplementing the culture media with 1 µg/mL doxycycline for 6 days. Cells were then split for screening either via *in situ* sequencing or by FACS.

For *in situ* screening, 500,000 cells were seeded into each well of a glass-bottom 6-well plate (CellVis). After two days of culture, *in situ* padlock detection and sequencing were carried out as above, with the modification that prior to sequencing-by-synthesis, cells were

immunostained to detect frameshift reporter activation by blocking and permeabilizing with 3% BSA + 0.5 % Triton X-100 for 5 minutes, incubating in rabbit anti-HA (1:1000 dilution in 3% BSA) for 30 minutes, washing with PBS-T and incubating with goat anti-rabbit F(ab')₂ fragment Alexa 488 (CST 4412S, 1:1000 dilution in 3% BSA) for 30 minutes. Samples were changed into imaging buffer (200 ng/mL DAPI in 2X SSC) and phenotype images were acquired.

FACS screening was carried out by fixing cells with 4% PFA, permeabilizing with 70% ice-cold ethanol, and immunostaining with the same anti-HA primary and secondary antibodies and dilutions used for *in situ* analysis. Cells were sorted into HA⁺ and HA⁻ populations (Sony SH800) and genomically integrated perturbations were sequenced as described above. The enrichment for each barcoded perturbation was defined as the HA⁺/HA⁻ ratio of normalized read counts.

Frameshift reporter screens in U2-OS, A375, HT1080, HCT116 and HEK293T cell types were performed by first transducing these cells with lentiCas9-blast (Addgene #52962) (Sanjana et al., 2014) and selecting with blasticidin for 4 days. Cas9-expressing cells were then transduced with pL_FR_Hygro and selected with hygromycin for 7 days to generate reporter lines. These reporter lines were transduced with a CROPseq-puro library consisting of the same 5 targeting and 5 non-targeting sgRNAs used above. Cell libraries were selected with puromycin for 4 days and cells were seeded onto glass-bottom dishes 2 days prior to *in situ* sequencing.

NF-κB activation screens

HeLa-TetR-Cas9 cells were transduced with pR14_p65-mNeonGreen, a C-terminal fusion of p65 with a bright monomeric green fluorescent protein (Allele Biotechnology). Fluorescent cells were sorted by FACS (Sony SH800) and re-sorted to select for cells with stable expression. This reporter cell line was further transduced with a 4,063-barcode sgRNA library (962 genes targeted and 1,000 barcodes assigned to non-targeting sgRNAs) in LentiGuide-BC-CMV. Cells were selected with puromycin for 4 days and library representation was validated by NGS.

Cas9 expression was induced with 1 μg/mL doxycycline and cells were seeded onto 6-well cover glass-bottom plates 2 days prior to translocation experiments. The total time between Cas9 induction and performing the NF-κB activation assay was 7 days. Cells were stimulated with either 30 ng/mL TNFα or 30 ng/mL IL-1β (Invivogen) for 40 minutes prior to fixation with 4% paraformaldehyde for 30 minutes and initiation of the *in situ* sequencing protocol. Translocation phenotypes were recorded after the post-fixation step by exchanging for imaging buffer (2X SSC + 200 ng/mL DAPI) and imaging the nuclear DAPI stain and p65-mNeonGreen. After phenotyping, the remainder of the *in situ* sequencing protocol (gap-fill and rolling circle amplification) and 12 bases of sequencing-by-synthesis were completed.

HeLa Cas9-blast, A549 Cas9-blast and HCT116 Cas9-blast cells were transduced with a CROPseq-puro library consisting of 5,738 sgRNAs (952 genes targeted, 100 non-targeting sgRNAs). Cells were selected for 4 days and seeded onto 6-well plates 2 days prior to

translocation experiments. The total time between library transduction and the screen was between 7 and 14 days. Cells were stimulated with TNF α or IL-1 β (30 ng/mL for HCT116, 3 ng/mL for HeLa and A549) for 40 minutes prior to fixation with 4% paraformaldehyde for 30 minutes. Cells were then processed for *in situ* sequencing as in the HeLa p65-mNeonGreen activation screen, except for the following differences. Cells were stained after the post-fixation step by incubating for 1 hour with a rabbit antibody against p65 (CST 8242S, 1:400 dilution in 3% BSA), washing with PBS-T, then incubating with goat anti-rabbit F(ab')₂ fragment Alexa 488 (CST 4412S, 1:1000 dilution in 3% BSA) for 45 minutes and washing with PBS-T before proceeding to the gap-fill step. After RCA, cells were exchanged into imaging buffer (2X SSC + 200 ng/mL DAPI) and imaged with DAPI and 488 (p65) filters. After phenotyping, sequencing primer was hybridized and 10 bases of sequencing-by-synthesis were acquired.

NF- κ B live-cell screens

HeLa-TetR-Cas9 stably expressing the p65-mNeonGreen reporter were transduced with the 952-gene CROPseq-puro library described above. Cells were selected with puromycin for 4 days and library representation was validated by NGS.

Cas9 expression was induced for 7 days with 1 μ g/mL doxycycline and cells were seeded onto 6-well cover glass-bottom plates 2 days prior to translocation experiments. The total time between the start of Cas9 induction and performing the NF- κ B live-cell assay was 14 days. Prior to the experiment, culture media was exchanged for imaging media: 0.1 ng/mL Hoechst 33342 in phenol red-free DMEM with HEPES and L-glutamine (Thermo Fisher Scientific 21063029), supplemented with 10% FBS and 100 U/mL penicillin-streptomycin. Cells were returned to the incubator for 2 hours, then stimulated with either 30 ng/mL TNF α or 30 ng/mL IL-1 β and immediately loaded onto an automated live-cell microscope with environmental control (Zeiss CellDiscoverer 7). Images of the Hoechst nuclear stain and p65-mNeonGreen were acquired at 5X magnification at 23 minute intervals over 6 hours. Immediately after stopping live-cell imaging, cells were fixed and the *in situ* sequencing protocol was carried out as above.

Pooled validation was conducted with a new CROPseq-puro library targeting 65 candidate genes from the primary screen (10 sgRNAs per gene, 100 non-targeting sgRNAs). The validation screen was conducted following the same protocol as the primary live-cell screen, except that imaging was carried out on a Nikon Ti2 inverted microscope equipped with a live-cell incubator (Okolab). Images were acquired at 10X magnification in 30 minute intervals.

NF- κ B arrayed validation

Top-ranking genes from the primary pooled screen were validated with individual sgRNAs. For each gene, 2 or 3 sgRNAs were tested, including at least one sgRNA not used in the primary screen. HeLa-TetR-Cas9 cells expressing p65-mNeonGreen were prepared and assayed as in the pooled screen, except that cells were transduced with the LentiGuide-Puro sgRNA expression vector (Addgene #52963) (Sanjana et al., 2014), and the translocation

assay was carried out in 96-well cover glass plates. Image acquisition and data analysis were performed with the same hardware and software settings as in the pooled screen.

Hit validation using clonal knockout cell lines

Hela-TetR-Cas9 cells expressing p65-mNeonGreen were transfected with synthetic crRNA/tracrRNA (IDT; *MED12* sgRNA: GGATCTTGAGCTACGAACAC; *MED24* sgRNA: GCGCTGGAGTGACTACCAAT) using Lipofectamine 2000 (Thermo Fisher Scientific). Two days after transfection, single cell clones were seeded at a density of 0.6 cells/well into 480 96-wells per target gene. After two weeks, live clones were identified by microscopy and were expanded to a 6-well format. Clones were genotyped by replacing media with direct lysis buffer (1 mM CaCl₂, 3 mM MgCl₂, 1 mM EDTA, 1% Triton X-100, 10 mM Tris pH 7.5, 0.2 mg/ml Proteinase K), heating to 65°C for 10 minutes, and incubating at 95°C for 15 minutes to inactivate Proteinase K. PCR amplification of target loci, deep sequencing (Illumina MiSeq), and data analysis using the [OutKnocker.org](https://www.outknocker.org) software were performed as described (Schmid-Burgk, Genome Research 2014). Clones bearing out-of-frame mutations in all alleles were further expanded, and analyzed by automated live cell microscopy (Nikon Ti2 inverted microscope with Okolab live-cell incubator) for 6 hours after stimulation with TNF α or IL-1 β as described above.

RNA-seq analysis of clonal knockout cell lines

Clonal cell lines and wild type control cells were plated in 24-well format at a density of 50,000 cells/well. The next day, cells were stimulated for 0, 1, or 4 hours with TNF α or IL-1 β (3 ng/mL) prior to lysis. Cells were washed with PBS, and lysed at room temperature in 150 μ l TCL buffer (Qiagen, supplemented with 1% beta-mercaptoethanol) per well, then stored at -80°C. Smart-seq2 was performed as described (Trombetta et al., 2014). Libraries were sequenced on a NextSeq 500 (Illumina) using the High Output Kit v2 75-cycle kit (paired end, 36 cycles forward, 36 cycles reverse).

QUANTIFICATION AND STATISTICAL ANALYSIS

Image analysis

Images of cell phenotype and *in situ* sequencing of perturbations were coarsely aligned during acquisition using nuclear masks to calibrate plate position, and finely aligned during analysis using cross-correlation of DAPI or *in situ* sequencing signal between cycles. Nuclei were detected using local thresholding and watershed-based segmentation. Cells were segmented using local thresholding of cytoplasmic background from *in situ* sequencing signal and assignment of pixels to the nearest nucleus by the fast-marching method. Frameshift reporter and NF- κ B translocation phenotypes were quantified by calculating pixel-wise correlations between the nuclear DAPI channel and 488 channel (HA stain or p65-mNeonGreen, respectively). Mitotic cells and cells with abnormally high or low reporter expression were filtered out based on maximum DAPI signal and mean p65-mNeonGreen signal (Figure S4A).

Sequencing reads were detected by applying a Laplacian-of-Gaussian linear filter (kernel width $\sigma = 1$ pixel) to subtract low-frequency background and enhance the point-like

sequencing spots. Peaks were detected by calculating the per-pixel, per-channel standard deviation over sequencing cycles, averaging over color channels, and finding local maxima in the resulting image. The base intensity at each cycle was defined as the maximum value in a 3×3 pixel window centered on the read. A linear transformation to correct for optical cross-talk and intensity differences between color channel was then estimated from the data and applied. Finally, each base was called according to the channel with maximum corrected intensity, and a per-base quality score was defined as the ratio of intensity for the maximum channel to total intensity for all channels. The output of the sequencing image analysis was a table recording each sequencing read along with the identity of the overlapping cell, quality score per base, and spatial location.

Data analysis functions were written in Python, using Snakemake for workflow control (Köster and Rahmann, 2012).

Frameshift reporter misidentification rate estimation

To estimate the rate at which cells are assigned an incorrect perturbation barcode, we first assumed all HA⁺ cells mapped to a non-targeting control sgRNA (4.7%) were false positive events due to incorrect barcode assignment (supported by the very low false positive rate (<0.001 %) of the frameshift reporter itself, measured for a single perturbation in arrayed format). However, as incorrect barcode assignments were equally likely to map an HA⁺ cell to a targeting or control sgRNA, we estimated the misidentification rate to be twice as large, or 9.4%.

NF- κ B activation screen analysis

Nuclei of individual cells were segmented by thresholding background-subtracted DAPI signal and separating the resulting regions using the watershed method. Cells with at least one read exactly matching a library barcode were retained for analysis. In order to remove mitotic cells and cells with abnormally high or low reporter expression, cells were further filtered based on nuclear area, maximum DAPI signal, and mean p65-mNeonGreen signal. Pixel-wise DAPI-mNeonGreen correlation within the segmented nuclear region, described above, was used to define the translocation score for each cell as it most clearly separated perturbations against known NF- κ B genes from non-targeting controls. The phenotypic effects of perturbations targeting known NF- κ B genes ranged from a large increase in fully untranslocated cells (e.g., *MAP3K7*) to more subtle negative shifts in the distribution of scores (e.g., *IKBKB*).

To capture a broad range of effect sizes, we calculated an sgRNA translocation defect in a given replicate by computing the difference in translocation score distribution compared to non-targeting controls (the shaded area in Fig. 4, C and E). We found this metric performed better at separating known genes from controls than the often-used Kolmogorov-Smirnov distance. We defined the gene translocation defect as the second-largest sgRNA translocation defect for sgRNAs targeting that gene. This statistic helps reduce the false positive rate due to clonal effects (integration of an sgRNA into a cell that is defective in translocation) which are independent among sgRNAs and screen replicates, as well as false negatives due to inefficient sgRNAs.

A permutation test was used to calculate p-values for the gene translocation defects. Random subsets of sgRNA translocation defects were sampled from non-targeting controls to build a null distribution (3 sgRNAs per replicate, repeated 100,000 times). The cumulative null distribution was used to determine p-values for the gene translocation defects. Hits at an estimated FDR <10% and <20% were identified using the Benjamini-Hochberg procedure (Table S3). KEGG-annotated genes were defined as members of KEGG pathway HSA04064 (NF-kappa B signaling pathway) between IL-1 β or TNF α and p65/p50.

Cellular morphology analysis

Cellular morphology was analyzed by calculating image features from DAPI intensity as well as cell and nuclear geometry. We summarized the gene-level (across all guides and cells) distribution of morphological features (cell area, nucleus area, DAPI max, DAPI mean, cell eccentricity and nucleus eccentricity) by its first, second and third quartiles, producing an 18-dimensional vector for each gene. We performed dimensionality reduction by Principal Components Analysis (PCA) and visualized the first two principal components, finding distinct clusters for regulators downstream of IKK (Figure S4).

Analysis of NF- κ B live-cell screens

Image analysis was conducted as for the initial NF- κ B screen, with additional preprocessing steps to (a) track cells through the live-cell time course, and (b) align the final time point of live-cell with the first cycle of sequencing-by-synthesis. For each cell, the translocation score at each time point was subtracted from a baseline translocation score, interpolated in time from control cells imaged in the same well. For each gene, an integrated translocation score was calculated by integrating each cell's baseline-subtracted translocation scores from 45 to 345 minutes post-stimulation. Statistical significance for deviation in the integrated translocation score between perturbed and control cells was quantified using the non-parametric Mann-Whitney U test.

NF- κ B arrayed validation

The translocation defect for each sgRNA and cytokine translocation was assessed by computing the difference in translocation score distribution compared to the average of at least 3 non-targeting control sgRNAs assayed on the same plate. For each cytokine, the translocation defects were standardized using the mean and standard deviation of the translocation defects for non-targeting control guides. These standardized values were averaged over replicate sgRNAs for a given gene and cytokine pair to obtain validation Z-scores (Table S3).

RNA-seq quantification

Transcript abundance was quantified using kallisto and differential gene expression was analyzed using DEBrowser as a graphical interface to edgeR, with default edgeR parameters: TMM-based normalization, dispersion estimated using the edgeR function estimateDisp, and p-values adjusted with the Benjamini-Hochberg procedure (Bray et al., 2016; Kucukural et al., 2019; Robinson et al., 2010). Highly induced NF- κ B genes were defined as genes with >50-fold increase after 4 hours of stimulation by either cytokine in

wild type cells (adjusted p-value $< 10^{-3}$). *MED12* or *MED24*-dependent genes were defined as genes with >8 -fold change between knockout and wild type conditions after 4 hours of stimulation with either cytokine. All pairwise comparisons were performed using the edgeR function exactTest ($n = 3$ or 4 biological replicates). Biological replicates were defined as replicate clonal knockout cell lines (for *MED12* and *MED24* knockouts) or replicate stimulations (for wild type control cells).

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

ACKNOWLEDGEMENTS

We thank Rhiannon Macrae, Iain Cheeseman, Aviv Regev, Fei Chen, Arjun Raj, Nir Hacohen, Benjamin Gewurz, Sara Jones, Emily Botelho, and members of the Blainey and Zhang labs for critical feedback and discussions. We thank Leslie Gaffney for assistance with figures. We gratefully acknowledge Linyi Gao for the gift of the integration-deficient co-packaging plasmid pR_LG. We thank Zohar Bloom-Ackermann for assistance with live-cell imaging and FuNien Tsai for preparing *in situ* sequencing of D458 cells.

This work was supported by a Simons Center Seed Grant from MIT, the Broad Institute through startup funding (to P.C.B) and the BN10 program, and two grants from the National Human Genome Research Institute (HG009283 and HG006193). P.C.B. is supported by a Career Award at the Scientific Interface from the Burroughs Wellcome Fund. F.Z. is a New York Stem Cell Foundation–Robertson Investigator. F.Z. is supported by NIH grants (1R01-HG009761, 1R01-MH110049, and 1DP1-HL141201); the Howard Hughes Medical Institute; the New York Stem Cell, Simons, Paul G. Allen Family, and Vallee Foundations; the Poitras Center for Affective Disorders Research at MIT; the Hock E. Tan and K. Lisa Yang Center for Autism Research at MIT; J. and P. Poitras; and R. Metcalfe. A.M. is supported by the Swedish Research Council (grant 2015–06403). R.J.C. is supported by a Fannie and John Hertz Foundation Fellowship and an NSF Graduate Research Fellowship. J.L.S.-B. is supported by an EMBO Long-Term Fellowship (ALTF 199–2017).

The Broad Institute and MIT may seek to commercialize aspects of this work, and related applications for intellectual property have been filed.

REFERENCES

- Adamson B, Norman TM, Jost M, Cho MY, Nuñez JK, Chen Y, Villalta JE, Gilbert LA, Horlbeck MA, Hein MY, et al. (2016). A Multiplexed Single-Cell CRISPR Screening Platform Enables Systematic Dissection of the Unfolded Protein Response. *Cell* 167, 1867–1882.e21. [PubMed: 27984733]
- Adamson B, Norman TM, Jost M, and Weissman JS (2018). Approaches to maximize sgRNA-barcode coupling in Perturb-seq screens. *BioRxiv* 298349.
- Bauernfeind F, Niepmann S, Knolle PA, and Hornung V. (2016). Aging-Associated TNF Production Primes Inflammation Activation and NLRP3-Related Metabolic Disturbances. *The Journal of Immunology* 197, 2900–2908. [PubMed: 27566828]
- Bononi A, Yang H, Giorgi C, Patergnani S, Pellegrini L, Su M, Xie G, Signorato V, Pastorino S, Morris P, et al. (2017). Germline BAP1 mutations induce a Warburg effect. *Cell Death Differ* 24, 1694–1704. [PubMed: 28665402]
- Bray NL, Pimentel H, Melsted P, and Pachter L. (2016). Near-optimal probabilistic RNA-seq quantification. *Nat. Biotechnol* 34, 525–527. [PubMed: 27043002]
- Buschmann T, and Bystrykh LV (2013). Levenshtein error-correcting barcodes for multiplexed DNA sequencing. *BMC Bioinformatics* 14, 272. [PubMed: 24021088]
- Chen ZJ (2005). Ubiquitin Signaling in the NF- κ B Pathway. *Nat. Cell Biol* 7, 758–765. [PubMed: 16056267]
- Chen KH, Boettiger AN, Moffitt JR, Wang S, and Zhuang X. (2015). Spatially resolved, highly multiplexed RNA profiling in single cells. *Science* 348, aaa6090.

- Chen X, Sun Y-C, Church GM, Lee JH, and Zador AM (2018). Efficient in situ barcode sequencing using padlock probe-based BaristaSeq. *Nucleic Acids Res* 46, e22.
- Chia N-Y, Chan Y-S, Feng B, Lu X, Orlov YL, Moreau D, Kumar P, Yang L, Jiang J, Lau M-S, et al. (2010). A genome-wide RNAi screen reveals determinants of human embryonic stem cell identity. *Nature* 468, 316–320. [PubMed: 20953172]
- Colleran A, Collins PE, O'Carroll C, Ahmed A, Mao X, McManus B, Kiely PA, Burstein E, and Carmody RJ (2013). Deubiquitination of NF- κ B by Ubiquitin-Specific Protease-7 promotes transcription. *Proc. Natl. Acad. Sci. U. S. A* 110, 618–623. [PubMed: 23267096]
- Collinet C, Stöter M, Bradshaw CR, Samusik N, Rink JC, Kenski D, Habermann B, Buchholz F, Henschel R, Mueller MS, et al. (2010). Systems survey of endocytosis by multiparametric image analysis. *Nature* 464, 243–249. [PubMed: 20190736]
- Dai F, Lee H, Zhang Y, Zhuang L, Yao H, Xi Y, Xiao Z-D, You MJ, Li W, Su X, et al. (2017). BAP1 inhibits the ER stress gene regulatory network and modulates metabolic stress response. *Proc. Natl. Acad. Sci. U. S. A* 114, 3192–3197. [PubMed: 28275095]
- Dang Y, Jia G, Choi J, Ma H, Anaya E, Ye C, Shankar P, and Wu H. (2015). Optimizing sgRNA structure to improve CRISPR-Cas9 knockout efficiency. *Genome Biol* 16, 280. [PubMed: 26671237]
- Datlinger P, Rendeiro AF, Schmidl C, Krausgruber T, Traxler P, Klughammer J, Schuster LC, Kuchler A, Alpar D, and Bock C. (2017). Pooled CRISPR screening with single-cell transcriptome readout. *Nat. Methods* 14, 297–301. [PubMed: 28099430]
- Dixit A, Parnas O, Li B, Chen J, Fulco CP, Jerby-Aron L, Marjanovic ND, Dionne D, Burks T, Raychowdhury R, et al. (2016). Perturb-Seq: Dissecting Molecular Circuits with Scalable Single-Cell RNA Profiling of Pooled Genetic Screens. *Cell* 167, 1853–1866.e17. [PubMed: 27984732]
- Doench JG, Hartenian E, Graham DB, Tothova Z, Hegde M, Smith I, Sullender M, Ebert BL, Xavier RJ, and Root DE (2014). Rational design of highly active sgRNAs for CRISPR-Cas9-mediated gene inactivation. *Nat. Biotechnol* 32, 1262–1267. [PubMed: 25184501]
- Essen D van, Engist B, Natoli G, and Sacconi S. (2009). Two Modes of Transcriptional Activation at Native Promoters by NF- κ B p65. *PLoS Biol* 7, e1000073.
- Feldman D, Singh A, Garrity AJ, and Blainey PC (2018). Lentiviral co-packaging mitigates the effects of intermolecular recombination and multiple integrations in pooled genetic screens. *BioRxiv* 262121.
- Floyd SR, Pacold ME, Huang Q, Clarke SM, Lam FC, Cannell IG, Bryson BD, Rameseder J, Lee MJ, Blake EJ, et al. (2013). The bromodomain protein Brd4 insulates chromatin from DNA damage signalling. *Nature* 498, 246–250. [PubMed: 23728299]
- Gewurz BE, Towfic F, Mar JC, Shinnars NP, Takasaki K, Zhao B, Cahir-McFarland ED, Quackenbush J, Xavier RJ, and Kieff E. (2012). Genome-wide siRNA screen for mediators of NF- κ B activation. *Proc. Natl. Acad. Sci. U. S. A* 109, 2467–2472. [PubMed: 22308454]
- Gilbert LA, Horlbeck MA, Adamson B, Villalta JE, Chen Y, Whitehead EH, Guimaraes C, Panning B, Ploegh HL, Bassik MC, et al. (2014). Genome-Scale CRISPR-Mediated Control of Gene Repression and Activation. *Cell* 159, 647–661. [PubMed: 25307932]
- Groot R. de, Lüthi J, Lindsay H, Holtackers R, and Pelkmans L. (2018). Large-scale image-based profiling of single-cell phenotypes in arrayed CRISPR-Cas9 gene perturbation screens. *Mol. Syst. Biol* 14, e8064.
- Guermah M, Malik S, and Roeder RG (1998). Involvement of TFIID and USA components in transcriptional activation of the human immunodeficiency virus promoter by NF- κ B and Sp1. *Mol. Cell. Biol* 18, 3234–3244. [PubMed: 9584164]
- Guo Y, Walther TC, Rao M, Stuurman N, Goshima G, Terayama K, Wong JS, Vale RD, Walter P, and Farese RV (2008). Functional genomic screen reveals genes involved in lipid-droplet formation and utilization. *Nature* 453, 657–661. [PubMed: 18408709]
- Gustafsdottir SM, Ljosa V, Sokolnicki KL, Wilson JA, Walpita D, Kemp MM, Seiler KP, Carrel HA, Golub TR, Schreiber SL, et al. (2013). Multiplex Cytological Profiling Assay to Measure Diverse Cellular States. *PLoS ONE* 8, e80999.
- Hayden MS, and Ghosh S. (2012). NF- κ B, the first quarter-century: remarkable progress and outstanding questions. *Genes Dev* 26, 203–234. [PubMed: 22302935]

- Heigwer F, Kerr G, and Boutros M. (2014). E-CRISP: fast CRISPR target site identification. *Nat. Methods* 11, 122–123. [PubMed: 24481216]
- Hill AJ, McFaline-Figueroa JL, Starita LM, Gasperini MJ, Matreyek KA, Packer J, Jackson D, Shendure J, and Trapnell C. (2018). On the design of CRISPR-based single-cell molecular screens. *Nat. Methods* 15, 271–274. [PubMed: 29457792]
- Jäättelä M, Mouritzen H, Elling F, and Bastholm L. (1996). A20 zinc finger protein inhibits TNF and IL-1 signaling. *J. Immunol. Baltim. Md 1950* 156, 1166–1173.
- Jaitin DA, Weiner A, Yofe I, Lara-Astiaso D, Keren-Shaul H, David E, Salame TM, Tanay A, van Oudenaarden A, and Amit I. (2016). Dissecting Immune Circuits by Linking CRISPR-Pooled Screens with Single-Cell RNA-Seq. *Cell* 167, 1883–1896.e15. [PubMed: 27984734]
- Johnson KM, Mahajan SS, and Wilson AC (1999). Herpes simplex virus transactivator VP16 discriminates between HCF-1 and a novel family member, HCF-2. *J. Virol* 73, 3930–3940. [PubMed: 10196288]
- Kanehisa M, and Goto S. (2000). KEGG: kyoto encyclopedia of genes and genomes. *Nucleic Acids Res* 28, 27–30. [PubMed: 10592173]
- Karlas A, Machuy N, Shin Y, Pleissner K-P, Artarini A, Heuer D, Becker D, Khalil H, Ogilvie LA, Hess S, et al. (2010). Genome-wide RNAi screen identifies human host factors crucial for influenza virus replication. *Nature* 463, 818–822. [PubMed: 20081832]
- Ke R, Mignardi M, Pacureanu A, Svedlund J, Botling J, Wählby C, and Nilsson M. (2013). In situ sequencing for RNA analysis in preserved tissue and cells. *Nat. Methods* 10, 857–860. [PubMed: 23852452]
- Köster J, and Rahmann S. (2012). Snakemake—a scalable bioinformatics workflow engine. *Bioinformatics* 28, 2520–2522. [PubMed: 22908215]
- Kucukural A, Yukselen O, Ozata DM, Moore MJ, and Garber M. (2019). DEBrowser: interactive differential expression analysis and visualization tool for count data. *BMC Genomics* 20, 6. [PubMed: 30611200]
- Larsson C, Grundberg I, Söderberg O, and Nilsson M. (2010). In situ detection and genotyping of individual mRNA molecules. *Nat. Methods* 7, 395–397. [PubMed: 20383134]
- Lawson MJ, Camsund D, Larsson J, Baltekin Ö, Fange D, and Elf J. (2017). In situ genotyping of a pooled strain library after characterizing complex phenotypes. *Mol. Syst. Biol* 13, 947. [PubMed: 29042431]
- Lee D-F, Kuo H-P, Liu M, Chou C-K, Xia W, Du Y, Shen J, Chen C-T, Huo L, Hsu M-C, et al. (2009). KEAP1 E3 ligase-mediated downregulation of NF-kappaB signaling by targeting IKKbeta. *Mol. Cell* 36, 131–140. [PubMed: 19818716]
- Lee JH, Daugharthy ER, Scheiman J, Kalhor R, Yang JL, Ferrante TC, Terry R, Jeanty SSF, Li C, Amamoto R, et al. (2014). Highly multiplexed subcellular RNA sequencing in situ. *Science* 343, 1360–1363. [PubMed: 24578530]
- Linhoff MW, Lauren J, Cassidy RM, Dobie FA, Takahashi H, Nygaard HB, Airaksinen MS, Strittmatter SM, and Craig AM (2009). An unbiased expression screen for synaptogenic proteins identifies the LRRTM protein family as synaptic organizers. *Neuron* 61, 734–749. [PubMed: 19285470]
- Lubeck E, Coskun AF, Zhiyentayev T, Ahmad M, and Cai L. (2014). Single-cell in situ RNA profiling by sequential hybridization. *Nat. Methods* 11, 360–361. [PubMed: 24681720]
- Machida YJ, Machida Y, Vashisht AA, Wohlschlegel JA, and Dutta A. (2009). The deubiquitinating enzyme BAP1 regulates cell growth via interaction with HCF-1. *J. Biol. Chem* 284, 34179–34188.
- Malik S, and Roeder RG (2005). Dynamic regulation of pol II transcription by the mammalian Mediator complex. *Trends Biochem. Sci* 30, 256–263. [PubMed: 15896744]
- Melnikov A, Murugan A, Zhang X, Tesileanu T, Wang L, Rogov P, Feizi S, Gnirke A, Callan CG, Kinney JB, et al. (2012). Systematic dissection and optimization of inducible enhancers in human cells using a massively parallel reporter assay. *Nat. Biotechnol* 30, 271–277. [PubMed: 22371084]
- Misaghi S, Ottosen S, Izrael-Tomasevic A, Arnott D, Lamkanfi M, Lee J, Liu J, O’Rourke K, Dixit VM, and Wilson AC (2009). Association of C-terminal ubiquitin hydrolase BRCA1-associated protein 1 with cell cycle regulator host cell factor 1. *Mol. Cell. Biol* 29, 2181–2192. [PubMed: 19188440]

- Moffat J, Grueneberg DA, Yang X, Kim SY, Kloepfer AM, Hinkle G, Piqani B, Eisenhaure TM, Luo B, Grenier JK, et al. (2006). A lentiviral RNAi library for human and mouse genes applied to an arrayed viral high-content screen. *Cell* 124, 1283–1298. [PubMed: 16564017]
- Nakatake Y, Fujii S, Masui S, Sugimoto T, Torikai-Nishikawa S, Adachi K, and Niwa H. (2013). Kinetics of drug selection systems in mouse embryonic stem cells. *BMC Biotechnol* 13, 64. [PubMed: 23919313]
- Neumann B, Walter T, Heriche J-K, Bulkescher J, Erfle H, Conrad C, Rogers P, Poser I, Held M, Liebel U, et al. (2010). Phenotypic profiling of the human genome by time-lapse microscopy reveals cell division genes. *Nature* 464, 721–727. [PubMed: 20360735]
- Orvedahl A, Sumpter R, Xiao G, Ng A, Zou Z, Tang Y, Narimatsu M, Gilpin C, Sun Q, Roth M, et al. (2011). Image-based genome-wide siRNA screen identifies selective autophagy factors. *Nature* 480, 113–117. [PubMed: 22020285]
- Pahl HL (1999). Activators and target genes of Rel/NF-kappaB transcription factors. *Oncogene* 18, 6853–6866. [PubMed: 10602461]
- Parnas O, Jovanovic M, Eisenhaure TM, Herbst RH, Dixit A, Ye CJ, Przybylski D, Platt RJ, Tirosh I, Sanjana NE, et al. (2015). A Genome-wide CRISPR Screen in Primary Immune Cells to Dissect Regulatory Networks. *Cell* 162, 675–686. [PubMed: 26189680]
- Robinson MD, McCarthy DJ, and Smyth GK (2010). edgeR: a Bioconductor package for differential expression analysis of digital gene expression data. *Bioinformatics* 26, 139–140. [PubMed: 19910308]
- Rubin AJ, Parker KR, Satpathy AT, Qi Y, Wu B, Ong AJ, Mumbach MR, Ji AL, Kim DS, Cho SW, et al. (2019). Coupled Single-Cell CRISPR Screening and Epigenomic Profiling Reveals Causal Gene Regulatory Networks. *Cell* 176, 361–376.e17. [PubMed: 30580963]
- Sack LM, Davoli T, Xu Q, Li MZ, and Elledge SJ (2016). Sources of Error in Mammalian Genetic Screens. *G3 Genes Genomes Genet* 6, 2781–2790.
- Sanjana NE, Shalem O, and Zhang F. (2014). Improved vectors and genome-wide libraries for CRISPR screening. *Nat. Methods* 11, 783–784. [PubMed: 25075903]
- Schwartz JJ, Lee C, and Shendure J. (2012). Accurate gene synthesis with tag-directed retrieval of sequence-verified DNA molecules. *Nat. Methods* 9, 913–915. [PubMed: 22886093]
- Shalem O, Sanjana NE, Hartenian E, Shi X, Scott DA, Mikkelsen T, Heckl D, Ebert BL, Root DE, Doench JG, et al. (2014). Genome-scale CRISPR-Cas9 knockout screening in human cells. *Science* 343, 84–87. [PubMed: 24336571]
- Sims JE, and Smith DE (2010). The IL-1 family: regulators of immunity. *Nat. Rev. Immunol* 10, 89–102. [PubMed: 20081871]
- Wang C, Lu T, Emanuel G, Babcock HP, and Zhuang X. (2019). Imaging-based pooled CRISPR screening reveals regulators of lncRNA localization. *Proc. Natl. Acad. Sci* 116, 10842–10851.
- Wang T, Wei JJ, Sabatini DM, and Lander ES (2014). Genetic screens in human cells using the CRISPR-Cas9 system. *Science* 343, 80–84. [PubMed: 24336569]
- Wang X, Allen WE, Wright MA, Sylwestrak EL, Samusik N, Vesuna S, Evans K, Liu C, Ramakrishnan C, Liu J, et al. (2018). Three-dimensional intact-tissue sequencing of single-cell transcriptional states. *Science* 361, eaat5691.
- Wroblewska A, Dhainaut M, Ben-Zvi B, Rose SA, Park ES, Amir E-AD, Bektesevic A, Baccarini A, Merad M, Rahman AH, et al. (2018). Protein Barcodes Enable High-Dimensional Single-Cell CRISPR Screens. *Cell* 175, 1141–1155.e16. [PubMed: 30343902]
- Xie S, Cooley A, Armendariz D, Zhou P, and Hon GC (2018). Frequent sgRNA-barcode recombination in single-cell perturbation assays. *PloS One* 13, e0198635.
- Yang X, Boehm JS, Yang X, Salehi-Ashtiani K, Hao T, Shen Y, Lubonja R, Thomas SR, Alkan O, Bhimdi T, et al. (2011). A public genome-scale lentiviral expression library of human ORFs. *Nat. Methods* 8, 659–661. [PubMed: 21706014]
- Zuker M. (2003). Mfold web server for nucleic acid folding and hybridization prediction. *Nucleic Acids Res* 31, 3406–3415. [PubMed: 12824337]

Highlights

- *In situ* sequencing of perturbations or barcodes enables image-based pooled screens
- p65 translocation is assayed by imaging in fixed and live cell pools
- Pooled live-cell screen identifies *MED12* and *MED24* as negative regulators of NF- κ B




Figure 1. Optical pooled genetic screens

(A) In pooled screens, a library of genetic perturbations is introduced, typically at a single copy per target cell. In existing approaches, cellular phenotypes are evaluated by bulk NGS of enriched cell populations or single-cell molecular profiling (e.g. single-cell RNA-seq). In optical pooled screens, high-content imaging assays are used to extract rich spatiotemporal information from the sample prior to enzymatic amplification and *in situ* detection of RNA barcodes, enabling linkage between the phenotype and perturbation genotype of each cell.

(B) Targeted *in situ* sequencing is used to read out RNA barcodes expressed from a single genomic integration. Barcode transcripts are fixed in place, reverse transcribed, and hybridized with single-stranded DNA padlock probes, which bind to common sequences flanking the barcode. The 3' arm of the padlock is extended and ligated, copying the barcode into a circularized ssDNA molecule, which then undergoes rolling circle amplification. The barcode sequence is then read out by multiple rounds of *in situ* sequencing-by-synthesis. See also Figure S1A.




Figure 2. Identification of perturbation barcodes by *in situ* sequencing

(A) Schematic of perturbation detection by *in situ* sequencing. Barcodes representing perturbations are expressed on a Pol II transcript and enzymatically converted into cDNA and amplified by RCA. RCA products serve as templates for sequencing-by-synthesis, in which barcodes are read out by multiple cycles of fluorescent nucleotide incorporation, imaging and dye cleavage.

(B) A 125-nt oligo pool encoding perturbations (sgRNAs) and associated 12-nt barcodes was cloned into a lentiviral vector (lentiGuide-BC) and delivered into HeLa cells. Expressed barcode sequences were read out by padlock detection, rolling circle amplification, and 12 cycles of sequencing-by-synthesis. A linear filter (Laplacian-of-Gaussian, kernel width $\sigma = 1$ pixel) was applied to sequencing channels to enhance spot-like features (scale bar 10 μm ; composite image of DAPI and four sequencing channels). See also Movie S1.

(C) >80% of barcodes map to 40 designed sequences out of 16.7 million possible 12-nt sequences. See also Figure S1B-D.

(D) Most cells contain multiple barcode reads that map to the designed library.

(E) Cellular read distribution further categorized by read identity for 77% of cells containing at least one read.

(F) The number of possible barcodes scales geometrically with barcode length. Sufficient 12-nt barcodes can be designed to cover a genome-scale perturbation library while maintaining the ability to detect and reject single or double sequencing errors (minimum pairwise Levenshtein distance $d = 2$ or 3, respectively).




Figure 3. Accuracy of phenotype-to-genotype mapping assessed with a fluorescent reporter

(A) Workflow for CRISPR-Cas9 knockout-based screening of a genetically-encoded frameshift reporter. A library of targeting and non-targeting guides was cloned into either LentiGuide-BC or the CROP-seq vector and transduced into cells at low MOI. Cas9 expression generates indels at the frameshift reporter target locus in cells with a targeting guide and leads to expression of a nuclear-localized HA epitope. HA expression was assayed by immunofluorescence and correlated with sgRNAs detected by *in situ* sequencing. Frameshift reporter accuracy was estimated using the relative abundances of HA⁺ cells mapped to targeting and non-targeting guides (X and Y, respectively). Scale bar is 30 μm.

(B) Targeting and control barcodes expressed from LentiGuide-BC in HeLa-TetR-Cas9 cells were well separated by fraction of HA⁺ cells.

(C) The same cell library was screened by flow sorting cells into HA⁺ and HA⁻ bins and performing next-generation sequencing of the genomically integrated barcode. See also Figure S3D.

(D) The experiment was repeated across a panel of cell lines using the CROP-seq library and an optimized padlock detection protocol, yielding a similar distribution of mapped reads (top) and frameshift reporter accuracies (bottom). Error bars indicate the range between two replicate sequencing experiments. Cell types are indicated by the same colors in both plots.




Figure 4. A screen for regulators of NF- κ B signaling

(A) Workflow for CRISPR-Cas9 knockout-based screening using a fluorescently tagged reporter cell line. Screen hits were identified by the failure of p65-mNeonGreen to translocate to the nucleus following stimulation with IL-1 β or TNF α cytokines.

(B) Known NF- κ B regulators were identified as high-ranking screen hits. Cells were assigned translocation scores based on the pixelwise correlation between mNeonGreen fluorescence and a DAPI nuclear stain; thus a score 1 indicates maximum translocation while a score of -1 indicates maximum cytoplasmic localization. The translocation defect for a gene was

defined based on the integrated difference in the distribution of translocation scores relative to non-targeting control sgRNAs across three replicate screens. See also Tables S1 and S3. (C) Cumulative distributions of translocation scores (second-ranked guide) of known $\text{NF-}\kappa\text{B}$ regulators in response to both cytokines. The shaded areas depict the difference between the translocation score distributions for targeting sgRNAs and non-targeting control sgRNAs (gray).

(D) $\text{NF-}\kappa\text{B}$ pathway map (KEGG HSA04064) color-coded as in (B). KEGG pathway members colored gray did not show a translocation defect when individually knocked out in HeLa cells.

(E) Top-ranked genes were validated with arrayed CRISPR-Cas9 knockouts (scale bar 10 μm). Histograms show the cumulative distributions of IL-1 β and TNF α -induced translocation scores (averaged over two guides) for each gene knockout compared to wild type cells (gray). See also Figure S4B.




Figure 5. A live-cell screen identifies a role for Mediator components in regulating p65 kinetics
 (A) The initial 952 gene translocation screen was repeated using live-cell imaging to monitor p65 translocation kinetics (n = 361,587 analyzed cells). Hierarchical clustering of the time-dependent translocation difference between each gene and the non-targeting controls grouped together KEGG-annotated regulators, as well as other positive and negative regulators with distinct cytokine-specific kinetic signatures. See also Table S4.
 (B) and (C) A validation live-cell screen was performed to improve sampling of regulators identified by the primary screen and helped group regulators with distinct p65 translocation kinetics. Individual traces represent different sgRNAs for each gene. See also Figure S5 and Table S4.

(D) Clonal knockouts of *MED12* and *MED24* recapitulated the increased retention time phenotype seen in the primary screen. Each trace represents a different knockout clone (scale bar 10 μm).

(E) Cytokine stimulation of Mediator clonal knockouts led to differential activation of $\text{NF-}\kappa\text{B}$ target genes, including increased expression of chemokines after stimulation with $\text{TNF}\alpha$ (1 ng/mL) or $\text{IL-1}\beta$ (30 ng/mL). Error bars show the range among knockout clones or wild type biological replicates, clones same as in (D).

(F) *IL1B* expression was induced in the Mediator clonal knockouts, but not in wild type cells, by $\text{TNF}\alpha$ stimulation.

KEY RESOURCES TABLE

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
rabbit anti-HA antibody	Cell Signaling Technologies	3724
rabbit anti-p65 antibody	Cell Signaling Technologies	8242
goat anti-rabbit IgG Alexa Fluor 488	Cell Signaling Technologies	4412
Bacterial and Virus Strains		
Endura Electrocompetent Cells	Lucigen	69242
Chemicals, Peptides, and Recombinant Proteins		
TNF α	Invivogen	rcyc-htnfa
IL-1 β	Invivogen	rcyec-hil1b
Critical Commercial Assays		
Revertaid H minus RT	Thermo Fisher Scientific	EP0452
Ribolock RNase inhibitor	Thermo Fisher Scientific	EO0384
RNase H	Enzymatics / Qiagen Beverly	Y9220L
TaqIT DNA polymerase	Enzymatics / Qiagen Beverly	P7620L
Stoffel Fragment	Blirt SA / DNA Gdansk	RP810
Ampligase	Lucigen	A3210K
Phi29 DNA polymerase	Thermo Fisher Scientific	EP0092
MiSeq 500 cycle Nano kit	Illumina	MS-103-1003
Deposited Data		
RNA-seq of clonal knockouts	Gene Expression Omnibus	GSE132704
Screening image data	Image Data Resource	Accession number pending
Experimental Models: Cell Lines		
HeLa-TetR-Cas9	Iain Cheeseman Lab, Whitehead Institute	N/A
Oligonucleotides		
oRT_LentiGuide-BC (for use with LentiGuide-BC): G+AC+GT+GT+GC+TT+AC+CCAAAGG	This paper	N/A
oPD_LentiGuide-BC (for use with LentiGuide-BC): /5Phos/ actggetattcattcgcCTCCTGTTCGA CAGTCAGCCGCATCTGCGTCTA1 1 IAGTGGAGCCCC TT Ggttcaatcaacattcc	This paper	N/A
oSBS_LentiGuide-BC (for use with oPD_LentiGuide- BC): TTCGACAGTCAGCCGCATCTGCGTCTATTAGTGG AGCCCTT Ggttcaatcaacattcc	This paper	N/A
oRT_CROPseq (for use with CROPseq-puro and CROPseq-zeo): G+AC+TA+GC+CT+TA+TT+TTAACTTGCTAT	This paper	N/A

REAGENT or RESOURCE	SOURCE	IDENTIFIER
oPD_CROPseq (for use with CROPseq-puro and CROPseq-zeo, optimized padlock): /5Phos/gttttagagctagaatagcaagCTCCTGTTCGACACC TACCCACCTCATCCCCTCTTCAaaaggacgaaacaccg	This paper	N/A
oSBS_CROPseq (for use with oPD_CROPseq): CACCTCATCCCCTCTTCAaaaggacgaaacaccg	This paper	N/A
oRT_CROPseq-v2 (for use with CROPseq-puro-v2): A+CT+CG+GT+GC +CA+CT+TTT	This paper	N/A
oPD_CROPseq-v2 (for use with CROPseq-puro-v2, optimized padlock):/ 5Phos/GTTTCAGAGCTATGCTGGCTCCTGT TCGCTTCTCCCTTACCTCCTTCCCTTCCATCCTATAT CCTCCACTCATAaaaggacgaaacaccg	This paper	N/A
oSBS_CROPseq-v2 (for use with oPD_CROPseq-v2): TCCATCCTATATCCTCCACTCATAaaaggacgaaacaccg	This paper	N/A
oPD_323 (previously used with CROPseq-puro; not optimized): /5Phos/ gttttagagctagaatagcCTCCTGTTCGACAGTCA GCCGCATCTGCGTCTATTTAGTGGAGCCCTTgaagga cgaaacaccg	This paper	N/A
oSBS_394 (previously used with oPD_323; not optimized): TCAGCCGCATCTGCGTCTATTTAGTGGAGCCCTTgaaggacgaaacaccg	This paper	N/A
Recombinant DNA		
LentiCas9-blast	Sanjana et al, 2014	Addgene Plasmid #52962
LentiGuide-Puro	Sanjana et al, 2014	Addgene Plasmid #52963
pXPR_011	Doench et al, 2014	Addgene Plasmid #59702
CROPseq-Guide-Puro	Datlinger et al., 2017	Addgene Plasmid #86708
pR_LG	Feldman et al., 2018	Addgene Plasmid #112895
LentiGuide-BC	This paper	Addgene Plasmid #127168
LentiGuide-BC-CMV-Puro	This paper	Addgene Plasmid #127169
LentiGuide-BC-CMV-Puro	This paper	Addgene Plasmid #127170
pL_FR_Hygro	This paper	Addgene Plasmid #127171
pR14_p65-mNeonGreen	This paper	Addgene Plasmid #127172
CROPseq-Zeo	This paper	Addgene Plasmid #127173
CROPseq-Puro-v2	This paper	N/A
Software and Algorithms		
Snakemake	Köster et al., 2012	https://snakemake.readthedocs.io/en/stable/
Sample <i>in situ</i> sequencing data and analysis pipeline from data to sequencing read table	This paper	https://github.com/blaineylab/OpticalPooledScreens
Other		
Microscope light source	Lumencor	Sola SE365 FISH
Camera	Hamamatsu	ORCA-Flash 4.0 v3
Objective Lens	Nikon	CFI Plan Apochromat Lambda 10X/ 0.45
DAPI filter set	Semrock	LED-DAPI-A-NTE-ZERO
GFP filter set	Semrock	GFP-1828A-NTE-ZERO
Cy3 filter set (MiSeq G)	Semrock	FF01-534/20-25 FF552-Di02-25×36 FF01-572/28-25
Alexa Fluor 594 filter set (MiSeq T)	Semrock	FF03-575/25-25 FF596-Di01-25×36 FF01-615/24-25

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Cy5 filter set (MiSeq A)	Semrock	FF01-635/18-25 FF652-DI01-25×36 FF01-680/42-25
Cy7 filter set (MiSeq C)	Semrock	FF01-661/20-25 FF695-DI01-25X36 FF01-732/68-25
Sample <i>in situ</i> sequencing data and analysis pipeline from data to sequencing read table	This paper	https://github.com/blaineylab/OpticalPooledScreens

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript