

Published in final edited form as:

Leukemia. 2019 December ; 33(12): 2884–2897. doi:10.1038/s41375-019-0478-9.

Spingosine-1-phosphate Signalling drives an Angiogenic Transcriptional Programme in Diffuse Large B Cell Lymphoma

Lauren Lupino¹, Tracey Perry¹, Sandra Margielewska¹, Robert Hollows¹, Maha Ibrahim^{1,2}, Matthew Care³, Jeremy Allegood⁴, Reuben Tooze³, Roger Sabbadini⁵, Gary Reynolds⁶, Roy Bicknell⁷, Zbigniew Rudzki⁸, Ye Lin Hock⁹, Ulises Zanetto¹⁰, Wenbin Wei^{1,11}, William Simmons¹, Sarah Spiegel⁴, Ciaran BJ Woodman¹, Martin Rowe¹, Katerina Vrzalikova^{1,†}, Paul G Murray^{1,12,†}

¹Institute of Cancer and Genomic Sciences, University of Birmingham, Birmingham, United Kingdom

²South Egypt Cancer Institute, Assiut University, Assiut, Egypt

³Section of Experimental Haematology, Leeds Institute of Cancer and Pathology, University of Leeds, Leeds, United Kingdom

⁴Department of Biochemistry and Molecular Biology and Massey Cancer, Virginia Commonwealth University School of Medicine, Richmond, VA, USA

⁵Department of Biology, San Diego State University, San Diego, CA, USA

⁶Institute of Immunology & Immunotherapy, University of Birmingham, Birmingham, United Kingdom

⁷Institutes of Cardiovascular Sciences and Biomedical Research, University of Birmingham, Birmingham, United Kingdom

⁸Department of Histopathology, Heartlands Hospital, Birmingham, United Kingdom

⁹Department of Histopathology, Walsall Manor Hospital, Walsall, United Kingdom

¹⁰Department of Histopathology, City Hospital, Birmingham, United Kingdom

¹¹Sheffield Institute of Translational Neuroscience, University of Sheffield, Sheffield, United Kingdom

¹²Department of Clinical and Molecular Pathology, Institute of Molecular and Translational Medicine, Faculty of Medicine and Dentistry, Palacky University, Olomouc, Czech Republic

Users may view, print, copy, and download text and data-mine the content in such documents, for the purposes of academic research, subject always to the full Conditions of use:http://www.nature.com/authors/editorial_policies/license.html#terms

†Joint corresponding authors: Paul Murray and Katerina Vrzalikova, Institute of Cancer and Genomic Sciences, University of Birmingham, Vincent Drive, Birmingham, B15 2TT, United Kingdom, Tel: (44-121) 414 4021, Fax: (44-121) 414 4486, p.g.murray@bham.ac.uk, k.vrzalikova@bham.ac.uk.

Authorship Contributions:

LL, CBJW, RS, MR, KV, RB, SS, PGM designed research; LL, KV, TP, SM, MI, GR, WS, ZR, YLH, UZ, JA performed research and analysed data; RH, MC, RT, WW, PGM contributed to statistical analysis; ZR, YLH, WS, UZ undertook pathology evaluation; LL, KV, SS, PGM wrote the manuscript.

Conflict of Interest Disclosures: Authors declare no competing financial interests.

Abstract

Although the over-expression of angiogenic factors is reported in diffuse large B-cell lymphoma (DLBCL), the poor response to anti-VEGF drugs observed in clinical trials suggests that angiogenesis in these tumours might be driven by VEGF-independent pathways. We show that sphingosine kinase-1 (SPHK1), which generates the potent bioactive sphingolipid sphingosine-1-phosphate (S1P), is over-expressed in DLBCL. A meta-analysis of over 2000 cases revealed that genes correlated with SPHK1 mRNA expression in DLBCL were significantly enriched for tumour angiogenesis meta-signature genes; an effect evident in both major cell of origin (COO) and stromal subtypes. Moreover, we found that S1P induces angiogenic signalling and a gene expression programme that is present within the tumour vasculature of SPHK1-expressing DLBCL. Importantly, S1PR1 functional antagonists, including Siponimod, and the S1P neutralising antibody, Sphingomab, inhibited S1P signalling in DLBCL cells *in vitro*. Furthermore, Siponimod, also reduced angiogenesis and tumour growth in an S1P-producing mouse model of angiogenic DLBCL. Our data define a potential role for S1P signalling in driving an angiogenic gene expression programme in the tumour vasculature of DLBCL and suggest novel opportunities to target S1P-mediated angiogenesis in patients with DLBCL.

Keywords

S1P; SPHK1; DLBCL; HUVEC; endothelial cells

Introduction

Despite the use of rituximab and chemotherapy (R-CHOP), the survival of patients with DLBCL remains poor with a significant proportion of patients being refractory or relapsing after therapy [1, 2]. Gene expression profiling has identified two clinically distinct DLBCL subgroups, the activated B cell (ABC) and germinal centre B cell (GCB) forms, characterised by different, therapeutically tractable, molecular abnormalities. Gene expression profiling has also identified tumour microenvironment-derived molecular signatures, including the Stromal-2 signature that is reported to be enriched for angiogenesis-associated genes [3].

Although there may be therapeutic benefits in targeting angiogenesis in DLBCL, the results of clinical trials of bevacizumab, a monoclonal antibody against VEGFA, have been disappointing. Although this drug was well tolerated when delivered as a single agent in relapsed DLBCL [4] or in combination with R-CHOP in first line treatment [5], a phase III study of RA-CHOP (MAIN trial) was stopped due to increased cardiotoxicity in the absence of any significant impact on progression-free survival [6]. One explanation for the limited clinical efficacy of bevacizumab in DLBCL is that angiogenesis in DLBCL does not depend on VEGF; a contention supported by studies showing that VEGF expression does not correlate with micro-vessel density in DLBCL [7, 8].

S1P is a bioactive sphingolipid metabolite which promotes cell growth and survival and has been shown to be a potent inducer of angiogenesis [9]. SPHK1, one of the two isoenzymes responsible for the production of S1P, is over-expressed in different cancers [10, 11].

Secreted S1P is a ligand for a family of five G-protein-coupled S1P receptors (S1PR1-5). Signalling through S1PR1 and/or S1PR3 has been shown to stimulate DNA synthesis, chemotactic motility and tube formation in endothelial cells and to induce angiogenesis *in vivo* [12–15]. As with other potent bioactive mediators, S1P levels are tightly regulated and controlled by the balance between its generation and its degradation by S1P lyase and S1P phosphatases [16]. Although it has been suggested that SPHK1 might play a role in haematological malignancies [17], its role in DLBCL remains to be established [18]. Furthermore, the effects of S1P signalling on the DLBCL microenvironment, including its influence on the tumour vasculature, have not been explored.

In the present study we have shown that the over-expression of SPHK1 correlates with an angiogenic transcriptional programme in DLBCL. We defined an endothelial cell transcriptional signature of S1P signalling and used this to show that the expression of S1P target genes in these cells was correlated with that of SPHK1 in primary DLBCL. Moreover, Siponimod, a small-molecule functional antagonist of S1PR1 [19], reversed S1P signalling and reduced angiogenesis and tumour growth in an S1P-producing mouse model of DLBCL. Our data suggest novel opportunities to target S1P-mediated angiogenesis in patients with DLBCL.

Materials and Methods

Cells and tissues

Tonsils and DLBCL samples were obtained with informed consent and ethical approval (REC_RG_HBRC_12-071). DLBCL cases were reviewed by haematopathologists (ZR, YLH, UZ). Isolation of tonsillar germinal centre (GC) and blood-derived B cells was described before [20–22]. Endothelial cells (EC) were isolated from umbilical cords (HUVEC) under informed consent (REC_RG_HBRC_14-180) using collagenase treatment [23] and cultured in M199 media supplemented with 5% fetal bovine serum (FBS), 1% glutamine, 1% penicillin/streptomycin (ThermoFisher Scientific, Waltham, MA, USA) and 1% EC-growth supplement (Caltag Medsystems, Buckingham, UK) at 37°C/5% CO₂. HT, Karpas-442, OCI-LY1, OCI-LY7, SUDHL4, SUDHL5, SUDHL6 are EBV-negative GC-DLBCL lines, Farage is an EBV-positive GC-DLBCL line. OCI-LY3 and U2932 are EBV-negative ABC-DLBCL lines. Lines were from DSMZ (Braunschweig, Germany), OCI (Ontario, Canada) or ATCC (Manassas, VA, USA) and were cultured in RPMI1640 or IMDM (OCI-LY1, OCI-LY7) media (ThermoFisher Scientific) supplemented with 10% FBS and 1% penicillin/streptomycin.

Mouse xenografts and flow cytometry

3x10⁶ SUDHL6 cells were injected subcutaneously into NSG mice (Charles River Laboratories, Wilmington, MA, USA). After 17 days (when tumour volume averaged 63mm³) mice were randomised into two groups (each n=4) and treated orally with either vehicle (0.1% DMSO in 10% 2-hydroxypropyl-β-cyclodextrin; Cayman Chemical, MI, USA) or 6mg/kg Siponimod (Selleckchem.com, Munich, Germany) every 48h. Mice were culled when average tumour volumes in control mice reached 400mm³ (28 days). Organs were weighed, minced and incubated with Liberase DL/Liberase TL and DNASEI (Roche,

Basel, Switzerland) [24]. Cell suspensions were labelled with mouse CD31 and CountBright absolute counting beads (ThermoFisher Scientific) and analysed by flow cytometry on LSRII and FACS diva 8 (BD, Franklin Lakes, NJ, USA). Details of the other mouse models tested are in Supplementary Materials and Methods. All mouse experiments were done according to UK Home Office guidelines.

S1P measurements

For intracellular S1P measurements, cell pellets were snap-frozen in liquid nitrogen. For secreted S1P measurements, SUDHL4 cells were cultured in serum-free RPMI (without phenol-red) supplemented with 1% tissue-culture grade fatty acid-free BSA (Sigma-Aldrich., St Louis, MO, USA) for indicated times. Supernatants were harvested into pre-chilled HPLC grade methanol (Sigma-Aldrich) supplemented with Halt™ Protease and Phosphatase Inhibitor Cocktail (ThermoFisher Scientific). S1P levels were quantified by liquid chromatography-electrospray ionization-tandem mass spectrometry (LC-ESI-MS/MS; 4000 QTRAP, AB Sciex, Framingham, MA, USA) as previously described [25].

Treatment of cells

S1P (Sigma-Aldrich) was prepared as in Supplementary Materials and Methods and as before [26]. Prior to treatments, HUVEC were cultured in full media depleted of EC-growth supplement for 16h and stimulated with 0.5µM S1P (or control/vehicle) for 5min to detect ERK1/2 activation or for 4h to detect S1P transcriptional targets. For S1P inhibition, Sphingomab or isotype/control antibodies (LPath Inc., San Diego, CA, USA) were incubated with S1P (at 150µg/ml per µM of S1P) for 1h prior to S1P stimulation. For S1PR1 inhibition, HUVEC were treated with 100nM S1PR1 functional antagonists Siponimod/BAF312, Ozanimod/RPC1063 or Ponesimod/ACT-128800 (Selleckchem.com) for 1h prior to S1P stimulation.

Quantitative RT-PCR (Q-PCR)

RNA was extracted with RNeasy Mini kit including genomic DNA removal with RNase-Free DNase Set (Qiagen, Manchester, UK). cDNA was generated with qScript™ cDNA SuperMix (QuantaBio, Beverly, MA, USA). Gene transcripts were quantified with commercial gene expression assays (ThermoFisher Scientific; Supplementary Table 1a) [21]. The 2^{-CT} method was used to quantify target expression relative to housekeeping control. The normalized values are shown relative to the reference sample that was set to a relative quantity value of 1.

Protein analysis

Immunoblotting was by standard methodology [20]. Detection was with enhanced chemiluminescence (ECL; GE Healthcare, Chalfont St Giles, UK) on a ChemiDoc MP (Bio-Rad, Watford, UK). Immunohistochemistry (IHC) was on 4µm FFPE sections and HUVEC grown on microscope slides, using either citrate or EDTA buffer antigen retrieval and 0.3% H₂O₂ and 5x casein blocking [26–28]. Slides were incubated either overnight at 4°C or for 1 hour at room temperature in primary antibodies in 0.05% PBS/Tween and visualised with species-specific ImmPRESS kits, followed by diaminobenzidine or ImmPACT NovaRED

(Vector Laboratories, Peterborough, UK) with haematoxylin counterstain. DLBCL were stained for BCL6, CD10 and IRF4, and defined as either GCB or non-GCB type by Hans algorithm [29]. A subset of DLBCL was also stained for SPHK1, S1PR1, CXCL12, SELE, COL1A1 and MAP1B. Mouse tumours were stained for Ki67, cPARP and CXCL12. For CXCL12, a Mouse-on-Mouse block (Vector Laboratories) was performed prior to incubation in casein. cPARP- and Ki67- positive cells were enumerated in at least 10 high power fields per tumour. Photomicrographs were acquired on a Nikon Eclipse E400 microscope with x20 or x40 objectives (NA 0.40 and 0.65) at room temperature and a Nikon DS-Fi-1 camera. Supplementary Table 1b lists antibodies used for immunoblotting and IHC.

RNAseq analysis

RNAseq was performed on 4 biological replicates of HUVEC treated with S1P or vehicle/control for 4h. Quality control (RNA integrity number >7), library construction and sequencing was performed by BGI Tech Solutions (Hong Kong). RNA was hybridised to Illumina HiSeq 2000 platform and data obtained using 10 M clean reads.

Analysis of gene expression

RNAseq data were aligned to the hg19 human genome using Rsubread aligner [30] and assigned to genes with featureCounts function. Read counts normalised between samples were converted to counts-per-million (CPM) reads using the edgeR package in R [31]. RNAseq data for 32 ABC and 54 GCB DLBCL were from the controlled access area of NIH database of genotypes and phenotypes (dbGaP; accession code phs000532.v5.p2) [32]. The RNAseq data for 4 GC B cell samples were from GEO (GSE45982) [33]. Differentially expressed genes were identified using edgeR with $p < 0.05$ and read CPM > 1 in at least half of the samples. Microarray data for 11 DLBCL and 10 GC B cell samples were from GEO (GSE12453)[34] and analysed with MAS5 algorithm of the Affymetrix Expression Console to generate expression levels for each probe set (GCOS Signal). The MAS5 TGT was set to 100. The series matrix expression and clinical data reported in Lenz et al [3] were downloaded from GEO (GSE10846). Meta-analysis of 11 different DLBCL datasets was performed as before [35]. Spearman test was used to correlate gene expression between samples.

Statistics

Statistical tests are indicated in relevant sections. All experiments were performed at least in triplicate, and for B cells on at least three separate donors. Tests were considered statistically significant if $p < 0.05$.

Results

SPHK1 is over-expressed in the tumour cells of DLBCL

To explore the contribution of S1P signalling to angiogenesis in DLBCL, we examined the expression of SPHK1, an enzyme which generates S1P from sphingosine. We measured SPHK1 mRNA in published datasets, comparing global gene expression in DLBCL with that in normal GC B cells [32, 34]. SPHK1 mRNA levels were significantly higher in

DLBCL compared to GC B cells (Figure 1a) and in both ABC- and GCB- DLBCL subtypes when analysed separately (Figure 1b).

We performed IHC using an antibody we previously showed is specific for SPHK1 [26]. We found that SPHK1 was present in normal GC B cells, and in the tumour cells of all 32 cases of DLBCL examined; including cases of both GCB and non-GCB type defined by the Hans algorithm [29] (Figure 1c; Supplementary Table 2). The intensity of SPHK1 expression in tumour cells was variable but was generally higher than in background of non-malignant cells, including macrophages and small reactive lymphocytes (not shown). Furthermore, in contrast to the strong staining observed in tumour cells, SPHK1 was either only weakly detectable or undetectable in EC of most (27/31) evaluable cases (Supplementary Table 2; Figure 1c).

SPHK1 mRNA and protein were also expressed in DLBCL lines (Figure 1d). Because the catalytic activity of SPHK1 is enhanced by its phosphorylation at Ser225 we used an antibody specific for this phosphorylation site. [36], [26]. Immunoblotting with this antibody revealed that SPHK1 was phosphorylated in all cell lines examined (Figure 1d; lower panel and Supplementary Figure 1A). S1P levels were significantly higher in all six DLBCL cell lines examined compared to normal B cells (Figure 1e). Because S1P is a ligand for S1P receptors present on the cell surface, it was of interest to determine whether DLBCL cells such as SUDHL4, which express SPHK1 and produce S1P intracellularly, can also secrete it. We found that DLBCL cells readily export S1P outside the cells reaching a maximum level of around 1nM within 2min (Figure 1f).

Finally, we stained the DLBCL cases for S1PR1. We found that S1PR1 was expressed on endothelial cells in all 32 cases and in tumour cells in 20/32 cases (Supplementary Table 2, Supplementary Figure 1B upper panels). In keeping with this, Q-PCR revealed that some, but not all, DLBCL cell lines expressed S1PR1 (Supplementary Figure 1B, lower panel). In primary tumours, there was no relationship between tumour cell expression of S1PR1 and disease subtype.

SPHK1 expression is associated with an angiogenic transcriptome in DLBCL

We next performed a meta-analysis of 11 primary DLBCL gene expression datasets comprising over 2000 cases of DLBCL [35]. For each data set, genes were ordered by their variance across the patient samples and the top 80% were used to calculate the Spearman's rank correlations for their expression against that of SPHK1. The resultant p values and correlation matrices were merged across the 11 datasets by taking the median values. A SPHK1-correlated gene set was created by taking all genes which were present in six or more datasets with a median $p < 0.05$. This identified 2236 genes positively correlated and 1658 genes negatively correlated with SPHK1 expression in DLBCL (Supplementary Table 3). We noted that genes positively correlated with SPHK1 included well known angiogenesis-associated genes, such as VEGF and VEGFR ($r=0.28$, $p=0.0001$; and $r=0.19$, $p=0.0042$, respectively). A gene ontology (GO) analysis [37] also showed a significant enrichment of angiogenesis and vasculature functions among genes positively correlated with SPHK1 (Figure 2a). To provide further confirmation of a relationship between SPHK1 expression and angiogenesis, we utilised a published 'tumour vascular gene signature'

generated from a meta-analysis of more than 1000 primary human cancers [38]. We found that genes positively correlated with SPHK1 expression in DLBCL were significantly enriched for tumour vascular signature genes (odds ratio (OR)=7.92, $p<0.0001$), whereas genes negatively correlated with SPHK1 were significantly depleted for tumour vascular signature genes (OR=0.37, $p<0.0001$; Figure 2b).

Repeating the meta-analysis, we found that tumour vascular signature genes were significantly enriched among genes positively correlated with SPHK1 expression in GCB and ABC subtypes when analysed separately (both $p<0.0001$; Supplementary Figure 2). As a further check on the association between SPHK1 and angiogenesis, we used two further gene signatures of angiogenesis (the “Hallmark” angiogenic signature, M5944, and the signature for the GO term ‘angiogenesis’). For both alternative angiogenic gene sets, we again found a significant overlap with genes positively correlated with SPHK1 (both $p<0.0001$, not shown). Finally, we directly compared the mRNA expression of SPHK1 with that of classical EC markers in a further published RNAseq dataset reported by Morin et al., [32] not included in our original meta-analysis. This revealed a significant positive correlation between the expression of SPHK1 and that of CD34, CDH5, PECAM1/CD31 and VWF (Figure 2c; Supplementary Figure 3).

We also used the same 11 primary DLBCL datasets described above to perform a meta-analysis of genes correlated with S1PR1 in DLBCL. In keeping with the observed strong expression of S1PR1 in the tumour vasculature of DLBCL, we found that genes positively correlated with S1PR1 in primary tumours were also significantly enriched for tumour vascular signature genes (OR=7.92, $p<0.0001$; not shown). Furthermore, we also found that S1PR1 was positively correlated with CD34, (Spearman test, $r=0.48$; $p<0.0001$), CDH5 ($r=0.42$; $p<0.0001$), PECAM1 ($r=0.46$; $p<0.0001$), and VWF ($r=0.44$; $p<0.0001$) in the dataset reported by Morin et al [32].

We conclude that the expression of both SPHK1 and S1PR1 is associated with an angiogenic transcriptome in DLBCL.

SPHK1 expression is associated with angiogenesis in both Stromal-1 and Stromal-2 DLBCL subtypes

Given that angiogenesis was reported to be associated with the Stromal-2 signature in DLBCL [3], we next explored the correlation of SPHK1 expression with the expression of stromal signature genes [3]. First, we studied the extent to which Stromal-2 genes overlapped with the angiogenic gene signatures. As expected, we found a significant overlap of Stromal-2 signature genes, but only with two of the three angiogenic gene sets (Figure 3a; upper panel). Consistent with this, we found that SPHK1 expression was positively correlated with Stromal-2 gene expression (calculated as the average expression across genes in the Stromal-2 signature [3] (Spearman correlation=0.46 and 0.47 for CHOP- and R-CHOP-treated patients, respectively; $p<0.0001$, not shown). However, when we repeated this analysis, this time using Stromal-1 signature genes [3], we were surprised to find that SPHK1 expression was even more highly correlated with Stromal-1 gene expression (Spearman correlation=0.80 and 0.77, respectively, $p<0.0001$) and included a highly significant overlap with all three angiogenic gene sets (Figure 3b; lower panel). In contrast,

we observed that S1PR1 expression was positively correlated only with Stromal-2 gene expression (Spearman correlation=0.34 and 0.45 for CHOP- and R-CHOP-treated patients, respectively; both $p < 0.0001$), and not with Stromal-1-gene expression (Spearman correlation $r = 0.02$, $p = 0.807$; and $r = 0.09$, $p = 0.212$, for CHOP- and R-CHOP-treated patients, respectively). We conclude that genes positively correlated with SPHK1 are enriched in both stromal signatures of DLBCL, most likely reflecting the contribution of SPHK1 to angiogenesis in both stromal subgroups.

An *in vitro* transcriptional signature of S1P signalling in endothelial cells

With the aim of revealing a more direct association between S1P signalling and angiogenesis in DLBCL, we next defined a transcriptional signature of S1P signalling in EC. To do this we used human umbilical vein endothelial cells (HUVEC) which we showed express the same pattern of S1P receptors as the EC of DLBCL, marked by the high expression of S1PR1 and the absence of S1PR2 and S1PR3 in most cases (Figure 4a). We showed that S1P treatment of HUVEC induced the robust phosphorylation of ERK1/2, a well-defined downstream target of S1P signalling in EC [12] (Figure 4b).

We next used RNAseq to describe the global transcriptional consequences of S1P signalling in HUVEC. Treatment of HUVEC with S1P was followed by the up-regulation of 115 genes and the down-regulation of 126 genes (Supplementary Table 4). We used Q-PCR to validate a subset of the genes which were both up-regulated by S1P and positively correlated with SPHK1 in our meta-analysis, including IL-8, ICAM1, SELE, PDGFA, ANGPTL4 and CXCL12 (Figure 4c).

To confirm the specificity of these effects, we repeated these experiments in the presence of Spingomab, a monoclonal antibody which depletes extracellular S1P [39]. Spingomab greatly suppressed S1P-induced phosphorylation of ERK1/2 (Figure 4d) and the up-regulation of its downstream transcriptional targets (Figure 4e). In contrast, S1P induced the phosphorylation of ERK1/2 and the up-regulation of its target genes in the presence of the isotype control antibody.

Genes up-regulated by S1P signalling and correlated with SPHK1 mRNA are expressed in the tumour vasculature of DLBCL

A gene ontology analysis revealed that genes up-regulated by S1P in HUVEC were significantly enriched for GO terms associated with angiogenesis, and for GO terms associated with the known effects of S1P in EC, including protection from apoptosis, leukocyte adhesion and migration (Figure 5a). Furthermore, genes up-regulated (OR=4.30, $p < 0.0001$), but not those down-regulated (OR=0.78, $p = 0.48$), by S1P were significantly enriched among genes positively correlated with SPHK1 in DLBCL (Figure 5b, left panel). Similarly, genes downregulated (OR=2.58, $p < 0.0001$), but not those up-regulated (OR=0.73, $p = 0.48$), by S1P were significantly enriched in genes negatively correlated with SPHK1 (Figure 5b, right panel). The enrichment of up-regulated S1P targets among genes positively correlated with SPHK1 was also evident when the GCB and ABC subtypes were analysed separately (Supplementary Figure 4).

To provide direct evidence of the expression of S1P target genes in the tumour vasculature of DLBCL we performed IHC for CXCL12 and SELE, as well as COL1A1 and MAP1B, two further genes up-regulated in S1P-treated HUVEC, positively correlated with SPHK1 expression in DLBCL, and that had available antibodies that work robustly in FFPE tissues. We found that all four markers were expressed in the EC of both subtypes of DLBCL. In contrast, tumour cells were either negative (CXCL12, COL1A1, SELE) or showed only weak expression (MAP1B) (Supplementary Table 5). In summary, these data provide evidence of an S1P-regulated transcriptional programme in the tumour vasculature of DLBCL.

S1PR1 inhibition reverses S1P signalling *in vitro* and reduces angiogenesis and tumour growth in a mouse model of angiogenic DLBCL

Finally, we explored the effects of inhibiting S1P signalling in a mouse model of DLBCL. We initially studied ERK1/2 phosphorylation *in vitro* following the stimulation of HUVEC with S1P, in the presence or absence of Siponimod, Ozanimod and Ponesimod, three functional antagonists of S1PR1 [19, 40, 41]. We found that all three drugs decreased ERK1/2 phosphorylation in HUVEC, confirming their ability to reduce the effects of S1P on downstream signalling (Figure 6a). We next studied the impact of inhibiting S1P signalling in a mouse model of angiogenic DLBCL, focussing on Siponimod, a potent functional antagonist of S1PR1 and S1PR5 (EC50:0.39nM and 0.98nM, respectively) with >1000-fold selectivity for S1PR1 versus S1PR2, S1PR3 and S1PR4 [42]. For these experiments we used SUDHL6 cells, not only because they produce high levels of S1P (Figure 1e), but also because as xenografts they induce levels of angiogenesis that are almost 8-fold-higher than those found in xenografts of OCI-LY1 cells, and nearly 60-fold higher than those observed in the A20 syngeneic mouse model of DLBCL (Supplementary Figure 5A). We treated established xenografts of SUDHL6 cells with Siponimod for up to two weeks before culling the animals and measuring mouse EC numbers in tumours by flow cytometry. We found that compared to controls, treatment of Siponimod significantly reduced the numbers of CD31-positive mouse EC in tumour tissues (Figure 6b) as well as reducing their expression of the S1P target gene, CXCL12 (Supplementary Figure 5B). Furthermore, Siponimod led to a significant reduction in tumour volumes in treated animals at the later time points (Figure 6c) that was associated with increased numbers of apoptotic cells as measured by immunohistochemistry for cleaved PARP (Supplementary Figure 5C). In contrast, we did not observe significantly reduced tumour volumes following Siponimod treatment of xenografts of the poorly angiogenic OCI-LY1 and A20 lines (Supplementary Figures 5D and 5E). We conclude that the inhibition of S1P signalling can reduce angiogenesis and tumour growth in a mouse model of angiogenic DLBCL.

Discussion

Although previous studies have shown that targeting VEGF or its receptors can decrease tumour vascularization and reduce the growth of DLBCL-derived cell lines *in vivo*, a phase III trial of patients with aggressive NHL receiving R-CHOP plus bevacizumab showed cardiotoxicity without improvement in progression-free survival [6]. We speculated that DLBCL-associated angiogenesis could be driven by VEGF-independent signalling; a

contention supported by other studies which reveal no correlation between lymphoma cell VEGF expression and micro-vessel density in primary DLBCL [7, 8]. Here we have focused on the potential contribution of the potent bioactive sphingolipid metabolite, S1P. We have shown that SPHK1, the major enzyme responsible for the production of S1P, which can be secreted, is over-expressed in DLBCL. SPHK1 is strongly implicated in tumour angiogenesis in other tumours [43], and is important for the generation of S1P that drives tumour-induced hemangiogenesis and lymphangiogenesis in murine models of breast cancer [44]. The underlying mechanisms responsible for the over-expression of SPHK1 in DLBCL are not known but do not usually involve amplification and/or mutation; our review of five separate cohorts of DLBCL has revealed that the frequency of amplification and/or mutation of the SPHK1 gene is very low (<1%; not shown; TCGA Research Network: <http://cancergenome.nih.gov/>; [32, 45–47]).

SPHK1 can be activated by growth factors and cytokines [48–52], and its catalytic activity is enhanced by ERK1/2-mediated phosphorylation at Ser225 [36]. The pSer225 antibody used in our study showed that SPHK1 was constitutively phosphorylated at Ser225 in DLBCL cell lines. In keeping with this, we showed that S1P production was increased in DLBCL cell lines compared to normal B cells, and furthermore that this S1P was secreted into the extracellular environment.

Our meta-analysis of DLBCL revealed a strong correlation between the expression of SPHK1 and that of angiogenesis meta-signature and EC-classifier genes. One explanation for this finding could be expression of SPHK1 in tumour-associated EC. However, we observed SPHK1 expression predominantly in cancer cells, consistent with the notion that SPHK1 expression is upregulated primarily in the tumour cell population [18].

Although the meta-analysis and our *in situ* expression data suggested that angiogenesis in DLBCL is driven by tumour-derived SPHK1, these approaches do not directly implicate S1P. Interactions between tumour and EC mediated by S1P have been shown to be important for angiogenesis in many solid cancers, including those of the breast, prostate, liver and kidney [53–55]. To investigate the involvement of S1P in DLBCL-associated angiogenesis, we defined a transcriptional signature of S1P signalling in HUVEC [23]. Gene ontology analysis revealed that genes up-regulated in this transcriptional signature were enriched for angiogenesis-associated GO terms, as well as for GO terms reflecting known functions of S1P [9, 12, 13, 56, 57]. We used this transcriptional signature to show that genes up-regulated by S1P were enriched among genes correlated with SPHK1, an association that was also evident separately for both COO subtypes. IHC revealed that S1P target genes were expressed in the tumour vasculature of SPHK1-expressing DLBCL, with little or no tumour cell expression, suggesting that the enrichment of S1P target genes observed in SPHK1-expressing DLBCL was primarily a consequence of their expression in tumour-associated EC.

Recent studies exploring the transcriptional landscape of DLBCL have defined tumour microenvironment-derived gene signatures predicting clinical outcome [2, 3]. The ‘Stromal-1’ signature, associated with a favourable outcome, is characterised by expression of extracellular matrix- and macrophage-associated genes. In contrast, the ‘Stromal-2’

signature has an unfavourable outcome, and is reportedly enriched for angiogenesis-associated genes [3]. We formally tested the relationship between angiogenesis and these stromal signatures [3]. As expected, we found a significant enrichment of angiogenesis-associated genes within the Stromal-2 signature. However, we were surprised to find an even stronger enrichment of angiogenesis genes within the Stromal-1 signature. Both stromal signatures were enriched for genes positively correlated with SPHK1 expression, suggesting that SPHK1 contributes to angiogenesis in both stromal subtypes.

Our data suggest novel therapeutic opportunities for patients with SPHK1-expressing DLBCL insofar as the S1PR1 functional antagonists, Siponimod, Ozanimod and Ponesimod, were all able to block S1P signalling in EC. Although we observed the same effect with the S1P-specific monoclonal antibody, Sphingomab, we suggest that S1PR1 is a preferred target; S1P-blocking agents will affect signalling through all S1P receptors. Potentially unwanted side-effects of blocking S1P could include the inhibition of S1PR2 signalling which we and others have reported to be tumour suppressive in DLBCL [58, 59] (Vockerodt et al, in press). Moreover, we showed that Siponimod, a highly potent functional antagonist of S1PR1 that is effective and safe in patients with multiple sclerosis [60], can reduce angiogenesis and tumour growth of SUDHL6 xenografts which produce S1P and which we used as a mouse model of angiogenic DLBCL. Targeting tumour cell over-expression of S1PR1, which is observed in a subset of poor prognosis DLBCL, is also potentially advantageous, since S1PR1 has been shown to be important for the proliferation and survival of DLBCL [61, 62]. S1P/S1PR1 signalling also promotes the accumulation of Treg while inhibiting CD8+ T cell recruitment and activation in xenograft models of breast cancer and melanoma [63]. Therefore, blocking S1PR1 might also promote anti-tumour immunity. Taken together our observations suggest that specific S1PR1 antagonists should be investigated for their therapeutic potential in DLBCL patients [40, 64, 65].

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This work was supported by Bloodwise (LL, TP, RH, MI, CJBW, KV, PGM) and the Traynor Foundation (LL), and in part by grants RVO:61989592 and NPS I LO1304 from the Czech Ministry of Education (PGM), NIH grant R01GM043880 (SS), and by Cancer Research UK Programme Grant C7845/A17723 (MC).

References

1. Coiffier B, Thieblemont C, Van Den Neste E, Lepage G, Plantier I, Castaigne S, et al. Long-term outcome of patients in the LNH-98.5 trial, the first randomized study comparing rituximab-CHOP to standard CHOP chemotherapy in DLBCL patients: a study by the Groupe d'Etudes des Lymphomes de l'Adulte. *Blood*. 2010; 116(12):2040–5. [PubMed: 20548096]
2. Sehn LH, Gascoyne RD. Diffuse large B-cell lymphoma: optimizing outcome in the context of clinical and biologic heterogeneity. *Blood*. 2015; 125(1):22–32. [PubMed: 25499448]
3. Lenz G, Wright G, Dave SS, Xiao W, Powell J, Zhao H, et al. Stromal Gene Signatures in Large-B-Cell Lymphomas. *New England Journal of Medicine*. 2008; 359(22):2313–23. [PubMed: 19038878]


4. Stopeck AT, Unger JM, Rimsza LM, Bellamy WT, Iannone M, Persky DO, et al. A phase II trial of single agent bevacizumab in patients with relapsed, aggressive non-Hodgkin lymphoma: Southwest oncology group study S0108. *Leuk Lymphoma*. 2009; 50(5):728–35. [PubMed: 19373598]
5. Ganjoo KN, An CS, Robertson MJ, Gordon LI, Sen JA, Weisenbach J, et al. Rituximab, bevacizumab and CHOP (RA-CHOP) in untreated diffuse large B-cell lymphoma: safety, biomarker and pharmacokinetic analysis. *Leuk Lymphoma*. 2006; 47(6):998–1005. [PubMed: 16840188]
6. Seymour JF, Pfreundschuh M, Trněný M, Sehn LH, Catalano J, Csinady E, et al. R-CHOP with or without bevacizumab in patients with previously untreated diffuse large B-cell lymphoma: final MAIN study outcomes. *Haematologica*. 2014; 99(8):1343–9. [PubMed: 24895339]
7. Gratzinger D, Zhao S, Tibshirani RJ, Hsi ED, Hans CP, Pohlman B, et al. Prognostic significance of VEGF, VEGF receptors, and microvessel density in diffuse large B cell lymphoma treated with anthracycline-based chemotherapy. *Lab Invest*. 2008; 88(1):38–47. [PubMed: 17998899]
8. Jorgensen JM, Sorensen FB, Bendix K, Nielsen JL, Olsen ML, Funder AM, et al. Angiogenesis in non-Hodgkin's lymphoma: clinico-pathological correlations and prognostic significance in specific subtypes. *Leuk Lymphoma*. 2007; 48(3):584–95. [PubMed: 17454603]
9. Takuwa Y, Du W, Qi X, Okamoto Y, Takuwa N, Yoshioka K. Roles of sphingosine-1-phosphate signaling in angiogenesis. *World J Biol Chem*. 2010; 1(10):298–306. [PubMed: 21537463]
10. Pyne NJ, Pyne S. Sphingosine 1-phosphate and cancer. *Nat Rev Cancer*. 2010; 10(7):489–503. [PubMed: 20555359]
11. Maceyka M, Harikumar KB, Milstien S, Spiegel S. Sphingosine-1-phosphate signaling and its role in disease. *Trends in Cell Biology*. 2012; 22(1):50–60. [PubMed: 22001186]
12. Lee OH, Kim YM, Lee YM, Moon EJ, Lee DJ, Kim JH, et al. Sphingosine 1-phosphate induces angiogenesis: its angiogenic action and signaling mechanism in human umbilical vein endothelial cells. *Biochem Biophys Res Commun*. 1999; 264(3):743–50. [PubMed: 10544002]
13. Wang F, Van Brocklyn JR, Hobson JP, Movafagh S, Zukowska-Grojec Z, Milstien S, et al. Sphingosine 1-phosphate stimulates cell migration through a G(i)-coupled cell surface receptor. Potential involvement in angiogenesis. *J Biol Chem*. 1999; 274(50):35343–50. [PubMed: 10585401]
14. English D, Welch Z, Kovala AT, Harvey K, Volpert OV, Brindley DN, et al. Sphingosine 1-phosphate released from platelets during clotting accounts for the potent endothelial cell chemotactic activity of blood serum and provides a novel link between hemostasis and angiogenesis. *FASEB J*. 2000; 14(14):2255–65. [PubMed: 11053247]
15. Liu F, Verin AD, Wang P, Day R, Wersto RP, Chrest FJ, et al. Differential regulation of sphingosine-1-phosphate- and VEGF-induced endothelial cell chemotaxis. Involvement of G(alpha2)-linked Rho kinase activity. *Am J Respir Cell Mol Biol*. 2001; 24(6):711–9. [PubMed: 11415936]
16. Spiegel S, Milstien S. Sphingosine-1-phosphate: an enigmatic signalling lipid. *Nat Rev Mol Cell Biol*. 2003; 4(5):397–407. [PubMed: 12728273]
17. Wallington-Beddoe CT, Bradstock KF, Bendall LJ. Oncogenic properties of sphingosine kinases in haematological malignancies. *Br J Haematol*. 2013; 161(5):623–38. [PubMed: 23521541]
18. Bayerl MG, Bruggeman RD, Conroy EJ, Hengst JA, King TS, Jimenez M, et al. Sphingosine kinase 1 protein and mRNA are overexpressed in non-Hodgkin lymphomas and are attractive targets for novel pharmacological interventions. *Leuk Lymphoma*. 2008; 49(5):948–54. [PubMed: 18452097]
19. Pan S, Gray NS, Gao W, Mi Y, Fan Y, Wang X, et al. Discovery of BAF312 (Siponimod), a Potent and Selective S1P Receptor Modulator. *ACS Med Chem Lett*. 2013; 4(3):333–7. [PubMed: 24900670]
20. Vockerodt M, Morgan SL, Kuo M, Wei W, Chukwuma MB, Arrand JR, et al. The Epstein-Barr virus oncoprotein, latent membrane protein-1, reprograms germinal centre B cells towards a Hodgkin's Reed-Sternberg-like phenotype. *J Pathol*. 2008; 216(1):83–92. [PubMed: 18566961]
21. Vrzalikova K, Vockerodt M, Leonard S, Bell A, Wei W, Schrader A, et al. Down-regulation of BLIMP1 α by the EBV oncogene, LMP-1, disrupts the plasma cell differentiation program and prevents viral replication in B cells: implications for the pathogenesis of EBV-associated B-cell lymphomas. *Blood*. 2011; 117(22):5907–17. [PubMed: 21411757]

22. Vrzalikova K, Leonard S, Fan Y, Bell A, Vockerodt M, Flodr P, et al. Hypomethylation and Over-Expression of the Beta Isoform of BLIMP1 is Induced by Epstein-Barr Virus Infection of B Cells; Potential Implications for the Pathogenesis of EBV-Associated Lymphomas. *Pathogens*. 2012; 1(2):83–101. [PubMed: 25436766]
23. Jaffe EA, Nachman RL, Becker CG, Minick CR. Culture of human endothelial cells derived from umbilical veins. Identification by morphologic and immunologic criteria. *J Clin Invest*. 1973; 52(11):2745–56. [PubMed: 4355998]
24. Cassetta L, Noy R, Swierczak A, Sugano G, Smith H, Wiechmann L, et al. Isolation of Mouse and Human Tumor-Associated Macrophages. *Adv Exp Med Biol*. 2016; 899:211–29. [PubMed: 27325269]
25. Hait NC, Allegood J, Maceyka M, Strub GM, Harikumar KB, Singh SK, et al. Regulation of histone acetylation in the nucleus by sphingosine-1-phosphate. *Science*. 2009; 325(5945):1254–7. [PubMed: 19729656]
26. Vrzalikova K, Ibrahim M, Vockerodt M, Perry T, Margielewska S, Lupino L, et al. S1PR1 drives a feedforward signalling loop to regulate BATF3 and the transcriptional programme of Hodgkin lymphoma cells. *Leukemia*. 2018; 32(1):214–23. [PubMed: 28878352]
27. Reynolds GM, Billingham LJ, Gray LJ, Flavell JR, Najafipour S, Crocker J, et al. Interleukin 6 expression by Hodgkin/Reed-Sternberg cells is associated with the presence of 'B' symptoms and failure to achieve complete remission in patients with advanced Hodgkin's disease. *Br J Haematol*. 2002; 118(1):195–201. [PubMed: 12100148]
28. Flavell JR, Baumforth KR, Wood VH, Davies GL, Wei W, Reynolds GM, et al. Down-regulation of the TGF-beta target gene, PTPRK, by the Epstein-Barr virus encoded EBNA1 contributes to the growth and survival of Hodgkin lymphoma cells. *Blood*. 2008; 111(1):292–301. [PubMed: 17720884]
29. Hans CP, Weisenburger DD, Greiner TC, Gascoyne RD, Delabie J, Ott G, et al. Confirmation of the molecular classification of diffuse large B-cell lymphoma by immunohistochemistry using a tissue microarray. *Blood*. 2004; 103(1):275–82. [PubMed: 14504078]
30. Liao Y, Smyth GK, Shi W. The Subread aligner: fast, accurate and scalable read mapping by seed-and-vote. *Nucleic Acids Res*. 2013; 41(10):e108. [PubMed: 23558742]
31. Robinson MD, McCarthy DJ, Smyth GK. edgeR: a Bioconductor package for differential expression analysis of digital gene expression data. *Bioinformatics*. 2010; 26(1):139–40. [PubMed: 19910308]
32. Morin RD, Johnson NA, Severson TM, Mungall AJ, An J, Goya R, et al. Somatic mutations altering EZH2 (Tyr641) in follicular and diffuse large B-cell lymphomas of germinal-center origin. *Nat Genet*. 2010; 42(2):181–5. [PubMed: 20081860]
33. Beguelin W, Popovic R, Teater M, Jiang Y, Bunting KL, Rosen M, et al. EZH2 is required for germinal center formation and somatic EZH2 mutations promote lymphoid transformation. *Cancer Cell*. 2013; 23(5):677–92. [PubMed: 23680150]
34. Brune V, Tiacchi E, Pfeil I, Döring C, Eckerle S, van Noesel CJM, et al. Origin and pathogenesis of nodular lymphocyte-predominant Hodgkin lymphoma as revealed by global gene expression analysis. *The Journal of Experimental Medicine*. 2008; 205(10):2251–68. [PubMed: 18794340]
35. Care MA, Westhead DR, Tooze RM. Gene expression meta-analysis reveals immune response convergence on the IFNgamma-STAT1-IRF1 axis and adaptive immune resistance mechanisms in lymphoma. *Genome Med*. 2015; 7(96):15–218. [PubMed: 25705260]
36. Pitson SM, Moretti PA, Zebol JR, Lynn HE, Xia P, Vadas MA, et al. Activation of sphingosine kinase 1 by ERK1/2-mediated phosphorylation. *EMBO J*. 2003; 22(20):5491–500. [PubMed: 14532121]
37. Dennis BTS, Douglas A, Hosack, Yang Jun, Michael W, Baseler H, Lane Clifford, Lempicki Richard A. DAVID: Database for Annotation, Visualization, and Integrated Discovery. *Genome Biology*. 2003; 4(5)
38. Masiero M, Simoes FC, Han HD, Snell C, Peterkin T, Bridges E, et al. A core human primary tumor angiogenesis signature identifies the endothelial orphan receptor ELTD1 as a key regulator of angiogenesis. *Cancer Cell*. 2013; 24(2):229–41. [PubMed: 23871637]

39. O'Brien N, Jones ST, Williams DG, Cunningham HB, Moreno K, Visentin B, et al. Production and characterization of monoclonal anti-sphingosine-1-phosphate antibodies. *Journal of Lipid Research*. 2009; 50(11):2245–57. [PubMed: 19509417]
40. Olsson T, Boster A, Fernandez O, Freedman MS, Pozzilli C, Bach D, et al. Oral ponesimod in relapsing-remitting multiple sclerosis: a randomised phase II trial. *J Neurol Neurosurg Psychiatry*. 2014; 85(11):1198–208. [PubMed: 24659797]
41. Scott FL, Clemons B, Brooks J, Brahmachary E, Powell R, Dedman H, et al. Ozanimod (RPC1063) is a potent sphingosine-1-phosphate receptor-1 (S1P1) and receptor-5 (S1P5) agonist with autoimmune disease-modifying activity. *Br J Pharmacol*. 2016; 173(11):1778–92. [PubMed: 26990079]
42. Hale JJ, Lynch CL, Neway W, Mills SG, Hajdu R, Keohane CA, et al. A rational utilization of high-throughput screening affords selective, orally bioavailable 1-benzyl-3-carboxyazetidide sphingosine-1-phosphate-1 receptor agonists. *J Med Chem*. 2004; 47(27):6662–5. [PubMed: 15615513]
43. Zhang L, Wang XE, Bullock AJ, Callea M, Shah H, Song JX, et al. Anti-S1P Antibody as a Novel Therapeutic Strategy for VEGFR TKI-Resistant Renal Cancer. *Clin Cancer Res*. 2015; 21(8):1925–34. [PubMed: 25589614]
44. Nagahashi M, Ramachandran S, Kim EY, Allegood JC, Rashid OM, Yamada A, et al. Sphingosine-1-phosphate produced by sphingosine kinase 1 promotes breast cancer progression by stimulating angiogenesis and lymphangiogenesis. *Cancer Res*. 2012; 72(3):726–35. [PubMed: 22298596]
45. Reddy A, Zhang J, Davis NS, Moffitt AB, Love CL, Waldrop A, et al. Genetic and Functional Drivers of Diffuse Large B Cell Lymphoma. *Cell*. 2017; 171(2):481–94 e15. [PubMed: 28985567]
46. Chapuy B, Stewart C, Dunford AJ, Kim J, Kamburov A, Redd RA, et al. Molecular subtypes of diffuse large B cell lymphoma are associated with distinct pathogenic mechanisms and outcomes. *Nat Med*. 2018; 24(5):679–90. [PubMed: 29713087]
47. Schmitz R, Wright GW, Huang DW, Johnson CA, Phelan JD, Wang JQ, et al. Genetics and Pathogenesis of Diffuse Large B-Cell Lymphoma. *N Engl J Med*. 2018; 378(15):1396–407. [PubMed: 29641966]
48. Vann LR, Payne SG, Edsall LC, Twitty S, Spiegel S, Milstien S. Involvement of sphingosine kinase in TNF-alpha-stimulated tetrahydrobiopterin biosynthesis in C6 glioma cells. *J Biol Chem*. 2002; 277(15):12649–56. [PubMed: 11815603]
49. Osawa Y, Banno Y, Nagaki M, Brenner DA, Naiki T, Nozawa Y, et al. TNF-alpha-induced sphingosine 1-phosphate inhibits apoptosis through a phosphatidylinositol 3-kinase/Akt pathway in human hepatocytes. *J Immunol*. 2001; 167(1):173–80. [PubMed: 11418646]
50. Olivera A, Edsall L, Poulton S, Kazlauskas A, Spiegel S. Platelet-derived growth factor-induced activation of sphingosine kinase requires phosphorylation of the PDGF receptor tyrosine residue responsible for binding of PLCgamma. *FASEB J*. 1999; 13(12):1593–600. [PubMed: 10463951]
51. Young KW, Bootman MD, Channing DR, Lipp P, Maycox PR, Meakin J, et al. Lysophosphatidic acid-induced Ca²⁺ mobilization requires intracellular sphingosine 1-phosphate production. Potential involvement of endogenous EDG-4 receptors. *J Biol Chem*. 2000; 275(49):38532–9. [PubMed: 10954727]
52. Meyer zu Heringdorf D, Lass H, Alemany R, Laser KT, Neumann E, Zhang C, et al. Sphingosine kinase-mediated Ca²⁺ signalling by G-protein-coupled receptors. *EMBO J*. 1998; 17(10):2830–7. [PubMed: 9582276]
53. Ader I, Gstalder C, Bouquerel P, Golzio M, Andrieu G, Zalvidea S, et al. Neutralizing S1P inhibits intratumoral hypoxia, induces vascular remodelling and sensitizes to chemotherapy in prostate cancer. *Oncotarget*. 2015; 6(15):13803–21. [PubMed: 25915662]
54. Lu ZP, Zhang WY, Gao S, Jiang QL, Xiao ZL, Ye LH, et al. MiR-506 suppresses liver cancer angiogenesis through targeting sphingosine kinase 1 (SPHK1) mRNA. *Biochem Bioph Res Co*. 2015; 468(1–2):8–13.
55. Mukhopadhyay P, Ramanathan R, Takabe K. S1P promotes breast cancer progression by angiogenesis and lymphangiogenesis. *Breast Cancer Manag*. 2015; 4(5):241–4. [PubMed: 27293484]

56. Hisano N, Yatomi Y, Satoh K, Akimoto S, Mitsumata M, Fujino MA, et al. Induction and suppression of endothelial cell apoptosis by sphingolipids: a possible in vitro model for cell-cell interactions between platelets and endothelial cells. *Blood*. 1999; 93(12):4293–9. [PubMed: 10361127]
57. Fernandez-Pisonero I, Duenas AI, Barreiro O, Montero O, Sanchez-Madrid F, Garcia-Rodriguez C. Lipopolysaccharide and sphingosine-1-phosphate cooperate to induce inflammatory molecules and leukocyte adhesion in endothelial cells. *J Immunol*. 2012; 189(11):5402–10. [PubMed: 23089395]
58. Flori M, Schmid CA, Sumrall ET, Tzankov A, Law CW, Robinson MD, et al. The hematopoietic oncoprotein FOXP1 promotes tumor cell survival in diffuse large B-cell lymphoma by repressing S1PR2 signaling. *Blood*. 2016; 127(11):1438–48. [PubMed: 26729899]
59. Muppidi JR, Schmitz R, Green JA, Xiao W, Larsen AB, Braun SE, et al. Loss of signalling via Galpha13 in germinal centre B-cell-derived lymphoma. *Nature*. 2014; 516(7530):254–8. [PubMed: 25274307]
60. Vockerodt M, Vrzalikova K, Ibrahim M, Nagy E, Margielewska S, Hollows R, Lupino L, Tooze R, Care M, Simmons W, Schrader A, Perry T, Abdullah M, Foster S, Reynolds G, Dowell A, Rudzki Z, Krappmann D, Kube D, Woodman C, Wei W, Taylor G, Murray PG, et al. Regulation of S1PR2 by the EBV oncogene LMP1 in aggressive ABC-subtype diffuse large B-cell lymphoma. *J Pathol*. 2019 Jan 21[Epub ahead of print]
61. Kappos L, Bar-Or A, Cree BAC, Fox RJ, Giovannoni G, Gold R, et al. Siponimod versus placebo in secondary progressive multiple sclerosis (EXPAND): a double-blind, randomised, phase 3 study. *Lancet*. 2018; 391(10127):1263–73. [PubMed: 29576505]
62. Koresawa R, Yamazaki K, Oka D, Fujiwara H, Nishimura H, Akiyama T, et al. Sphingosine-1-phosphate receptor 1 as a prognostic biomarker and therapeutic target for patients with primary testicular diffuse large B-cell lymphoma. *Br J Haematol*. 2016; 174(2):264–74. [PubMed: 27061580]
63. Liu Y, Deng J, Wang L, Lee H, Armstrong B, Scuto A, et al. S1PR1 is an effective target to block STAT3 signaling in activated B cell-like diffuse large B-cell lymphoma. *Blood*. 2012; 120(7):1458–65. [PubMed: 22745305]
64. Priceman SJ, Shen S, Wang L, Deng J, Yue C, Kujawski M, et al. S1PR1 is crucial for accumulation of regulatory T cells in tumors via STAT3. *Cell Rep*. 2014; 6(6):992–9. [PubMed: 24630990]
65. Kappos L, Li DK, Stuve O, Hartung HP, Freedman MS, Hemmer B, et al. Safety and Efficacy of Siponimod (BAF312) in Patients With Relapsing-Remitting Multiple Sclerosis: Dose-Blinded, Randomized Extension of the Phase 2 BOLD Study. *JAMA Neurol*. 2016; 73(9):1089–98. [PubMed: 27380540]
66. Tran JQ, Hartung JP, Peach RJ, Boehm MF, Rosen H, Smith H, et al. Results From the First-in-Human Study With Ozanimod, a Novel, Selective Sphingosine-1-Phosphate Receptor Modulator. *J Clin Pharmacol*. 2017; 57(8):988–96. [PubMed: 28398597]

Figure 1

**Figure 1. SPHK1 is over-expressed in primary DLBCL.**

(A) SPHK1 mRNA expression in the re-analysis of a published microarray dataset [34] and (B) SPHK1 mRNA expression in the re-analysis of published RNAseq data [32]. (C) IHC for SPHK1 protein expression in normal GC B cells and also in the tumour cells of representative cases of DLBCL. Arrows indicate tumour-associated EC which did not express SPHK1 (lower middle and lower right panels). GC=germinal centre, MZ=mantle zone. Original magnifications x200 and x600. D) Upper panel: qPCR analysis for the expression of SPHK1 mRNA in DLBCL cell lines. Data are in triplicate and each is

representative of three separate biological replicates. Lower panel: Immunoblotting of DLBCL cell lines for phosphorylated and total SPHK1 (pSPHK1, tSPHK1). β -tubulin is a loading control. E) Intracellular S1P levels in primary B cells and DLBCL cell lines. Data show means (\pm SEM) of four biological replicates. Statistics were based on a comparison of individual cell lines with B cells#2. *denotes $p<0.05$, ** $p<0.01$, *** $p<0.001$ (Student's t test). F) S1P levels secreted by SUDHL4 cells incubated in serum-free media for indicated times measured by mass spectrometry. Data show means (\pm SEM) of three biological replicates. *denotes $p<0.05$.




Figure 2. SPHK1 expression is associated with an angiogenic transcriptome in primary DLBCL. (A) Enrichment of angiogenesis-related terms (red arrows) in a GO analysis of genes positively correlated with SPHK1 expression in primary DLBCL. The top 10 significant GO categories are shown ordered by p value. (B) Tumour vascular signature genes [38] were significantly enriched among genes positively correlated with SPHK1 expression in primary DLBCL (left panel), but significantly depleted among those negatively correlated with SPHK1 expression in primary DLBCL (right panel) (Fisher's exact test for both comparisons). (C) Re-analysis of gene expression from primary DLBCL [32] reveals a

statistically significant positive correlation between SPHK1 expression and the expression of EC marker genes CD34, CDH5, PECAM1 and VWF (r =Spearman rank correlation coefficient).




Figure 3. Both Stromal-1 and Stromal-2 gene signatures are enriched for angiogenic genes. (A) Overlap between Stromal-2 genes and genes comprising the three different angiogenic gene signatures. (B) Overlap between Stromal-1 genes and the angiogenic gene signatures.




Figure 4. Identification of S1P target genes in endothelial cells.

(A) Representative images of IHC for S1PR1, S1PR2 and S1PR3 in primary DLBCL and HUVEC. DLBCL-associated EC (left and middle panels) were positive for S1PR1 and negative for S1PR2 and S1PR3 in most cases (black arrows). Red blood cells stained strongly for S1PR2 (red arrows) and served as an internal positive control. S1PR3-positive tumour cells (bottom left) and/or squamous epithelium (not shown) were internal positive controls for S1PR3 expression. HUVEC (right panel) were positive for S1PR1 and negative for S1PR2 and S1PR3. Original magnifications x200 (tissues) and x600 (cells). (B) S1P

treatment of HUVEC increased ERK1/2 phosphorylation, shown here for 4 separate donors compared with vehicle only treated cells (H1-H4). β -tubulin is a loading control. The same protein lysates were used to detect phosphorylated and total ERK1/2 (pERK1/2, tERK1/2) in separate blots. (C) Q-PCR for S1P transcriptional targets in HUVEC of four donors following treatment with S1P (dark grey bars) compared with vehicle (light grey bars) normalised to 1. *denotes $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$ (Student's t test). (D) Immunoblotting for pERK1/2, tERK1/2 and β -tubulin in HUVEC representative of three donors treated with S1P in the presence of Sphingomab (SmAb) or isotype/control antibody (CmAb). The same protein lysates were used to detect pERK1/2 and tERK1/2 in separate blots. (E) Q-PCR for S1P targets following treatment of HUVEC with S1P in the presence of either SmAb or CmAb. Data shown are representative of three donors. *denotes $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$ (Student's t test).




Figure 5. Genes upregulated by S1P and correlated with SPHK1 mRNA are expressed in DLBCL tumour vasculature.

(A) GO analysis of genes upregulated by S1P in HUVEC revealed a significant enrichment of GO terms associated with angiogenesis. (B) Genes upregulated (left panel), but not those downregulated (not shown) in S1P-treated HUVEC were enriched among genes positively correlated with SPHK1 expression in primary DLBCL. Genes downregulated (right panel), but not those upregulated (not shown) in S1P-treated HUVEC were enriched among genes negatively correlated with SPHK1 expression in primary DLBCL. (C) Representative

examples of the expression of the S1P target genes, CXCL12, SELE, COL1A1 and MAP1B, in the tumour vasculature of DLBCL. Original magnifications x200.




Figure 6. S1PR1 inhibition decreases S1P-induced signalling and reduces angiogenesis in a mouse model of DLBCL.

(A) Immunoblot of HUVEC pre-treated for 1h with 100nM of S1PR1 functional antagonists, Siponimod, Ozanimod and Ponesimod and stimulated with S1P or control (S1P vehicle) for 5min. S1P or control only treated cells received the equivalent dose of DMSO (drug solvent) to S1PR1 inhibitor treated cells (0.01µl DMSO per 1mL of media). β-tubulin was used as a loading control. The same protein lysates were used to detect pERK1/2 and tERK1/2 in separate blots. Data are representative of three separate donors. (B) Flow cytometry analysis of numbers of CD31-positive mouse cells/per mg tumour of SUDHL6 xenografts in NSG

mice treated with Siponimod compared with drug vehicle (4 mice for each group). (C) Tumour volumes in days after injection with SUDHL6 cells in Siponimod-treated versus vehicle-treated animals. Arrows indicate treatment days. *denotes $p < 0.05$ in Student's t test.