



Correction to: Short-term progression of cardiometabolic risk factors in relation to age at type 2 diabetes diagnosis: a longitudinal observational study of 100,606 individuals from the Swedish National Diabetes Register

Andri O. Steinarsson¹ · Araz Rawshani² · Soffia Gudbjörnsdóttir^{2,3} · Stefan Franzén³ · Ann-Marie Svensson³ · Naveed Sattar⁴

Published online: 2 September 2019
© Springer-Verlag GmbH Germany, part of Springer Nature 2019

Correction to: Diabetologia

<https://doi.org/10.1007/s00125-017-4532-8>

Unfortunately, the symbols in Fig. 1 were incorrectly described in the legend. The corrected figure is given here.

The online version of the original article can be found at <https://doi.org/10.1007/s00125-017-4532-8>

- ✉ Soffia Gudbjörnsdóttir
soffia.gudbjornsdottir@medic.gu
- ✉ Naveed Sattar
naveed.sattar@glasgow.ac.uk

¹ School of Health Science, Faculty of Medicine, University of Iceland, Reykjavík, Iceland

² Department of Molecular and Clinical Medicine, Institute of Medicine, University of Gothenburg, Gothenburg, Sweden

³ The Swedish National Diabetes Register, Västra Götalandsregionen, Gothenburg, Sweden

⁴ BHF Glasgow Cardiovascular Research Centre, Institute of Cardiovascular and Medical Sciences, University of Glasgow, 126 University Place, Glasgow G12 8TA, UK

Fig. 1 Yearly averages (95% CI) stratified by age group for (a) BMI, (b) HbA_{1c}, (c) triacylglycerol, (d) HDL-cholesterol and (e) LDL-cholesterol. Circles, 18–44 years old; squares, 45–59 years old; diamonds, 60–74 years old; triangles, ≥75 years old. All analyses, $p < 0.05$ where 95% CI do not overlap

