CORRECTION



Correction to: c-Jun Amino-Terminal Kinase is Involved in Valproic Acid-Mediated Neuronal Differentiation of Mouse Embryonic NSCs and Neurite Outgrowth of NSC-Derived Neurons

Lu Lu¹ · Hengxing Zhou¹ · Bin Pan¹ · Xueying Li² · Zheng Fu² · Jun Liu¹ · Zhongju Shi¹ · Tianci Chu³ · Zhijian Wei¹ · Guangzhi Ning¹ · Shiqing Feng¹

Published online: 20 August 2019
© Springer Science+Business Media, LLC, part of Springer Nature 2019

Correction to:

Neurochemical Research (2017) 42(4):1254–1266 https://doi.org/10.1007/s11064-016-2167-7

In the original version of this article, unfortunately, the images in Fig. 4 and 7 are mixed. The correct version of the Figs. 4 and 7 is given below.

The original article can be found online at https://doi.org/10.1007/s11064-016-2167-7.


- ² Key Laboratory of Immuno Microenvironment and Disease of the Educational Ministry of China, Department of Immunology, Tianjin Medical University, No. 22 Qixiangtai Road, Heping District, Tianjin 300070, People's Republic of China
- ³ Kosair Children's Hospital Research Institute at the Department of Pediatrics, University of Louisville School of Medicine, Louisville, KY 40202, USA



Shiqing Feng fshiqing 1967@hotmail.com

Department of Orthopaedics, Tianjin Medical University General Hospital, No. 154 Anshan Road, Heping District, Tianjin 300052, People's Republic of China

Fig. 4 Mouse embryonic NSCs were treated with 1 mM VPA for up to 7 days and immunostained with anti- β -tubulin-III, anti-MAP-2, anti-CNPase, and anti-GFAP. *Scale bar* 100 μ m








Fig. 7 Mouse embryonic NSCs were treated with 1 mM VPA for up to 7 days in the absence or presence of 10 μM SP600125 and immunostained with anti-β-tubulin-III, anti-MAP-2, anti-CNPase, and anti-GFAP. *Scale bar* 100 μm

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

