



Published in final edited form as:

Clin Cancer Res. 2019 June 01; 25(11): 3229–3238. doi:10.1158/1078-0432.CCR-18-2675.

A Phase 2 Study of Alisertib in Children with Recurrent/ Refractory Solid Tumors or Leukemia: Children's Oncology Group Phase 1 and Pilot Consortium (ADV L0921)

Yael P. Mossé, M.D.¹, Elizabeth Fox, M.D.¹, David T. Teachey, M.D.¹, Joel M. Reid, PhD²,
Stephanie L. Safgren, PhD², Hernan Carol³, Richard B Lock³, Peter J Houghton⁴, Malcolm
A Smith⁵, David Hall, MS⁶, Donald A Barkauskas, PhD⁶, Mark Krailo, PhD⁷, Stephan D.
Voss, MD, PhD⁸, Stacey L. Berg, M.D.⁹, Susan M. Blaney, M.D.⁹, Brenda J. Weigel, M.D.¹⁰

¹Division of Oncology and Center for Childhood Cancer Research, Children's Hospital of Philadelphia; Department of Pediatrics, Perelman School of Medicine at the University of Pennsylvania, Philadelphia, PA, USA ²Mayo Clinic, Rochester, MN, USA ³Children's Cancer Institute, Lowy Cancer Research Centre, UNSW Sydney, Sydney, Australia ⁴Greehey Children's Cancer Research Institute, University of Texas Health Science Center San Antonio, San Antonio, Texas ⁵Cancer Therapy Evaluation Program, NCI, Bethesda, Maryland ⁶Children's Oncology Group, Monrovia, CA, USA ⁷Department of Preventive Medicine, University of Southern California, Los Angeles, CA ⁸Dana-Farber/Boston Children's Cancer and Blood Disorders Center, Boston, MA, USA ⁹Texas Children's Cancer Center, Baylor College of Medicine, Houston, TX, USA ¹⁰Division of Pediatric Oncology, University of Minnesota Cancer Center, Minneapolis, MN, USA

Abstract

Background—Aurora A kinase (AAK) plays an integral role in mitotic entry, DNA damage checkpoint recovery, and centrosome and spindle maturation. Alisertib (MLN8237) is a potent and selective AAK inhibitor. In pediatric preclinical models, antitumor activity was observed in neuroblastoma, acute lymphoblastic leukemia, and sarcoma xenografts. We conducted a phase 2 trial of alisertib in pediatric patients with refractory or recurrent solid tumors or acute leukemias ().

Procedures—Alisertib (80 mg/m²/dose) was administered orally, daily for 7 days every 21 days. Pharmacogenomic (PG) evaluation for polymorphisms in the AURK gene and drug metabolizing enzymes (UGT1A1*28), and plasma pharmacokinetic studies (PK) were performed. Using a two-

Corresponding Author: Yael P. Mossé, MD, Children's Hospital of Philadelphia, Colket Translational Research Building, 3501 Civic Center Blvd- Office 3056, Philadelphia, PA 19104-4318, Tel: 215-590-0965, Fax: 267-426-0685, mosse@chop.edu.

Author Contributions:

Conception and design (YPM, DTT, JMR, MK, SMB, BJW, MAS, RBL)

Acquisition of data (YPM, EF, DTT, JMR, MK, SDV, SB, SMB, BJW, HC)

Analysis & interpretation data (YPM, EF, DTT, JMR, SLS, DH, DAB, MK, SDV, SB, SMB, BJW, MAS, RBL, HC)

Writing, review and/or revision of the manuscript (YPM, EF, DTT, JMR, SLS, DH, DAB, MK, SDV, SB, SMB, BJW, MAS, RBL, PJH, HC)

Study supervision (YPM, EF, JMR, MK, SDV, SB, SMB, BJW)

Conflict of Interest Disclosure Statement: The authors declare no potential conflict of interest.

stage design, patients were enrolled to 12 disease strata (10 solid tumor and 2 acute leukemia). Response was assessed after cycle 1, then every other cycle.

Results—A total of 139 children and adolescents (median age 10 years) were enrolled, 137 were evaluable for response. Five objective responses were observed (2 complete responses and 3 partial responses). The most frequent toxicity was myelosuppression. The median alisertib trough concentration on day 4 was 1.3 μM , exceeding the 1 μM target trough concentration in 67% of patients. No correlations between PG or PK and toxicity were observed.

Conclusion—Despite alisertib activity in pediatric xenograft models and cogent pharmacokinetic-pharmacodynamic relationships in preclinical models and adults, the objective response rate in children and adolescents receiving single agent alisertib was less than 5%.

Keywords

Aurora A Kinase; Clinical Trial; Childhood Cancer; Pharmacokinetics; Pharmacogenomics

Introduction

The Aurora kinase family is essential in the regulation of chromosome segregation and cytokinesis during mitosis¹. Aurora A kinase (AAK) plays an integral role in mitotic entry, DNA damage checkpoint recovery, and centrosome and spindle maturation¹. Aurora A is overexpressed in many adult tumors including bladder, breast, lung and head and neck cancers; ¹⁻⁴ as well as pediatric malignancies.⁵⁻⁸ The Aurora A kinase gene (AURK) has 2 two common polymorphisms; the phe31Ile polymorphism, which alters the kinase function and is associated with tumorigenesis or advanced cancers;^{9,10} and the Va571Ile polymorphism that in combination with phe31Ile may be associated with an increased risk of cancer or treatment related adverse events.¹¹⁻¹⁴

Alisertib (MLN8237) is a potent and selective AAK inhibitor previously investigated alone and in combination with chemotherapy as a potential treatment for patients with relapsed/refractory peripheral T-cell lymphoma as well as advanced solid tumors¹⁵⁻¹⁸. In preclinical models, antitumor activity and maximum pharmacodynamic effect were associated with alisertib concentrations exceeding 1 μM .^{19,20} In adults, the recommended dose of alisertib is 50 mg twice daily for 7 days with associated dose limiting toxicities (DLT) of neutropenia and stomatitis.²¹ The maximum concentration (C_{max}) and area under concentration time curve (AUC) following administration of 50 mg enteric coated tablets were 2.9 μM and 20.9 $\mu\text{M}\cdot\text{h}$, respectively; and the steady state trough concentration (C_{min}) exceeded 1 μM .²² Pharmacokinetic parameters derived from a population pharmacokinetic (PK) model based on data from 363 adults enrolled on 7 alisertib single agent trials showed a terminal half-life of 19.3 h and an apparent clearance (CL/F) of 4.25 L/h²³. Although the major metabolic pathway of alisertib is glucuronidation via the UDP-glucuronosyltransferase, UGT1A1, alisertib CL/F was not altered in adult subjects with UGT1A1*28 polymorphisms. Single agent phase 2 trials of alisertib demonstrated modest activity in adults with ovarian cancer²⁴, acute myeloid leukemia (AML),²⁵ and T- or B- cell lymphoma.²⁶

In the Pediatric Preclinical Testing Program (PPTP) alisertib was active in neuroblastoma (NBL) and acute lymphoblastic leukemia (ALL) xenografts with maintained complete responses observed in 3 of 7 NBL xenografts and 6 out of 6 ALL xenografts; sustained concentrations $1\mu\text{M}$ were associated with response.²⁷ Statistically significant improvement in event free survival in Wilms tumor (WT), rhabdomyosarcoma (RMS), and osteosarcoma (OS) xenograft models was also observed.²⁸ In these preclinical studies, alisertib was administered 5 days per week for 3 and 6 consecutive weeks for ALL and solid tumor models, respectively. In addition, alisertib was active in p53-wildtype, therapy-refractory NBL cell lines,²⁹ as a result of disruption of the Aurora-A/N-Myc complex resulting in inhibition of N-Myc dependent transcription.³⁰ Alisertib has also been shown to induce cell death and augment radiation sensitivity in atypical teratoid rhabdoid (ATRT) cell lines that overexpress AAK and have mutations in SMARCB1 (SNF5/INI1), a tumor suppressor and component of chromatin remodeling.⁷

Based on the preclinical anti-tumor activity in pediatric cell lines and xenograft models and the clinical anti-tumor activity in adult trials, a Children's Oncology Group phase 1 trial of alisertib in children and adolescents with relapsed or refractory solid tumors evaluated both once daily and twice daily schedules³¹. The recommended phase 2 dose and schedule was 80 mg/m^2 orally, once daily for 7 days. DLTs included myelosuppression, mood alterations, somnolence, mucositis, fatigue, alopecia, elevated hepatic transaminases, agitation and euphoria. In contrast to adults, the twice daily schedule in children resulted in a higher frequency of neutropenia and palmar-plantar erythrodysesthesia. There was marked inter-patient variability in the alisertib PK parameters in children. At the recommended dose, the C_{max} and AUC were $7.5 \pm 0.1\ \mu\text{M}$ and $75 \pm 13.5\ \mu\text{M}\cdot\text{h}$, respectively. The alisertib trough 24 hours after the first dose at the 80 mg/m^2 dose level was $1.1\ \mu\text{M}$. Of 33 response evaluable children in the phase 1 trial, one with hepatoblastoma (HBL) had a PR and 8 [NBL (n=4) and sarcoma (n=4)] had stable disease for 5–35 cycles.³¹

Based on the mechanism of action and preclinical activity of alisertib, a phase 2 trial was conducted to evaluate the objective response rate of alisertib in children and adolescents with relapsed/refractory solid tumors or acute leukemia. In parallel, and subsequent to the establishment of a dose and schedule for alisertib in a pediatric population, we selected two ALL xenografts against which we have previously reported single-agent alisertib efficacy at a dose resulting in drug plasma levels that are achievable in humans for additional testing to compare dosing schedules.

Methods

Patient Population

Patients with relapsed or refractory cancer were enrolled in one of 12 strata based on histology, including two strata for neuroblastoma (NBL); one for those with measurable disease by computed tomography (CT) or magnetic resonance imaging (MRI) and another for those with disease evaluable by MIBG scintigraphy, but no measurable disease. Patients with NBL limited to the bone marrow were not eligible. Patients with other solid tumors were required to have measurable disease as defined by Response Evaluation Criteria in Solid Tumors (RECIST 1.1).³² Strata for patients with sarcoma included

rhabdomyosarcoma (RMS), non-RMS soft tissue sarcoma (NRSTS), osteosarcoma (OS), or Ewing sarcoma/ peripheral PNET (EWS). Additional solid tumor strata included enrollment of patients with Wilms tumor (WT), hepatoblastoma (HBL), malignant germ cell tumors (GCT), and rhabdoid tumors (central nervous system atypical teratoid rhabdoid or other malignant rhabdoid tumors) with loss of INI1 by immunohistochemistry or molecular analysis. Patients with hematological malignancies without CNS involvement who were refractory or recurrent after at least 2 prior induction chemotherapy regimens, including those with acute myeloid leukemia (AML) and at least 5% myeloblasts in the bone marrow or those with acute lymphoblastic leukemia (ALL) and greater than 25% blasts (M3) in bone marrow, were also eligible. Subjects enrolled on the COG phase 1 trial (ADVL0812) who received alisertib at the recommended phase 2 dose and who met criteria for inclusion in one of the 12 disease stratum defined in this trial were included in this study population by prospective design.³¹

Patients were required to swallow alisertib tablets intact. Other inclusion criteria included age > 12 months and < 22 years; performance status of ≥ 50 by the Karnofsky scale for patients > 16 years or by the Lansky scale if ≤ 16 years; adequate renal function (normal serum creatinine for age and gender); and hepatic function (total bilirubin ≤ 1.5 -fold greater than the upper limit of normal, alanine aminotransferase (ALT) less than 225 U/L, and serum albumin of at least 2 g/dL) was required. In patients with solid tumors, bone marrow function for patients without known tumor infiltration of bone marrow included an absolute neutrophil count (ANC) $\geq 1000/\mu\text{L}$, platelet count $\geq 100,000/\mu\text{L}$, and hemoglobin ≥ 8 gm/dL; for patients with solid tumors and known bone marrow metastatic disease an ANC $\geq 750/\mu\text{L}$, platelet count $\geq 50,000/\mu\text{L}$, and hemoglobin ≥ 8 gm/dL were required. Patients with leukemia could enroll if they were not refractory to red blood cell or platelet transfusions.

Patients were required to have recovered from the acute toxic effects of all prior treatment. Requirements for the interval of time from prior therapy were standard.³¹ Exclusion criteria included uncontrolled infection; pregnancy; lactation; concurrent administration of selected P-glycoprotein substrates (digoxin, cyclosporine, tacrolimus or sirolimus); or use of daily benzodiazepines, because of the potential benzodiazepine-like effects of alisertib.

This study was conducted in compliance with the Declaration of Helsinki, the International Conference on Harmonization, Guideline for Good Clinical Practice, and applicable national and local regulatory requirements. Institutional Review Boards at participating institutions approved the study. Informed consent was obtained from patients, ages 18 years or older, or from parents/legal guardians of children aged less than 18 years, with child assent when appropriate, according to institutional policies.

Treatment Program

Alisertib (Millennium Pharmaceuticals, Inc. Cambridge, MA) was distributed by the NCI Cancer Therapy Evaluation Program as 10 mg enteric-coated tablets. Alisertib (80 mg/m^2) was administered orally once daily for seven consecutive days. Cycle duration was 21 days. The dose was reduced to 60 mg/m^2 for reversible toxicity as outlined in the protocol. The maximum daily dose of alisertib was 160 mg. Adherence was monitored using daily dosing diaries completed by the patient or parent/guardian. In the absence of progressive disease or

unacceptable toxicity, the maximum total duration of protocol therapy was 35 cycles, approximately 2 years.

Toxicity Monitoring and Dose Modifications

The Common Terminology and Adverse Events (CTCAE v 4.0) criteria were used to grade toxicity. Prior to each cycle, physical examination, complete blood count (CBC), serum electrolytes, creatinine, and liver function tests were performed. During cycle 1, physical examinations and serum chemistries were performed weekly; CBCs were performed twice weekly. In subsequent cycles, CBCs were performed weekly or more frequently if hematological toxicity occurred.

Response

Disease evaluations were performed at baseline, the end of cycle 1 and after completion of every other cycle of protocol therapy. In patients with non-CNS solid tumors, response was assessed using RECIST version 1.1 criteria.³² Response in NBL subjects with non-measurable but MIBG evaluable disease was assessed using the Curie Score criteria.³³ Response for subjects with AML was assessed using the International Working Group Criteria³⁴ and for those with ALL response was defined by morphology. Response in patients with central nervous system ATRT was assessed based on the sum of the products of the longest diameter × perpendicular diameter. All objective responses were confirmed by central review.

Any eligible patient who received at least one dose of alisertib was considered evaluable for response provided: (1) the patient was observed on protocol therapy for at least one cycle and the tumor was not removed surgically prior to the time an objective response was confirmed; or (2) the patient demonstrated a complete or partial response as confirmed by central review; or (3) the patient demonstrated progressive disease or died while on protocol therapy. All other evaluable patients with solid tumors were considered to be non-responders. The maximum evaluation period for determination of the overall best response was six treatment cycles for ADVL0921; the evaluation period for determination of overall best response for ADVL0812 was the time from enrollment to termination of protocol therapy.

Pharmacokinetics and Pharmacogenomics

To characterize the PKs of alisertib in children and adolescents, blood samples (3 mL, EDTA) were required in all participants during cycle 1 prior to alisertib administration on days 1, 4±1, and 7±1. If consent was provided, optional sampling was performed at 1–2, 3–4, and 6–8 hours after the first dose. Plasma was stored at –80°C until analysis. Alisertib concentrations were measured as previously described and PK parameters were calculated using non-compartmental analyses.^{31,35}

Consenting patients provided whole blood in EDTA tubes prior to day 7 of the first cycle for genotyping of patients for germline polymorphisms in UGT1A1 or aurora AAK gene (AURK, Phe31Ile and Val57Ile). DNA was extracted by QIAamp DNA Blood Mini Kit as per manufacturer's instructions. Methods were validated with a panel of 60 Caucasian DNA

samples from the Coriell Institute. Positive and negative controls were included for each analysis. For *UGT1A1* *28 (rs8175347), the number of TA repeats in the promoter region were detected and quantified by a modification of the method described by Akaba et al.³⁶ *UGT1A1* polymorphisms rs4124874 and rs10929302 were evaluated with PCR amplification and dye-terminator sequencing. Specific primers were designed and validated to amplify the region for both SNPs. Forward and reverse primers are AGTTCTCTTCACCTCCTCCT and AATAAA CCCACCTCACCAC, respectively. For *AURKA*, genotyping for the G>A polymorphism (rs1047972 in codon 57) and T>A polymorphism (rs2273535 in codon 31) was performed by amplification and detected on a Bio-Rad CFX384 Real-Time PCR detection system (Hercules CA). The real time PCR methods were validated against a standard PCR reaction with sequence detection of the polymorphisms. Primer and probe sequences were provided by Millennium Pharmaceuticals, Inc. (Cambridge, MA). The forward and reverse primer sequences for rs2273535 were CTGGCCACTATTTACAGGTAATGGA and TGGAGGTCCAAAACGTGTTCTC, respectively with probe/reporter 1 (VIC-labeled) sequence ACTCAGCAATTTCCCTT and probe/reporter 2 (FAM-labeled) sequence CTCAGCAAATTCCTT. The forward and reverse primer sequences for rs1047972 were CGGCTTGTGACTGGAGACA and GGGTCTTGTGTCCTTCAAATTCTTC, respectively with probe/reporter 1 (VIC-labeled) sequence CAGCGCGTTCCTT and probe/reporter 2 (FAM-labeled) sequence CAGCGCATTCCTT. The *AURKA* haplotypes were determined using the Phe31Ile and val57Ile SNPs as described by Ishikawa et al.¹⁴

Statistical Analysis Plan

A two-stage design was used to evaluate alisertib anti-tumor activity in seven primary strata: NBL with RECIST measurable disease, NBL with MIBG only evaluable disease, OS, EWS, RMS, ALL, and AML. For each of the two NBL strata, 14 patients were enrolled at the first stage. If no patients experienced a complete or partial response, alisertib was considered inactive in that stratum, and further enrollment to that stratum was terminated. If one or more patients achieved an objective response, 10 additional patients were enrolled to the stratum. Alisertib was active if 4 of 24 patients in an expanded stratum experienced an objective response. With this design, alisertib was identified as inactive with probability 0.96 if the true response rate was 5% and as active with probability 0.91 if the true response rate was 25%. For the other five secondary strata (NRSTS, HBL, GCT, WT and rhabdoid tumors), at the first stage for each stratum, 10 patients were enrolled. If no patients experienced an objective response, alisertib was considered inactive in that stratum and enrollment was terminated. If one or more patients experienced an objective response, 10 additional patients were enrolled to that stratum. Alisertib was considered active if 3 of 20 patients in an expanded stratum experienced an objective response. With this design, alisertib would be identified as inactive with probability 0.93 if the true response rate was 5%, and was identified as active with probability 0.88 if the true response rate was 25%. Because of the rarity of tumors in the secondary strata, enrollment to the study was designed to be closed, irrespective of enrollment numbers, when the evaluation of the seven primary strata was completed. If sufficient enrollment was obtained, the two-stage design used for the non-NBL stratum was applied to the secondary stratum.

Xenograft studies

Subsequent to the prior preclinical evaluation in which alisertib was administered on a twice-daily schedule for 5 days and repeated for 3-weeks,²⁷ the maximum tolerated dose in pediatric patients was determined to be once daily for 7 days.³¹ An experiment was designed to compare the preclinical and clinical schedules. ALL xenografts were generated as described previously.^{28,37} ALL-8 and ALL-19 xenograft cells, generated from 2 patients with relapsed ALL, were inoculated into NOD/SCID mice and engraftment monitored by weekly flow cytometric enumeration of the percentage of human CD45 (%huCD45) cells in murine peripheral blood (PB). When the %huCD45 cells reached a median of >1% for the entire cohort, mice were randomized to receive treatment with alisertib or vehicle control. Alisertib was administered using two alternative schedules: Schedule A) twice daily for 7 days; or Schedule B) twice daily for 5 days repeated for 3 weeks. In both cases the dose used was 10.4 mg/kg, administered by oral gavage as a suspension in 10% cyclodextrin. Groups of 4–6 mice were euthanized at Days 0, 7 and 21 post treatment initiation and at the end of the evaluation period (Day 42) to assess leukemic infiltration of PB, bone marrow and spleens. An additional experimental endpoint for each mouse was when the %huCD45 cells in PB reached 25% (deemed an event). Mice were euthanized if morbid or if they experienced weight loss ≥20%. Event free survival (EFS), Treated-Control (T-C), T/C and overall response measure (ORM) estimations were carried out according to established methodology.³⁷ Individual mice were assigned an ORM depending on the leukemic growth characteristics observed in the 42 days following treatment according to the established criteria used for evaluating single agents, and the median ORM was used to obtain the group score.

Results

Patients

Characteristics for all patients are presented in Table I. All patients (n=139) were eligible. Two patients, one with RECIST-measurable NBL and one with AML, were not evaluable for response due to rapid progression of disease prior to the start of protocol therapy. The median number of treatment cycles for 137 response-evaluable patients was 2 (range 1–35). A total of 500 cycles of alisertib were delivered. Three patients completed 35 cycles (24 months) of protocol therapy.

Toxicity

During cycle 1, 18 patients (13%) experienced dose limiting toxicity including myelosuppression, mucositis, febrile neutropenia, enterocolitis, diarrhea, depression, hypersomnia, photophobia, tumor lysis syndrome, hyperbilirubinemia and/or electrolyte abnormalities. The frequency of alisertib related grade 3 and 4 toxicities is shown in Table 2. Alisertib-related grade 3 or 4 toxicities that occurred in ≥10% of delivered cycles (n=500) were anemia (13.6%), lymphopenia (12.2%), neutropenia (51.8%), thrombocytopenia (20.8%) and leukopenia (33%). During cycle 1, two patients had fatal adverse events possibly related to alisertib: a patient with pelvic soft tissue sarcoma experienced grade 5 respiratory failure and a patient with hepatoblastoma experienced a fatal hepatic hemorrhage.

Response

Five objective responses were observed. Two patients had complete responses, one patient with MIBG-only evaluable NBL (Figure 1) and one patient with Wilms tumor. Three patients had partial responses, one each with RECIST-measurable NBL, MIBG evaluable NBL and HBL. The patient with HBL and partial response was previously reported in the phase 1 trial.³¹ Unlike the phase 1 study (ADVL0812), prolonged stable disease was not considered as a response in this trial and was not centrally reviewed. No responses were achieved in the other primary disease strata (OS, EWS, RMS, ALL, AML). The objective responses are summarized in Table 3 and includes the number of cycles of alisertib administered for patients with a best response of stable disease. Accrual to secondary strata (HBL, WT, GCT, NRSTS) was discontinued due to insufficient response in the primary strata. At the time the study was closed, accrual to the first stage was not completed for HBL, GCT or rhabdoid tumors.

Pharmacokinetics and Pharmacogenomics

Forty-five patients provided consent for optional PK samples on ADVL0912 and 2 patients from ADVL0812 had trough levels obtained on day 4. The alisertib PK parameters were highly variable (Supplemental Table 1). There was no correlation between age or gender and alisertib C_{min} or AUC. The alisertib C_{max} exceeded $1\mu\text{M}$ in 98% (44/45) of patients participating in detailed PK studies on day 1. The median C_{min} on day 4 was $1.6\mu\text{M}$, exceeding $1\mu\text{M}$ target trough concentration in 67% (26/39) of patients. The median C_{min} on day 7 was $0.9\mu\text{M}$, exceeding $1\mu\text{M}$ in 41% of patients (11/27).

A total of 87 patients underwent genotyping for AURKA and UGT1A1 (Table 4A). There was no relationship between cycle 1 toxicity and either the Phe31Ile or Val57Ile single nucleotide polymorphisms (SNP) in the AURKA genotype or in the AURKA haplotypes (Table 4B). There was no relationship between grade 2 toxicities and AURKA genotype for the Ile31Phe SNP or the AURKA haplotypes. However, patients that were heterozygous (WV) for the Val57Ile SNP appeared to have fewer grade 2 toxicities (Table 4B). Given the small patient cohort, it is not possible to determine if this is clinically significant. In addition, there were no relationships between treatment response and AURKA genotype for either SNP.

Paired data for UGT1A1 phenotype and day 4 C_{min} was obtained in 32 patients. There did not appear to be a difference in Day 4 C_{min} between the intermediate (IM) and poor metabolizer (PM) phenotypes, therefore, the data for these patients were pooled (Table 4C). The mean \pm SE alisertib trough concentration for the extensive metabolizer group (EM, $n=16$) was $1.01 \pm 0.23\mu\text{M}$ (95% C.I. of $0.53 - 1.49\mu\text{M}$) and for IM/PM ($n=16$) was $2.06 \pm 0.30\mu\text{M}$ (95% C.I. of $1.42 - 2.70\mu\text{M}$). The difference in the population means, $1.05\mu\text{M}$ (95% C.I. of $0.28 - 1.82\mu\text{M}$), was statistically significant (p-value of pooled t-test, 0.0091). The mean alisertib trough concentration for patients with and without grade 2 toxicities were $1.50 \pm 0.25\mu\text{M}$ (95% C.I. of $0.96 - 2.04\mu\text{M}$) and $1.58 \pm 0.35\mu\text{M}$ (95% C.I. of $0.83 - 2.32\mu\text{M}$), respectively. The difference in the population means, $0.08\mu\text{M}$ (95% C.I. of $-0.79 - 0.94\mu\text{M}$), was not statistically significant. While alisertib trough concentrations were statistically significantly higher for IM/PM patients, there did not appear to be a relationship

with the occurrence of grade 2 toxicity (p-value = 0.86). Furthermore, we found no significant interaction between EM (yes/no) and grade 2 adverse events (yes/no) (p-value = 1.0) (Table 4C).

Xenograft studies

For the T-lineage ALL-8, based on serial peripheral blood parameters, leukemia progression was significantly delayed compared to vehicle control for both of the treatment schedules, resulting in increased EFS (Figure 2 and Table 5). Leukemia progression was delayed by an additional 12.1 days in mice treated with Schedule B compared with Schedule A ($p=0.004$), with T/C values 3.6 and 2.2, respectively. However, neither of the treatment schedules induced an objective response. Data are summarized in Table 5 and Supplementary Table 2. Engraftment levels for ALL-8 engrafted mice detected in the three compartments analyzed at autopsy (blood, bone marrow and spleen) are shown in Supplementary Figure 1, Figure 2 and Figure 3, respectively. Treatment with both alisertib schedules limited leukemia progression to a similar extent by Day 7 in all organs analyzed. These effects were not complete, with 5–10 % human cells in the spleen and approximately 20% human cells in the bone marrow as the lowest levels achieved. Schedule B was more effective than Schedule A in reducing leukemia levels measured at Day 21 in the three compartments analyzed, but the differences between the treatments were transient and after drug treatments ceased there was a rapid progression of the disease (Day 42).

For the B-lineage ALL-19, based on serial peripheral blood parameters, leukemia progression was not significantly different from that of controls for Schedule A (Figure 3 and Table 5), However, Schedule B significantly delayed ALL-19 progression by 24.6 days relative to controls, which was 17.6 days greater than Schedule A ($p=0.048$) with T/C values 4.7 and 2.0, respectively. Furthermore, while treatment on Schedule A resulted in progressive disease (PD), Schedule B induced a complete response (CR). Data are summarized in Table 5 and Supplementary Table S2. Engraftment levels for ALL-19 engrafted mice detected in the three compartments analyzed at autopsy (blood, bone marrow and spleen) are shown in Supplementary Figure 1, Figure 2 and Figure 3, respectively. They followed a similar pattern to those of ALL-8, with the exception of being higher at Day 7, particularly in the bone marrow.

Discussion

In this phase 2 study, anti-tumor activity of alisertib was evaluated in 137 children and adolescents in 7 primary and 5 secondary disease strata. We demonstrated that children achieved target concentrations established in adults and preclinical models. The higher alisertib trough concentration in patients with intermediate and poor metabolizer UGT1A1 phenotypes compared to the extensive metabolizer phenotype was statistically significant. However, we did not find a difference in frequency of toxicity among these groups. Evaluation of AURK somatic mutation status and AAK expression in archival tumor specimens from children enrolled on this study was not performed, therefore, the impact of enrollment stratification by somatic mutation or expression in tumors of children enrolled on this trial is unknown. Pharmacogenomic profiling of germline AURKA in adults has focused

on cancer susceptibility and early adverse reactions.¹⁴ In this study, we evaluated germline AURKA SNPs and did not find correlation with toxicity during cycle 1 or response.

Despite striking efficacy in pediatric xenograft models in which objective responses were reported in 80% of solid tumor pediatric solid tumor models and all leukemia models²⁸, the objective response rate in children and adolescents receiving single agent alisertib on this trial was less than 5%. In patients receiving 50 mg BID, the C_{max} and AUC_{0–24 h} were 1.3 and 40µM h, respectively²⁷. At the recommended phase 2 dose of 50 mg BID for 7 days, average trough concentrations exceeded 1µM, the efficacious concentration estimated in previous preclinical work. In mice receiving alisertib at 10 mg/kg, the C_{max} and AUC_{0–24 h} were 16 and 39µM h, respectively, with the 12 h level being 1.2 µM²⁷. These data suggest that continuous drug exposure above 1µM throughout each 24-hour dosing period which can only be achieved with twice-daily dosing in mice, is crucial for anti-tumor activity. The initial preclinical studies evaluating alisertib in pediatric patient-derived xenograft models used a dose and schedule that was employed in preclinical assessment of adult cancer models in an attempt to mirror pharmacokinetic-pharmacodynamic relationships expected to be tolerated in humans. However, the continuous treatment schedule was too myelosuppressive in adult phase 1 studies and a 7-day on, 14-day off regimen was adopted to minimize toxicity. Thus, we hypothesized that the dose and schedule used in our trial might account for the discordance between the preclinical and clinical activity observed.

Concurrent with this trial, we tested alisertib in ALL xenograft models utilizing an intermittent dose and schedule, and found significantly less efficacy compared with the previously tested schedule. This demonstrates an important point in the clinical translation of new agents. While careful consideration is made to maximize the clinical relevance and translatability of pre-clinical oncology studies, numerous variables can influence the extent to which a drug against a given target will cause toxicity. This study highlights the critical importance of performing reverse translational studies to rigorously reproduce results when dose or schedule significantly change as a result of early phase human clinical trials. The continuous treatment schedule of alisertib, which was shown to be more effective in preclinical models than the 1-week administration schedule, was not feasible due to toxicity to pursue in the clinical setting, providing a potential explanation for the differential observed between preclinical and clinical anti-tumor activity. To increase our confidence in the translation of results from preclinical studies, there needs to be continued efforts to redesign preclinical experiments as we learn from the corresponding human experience.

Identifying applicable preclinical cancer models remains a major challenge in augmenting the effectiveness of drug development and predicting success in the clinic. All models are limited and interrogating the complexity of human cancers in the laboratory remains a challenge that contributes appreciably to attrition in drug development. In recent years, patient-derived xenografts obtained by direct implants of human tumors in immunodeficient mice and then passaged directly from mouse to mouse have emerged as an important platform for translational oncology research³⁸. The ability of these models to predict clinical outcomes is being optimized through murine humanization strategies to improve the reach of these models as reliable tools for exploring tumor intrinsic and extrinsic heterogeneity, clonal evolution under the selective pressure of our therapies, discovery of integral

biomarkers and predictability of drug response in the clinic³⁸. Numerous challenges and limitations remain, including the lack of a proper anatomical and metastatic niche, engraftment failure of certain tumor subtypes, access to imaging technologies for robust tumor visualization, and hurdles to achieve complete human immune system reconstitution³⁸.

To improve the efficiency of this two-stage phase 2 trial, this trial was prospectively designed to include data from 21 patients on the phase 1 single agent alisertib trial (ADVL0812), who were treated at the recommended dose and met eligibility criteria for this trial, including 5 patients with NBL- measurable disease, 12 patients with NBL-MIGB evaluable disease, 2 patients with non-RMS soft tissue sarcoma and 2 patients with hepatoblastoma. Given the rarity of relapsed pediatric cancers, trial designs that improve efficiency are essential.

Clinical trials evaluating alisertib in combination with cytotoxic agents have shown anti-tumor activity in children and adolescents with relapsed solid tumors including neuroblastoma^{39,40}. Given the lack of objective response observed in this comprehensive single agent clinical trial of alisertib as well as dose limiting myelosuppression, alternative strategies, including novel: novel combinations simultaneously targeting other oncogenic signaling pathways or exploiting the pro-apoptotic machinery⁴¹, should be explored to harness this pathway and simultaneously minimize toxicity.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Financial Support: UM1CA097452 Phase 1 and Pilot Consortium Grant; National Institutes of Health National Cancer Institute NOI-CM-42216 and NOI-CM-91001-03; St Baldrick's Foundation; and Cookies for Kids Foundation

Abbreviations Key Table

AAK	Aurora A Kinase
ALL	Acute Lymphoblastic Leukemia
ALT	Alanine Aminotransferase
ANC	Absolute Neutrophil Count
AML	Acute Myeloid Leukemia
ATRT	Atypical Teratoid Rhabdoid Tumor
AUC	Area Under the Concentration × Time Curve
AURK	Aurora A Kinase Gene
CBC	Complete Blood Count with Differential

CL/F	Apparent Clearance
C_{max}	Maximum Concentration, peak
C_{min}	Minimum Concentration, trough
CR	Complete Remission, Complete Response
CT	Computed Tomography Imaging
CTCAE	Common Terminology Criteria for Adverse Events
dL	Deciliter
DLT	Dose Limiting Toxicity
EDTA	Ethylenediaminetetraacetic acid
EWS	Ewing Sarcoma Family of Tumors
GCT	Germ Cell Tumor (Malignant)
GGT	Gamma-Glutamyl Transferase
g	Gram
h	Hour
HBL	Hepatoblastoma
INR	International Normalized Ratio
L	Liter
NBL	Neuroblastoma
NRSTS	Non- Rhabdo Soft Tissue Sarcoma
m²	Square meters of body surface area
mg	Milligram
MIBG	Metaiodobenzylguanidine
mm³	Cubic millimeters
MRI	Magnetic Resonance Imaging
OS	Osteosarcoma
PB	Peripheral blood
PD	Progressive Disease
PK	Pharmacokinetics
PG	Pharmacogenomics

PPTP	Pediatric Preclinical Testing Program
PR	Partial Response, Partial Remission
RECIST	Response Criteria in Solid Tumors
RMS	Rhabdomyosarcoma
SD	Stable Disease
UGT1A1	UDP-glucuronosyltransferase 1A1
ULN	Upper Limit of Normal
WBC	White Blood Cell Count
WT	Wilms Tumor
μL	Microliter
μM	micromolar

References

- Gautschi O, Heighway J, Mack PC, et al.: Aurora kinases as anticancer drug targets. *Clinical cancer research : an official journal of the American Association for Cancer Research* 14:1639–48, 2008 [PubMed: 18347165]
- Nigg EA: Mitotic kinases as regulators of cell division and its checkpoints. *Nat Rev Mol Cell Biol* 2:21–32, 2001 [PubMed: 11413462]
- Lapenna S, Giordano A: Cell cycle kinases as therapeutic targets for cancer. *Nature reviews. Drug discovery* 8:547–66, 2009 [PubMed: 19568282]
- Sen S, Zhou H, Zhang RD, et al.: Amplification/overexpression of a mitotic kinase gene in human bladder cancer. *Journal of the National Cancer Institute* 94:1320–9, 2002 [PubMed: 12208897]
- Borges KS, Moreno DA, Martinelli CE Jr., et al.: Spindle assembly checkpoint gene expression in childhood adrenocortical tumors (ACT): Overexpression of Aurora kinases A and B is associated with a poor prognosis. *Pediatr Blood Cancer* 60:1809–16, 2013 [PubMed: 23788275]
- Hartsink-Segers SA, Zwaan CM, Exalto C, et al.: Aurora kinases in childhood acute leukemia: the promise of aurora B as therapeutic target. *Leukemia* 27:560–8, 2013 [PubMed: 22940834]
- Venkataraman S, Alimova I, Tello T, et al.: Targeting Aurora Kinase A enhances radiation sensitivity of atypical teratoid rhabdoid tumor cells. *Journal of neuro-oncology* 107:517–26, 2012 [PubMed: 22246202]
- Romain C, Paul P, Kim KW, et al.: Targeting Aurora kinase-A downregulates cell proliferation and angiogenesis in neuroblastoma. *J Pediatr Surg* 49:159–65, 2014 [PubMed: 24439602]
- Ewart-Toland A, Briassouli P, de Koning JP, et al.: Identification of Stk6/STK15 as a candidate low-penetrance tumor-susceptibility gene in mouse and human. *Nature genetics* 34:403–12, 2003 [PubMed: 12881723]
- Dicioccio RA, Song H, Waterfall C, et al.: STK15 polymorphisms and association with risk of invasive ovarian cancer. *Cancer epidemiology, biomarkers & prevention : a publication of the American Association for Cancer Research, cosponsored by the American Society of Preventive Oncology* 13:1589–94, 2004
- Egan KM, Newcomb PA, Ambrosone CB, et al.: STK15 polymorphism and breast cancer risk in a population-based study. *Carcinogenesis* 25:2149–53, 2004 [PubMed: 15271853]
- Tang W, Qiu H, Jiang H, et al.: Aurora-A V57I (rs1047972) polymorphism and cancer susceptibility: a meta-analysis involving 27,269 subjects. *PloS one* 9:e90328, 2014 [PubMed: 24598702]

13. Xu L, Zhou X, Jiang F, et al.: STK15 rs2273535 polymorphism and cancer risk: a meta-analysis of 74,896 subjects. *Cancer epidemiology* 38:111–7, 2014 [PubMed: 24252226]
14. Ishikawa A, Suga T, Shoji Y, et al.: Genetic variants of NPAT-ATM and AURKA are associated with an early adverse reaction in the gastrointestinal tract of patients with cervical cancer treated with pelvic radiation therapy. *Int J Radiat Oncol Biol Phys* 81:1144–52, 2011 [PubMed: 21050672]
15. Barr PM, Li H, Spier C, et al.: Phase II Intergroup Trial of Alisertib in Relapsed and Refractory Peripheral T-Cell Lymphoma and Transformed Mycosis Fungoides: SWOG 1108. *J Clin Oncol* 33:2399–404, 2015 [PubMed: 26077240]
16. Melichar B, Adenis A, Lockhart AC, et al.: Safety and activity of alisertib, an investigational aurora kinase A inhibitor, in patients with breast cancer, small-cell lung cancer, non-small-cell lung cancer, head and neck squamous-cell carcinoma, and gastro-oesophageal adenocarcinoma: a five-arm phase 2 study. *Lancet Oncol* 16:395–405, 2015 [PubMed: 25728526]
17. Friedberg JW, Mahadevan D, Cebula E, et al.: Phase II study of alisertib, a selective Aurora A kinase inhibitor, in relapsed and refractory aggressive B- and T-cell non-Hodgkin lymphomas. *J Clin Oncol* 32:44–50, 2014 [PubMed: 24043741]
18. Manfredi MG, Ecsedy JA, Chakravarty A, et al.: Characterization of Alisertib (MLN8237), an investigational small-molecule inhibitor of aurora A kinase using novel in vivo pharmacodynamic assays. *Clin Cancer Res* 17:7614–24, 2011 [PubMed: 22016509]
19. Manfredi MG, Ecsedy JA, Chakravarty A, et al.: Characterization of Alisertib (MLN8237), an investigational small-molecule inhibitor of aurora A kinase using novel in vivo pharmacodynamic assays. *Clinical cancer research : an official journal of the American Association for Cancer Research* 17:7614–24, 2011 [PubMed: 22016509]
20. Yang JJ, Li Y, Chakravarty A, et al.: Preclinical drug metabolism and pharmacokinetics, and prediction of human pharmacokinetics and efficacious dose of the investigational Aurora A kinase inhibitor alisertib (MLN8237). *Drug metabolism letters* 7:96–104, 2014 [PubMed: 24484538]
21. Cervantes A, Elez E, Roda D, et al.: Phase I pharmacokinetic/pharmacodynamic study of MLN8237, an investigational, oral, selective aurora a kinase inhibitor, in patients with advanced solid tumors. *Clinical cancer research : an official journal of the American Association for Cancer Research* 18:4764–74, 2012 [PubMed: 22753585]
22. Falchook G, Kurzrock R, Gouw L, et al.: Investigational Aurora A kinase inhibitor alisertib (MLN8237) as an enteric-coated tablet formulation in non-hematologic malignancies: phase 1 dose-escalation study. *Investigational new drugs* 32:1181–7, 2014 [PubMed: 24879333]
23. Venkatakrishnan K, Zhou X, Ecsedy J, et al.: Dose selection for the investigational anticancer agent alisertib (MLN8237): Pharmacokinetics, pharmacodynamics, and exposure-safety relationships. *J Clin Pharmacol* 55:336–47, 2015 [PubMed: 25302940]
24. Matulonis UA, Sharma S, Ghamande S, et al.: Phase II study of MLN8237 (alisertib), an investigational Aurora A kinase inhibitor, in patients with platinum-resistant or -refractory epithelial ovarian, fallopian tube, or primary peritoneal carcinoma. *Gynecologic oncology* 127:63–9, 2012 [PubMed: 22772063]
25. Goldberg SL, Fenaux P, Craig MD, et al.: An exploratory phase 2 study of investigational Aurora A kinase inhibitor alisertib (MLN8237) in acute myelogenous leukemia and myelodysplastic syndromes. *Leukemia research reports* 3:58–61, 2014 [PubMed: 25068104]
26. Friedberg JW, Mahadevan D, Cebula E, et al.: Phase II study of alisertib, a selective Aurora A kinase inhibitor, in relapsed and refractory aggressive B- and T-cell non-Hodgkin lymphomas. *Journal of clinical oncology : official journal of the American Society of Clinical Oncology* 32:44–50, 2014 [PubMed: 24043741]
27. Carol H, Boehm I, Reynolds CP, et al.: Efficacy and pharmacokinetic/pharmacodynamic evaluation of the Aurora kinase A inhibitor MLN8237 against preclinical models of pediatric cancer. *Cancer chemotherapy and pharmacology* 68:1291–304, 2011 [PubMed: 21448591]
28. Maris JM, Morton CL, Gorlick R, et al.: Initial testing of the aurora kinase a inhibitor MLN8237 by the Pediatric Preclinical Testing Program (PPTP). *Pediatr Blood Cancer*, 2010
29. Michaelis M, Selt F, Rothweiler F, et al.: Aurora kinases as targets in drug-resistant neuroblastoma cells. *PLoS one* 9:e108758, 2014 [PubMed: 25268132]

30. Brockmann M, Poon E, Berry T, et al.: Small molecule inhibitors of aurora-a induce proteasomal degradation of N-myc in childhood neuroblastoma. *Cancer cell* 24:75–89, 2013 [PubMed: 23792191]
31. Mosse YP, Lipsitz E, Fox E, et al.: Pediatric phase I trial and pharmacokinetic study of MLN8237, an investigational oral selective small-molecule inhibitor of Aurora kinase A: a Children's Oncology Group Phase I Consortium study. *Clinical cancer research : an official journal of the American Association for Cancer Research* 18:6058–64, 2012 [PubMed: 22988055]
32. Eisenhauer EA, Therasse P, Bogaerts J, et al.: New response evaluation criteria in solid tumours: revised RECIST guideline (version 1.1). *Eur J Cancer* 45:228–47, 2009 [PubMed: 19097774]
33. Ady N, Zucker JM, Asselain B, et al.: A new 123I-MIBG whole body scan scoring method-- application to the prediction of the response of metastases to induction chemotherapy in stage IV neuroblastoma. *European journal of cancer* 31A:256–61, 1995 [PubMed: 7718334]
34. Cheson BD, Bennett JM, Kopecky KJ, et al.: Revised recommendations of the International Working Group for Diagnosis, Standardization of Response Criteria, Treatment Outcomes, and Reporting Standards for Therapeutic Trials in Acute Myeloid Leukemia. *Journal of clinical oncology : official journal of the American Society of Clinical Oncology* 21:4642–9, 2003 [PubMed: 14673054]
35. Lipsitz E, Moorthy G, Mosse Y, et al.: A sensitive and selective liquid chromatography/tandem mass spectrometry method for determination of MLN8237 in human plasma. *Journal of chromatography. B, Analytical technologies in the biomedical and life sciences* 878:2369–73, 2010 [PubMed: 20685183]
36. Akaba K, Kimura T, Sasaki A, et al.: Neonatal hyperbilirubinemia and mutation of the bilirubin uridine diphosphate-glucuronosyltransferase gene: a common missense mutation among Japanese, Koreans and Chinese. *Biochem Mol Biol Int* 46:21–6, 1998 [PubMed: 9784835]
37. Houghton PJ, Morton CL, Tucker C, et al.: The pediatric preclinical testing program: description of models and early testing results. *Pediatr Blood Cancer* 49:928–40, 2007 [PubMed: 17066459]
38. Byrne AT, Alferez DG, Amant F, et al.: Interrogating open issues in cancer precision medicine with patient-derived xenografts. *Nat Rev Cancer* 17:254–268, 2017 [PubMed: 28104906]
39. DuBois SG, Marachelian A, Fox E, et al.: Phase I Study of the Aurora A Kinase Inhibitor Alisertib in Combination With Irinotecan and Temozolomide for Patients With Relapsed or Refractory Neuroblastoma: A NANT (New Approaches to Neuroblastoma Therapy) Trial. *J Clin Oncol* 34:1368–75, 2016 [PubMed: 26884555]
40. Dubois SG, Mosse YP, Fox E, et al.: Phase 2 Trial of Alisertib in Combination with Irinotecan and Temozolomide for Patients with Relapsed or Refractory Neuroblastoma. *Clin Cancer Res*, 2018
41. Ham J, Costa C, Sano R, et al.: Exploitation of the Apoptosis-Primed State of MYCN-Amplified Neuroblastoma to Develop a Potent and Specific Targeted Therapy Combination. *Cancer Cell* 29:159–72, 2016 [PubMed: 26859456]

Statement of Translational Relevance

The Aurora kinases are a family of serine-threonine kinases that play an essential role in regulating chromosome assembly and segregation during mitosis and are critical for cell proliferation. Aurora A dysregulation has been implicated in cancer, and as such, is a rational therapeutic target. This phase 2 study of alisertib (MLN2837), an oral small molecular inhibitor of Aurora A kinase, was evaluated in 137 pediatric patients with relapsed/refractory solid tumors or acute leukemia. The lack of robust objective responses suggests that alisertib, as a single agent, has limited anti-tumor activity and alternative strategies, including novel combinations simultaneously targeting other oncogenic signaling pathways, should be explored to harness this pathway and simultaneously minimize toxicity. The continuous treatment schedule of alisertib, which was shown to be more effective in preclinical models than the 1-week administration schedule, was not feasible in the clinical setting, providing a potential explanation for the differential observed between preclinical and clinical anti-tumor activity.




Figure 1.

Sagittal and axial SPECT/CT images from ^{123}I -MIBG examination performed at baseline, reporting periods (RP) 12 and 24; ^{123}I -MIBG anterior projection planar images at baseline, RP10 and RP 20. Each reporting period is one cycle. Arrows indicate MIBG-avid neuroblastoma in the L3 vertebral body and proximal right femur. The other areas of MIBG positivity on planar images include physiologic uptake in liver, salivary glands, renal collecting system, GI tract, and excretion in bladder. Uptake projecting of the thorax at RP 10 and RP 20 is residual tracer at the port injection site.




Figure 2. Percentage of huCD45 cells in peripheral blood over time (A), and event-free survival curves (B) for ALL-8 engrafted NOD/SCID mice treated with alisertib at 10.4 mg/kg twice daily for 7 days (Schedule A, red), or twice daily for 5 days repeated for 3 weeks (Schedule B, blue) in relation to vehicle-treated controls (dotted line).




Figure 3. Percentage of huCD45+ cells in peripheral blood over time (A), and event-free survival curves (B) for ALL-19 engrafted NOD/SCID mice treated with alisertib at 10.4 mg/kg twice daily for 7 days (Schedule A, red), or twice daily for 5 days repeated for 3 weeks (Schedule B, blue) in relation to vehicle-treated controls (dotted line).

Table 1.

Patient Characteristics at Enrollment

	Total	NBL_Measurable	NBL_Evaluable	AML	ALL	EWS	RMS	NR-ST5	OS	WT	HBL	GCT	MRT
Eligible (N)	139	25	24	11	10	10	10	10	10	10	8	7	4
Age (y) *	10 (2–21)	9 (3–20)	8 (4–19)	10 (3–15)	13 (7–19)	14 (6–20)	12 (4–21)	15 (9–20)	17 (12–21)	10 (3–18)	6 (3–14)	11 (3–19)	5 (2–7)
Female (%)	45%	48%	29%	55%	30%	30%	70%	50%	30%	70%	38%	57%	50%
Race n(%)													
Caucasian	80 (58%)	19 (76%)	15 (63%)	2 (18%)	5 (50%)	8 (80%)	7 (70%)	4 (40%)	7 (70%)	5 (50%)	4 (50%)	4 (57%)	-
Black	21 (15%)	2 (8%)	4 (17%)	2 (18%)	2 (20%)	-	1 (10%)	1 (10%)	2 (20%)	2 (20%)	2 (20%)	-	3 (75%)
Other	6 (4%)	0 (0%)	2 (8%)	1 (9%)	-	-	-	-	1 (10%)	-	1 (10%)	1 (14%)	-
Not Specified	32 (23%)	4 (16%)	3 (12%)	6 (55%)	3 (30%)	2 (20%)	2 (20%)	5 (50%)	-	3 (30%)	1 (10%)	2 (29%)	1 (25%)
Ethnicity N (%)													
Hispanic	30 (22%)	3 (12%)	1 (4%)	4 (36%)	2 (20%)	1 (10%)	1 (10%)	3 (30%)	3 (30%)	4 (40%)	3 (38%)	4 (57%)	1 (25%)
Non-Hispanic	97 (70%)	19 (76%)	22 (92%)	5 (46%)	8 (80%)	8 (80%)	7 (70%)	6 (60%)	7 (70%)	5 (50%)	5 (62%)	2 (29%)	3 (75%)
Not Reported	12 (8%)	3 (12%)	1 (4%)	2 (18%)	-	1 (10%)	2 (20%)	1 (10%)	-	1 (10%)	-	1 (14%)	-

* Median (range)

NBL –neuroblastoma; AML- Acute Myeloid Leukemia; ALL- Acute Lymphoblastic Leukemia; EWS-Ewing Sarcoma; RMS- Rhabdomyosarcoma; NR-ST5- Non-Rhabdo Soft Tissue Sarcoma; OS- Osteosarcoma; WT- Wilms Tumor; HBL- Hepatoblastoma; GCT- Malignant Germ Cell Tumor; MRT- Malignant Rhabdoid Tumor (2 patients with CNS atypical teratoid rhabdoid tumors and 2 patients with extracranial malignant rhabdoid tumors).

Table 2:

Frequency of Alisertib Related Grade 3 or 4 Toxicity in All Cycles (n=500)

CTCAE Class	Toxicity	Grade 3 n (%)	Grade 4 n (%)	Grade 5 n (%)
Hematologic	Anemia	63 (12.6%)	5 (1%)	
	Febrile neutropenia	18 (3.6%)		
	Lymphopenia	47 (9.4%)	14 (2.8%)	
	Neutropenia	124 (24.8%)	137 (27.4%)	
	Thrombocytopenia	54 (10.8%)	50 (10%)	
	Serum amylase increased	1 (0.2%)		
	Leukopenia	117 (23.4%)	48 (9.6%)	
Eye Disorders	Photophobia	1 (0.2%)		
Gastrointestinal	Diarrhea	2 (0.4%)		
	Enterocolitis	1 (0.2%)		
	Oral mucositis	19 (3.8%)		
	Oral pain	5 (1%)		
	Nausea	2 (0.4%)		
	Vomiting	2 (0.4%)		
Investigations (Laboratory)	ALT increased	17 (3.4%)		
	AST increased	10 (2%)		
	Hyperbilirubinemia	3 (0.6%)		
	GGT increased	1 (0.2%)		
	INR increased	1 (0.2%)		
Infection	Infection	1 (0.2%)		
	Pneumonia	1 (0.2%)		
	Urinary tract infection	1 (0.2%)		
Metabolism/Nutrition	Anorexia	1 (0.2%)		
	Dehydration	6 (1.2%)		
	Hyperuricemia		1 (0.2%)	
	Hypoalbuminemia	1 (0.2%)		
	Hypocalcemia	1 (0.2%)		
	Hypokalemia	4 (0.8%)	1 (0.2%)	
	Hyponatremia	3 (0.6%)		
	Hypophosphatemia	1 (0.2%)		
Tumor lysis syndrome		1 (0.2%)		
Psychiatric	Depression		1 (0.2%)	
Neurological	Dizziness	14 (2.8%)		
	Hypersomnia	1 (0.2%)		
Hepatobiliary	Hepatic hemorrhage			1 (0.2%)
Respiratory	Respiratory failure			1 (0.2%)
Skin/Dermatological	Palmar-plantar erythrodysesthesia	2 (0.4%)		

Table 3.

Responses to Alisertib

Stratum	Response Evaluable	Responder		Non-responders		
		Complete Response	Partial Response	Stable Disease n patients, median (range) cycles administered	Non-responders	Progressive Disease
Neuroblastoma (Measurable)	24		1	2 (6, 13 cycles)	5	16
Neuroblastoma (MIBG Evaluable)	24	1	1	9 13 (5–35) cycles	4	9
Acute Lymphoblastic Leukemia	10			3 (1,2,2 cycles)	1	6
Acute Myeloid Leukemia	10				10	
Ewing Sarcoma	10			3 (4, 5, 5 cycles)	2	5
Rhabdomyosarcoma	10			1 15 cycles	2	7
Non-Rhabdo Soft Tissue Sarcoma	10			1 5 cycles	2	7
Osteosarcoma	10				2	8
Wilms tumor	10	1		1 31 cycles	1	7
Hepatoblastoma	8		1*	1 5 cycles	2	4
Germ Cell Tumor	7			2 4 and 5 cycles	1	4
Rhabdoid Tumors	4				1	3
Total	137	2	3	23	33	76

* Reported previously in ADVL0812 publication.³¹

Table 4A:

AURKA, UGT1A1*28 and UGT1A1 PBREM genotype distribution.

Gene	RsSNP ID	Description	Genotype ^a (WW/WV/VV)
AURKA	rs2273535	91 A>T, Ile31Phe	51/27/8 ^b
	rs1047972	169 G>A, Val57Ile	63/21/2 ^b
UGT1A1*28	rs8175347	TA repeat (5,6,7,8)	39/32/11
UGT1A1 PBREM	rs4124874	-3279 T>G	23/42/22
	rs10929302	-3156 G>A	46/33/8

^aW = Wild Type Allele, V = Variant allele; For UGT1A1*28- WW = 66 or 57, WV = 67 or 68, and VV = 77 or 78

^bOne sample could not be genotyped with the available DNA
Five samples could not be genotyped with the available DNA

Table 4B.

Toxicity versus AURKA genotype

AURKA Genotype	Cycle 1 DLT		Grade 2+ Toxicity		
	Yes	No	Yes	No	
91 A>T	Total	13	73	72	14
	WW	8	43	42	9
	WV	5	22	22	5
	VV	0	8	8	0
169G>A	Total	13	73	72	14
	WW	10	53	54	9
	WV	3	18	17	4
	VV	0	2	1	1
Haplotype	Total	13	73	72	14
	H1	5	25	26	4
	H2	8	48	46	10

Toxicity versus UGT1A1 metabolizer phenotype and alisertib trough concentration (C_{\min}) on day 4.

Table 4C:

	EM		I/E, I, P	
	Grade 2; No (n=7)	Grade 2; Yes (n=9)	Grade 2; No (n=8)	Grade 2; Yes (n=8)
Alisertib C_{\min} (μM) Mean \pm SD	0.81 \pm 0.43	1.17 \pm 1.15	2.25 \pm 1.53	1.87 \pm 0.82
Alisertib C_{\min} (μM) Median (range)	0.99 (0.2–1.24)	0.53 (0.08–3.51)	2.22 (0.6–5.65)	1.80 (0.97–3.10)

Leukemic growth delay summary and clinical scoring for the two treatment schedules for xenografts ALL-8 and ALL-19.

Table 5.

	Treatment Schedule*	Median EFS (days)		EFS T-C (days)	EFS T/C	p value**	Median Response Score
		Control	Treated				
ALL-8	Schedule A	8.9	19.8	10.9	2.2	0.002	PD2
	Schedule B	8.9	31.9	23.0	3.6	0.002	SD
ALL-19	Schedule A	6.6	13.6	7.0	2.0	0.584	PD2
	Schedule B	6.6	31.2	24.6	4.7	0.002	CR

* Schedule A, twice daily \times 7 days; Schedule B, twice daily \times 5 days \times 3 weeks

** Statistically significant results are in bold.