

## RESEARCH ARTICLE

# *Hermetia illucens* in diets for zebrafish (*Danio rerio*): A study of bacterial diversity by using PCR-DGGE and metagenomic sequencing

Andrea Osimani<sup>1</sup>, Vesna Milanović<sup>1</sup>, Andrea Roncolini<sup>1\*</sup>, Paola Riolo<sup>1</sup>, Sara Ruschioni<sup>1</sup>, Nunzio Isidoro<sup>1</sup>, Nino Loreto<sup>1</sup>, Elena Franciosi<sup>2</sup>, Kieran Tuohy<sup>2</sup>, Ike Olivotto<sup>3</sup>, Matteo Zarantonello<sup>3</sup>, Federica Cardinali<sup>1</sup>, Cristiana Garofalo<sup>1</sup>, Lucia Aquilanti<sup>1\*</sup>, Francesca Clementi<sup>1</sup>

**1** Dipartimento di Scienze Agrarie, Alimentari ed Ambientali, Università Politecnica delle Marche, Ancona, Italy, **2** Food Quality and Nutrition Department (DQAN), Research and Innovation Center, Fondazione Edmund Mach (FEM), San Michele all'Adige, Italy, **3** Dipartimento di Scienze della Vita e dell'Ambiente, Università Politecnica delle Marche, Ancona, Italy

\* [l.aquilanti@univpm.it](mailto:l.aquilanti@univpm.it)(LA); [a.roncolini@pm.univpm.it](mailto:a.roncolini@pm.univpm.it)(AR)



## Abstract

In the present research, bacterial diversity was studied during a 6-month feeding trial utilizing zebrafish (*Danio rerio*) fed *Hermetia illucens* reared on different substrates with an emphasis on fish gut bacterial diversity. A polyphasic approach based on viable counting, PCR-DGGE and metagenomic 16S rRNA gene amplicon target sequencing was applied. Two different *H. illucens* groups were reared on coffee by-products (C) or a mixture of vegetables (S). Viable counts showed a wide variability based on substrate. PCR-DGGE and Illumina sequencing allowed the major and minor bacterial taxa to be detected. Both samples of larvae and their frass reared on the S substrate showed the highest richness and evenness of bacterial communities, whereas zebrafish (ZHC) fed *H. illucens* reared on substrate C and zebrafish (ZHS) fed *H. illucens* reared on substrate S had the lowest bacterial richness and evenness. A stimulating effect of bioactive compounds from coffee by-products on the occurrence of Lactobacillaceae and Leuconostocaceae in *H. illucens* reared on substrate C has been hypothesized. Zebrafish gut samples originating from the two feeding trials showed complex microbial patterns in which Actinobacteria and Alteromonadales were always detected, irrespective of the diet used. Enterobacteriaceae in fish guts were more abundant in ZHS than in ZHC, thus suggesting an influence of the bioactive compounds (chlorogenic and caffeic acids) in the substrate on Enterobacteriaceae in fish guts. ZHC showed a higher abundance of Clostridia than did ZHS, which was likely explained by stimulating activity on the bacteria in this class by the bioactive compounds contained in *H. illucens* reared on substrate C. An influence of the microbiota of *H. illucens* or insect-derived bioactive compounds on the gut microbiota of zebrafish has been suggested. The presence of bacteria consistently associated with zebrafish guts has been found irrespective of the diet, thus attesting to the likely stability of the core fish microbiota.

## OPEN ACCESS

**Citation:** Osimani A, Milanović V, Roncolini A, Riolo P, Ruschioni S, Isidoro N, et al. (2019) *Hermetia illucens* in diets for zebrafish (*Danio rerio*): A study of bacterial diversity by using PCR-DGGE and metagenomic sequencing. PLoS ONE 14(12): e0225956. <https://doi.org/10.1371/journal.pone.0225956>

**Editor:** Patrizia Falabella, Università degli Studi della Basilicata, ITALY

**Received:** August 12, 2019

**Accepted:** November 16, 2019

**Published:** December 10, 2019

**Copyright:** © 2019 Osimani et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Data Availability Statement:** All relevant data are within the manuscript.

**Funding:** The authors received no specific funding for this work.

**Competing interests:** The authors have declared that no competing interests exist.

## Introduction

The use of edible insects as food and feed could represent a suitable source of alternative protein due to their high-quality nutritional values and the environmental sustainability of their rearing [1, 2]. Indeed, edible insects can grow on substrates deriving from food industry by-products, thus adding value to these organic matrices. Moreover, edible insects are already reared in Asia, Africa, Oceania and South America, thus representing a growing source of income [3, 4]. In the European Union (EU), stakeholders have recently begun paying attention to the applications of edible insects as food or feed. In 2015, the European Food Safety Authority (EFSA) listed a few insect species with potential use as food and feed in the EU. The EFSA list included the following species: *Musca domestica*, *Hermetia illucens*, *Tenebrio molitor*, *Zophobas atratus*, *Alphitobius diaperinus*, *Galleria mellonella*, *Achroia grisella*, *Bombyx mori*, *Acheta domesticus*, *Gryllobates sigillatus*, *Locusta migratoria migratorioides*, and *Schistocerca americana* [5]. The EFSA also highlighted the need for in-depth studies to assess the safety of edible insects to achieve large-scale production [6]. Since insects can be natural reservoirs of microorganisms that can be transferred to the environment and to living animals used as food sources, their microbiological assessment must be properly conducted [7].

More recently, the European Commission issued Regulation No. 2017/893 of 24 May 2017 that allowed the use of insects as processed animal proteins (PAPs) in aquaculture [8]; such a Regulation paved the way for a more concrete exploitation of edible insects in the EU. In such a context, *H. illucens* (Diptera:Stratiomyidae) (commonly known as black soldier fly) represents one of the most promising species to be used in aquaculture [4, 9–13]. Indeed, *H. illucens* is economically among the most important farmed insects in the world [14]. It is noteworthy that, although many studies have already been carried out to optimize the rearing conditions and to increase the yield of the bioconversion process, data on microbiological aspects of this insect are still scarce [15]. Recent studies have highlighted that *H. illucens* larvae can be carriers of microorganisms whose presence may be influenced in part by the rearing substrate. De Smet et al. [16] recently reviewed the composition of microbial communities of *H. illucens* larvae reared on different substrates, such as food waste, cooked rice, calf forage and the so-called Gainesville housefly diet. The authors reported that the presence of a core microbiota in the insect gut can be conceivable, although the different rearing substrates affected the relative abundances of the detected microbial taxa [16]. Similarly, Wynants et al. [17], highlighted that *H. illucens* larvae reared at several locations showed the presence of a core microbiota among which the phyla Firmicutes and Proteobacteria were dominant. The occurrence of interacting factors, both abiotic and biotic, capable of affecting the larval microbiota, was also considered by the same authors [17]. Interestingly, Bruno et al. [15] reported a variable influence of ingested food on the microbiota of different gut regions of *H. illucens* larvae. The same study reported that an excessively high protein content in the feeding substrate seemed to favor midgut dysbiosis. Furthermore, an influence of the rearing substrate on *H. illucens* larval microbiota was also highlighted by [18].

In the present research bacterial diversity was studied during a 6-month feeding trial with zebrafish (*Danio rerio*) fed *H. illucens* reared on different substrates, with emphasis on fish gut bacterial diversity. To this end, a polyphasic approach based on viable counting, Polymerase Chain Reaction-Denaturing Gradient Gel Electrophoresis (PCR-DGGE) and metagenomic 16S rRNA gene amplicon target sequencing (Illumina sequencing) was applied.

## Materials and methods

### Insect rearing

A *Hermetia illucens* laboratory colony was established from commercially available larvae (Smart Bugs, Ponzano Veneto, Italy). The rearing was maintained under controlled conditions

(27°C ±2°C, 70% ±10% RH, and 14:10 light:dark photoperiod). Larvae were placed in separate sterile plastic boxes (57 × 39 × 28 cm) until pupal stage was reached. Newly emerged adults were placed in two different mating and oviposition cages (2 m<sup>3</sup>). After two days, coffee by-products (C) (coffee silverskin obtained from the roasting process at Saccaria Caffè SRL, Marina di Montemarciano, Italy) or vegetable substrate (S) composed of 50% corn meal and fruit and 50% vegetable mixture were provided, in small sterile plastic boxes (16.5 × 22 cm), as egg-laying substrates. The plastic boxes were changed every 2 days and the eggs collected using brushes. Four thousand 4-day-old larvae (individually counted using tweezers and brushes) were transferred in sterile plastic boxes (57 × 39 × 28 cm) and fed with C (namely, HC larvae) or S (namely, HS larvae). The feeding substrate was supplied *ad libitum* to the developing larvae. The experiments were performed in triplicate for each rearing condition. Larval rearing boxes were covered with a lid provided with a single ventilation hole and equipped with a ramp (35°) and a collection device. When the larvae reached the prepupal stage, they left the feeding substrate and climbed up the ramp into the collecting vessel. Both prepupal groups (HC and HS) were collected, freeze-dried, and processed to obtain full-fat prepupae fish meals. Moreover, HC, HS, C, S and frass samples, composed of excrement from larvae mixed with substrate residues and exuviae, (namely, HCF from HC or HSF from HS) were analyzed as described below.

### Zebrafish rearing

*Danio rerio* fed with either 100% HC or 100% HS were produced as already reported by Zarantoniello et al. [13]. *D. rerio* were spawned and maintained under the rearing conditions already reported by Zarantoniello et al. [13]. Embryos were then collected, counted under a stereomicroscope (Leica Wild M3B, Leica Microsystems, Nussloch, Germany) and randomly divided into two experimental groups (in triplicate) according to the two test diets [13]. Fish (1500 larvae per dietary treatment; 3 × 20 L tanks containing 500 larvae each) were daily fed (2% body weight) the two different insect meals (HC; HS) for 6 months according to Vargas et al. [12] and Zarantoniello et al. [19]. At the end of the experimental period fish were anesthetized with a lethal dose of MS222 (1 g/L, Sigma Aldrich, Saint Louis, Missouri, USA). All efforts were made to minimize animal suffering [16]. The microbiota of ZHC (gut contents of zebrafish fed HC) and ZHS (gut contents of zebrafish fed HS) were then analyzed.

### Microbiological analyses

Microbiological enumeration was carried out on samples C, HC, HCF, S, HS and HSF. Ten grams of each sample were aseptically diluted in 90 mL of sterile peptone water (bacteriological peptone 1 g/L) and homogenized in a Stomacher 400 Circulator apparatus (VWR International PBI, Milan, Italy). Homogenate ten-fold dilutions were inoculated on specific solid media for the enumeration of the following microorganisms: total mesophilic aerobes counted according to the UNI EN ISO 4833:2004 standard method; Enterobacteriaceae counted in according to the ISO 21528–2:2004 standard method; and bacterial spores, for which homogenized samples were subjected to thermal treatment at 80°C for 15 min followed by cooling in iced water to inactivate the vegetative cells [20] and then grown in Standard Plate Count Agar (Oxoid, Basingstoke, UK) incubated at 30°C for 48 h. Presumptive lactic acid bacteria counts were carried out on MRS agar (Oxoid) with cycloheximide (250 mg/L) added, incubated at 37°C for 72 h under anaerobiosis.

The presence of *Listeria monocytogenes* and *Salmonella* spp. was assessed in accordance with AFNOR BIO 12/11-03/04 and AFNOR BIO 12/16-09/05 standard methods, respectively.

Analyses were carried out in duplicate. The results were subjected to ANOVA using the software JMP v.11.0.0 (SAS Institute Inc., Cary, NC) and expressed as the mean of log colony forming units (cfu) per gram  $\pm$  standard deviation of independent duplicate biological experiments.

## Molecular analyses

To gain insight into the microbial species occurring in the analyzed samples, C, HC, HCF, ZHC, S, HS, HSF and ZHS were subjected to culture-independent analyses via PCR-DGGE and Illumina sequencing. To this aim, DNA extraction from samples was carried out as already described by Garofalo et al. [21]. DNA extracts were standardized to 25 ng/ $\mu$ L, and equal portions derived from the biological replicates of each sample were pooled together and vortexed vigorously.

PCR DGGE analysis was performed as described by Milanović et al. [22] through PCR amplification of the V3 region of the 16S rRNA gene using universal prokaryotic primers 338f (5' -ACT CCT ACG GGA GGC AGC AGC AG-3') added with GC clamp and 518r (5' -ATT ACC GCG GCT GCT GG-3') [23]. After the DGGE, bands were excised and reamplified and amplicons sent for sequencing to Genewiz (Leipzig, Germany). The output sequences in FASTA format were compared with those deposited in the GenBank DNA database using the basic BLAST search tools [24].

Samples were further subjected to Illumina analysis. In more detail, a 464-nucleotide sequence of the bacterial V3-V4 region [25, 26] of the 16S rRNA gene (*Escherichia coli* positions 341 to 805) was amplified and processed as already described by Osimani et al. [27].

Unique barcodes were attached before the forward primers; the amplicons were cleaned using the Agencourt AMPure kit (Beckman coulter). The DNA concentration of amplicons was determined using the Quant-iT PicoGreen dsDNA kit (Invitrogen). The quality of the generated amplicon libraries was evaluated with a Bioanalyzer 2100 (Agilent, Palo Alto, CA, USA) using the High Sensitivity DNA Kit (Agilent). Raw paired-end FASTQ files were demultiplexed using idemp (<https://github.com/yhwu/idemp/blob/master/idemp.cpp>) and imported into Quantitative Insights Into Microbial Ecology (QIIME2, version 2018.2) [28]. Sequences were quality filtered, trimmed, denoised, and merged using DADA2 [29]. Chimeric sequences were identified and removed via the consensus method in DADA2. Representative sequences were aligned with MAFFT and used for phylogenetic reconstruction in FastTree using plugins alignment and phylogeny [30, 31]. Alpha and beta diversity metrics were calculated using the core-diversity plugin within QIIME2 and emperor [32]. Taxonomic and compositional analyses were conducted with plugins feature-classifier (<https://github.com/qiime2/q2-feature-classifier>). A pretrained Naive Bayes classifier based on the Greengenes 13\_8 97% operational taxonomic units (OTUs) database (<http://greengenes.secondgenome.com/>), which had been previously trimmed to the V4 region of 16S rDNA, bound by the 341F/805R primer pair, was applied to paired-end sequence reads to generate taxonomy tables.

The data generated by Illumina sequencing were deposited in the NCBI Sequence Read Archive (SRA) and are available under Ac. PRJNA546499.

## Results and discussion

*H. illucens* larvae are able to efficiently convert organic by-products into biomass to be used for feed or other purposes. Hence, *H. illucens* larvae constitute a potentially suitable ingredient for fish feed, as well as a good marketing opportunity for overcoming concerns regarding the use of fishmeal in aquaculture [4, 11]. Notwithstanding, studies on microbiological aspects of the use of *H. illucens* as feed are surprisingly very limited [15].

As reported in Table 1, except for HCF, the samples derived from coffee by-products (C and HC) were characterized by high counts for all the microbial groups assessed. Of note, Enterobacteriaceae counts in samples HC and HCF were lower than that in sample C, reflecting the previously described inhibitory effect of *H. illucens* larvae on Enterobacteriaceae [33]. Surprisingly, the inhibitory effect was not observed in HS or HSF, which exhibited higher Enterobacteriaceae loads than in S samples. The data collected on mesophilic aerobes, bacterial spores, and LAB were consistent with those reported by Wynants et al. [17] that reflected a wide variability based on the substrate, the rearing methods and the timing of larval harvest. In frass samples, viable counts were generally higher, probably because of the accumulation effect during the rearing period [27]. Finally, no *Listeria monocytogenes* or *Salmonella* spp. were detected in any of the analyzed samples. Raw data used for statistical analysis of viable counts are reported in S1 Table.

The data on the overall microbiota collected highlighted very complex populations in almost all the samples where, thanks to the combination of PCR-DGGE and Illumina sequencing, major and minor taxa were detected.

Regarding PCR-DGGE analysis, the closest relatives, the percent identities, and the accession numbers of the sequences obtained are shown in Table 2, whereas Fig 1 shows the DGGE profiles. The raw original image of the DGGE gel is shown in S1 Fig.

Moreover, the results of Illumina sequencing are reported in Fig 2; in addition, raw values of relative abundance (% of bacterial community) in the analyzed samples identified by analysis of V3-V4 regions of the 16S rRNA gene through MiSeq Illumina are reported in S2 Table. In more detail, the DNA extracted from the samples successfully amplified the bacterial V3-V4 16S rRNA gene region. After splitting and quality trimming the raw data, 453,815 reads remained for subsequent analysis. After alignment, the remaining operational taxonomic units (OTUs) were clustered at a 3% distance. To analyze the bacterial community richness of each sample, the number of OTUs, the coverage estimator, the Shannon diversity index and the Chao1 richness estimator were determined using QIIME at 97% similarity levels (Table 3). Good's estimator of coverage was over 99% for all the samples, which indicated that most of the bacterial phylotypes were detected. Based on the Shannon and Chao indices, pooled samples HC, HS and HSF showed the highest richness and evenness of bacterial communities, while pooled samples C, ZHC and S exhibited the lowest.

It is notable that, only for pooled S samples, vegetable DNA and a few saprophytic microbial species were observed; these data are consistent with those recently reported by Osimani et al.

**Table 1. Viable counts from the analyzed samples.**

| Samples         | Total mesophilic aerobes | Bacterial spores       | Lactic acid bacteria   | Enterobacteriaceae     |
|-----------------|--------------------------|------------------------|------------------------|------------------------|
| Rearing chain 1 |                          |                        |                        |                        |
| C               | 8.13±0.03 <sup>a</sup>   | 7.45±0.02 <sup>b</sup> | 7.42±0.01 <sup>a</sup> | 5.35±0.06 <sup>a</sup> |
| HC              | 6.08±0.00 <sup>c</sup>   | 5.92±0.01 <sup>c</sup> | 3.89±0.10 <sup>c</sup> | 3.30±0.02 <sup>b</sup> |
| HCF             | 7.77±0.03 <sup>b</sup>   | 7.71±0.04 <sup>a</sup> | 6.29±0.08 <sup>b</sup> | <1 <sup>c</sup>        |
| Rearing chain 2 |                          |                        |                        |                        |
| S               | 5.83±0.04 <sup>c</sup>   | 2.53±0.04 <sup>c</sup> | 3.93±0.04 <sup>c</sup> | 4.18±0.08 <sup>c</sup> |
| HS              | 7.13±0.03 <sup>b</sup>   | 3.36±0.08 <sup>b</sup> | 7.00±0.00 <sup>b</sup> | 6.90±0.01 <sup>b</sup> |
| HSF             | 10.50±0.06 <sup>a</sup>  | 4.14±0.01 <sup>a</sup> | 7.46±0.01 <sup>a</sup> | 7.48±0.01 <sup>a</sup> |

Results are expressed as log colony forming units (cfu) g<sup>-1</sup>. For microbial counts, means ± standard deviations of triplicate independent experiments are shown. For each rearing chain, within each column, means with different superscript letters are significantly different ( $P < 0.05$ ). C, coffee by-product; HC, *H. illucens* reared on coffee by-product; HCF, frass of *H. illucens* reared on coffee by-product; S, vegetable substrate; HS, *H. illucens* reared on vegetable substrate; HSF, frass of *H. illucens* reared on vegetable substrate.

<https://doi.org/10.1371/journal.pone.0225956.t001>

**Table 2. Sequencing results of the bands excised from the DGGE gel obtained from the amplified fragments of bacterial DNA extracted directly from the analyzed samples.**

| Samples | Bands <sup>a</sup> | Closest relative                    | % Ident. <sup>b</sup> | Acc. No. <sup>c</sup> |
|---------|--------------------|-------------------------------------|-----------------------|-----------------------|
| C       | 1                  | <i>Desulfomonile limimaris</i>      | 99                    | NR_025079             |
|         | 2                  | <i>Agromyces cerinus</i>            | 97                    | NR_036873             |
|         | 3                  | <i>Aliihoeflea</i> sp.              | 97                    | JN210898              |
| HC      | 4                  | <i>Streptomyces</i> sp.             | 97                    | MK702081              |
|         | 5                  | <i>Lysobacter rhizophilus</i>       | 97                    | NR_152715             |
|         | 6                  | <i>Streptomyces</i> sp.             | 95                    | MK702081              |
|         | 7                  | <i>Corynebacterium jeikeium</i>     | 97                    | NR_118863             |
|         | 8                  | <i>Kytococcus sedentarius</i>       | 97                    | NR_074714             |
|         | 9                  | <i>Roseomonas oryzae</i>            | 98                    | NR_137403             |
|         | 10                 | <i>Deferribacter desulfuricans</i>  | 97                    | NR_075025             |
| HCF     | 11                 | <i>Hymenobacter coalescens</i>      | 97                    | NR_151971             |
|         | 12                 | <i>Massilia alkalitolerans</i>      | 97                    | NR_043094             |
|         | 13                 | <i>Blastochloris viridis</i>        | 98                    | NR_041712             |
| ZHC     | 14                 | <i>Blastochloris viridis</i>        | 97                    | NR_041712             |
|         | 15                 | <i>Mobilicoccus pelagius</i>        | 98                    | NR_113143             |
|         | 16                 | <i>Pseudonocardia xishanensis</i>   | 98                    | NR_108411             |
|         | 17                 | <i>Saccharopolyspora</i> sp.        | 96                    | MH777902              |
|         | 18                 | <i>Corynebacterium</i> sp.          | 97                    | MH400695              |
|         | 19                 | <i>Phycococcus endophyticus</i>     | 97                    | NR_148775             |
|         | 20                 | <i>Saccharomonospora</i> sp.        | 97                    | KP639601              |
| S       | 21                 | <i>Corynebacterium halotolerans</i> | 97                    | NR_102500             |
|         | 22                 | <i>Phycococcus endophyticus</i>     | 97                    | NR_148775             |
|         | 23                 | Failed                              | -                     | -                     |
|         | 24                 | <i>Saccharomonospora</i> sp.        | 96                    | KP639601              |
| HS      | 25                 | <i>Lysobacter</i> sp.               | 98                    | MG237861              |
|         | 26                 | <i>Rhodocyclus purpureus</i>        | 97                    | NR_044679             |
|         | 27                 | <i>Aliihoeflea</i> sp.              | 97                    | JN210898              |
| HSF     | 28                 | <i>Lysobacter</i> sp.               | 97                    | MG237861              |
|         | 29                 | <i>Pseudomonas</i> sp.              | 98                    | KX079778              |
|         | 30                 | <i>Bacillus</i> sp.                 | 97                    | KX681796              |
|         | 31                 | <i>Phycococcus endophyticus</i>     | 97                    | NR_148775             |
|         | 32                 | <i>Janibacter</i> sp.               | 97                    | DQ268766              |
| ZHS     | 33                 | <i>Microbacterium</i> sp.           | 98                    | JN935776              |
|         | 34                 | <i>Shewanella</i> sp.               | 97                    | MF155923              |
|         | 35                 | <i>Bacillus</i> sp.                 | 98                    | KX785165              |
|         | 36                 | <i>Lysobacter</i> sp.               | 97                    | MG237861              |
|         | 37                 | <i>Phycococcus endophyticus</i>     | 97                    | NR_148775             |
|         | 38                 | <i>Shewanella</i> sp.               | 97                    | MF155923              |
|         | 39                 | <i>Microbacterium</i> sp.           | 98                    | JN935776              |


<sup>a</sup> Bands are numbered as indicated in Fig 1.

<sup>b</sup> Percentage of identical nucleotides in the sequence obtained from the isolates and the sequence of the closest relative found in the GenBank database.

<sup>c</sup> Accession number of the sequence of the closest relative found by BLAST search.

C, coffee by-product; HC, *H. illucens* reared on coffee by-product; HCF, frass of *H. illucens* reared on coffee by-product; S, vegetable substrate; HS, *H. illucens* reared on vegetable substrate; HSF, frass of *H. illucens* reared on vegetable substrate; ZHC, gut content of zebrafish fed HC; ZHS, gut content of zebrafish fed HS.

<https://doi.org/10.1371/journal.pone.0225956.t002>




**Fig 1. Bacterial DGGE profiles of the DNA extracted directly from the analyzed samples and amplified with primers 338fGC and 518r.** The bands indicated by the letters were excised, reamplified and subjected to sequencing. The identities of the bands are given in Table 2. C, coffee by-product; HC, *H. illucens* reared on coffee by-product; HCF, frass of *H. illucens* reared on coffee by-product; S, vegetable substrate; HS, *H. illucens* reared on vegetable substrate; HSF, frass of *H. illucens* reared on vegetable substrate; ZHC, gut content of zebrafish fed HC; ZHS, gut content of zebrafish fed HS.

<https://doi.org/10.1371/journal.pone.0225956.g001>

[27] for wheat meal used as insect feed. Pooled C samples revealed the dominance of Enterobacteriaceae, Brevibacteriaceae, *Corynebacterium*, Xanthomonadaceae and Rhizobiales (within this latter family, *Aliihoeflea* sp. was also detected via PCR-DGGE). The latter two taxa showed elevated values of percent relative abundance in pooled HCF samples. Pooled HC samples were characterized by the dominance of Leuconostocaceae, Lactobacillaceae and Enterococcaceae, which were minorities in C, thus suggesting a positive selective pressure for these taxa in the insect gut. As reported by Costa et al. [34], chlorogenic acid and caffeic acid represent important compounds in coffee by-products (such as silverskin) with potentially inhibitory effects on bacterial growth. Interestingly, different authors [35–37] have reported that such compounds may also stimulate the growth of lactic acid bacteria (e.g., *Lactobacillus collinoides*, *Lactobacillus brevis*, etc), thus likely explaining the massive presence of Lactobacillaceae or Leuconostocaceae in HC samples. It is noteworthy that lactic acid bacteria have been widely detected in different edible insect species with potential symbiotic or probiotic activities [38].

Among Actinobacteria detected in HC, PCR-DGGE allowed the presence of *Streptomyces*, *Corynebacterium*, and *Kytococcus* to be discovered. It is noteworthy that among *Streptomyces*, the species *Streptomyces griseus* and *Streptomyces coelicolor* is a chitinolytic bacterial species that can improve chitin digestibility in insect-based feeds [39, 40]. *Corynebacterium* has very recently been found in the gut of *H. illucens*, with functions in hydrogen metabolism, nitrogen cycling, and sulfur compound metabolism [41]. *Kytococcus* comprises Gram positive species



**Fig 2. Relative abundances (%) of bacterial community in the analyzed samples identified by analysis of V3-V4 region of the 16S rRNA gene through MiSeq Illumina.** C, coffee by-product; HC, *H. illucens* reared on coffee by-product; HCF, frass of *H. illucens* reared on coffee by-product; S, vegetable substrate; HS, *H. illucens* reared on vegetable substrate; HSF, frass of *H. illucens* reared on vegetable substrate; ZHC, gut content of zebrafish fed HC; ZHS, gut content of zebrafish fed HS.

<https://doi.org/10.1371/journal.pone.0225956.g002>

of environmental origin [42]; although found as minority species, to the authors’ knowledge this is the first report of *Kytococcus* in the microbiota of edible insects.

As already described for viable counts, reductions in Enterobacteriaceae relative abundance were observed in both HC and HCF. It is noteworthy that, regarding Enterobacteriaceae, HC was also characterized by lower abundance than in HS. It is likely that the presence of bioactive compounds from the coffee by-product, as chlorogenic or caffeic acids, could have modulated the growth of this bacterial family. Indeed, Zhang et al. [43] recently reported a decrease in the population of *Escherichia coli* in the colons of pigs fed diets supplemented with chlorogenic acid. Moreover, De Smet et al. [16] reported that the effectiveness of pathogen reduction by *H. illucens* can be affected by the composition of the substrate, thus supporting the results obtained in the present study.

Pooled HCF samples were also characterized by the presence of a rich microbiota mainly composed of Alteromonadales, Rhizobiales, Bacteroidetes, Rhodothermales, and Sphingobacteriaceae. To the authors’ knowledge the microbiota of *H. illucens* frass is actually poorly



**Table 3. Number of sequences analyzed (N reads), diversity richness (Chao 1), observed operational taxonomic units (OTUs), estimated sample coverage for 16S (Coverage) and diversity index (Shannon) for samples from the two rearing cycles.**

| Sample | N reads | Chao 1 | OTUs  | Coverage | Shannon |
|--------|---------|--------|-------|----------|---------|
| C      | 60,235  | 447    | 417   | 99.95    | 8.059   |
| HC     | 65,288  | 1,164  | 931   | 99.77    | 8.774   |
| HCF    | 60,530  | 614    | 565   | 99.92    | 8.361   |
| ZHC    | 44,728  | 347    | 346   | 99.97    | 7.587   |
| S      | 50,489  | 460    | 442   | 99.95    | 7.090   |
| HS     | 49,096  | 1,113  | 914   | 99.65    | 8.857   |
| HSF    | 63,101  | 1,430  | 1,151 | 99.79    | 9.079   |
| ZHS    | 60,348  | 856    | 761   | 99.82    | 8.322   |

C, coffee by-product; HC, *H. illucens* reared on coffee by-product; HCF, frass of *H. illucens* reared on coffee by-product; S, vegetable substrate; HS, *H. illucens* reared on vegetable substrate; HSF, frass of *H. illucens* reared on vegetable substrate; ZHC, gut content of zebrafish fed HC; ZHS, gut content of zebrafish fed HS.

<https://doi.org/10.1371/journal.pone.0225956.t003>

studied; hence, no further comparison is possible. It is noteworthy that, as reported by Mitchell and Hanks [44], frass can represent a vehicle for transmission of bacterial plant pathogens as well as a rich culturing media for saprophytic microorganisms. Moreover, the frass microbiota can synthesize chemical compounds that can be perceived by other insects, thus affecting the oviposition of some species [45]. Finally, insect frass used as fertilizer can also represent a source of safety risks for consumers due to the possible presence of human pathogens [46]; hence, a careful microbiological assessment of such a by-product must always be included in risk assessment.

In pooled HS samples the dominance of Enterobacteriaceae, Sphingobacteriaceae, Rhizobiales (among which closest relatives to *Aliihoeflea*) and Xanthomonadaceae (among which closest relatives to *Lysobacter*) was highlighted.

Sphingobacteriaceae have already been found in the microbiota of crambid butterflies, whereas the occurrence of species belonging to Xanthomonadaceae has already been reported in butterflies, beetles and crickets, thus confirming the association of these two bacterial families with the microbiota of insects [47, 48]. Finally, Rhizobiales have recently been detected in flea microbiota by Nziza et al. [49]. Members of Rhizobiales (e.g., *Bartonella*) can represent facultative intracellular parasites able to cause infection in human or wild and domestic animals [50].

In pooled HS samples, a low occurrence of Pseudomonadaceae was also observed; this bacterial family was in turn highly represented in the pooled HSF samples, together with Sphingobacteriaceae and Enterobacteriaceae.

The microbiota of insects can be strongly influenced by feeding substrates and vertical transmission from mother to offspring [27]. In the present study, the coffee by-product may have exerted a selective pressure on the microbiota of HC due to the presence of bioactive compounds such as chlorogenic acid and caffeine [51], thus perhaps explaining the differences in microbial composition between HC and HS. The results obtained in the present study are consistent with those very recently reported by Bruno et al. [15] which demonstrated that feed substrate can affect *H. illucens* midgut microbiota composition.

Zebrafish gut samples (ZHC and ZHS), obtained from the two feeding trials, were characterized by specific microbial patterns in which Actinobacteria and Alteromonadales were always detected, irrespective of the diet used.

Among Actinobacteria, PCR-DGGE allowed the presence of closest relatives to *Mobilicoccus*, *Pseudonocardia*, *Saccharopolyspora*, *Corynebacterium*, and *Phycoccus* to be discovered in pooled ZHC samples. Moreover, among Alteromonadales, closest relatives to *Shewanella* were

detected by PCR-DGGE in pooled ZHS samples, together with Actinobacteria genera (*Phycoccus* and *Microbacterium*).

Actinobacteria have already been detected in the gut microbiota of zebrafish, thus constituting one of the major microbial taxa involved in the function of the intestinal barrier of the fish and playing an essential role in the synthesis of antibiotics against fish pathogens [52]. Hence, the detection of Actinobacteria in both ZHC and ZHS pooled samples suggests healthy status of their gut systems.

High occurrence of Alteromonadales has been associated with good fish health in recirculating aquaculture systems [53]. Moreover, the presence of *Shewanella* detected in the present study is consistent with observations of Rimoldi et al. [54], who described this bacterial genus as part of the core microbiota detected in the gut of rainbow trout (*Oncorhynchus mykiss*) fed *H. illucens*-based diet.

Interestingly, the presence of Enterobacteriaceae in fish guts was higher in ZHS samples than in ZHC, thus suggesting a different contribution of the two feeding substrates (HC or HS) in the modulation of Enterobacteriaceae in the fish gut. Moreover, fish fed HC were characterized by the abundant presence of Actinomycetaceae, which, as recently reported by Nurul et al. [55], can establish a symbiotic association in fish with benefits for the bacteria and the host.

Finally, a higher relative abundance of Clostridia was detected in ZHC than in ZHS. This finding could likely be explained by the stimulating activity of chlorogenic acid and caffeine, contained in the zebrafish diet (HC), on this bacterial taxon [56].

## Conclusions

Overall, the data collected suggest that the microbiota of *H. illucens* may have been influenced by the feeding substrates. Similarly, zebrafish gut microbiota may have been influenced: i) by the microbiota of *H. illucens*-based feed and ii) eventually by insect-derived bioactive compounds contained in the experimental diets. Indeed, as recently reported by Bruni et al. [57], the fish gut microbial community is plastic and can be influenced by insect-based feed. Moreover, the results of both PCR-DGGE and Illumina sequencing showed the presence of a core microbiota constantly associated with the zebrafish gut, irrespective of the diet.

As highlighted by Huyben et al. [58], a high bacterial diversity can lead to improved health fish guts. Indeed, a rich microbial community could increase host resistance to pathogen invasion and intestinal infection. Hence, the low microbial richness exhibited by ZHC should be carefully considered when diets including HC are suggested.

Further research is needed to understand the interactions among *H. illucens* microbiota, *H. illucens* bioactive compounds (i.e., chitin, polyphenols, etc.) and the zebrafish gut microbiota.

Moreover, data collected on insect frass can be useful to widen the knowledge on such a poorly investigated insect rearing by-product for its further use.

## Compliance with ethical standards

All procedures involving animals were conducted in compliance with Italian legislation on experimental animals and were approved by the Ethics Committee of the Università Politecnica delle Marche (Ancona, Italy) (84/94-A). All efforts were made to minimize animal suffering by using an anesthetic (MS222; Sigma Aldrich, Saint Louis, Missouri, USA).

## Supporting information

**S1 Table. Raw data used for statistical analysis of viable counts.**  
(XLSX)

**S2 Table. Raw data of relative abundances (%) of bacterial community in the analyzed samples identified by analysis of V3-V4 region of the 16S rRNA gene by using MiSeq Illumina.**

(XLSX)

**S1 Fig. Raw original image of bacterial DGGE profiles of the DNA extracted directly from the analyzed samples and amplified with primers 338fGC and 518r.**

(PDF)

## Author Contributions

**Conceptualization:** Andrea Osimani, Paola Riolo, Ike Olivotto.

**Formal analysis:** Vesna Milanović, Andrea Roncolini, Sara Ruschioni, Nino Loreto, Elena Franciosi, Kieran Tuohy, Matteo Zarantoniello, Federica Cardinali, Cristiana Garofalo.

**Resources:** Nunzio Isidoro, Lucia Aquilanti, Francesca Clementi.

**Writing – original draft:** Andrea Osimani, Vesna Milanović, Paola Riolo, Ike Olivotto.

**Writing – review & editing:** Andrea Osimani, Vesna Milanović, Paola Riolo, Ike Olivotto.

## References

1. Oonincx DG, de Boer IJ. Environmental impact of the production of mealworms as a protein source for humans—a life cycle assessment. *PLoS One*. 2012; 7: e51145. <https://doi.org/10.1371/journal.pone.0051145> PMID: 23284661
2. Oonincx DG, van Itterbeek J, Heetkamp MJ, van den Brand H, van Loon JJ, van Huis A. An exploration on greenhouse gas and ammonia production by insect species suitable for animal or human consumption. *PLoS One*. 2010; 5: e14445. <https://doi.org/10.1371/journal.pone.0014445> PMID: 21206900
3. Rumpold BA, Schlüter OK. Nutritional composition and safety aspects of edible insects. *Molecul Nutr Food Res*. 2013; 57: 802–823.
4. van Huis A, Van Itterbeek J, Klunder H, Mertens E, Halloran A, Vantomme P. Edible insects: Future prospects for food and feed security. Rome: Food and Agriculture Organization of the United Nations. FAO Forestry Paper, FAO. 2013; 187 pp. Eds. <http://www.fao.org/docrep/018/i3253e/i3253e14.pdf>
5. EFSA Scientific Committee, 2015. Scientific opinion on a risk profile related to production and consumption of insects as food and feed. *EFSA J*. 13(10) (4257), 60. <http://dx.doi.org/10.2903/j.efsa.2015.4257>
6. ANSES (2015) Opinion of the French Agency for Food. In: Environmental and Occupational Health & Safety on “the Use of Insects as Food and Feed and the Review of Scientific Knowledge on the Health Risks Related to the Consumption of Insects” Maisons-Alfort, France.
7. Garofalo C, Milanović V, Cardinali F, Aquilanti L, Clementi F, Osimani A. Current knowledge on the microbiota of edible insects intended for human consumption: A state-of-the-art review. *Food Res Int*. 2019; 125: 108527 <https://doi.org/10.1016/j.foodres.2019.108527> PMID: 31554102
8. Commission Regulation (EU) 2017/893 of 24 May 2017 amending Annexes I and IV to Regulation (EC) No 999/2001 of the European Parliament and of the Council and Annexes X, XIV and XV to Commission Regulation (EU) No 142/2011 as regards the provisions on processed animal protein. *Official Journal of the European Union*, L 138/92-116. Available at: <https://eur-lex.europa.eu/legal-content/EN/TXT/PDF/?uri=CELEX:32017R0893&from=IT>
9. Cardinaletti G, Randazzo B, Messina M, Zarantoniello M, Giorgini E, Zimbelli A, et al. Effects of Graded Dietary Inclusion Level of Full-Fat *Hermetia illucens* Prepupae Meal in Practical Diets for Rainbow Trout (*Oncorhynchus mykiss*). *Animals*. 2019; 9, pii:E25. <https://doi.org/10.3390/ani9010025>
10. Cutrignelli MI, Messina M, Tulli F, Randazzo B, Olivotto I, Gasco L, et al. Evaluation of an insect meal of the Black Soldier Fly (*Hermetia illucens*) as soybean substitute: Intestinal morphometry, enzymatic and microbial activity in laying hens. *Res Vet Sci*. 2018; 117: 209–215. <https://doi.org/10.1016/j.rvsc.2017.12.020> PMID: 29304440
11. Diener S, Zurbrugg C, Tockner K. Conversion of organic material by black soldier fly larvae: Establishing optimal feeding rates. *Waste Manage Res*. 2009; 27: 603–610.

12. Vargas A, Randazzo B, Riolo P, Truzzi C, Gioacchini G, Giorgini E, et al. Rearing Zebrafish on Black Soldier Fly (*Hermetia illucens*): Biometric, Histological, Spectroscopic, Biochemical, and Molecular Implications. *Zebrafish*. 2018; 15: 404–419. <https://doi.org/10.1089/zeb.2017.1559> PMID: 29589997
13. Zarantoniello M, Randazzo B, Truzzi C, Giorgini E, Marcellucci C, Vargas-Abúndez JA, et al. A six-months study on Black Soldier Fly (*Hermetia illucens*) based diets in zebrafish. *Sci Rep*. 2019; 9: 8598. <https://doi.org/10.1038/s41598-019-45172-5> PMID: 31197206
14. Vogel H, Müller A, Heckel DG, Gutzeit H, Vilcinskas A. Nutritional immunology: Diversification and diet-dependent expression of antimicrobial peptides in the black soldier fly *Hermetia illucens*. *Dev Comp Immunol*. 2018; 78: 141–148. <https://doi.org/10.1016/j.dci.2017.09.008> PMID: 28966127
15. Bruno D, Bonelli M, De Filippis F, Di Lelio I, Tettamanti G, Casartelli M, et al. The Intestinal Microbiota of *Hermetia illucens* Larvae Is Affected by Diet and Shows a Diverse Composition in the Different Midgut Regions. *Appl Environ Microbiol*. 2019; 85, pii: e01864–18. <https://doi.org/10.1128/AEM.01864-18> PMID: 30504212
16. De Smet J, Wynants E, Cos P, Van Campenhout L. Microbial Community Dynamics during Rearing of Black Soldier Fly Larvae (*Hermetia illucens*) and Impact on Exploitation Potential. *Appl Environ Microbiol*. 2018; 84, pii: e02722–17. <https://doi.org/10.1128/AEM.02722-17> PMID: 29475866
17. Wynants E, Frooninckx L, Crauwels S, Verreth C, De Smet J, Sandrock C, et al. Assessing the Microbiota of Black Soldier Fly Larvae (*Hermetia illucens*) Reared on Organic Waste Streams on Four Different Locations at Laboratory and Large Scale. *Microb Ecol*. 2019; 77: 913–930. <https://doi.org/10.1007/s00248-018-1286-x> PMID: 30430196
18. Varotto Boccazzi I, Ottoboni M, Martin E, Comandatore F, Vallone L, Spranghers T, et al. A survey of the mycobiota associated with larvae of the black soldier fly (*Hermetia illucens*) reared for feed production. *PLoS One*. 2017; 12: e0182533. <https://doi.org/10.1371/journal.pone.0182533> PMID: 28771577
19. Zarantoniello M, Bruni L, Randazzo B, Vargas A, Gioacchini G, Truzzi C, et al. Partial Dietary Inclusion of *Hermetia illucens* (Black Soldier Fly) Full-Fat Prepupae in Zebrafish Feed: Biometric, Histological, Biochemical, and Molecular Implications. *Zebrafish*. 2018; 15: 519–532. <https://doi.org/10.1089/zeb.2018.1596> PMID: 29912648
20. Osimani A, Cardinali F, Aquilanti L, Garofalo C, Roncolini A, Milanović V, et al. Occurrence of transferable antibiotic resistances in commercialized ready-to-eat mealworms (*Tenebrio molitor* L.). *Int J Food Microbiol*. 2017; 263: 38–46. <https://doi.org/10.1016/j.ijfoodmicro.2017.10.009> PMID: 29028569
21. Garofalo C, Silvestri G, Aquilanti L, Clementi F. PCR-DGGE analysis of lactic acid bacteria and yeast dynamics during the production processes of three varieties of Panettone. *J Appl Microbiol*. 2008; 105: 243–254. <https://doi.org/10.1111/j.1365-2672.2008.03768.x> PMID: 18312562
22. Milanović V, Osimani A, Garofalo C, De Filippis F, Ercolini D, Cardinali F, et al. Profiling white wine seed vinegar bacterial diversity through viable counting, metagenomic sequencing and PCR-DGGE. *Int J Food Microbiol*. 2018; 286: 66–74. <https://doi.org/10.1016/j.ijfoodmicro.2018.07.022> PMID: 30048915
23. Ampe F, Ben Omar N, Moizan C, Wachter C, Guyot JP. Polyphasic study of the spatial distribution of microorganisms in Mexican pozol, a fermented maize dough, demonstrates the need for cultivation-independent methods to investigate traditional fermentations. *Appl Environ Microbiol*. 1999; 65: 5464–5473. PMID: 10584005
24. Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. Basic local alignment search tool. *J Mol Biol*. 1990; 215: 403–410. [https://doi.org/10.1016/S0022-2836\(05\)80360-2](https://doi.org/10.1016/S0022-2836(05)80360-2) PMID: 2231712
25. Baker GC, Smith JJ, Cowan DA. Review and re-analysis of domain-specific 16S primers. *J Microbiol Method*. 2003; 55: 541–555.
26. Claesson MJ, Wang Q, O'Sullivan O, Greene-Diniz R, Cole JR, Ross RP, et al. Comparison of two next-generation sequencing technologies for resolving highly complex microbiota composition using tandem variable 16S rRNA gene regions. *Nucleic Acids Res*. 2010; 38: e200–e200. <https://doi.org/10.1093/nar/gkq873> PMID: 20880993
27. Osimani A, Milanović V, Cardinali F, Garofalo C, Clementi F, Pasquini M, et al. The bacterial biota of laboratory-reared edible mealworms (*Tenebrio molitor* L.): from feed to frass. *Int J Food Microbiol*. 2018; 272: 49–60. <https://doi.org/10.1016/j.ijfoodmicro.2018.03.001> PMID: 29525619
28. Caporaso JG, Kuczynski J, Stombaugh J, Bittinger K, Bushma FD, Costello E.K., et al. QIIME allows analysis of highthroughput community sequencing data. *Nat Methods*. 2010; 7: 335–336. <https://doi.org/10.1038/nmeth.f.303> PMID: 20383131
29. Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJA, Holmes S.P. DADA2: high-resolution sample inference from Illumina amplicon data. *Nat Methods*. 2016; 13: 581. <https://doi.org/10.1038/nmeth.3869> PMID: 27214047
30. Katoh K, Standley DM. MAFFT multiple sequence alignment software version 7: improvements in performance and usability. *Mol Biol Evol*. 2013; 30: 772–780. <https://doi.org/10.1093/molbev/mst010> PMID: 23329690

31. Price MN, Dehal PS, Arkin AP. FastTree: computing large minimum evolution trees with profiles instead of a distance matrix. *Mol Biol Evol.* 2009; 26: 1641–165030. <https://doi.org/10.1093/molbev/msp077> PMID: 19377059
32. Vazquez-Baeza Y, Pirrung M, Gonzalez A, Knight R. EMPERor: a tool for visualizing high-throughput microbial community data. *GigaScience.* 2013; 2: 16. <https://doi.org/10.1186/2047-217X-2-16> PMID: 24280061
33. Lalander C, Diener S, Magri ME, Zurbrügg C, Lindström A, Vinnerås B. Faecal sludge management with the larvae of the black soldier fly (*Hermetia illucens*)—From a hygiene aspect. *Sci Total Environ.* 2013; 458–460: 312–318. <https://doi.org/10.1016/j.scitotenv.2013.04.033> PMID: 23669577
34. Costa ASG, Alves RC, Vinha AF, Costa E, Costa S.G, Nunes MA, et al. Nutritional, chemical and anti-oxidant/pro-oxidant profiles of silverskin, a coffee roasting by-product. *Food Chem.* 2018; 267: 28–35. <https://doi.org/10.1016/j.foodchem.2017.03.106> PMID: 29934169
35. Filannino P, Bai YP, Di Cagno R, Gobetti M, Gänzle MG. Metabolism of phenolic compounds by *Lactobacillus* spp. during fermentation of cherry juice and broccoli puree. *Food Microbiol.* 2015; 46: 272–279. <https://doi.org/10.1016/j.fm.2014.08.018> PMID: 25475296
36. Fritsch C, Heinrich V, Vogel RF, Toelstede S. Phenolic acid degradation potential and growth behavior of lactic acid bacteria in sunflower substrates. *Food Microbiol.* 2016; 57: 178–86. <https://doi.org/10.1016/j.fm.2016.03.003> PMID: 27052717
37. Parkar SG, Stevenson DE, Skinner MA. The potential influence of fruit polyphenols on colonic microflora and human gut health. *Int J Food Microbiol.* 2008; 124: 295–298. <https://doi.org/10.1016/j.ijfoodmicro.2008.03.017> PMID: 18456359
38. Osimani A, Milanović V, Garofalo C, Cardinali F, Roncolini A, Sabbatini R, et al. Revealing the microbiota of marketed edible insects through PCR-DGGE, metagenomic sequencing and real-time PCR. *Int J Food Microbiol.* 2018; 276: 54–62. <https://doi.org/10.1016/j.ijfoodmicro.2018.04.013> PMID: 29665523
39. Berini F, Katz C, Gruzdev N, Casartelli M, Tettamanti G, Marinelli F. Microbial and viral chitinases: Attractive biopesticides for integrated pest management. *Biotechnol Adv.* 2018; 36: 818–838. <https://doi.org/10.1016/j.biotechadv.2018.01.002> PMID: 29305895
40. Kroeckel S, Harjes AGE, Roth I, Katz H, Wuertz S, Susenbeth A, et al. When a turbot catches a fly: Evaluation of a pre-pupae meal of the Black Soldier Fly (*Hermetia illucens*) as fish meal substitute—Growth performance and chitin degradation in juvenile turbot (*Psetta maxima*). *Aquaculture.* 2012; 364–365: 345–352.
41. Jiang CL, Ji W.Z, Tao XH, Zhang Q, Zhu J, Feng SY, et al. Black soldier fly larvae (*Hermetia illucens*) strengthen the metabolic function of food waste biodegradation by gut microbiome. *Microb Biotechnol.* 2019; 12: 528–543. <https://doi.org/10.1111/1751-7915.13393> PMID: 30884189
42. Kämpfer P, Martin K, Schäfer J, Schumann P. *Kytococcus aerolatus* sp. nov., isolated from indoor air in a room colonized with moulds. *Syst Appl Microbiol.* 2009; 32: 301–305. <https://doi.org/10.1016/j.syapm.2009.05.004> PMID: 19541443
43. Zhang Y, Wang Y, Chen D, Yu B, Zheng P, Mao X, et al. Dietary chlorogenic acid supplementation affects gut morphology, antioxidant capacity and intestinal selected bacterial populations in weaned piglets. *Food Funct.* 2018; 9: 4968–4978. <https://doi.org/10.1039/c8fo01126e> PMID: 30183786
44. Mitchell RF, Hanks LM. Insect frass as a pathway for transmission of bacterial wilt of cucurbits. *Environ Entomol.* 2009; 38: 395–403. <https://doi.org/10.1603/022.038.0212> PMID: 19389288
45. Anbutu H, Togashi K. Oviposition deterrence associated with larval frass of the Japanese pine sawyer, *Monoctonus alternatus* (Coleoptera: Cerambycidae). *J Insect Physiol.* 2002; 48: 459–465. [https://doi.org/10.1016/s0022-1910\(02\)00067-7](https://doi.org/10.1016/s0022-1910(02)00067-7) PMID: 12770095
46. Yang CC, Kim MS, Millner P, Chao K, Cho BK, Mo C, et al. Development of multispectral imaging algorithm for detection of frass on mature red tomatoes. *Postharvest Biol Technol.* 2014; 93: 1–8.
47. Grabowski NT, Klein G. Bacteria encountered in raw insect, spider, scorpion, and centipede taxa including edible species, and their significance from the food hygiene point of view. *Trends Food Sci Technol.* 2017; 63: 80–90.
48. Milanović V, Cardinali F, Aquilanti L, Garofalo C, Roncolini A, Sabbatini R, et al. A Glimpse into the Microbiota of Marketed Ready-to-Eat Crickets (*Acheta domesticus*). *Indian J Microbiol.* 2019; <https://doi.org/10.1007/s12088-019-00817-x>
49. Nziza J, Tumushime JC, Cranfield M, Ntwari AE, Modry D, Mudakikwa A, et al. Fleas from domestic dogs and rodents in Rwanda carry *Rickettsia asembonensis* and *Bartonella tribocorum*. *Med Vet Entomol.* 2019; 33: 177–184. <https://doi.org/10.1111/mve.12340> PMID: 30390316

50. Breitschwerdt EB, Maggi RG, Chomel BB, Lappin MR. Bartonellosis: an emerging infectious disease of zoonotic importance to animals and human beings. *J Vet Emerg Crit Care (San Antonio)*. 2010; 20: 8–30.
51. Fernandez-Gomez B, Lezama A, Amigo-Benavent M, Ullate M, Herrer M., Martín MA, et al. Insights on the health benefits of the bioactive compounds of coffee silverskin extract. *J Funct Foods*. 2016; 25: 197–207.
52. Qiao R, Deng Y, Zhang S, Wolosker MB, Zhu Q, Ren H, et al. Accumulation of different shapes of microplastics initiates intestinal injury and gut microbiota dysbiosis in the gut of zebrafish. *Chemosphere*. 2019; 236: 124334. <https://doi.org/10.1016/j.chemosphere.2019.07.065> PMID: 31310986
53. Xue S, Xu W, Wei J, Sun J. Impact of environmental bacterial communities on fish health in marine recirculating aquaculture systems. *Vet Microbiol*. 2017; 203: 34–39. <https://doi.org/10.1016/j.vetmic.2017.01.034> PMID: 28619164
54. Rimoldi S, Gini E, Iannini F, Gasco L, Terova G. The Effects of Dietary Insect Meal from *Hermetia illucens* Prepupae on Autochthonous Gut Microbiota of Rainbow Trout (*Oncorhynchus mykiss*). *Animals (Basel)*. 2019; 9, pii: E143.
55. Nurul ANA, Muhammad DD, Okomoda VT, Nur AAB. 16S rRNA-Based metagenomic analysis of microbial communities associated with wild *Labroides dimidiatus* from Karah Island, Terengganu, Malaysia. *Biotechnol Rep (Amst.)*. 2019; 21: e00303.
56. Mills CE, Tzounis X, Oruna-Concha MJ, Mottram DS, Gibson GR, Spencer JP. In vitro colonic metabolism of coffee and chlorogenic acid results in selective changes in human faecal microbiota growth. *Br J Nutr*. 2015; 113: 1220–1227. <https://doi.org/10.1017/S0007114514003948> PMID: 25809126
57. Bruni L, Pastorelli R, Viti C, Gasco L, Parisi G. Characterisation of the intestinal microbial communities of rainbow trout (*Oncorhynchus mykiss*) fed with *Hermetia illucens* (black soldier fly) partially defatted larva meal as partial dietary protein source. *Aquaculture*. 2018; 487: 56–63.
58. Huyben D, Vidaković A, Werner Hallgren S, Langeland M. High-throughput sequencing of gut microbiota in rainbow trout (*Oncorhynchus mykiss*) fed larval and pre-pupae stages of black soldier fly (*Hermetia illucens*), *Aquaculture*. 2019; 500: 485–491.