



Published in final edited form as:

Curr Treatm Opt Rheumatol. 2019 June 15; 5: 168–178. doi:10.1007/s40674-019-00123-z.

Self-Efficacy and the Role of Non-Pharmacologic Treatment Strategies to Improve Pain and Affect in Arthritis

Dana DiRenzo, MD, MHS¹, Patrick Finan, PhD¹

¹The Johns Hopkins University, Baltimore, MD, USA

Abstract

Purpose of Review: There is increasing evidence that adjunctive, non-pharmacologic treatment programs are beneficial in the management of arthritis when added to traditional disease-modifying medications. This review focuses on non-pharmacologic management strategies that impact pain and affect, with a focus on self-efficacy, for those with osteoarthritis (OA) and rheumatoid arthritis (RA).

Recent Findings: We reviewed both office-based and internet-based self-management strategies, mindfulness based interventions (MBIs), and cognitive behavioral therapies (CBTs) for patients with arthritis. These behavioral strategies have shown to improve pain, mood disturbance, and physical function in those with both osteoarthritis and rheumatoid arthritis. Improvements in self-efficacy and coping capacity are associated with improvements in patient-reported outcomes (PROs) related to pain and functioning.

Summary: Self-management programs, MBIs, and CBTs are more effective at improving pain and mood disturbance compared to usual care for patients with arthritis although high quality randomized controlled trials are lacking. Non-pharmacologic management programs are increasingly available via the internet and mobile applications.

Keywords

Arthritis; Self-Efficacy; Anxiety; Depression; Pain; Mindfulness

Introduction.

Doctor-diagnosed arthritis, according to the National Health Interview Survey (2013–2015), is present in a large proportion of American adults (54.5 million individuals) of which 43.5% have arthritis-attributable activity limitations.(1) Osteoarthritis (OA) is the most

Corresponding Author: Dana DiRenzo, MD. 5200 Eastern Avenue, MFL Building, Center Tower, Suite 4100, Baltimore, MD, 21224. ddirenz1@jhmi.edu. Phone: 410-550-1925.

Publisher's Disclaimer: This Author Accepted Manuscript is a PDF file of a an unedited peer-reviewed manuscript that has been accepted for publication but has not been copyedited or corrected. The official version of record that is published in the journal is kept up to date and so may therefore differ from this version.

Conflicts of Interest:

The authors declare that there is no conflict of interest regarding the publication of this paper.

Human and Animal Rights and Informed Consent

This article does not contain any studies with human or animal subjects performed by any of the authors.

common of the arthritides, characterized by degenerative changes of the joints and affected by aging and obesity.(2) Rheumatoid Arthritis (RA) is the most common inflammatory arthritis which is characterized by joint pain, stiffness, and articular damage; treatment most likely requires immunosuppression.(3,4)

Anxiety and depression are prevalent in those with both OA and RA.(5,6) A large meta-analysis of those with OA (n=15,855) noted that approximately 20% of individuals had anxiety and depression, respectively.(7) For those with RA, the levels of anxiety and depression appear to be similar and present in approximately 20% of individuals.(8) Unfortunately, arthritis coupled with anxiety and depression has considerable impacts on physical function and quality of life.(9–11) In RA, patients with anxiety and depression at baseline and 2 year follow-up had more disability and disease activity compared to those without anxiety and depression.(12,13) Likewise, OA patients with anxiety and depression had worse physical function and pain severity compared to those without mood disturbance.(14,15)

The detrimental impacts of anxiety on functionality and patient perception of disease activity may be exacerbated by low self-efficacy, or the inner belief in one's ability to succeed in specific situations and tasks.(16,17) In RA patients, anxiety level has a strong inverse correlation with the degree of self-efficacy(18–20) and low self-efficacy at baseline is a strong predictor for declining health outcomes up to two-years later.(21) In patients with chronic pain, high degrees of self-efficacy had enhancing effects on the perception of quality of life and general health.(22) Self-efficacy was found to fully mediate pain catastrophizing and physical disability in obese individuals with osteoarthritis.(23)

Importantly, arthritis patients with high levels of self-efficacy report lower levels of pain, fatigue, physical disability, and psychological distress.(20,24–26) Self-efficacy appears closely related to coping capacity and in arthritis patients, this may protect against pain-related anxiety.(27) A recent systematic review of the role of self-efficacy in patients with RA similarly noted an association between high self-efficacy and positive affect, physical function, and ability to participate in social roles and activities.(28)

In recent years, several types of non-pharmacologic treatment strategies to improve self-efficacy have emerged including 1) self-management with a focus on exercise, 2) mindfulness based interventions (MBIs), and 3) cognitive behavioral therapies (CBTs) (Table 1). Across modalities, use of internet and smartphone-based platforms for intervention delivery is on the rise.

Current Treatment Options.

Self-Management Strategies.

Non-pharmacologic treatment strategies that positively impact mood disturbance and pain via improvements in self-efficacy have been explored for several decades.(24) Beginning in the late 1980s, arthritis self-management strategies such as the Arthritis Self-Management Program (ASMP) and a tailored print intervention (SMART) were found to improve self-efficacy and positively impact health outcomes including pain, anxiety, and depression.(29–

31) The ASMP was a 6-week, community-based program that focused on education, improving function through cognitive restructuring techniques, and problem-solving for health-related problems. The mailed intervention, SMART, included a tailored action plan by a physician, two arthritis self-help books, and relaxation tape. Compared to the ASMP, SMART had greater improvements in disability and self-efficacy after 1-year.(31)

The popularity of self-management strategies to improve self-efficacy grew throughout the early 2000s into present day and now additionally has focused on exercise. Self-management strategies may mediate the association of self-efficacy with physical activity.(32) A recent qualitative study evaluating patients with physician-diagnosed arthritis (n=197), found that the exercise, self-management program, First Step to Active Health™ (FSAH), was well-received by participants and improved motivation, accountability, and quality of life.(33) Similarly, the PLE²NO program, a self-management and exercise group for older individuals with knee osteoarthritis, found improvements in physical function, pain, and symptoms compared to an education group.(34)

For patients with RA, a 5-week targeted motivation and self-regulation program was found to increase physical activity, self-efficacy, and autonomous motivation compared to group based education.(35) For those in the self-regulation group, the percentage of RA patients meeting the daily physical activity recommendation (30 mins x 5 days per week) was 67% compared to 23% of controls (p<0.001). Self-management strategies tailored to the individual were also noted to improve pain, symptoms, affect (Arthritis Impact Measurement Scale), and self-efficacy (Arthritis Self-Efficacy Scale) for those with RA compared to a generalized education program.(36)

In more recent years, self-management programs have shifted to include internet and smartphone-based delivery systems.(37) A recent observational study of 200 patients with chronic pain, including those with arthritis, found moderate improvements in pain, anxiety, depression, and quality life after participating in an internet-based self-management program for six months.(38) The self-management program consisted of visiting two websites for 15 minutes per day and perusing information on exercise, nutrition, mindfulness, relaxation, and educational postings. Similarly, a qualitative study of a mobile internet service to increase physical activity in patients with RA found the internet-based program improved goal-setting, self-regulation, and self-efficacy after six-weeks of use.(39) Social media may be used to effectively disseminate self-management strategies to patients with arthritis via healthcare professionals.(40)

Mindfulness Based Interventions (MBIs).

Mindfulness is a mental state that is defined as non-judgmental awareness of the present moment and involves self-regulation of attention guided by curiosity, openness, and acceptance.(41) There are several traditional office-based mindfulness programs including Mindfulness Based Stress Reduction (MBSR) and Mindfulness Based Cognitive Therapy (MBCT).(42,43) MBSR and MBCT typically follow an 8-week format; participants meet once weekly for approximately 2.5 hours.(41) Other MBIs that have shown benefit in health-related quality of life (HRQOL) in those with RA include the Vitality Training Programme

(VTP), Mindful Awareness and Acceptance Therapy (MAAT), and Internal Family Systems (IFS).(44)

MBSR is the most well-known of the MBIs that focuses on learning how to mindfully respond to bodily sensations and emotions through guided meditative practices including sitting meditation, body scans, stretching, and gentle Hatha yoga.(42) The MBSR program strongly encourages daily home practice to solidify techniques learned during the weekly sessions. Patients receive printed materials and audiovisuals to aid their home practice; a portion of the MBSR clinic session is spent reviewing progress.(45) MBSR is an effective non-pharmacologic treatment strategy for anxiety and depression in the general population and provide benefit to those with RA through improvements in self-efficacy via emotion regulation and positive reappraisal.(41,46–51) Similarly, RA patients who completed the VTP and MAAT programs, which focus on symptom and emotion management, noted significant improvements in self-efficacy (pain, symptoms) after course completion.(52,53) A trial evaluating the effectiveness of MBSR for those with OA is currently ongoing (NCT03527849). While no trials have specifically evaluated the efficacy of MBIs for those with OA, patients who had higher total mindfulness scores (according to Five Facet Mindfulness Questionnaire) after a Tai Chi program were 38% more likely to meet Osteoarthritis Research Society International response criteria than those treated with physical therapy.(54)

Internet and smartphone mindfulness interventions are also gaining traction for use in research and clinical practice. Preliminary evidence demonstrates improvements in psychological well-being, perceived stress and pain severity for those with chronic disease; lasting effects have not been well-established.(55,56) Internet and smartphone mindfulness interventions characteristically are much shorter in length per session and can be utilized anywhere by the participant.(57)

Cognitive Behavioral Therapy.

Cognitive Behavioral Therapy (CBT) is a well-established treatment for anxiety and depression in patients with chronic pain and has been implemented in multi-modal management strategies.(58,59) In patients with chronic pain who were treated with CBT, improvements in self-efficacy for managing symptoms were noted.(58,60) A randomized controlled trial (RCT) for Pain Coping Skills Training (PCST), a form of CBT implemented in those with chronic hip and knee OA (n=256), found that those with moderate to high expectations of benefit, moderate to high osteoarthritis disease severity, advanced age, and who were highly educated benefited the most from the intervention.(61) Likewise, in patients with early RA, defined as disease duration less than 8 years, tailored CBT programs have been effective in improving fatigue, depression, helplessness, coping with stress, and medication compliance.(62)

A recent meta-analysis found improvements in physical function, pain intensity, and depression for patients with chronic pain treated with CBT, but no evidence of a significant difference compared to those treated with MBSR. (63) In a RCT that compared CBT, relaxation response training, and arthritis education in those with RA, significant improvements in symptoms, but not in pain, were noted at 12 month follow up.(64) There

was not a significant change in levels of anxiety and depression as measured by the Rand Mental Health Inventory (MHI) in the CBT group.

Like MBIs, CBT programs implemented via internet-based platforms are being utilized at increased frequency with beneficial effects on anxiety and depression.(65) More specifically, a RCT of RA patients (n=133) found significant improvements in depressed mood, anxiety, and fatigue in those who completed an internet-based CBT program with an average course duration of 17 weeks.(66) For older adults with knee OA, internet-based CBT was superior to usual care in improving depression (measured via Patient Health Questionnaire-9 (PHQ-9)).(67) Patients with hip and knee OA treated with an internet-based PCST noted improvements in pain, self-efficacy, pain-related anxiety, pain-related interference with physical function, and negative and positive affect.(68,69) Similarly, an internet-delivered multimodal program including exercise, education, and PCST for patients with knee OA, found significant improvements in pain at 3 months compared to internet education alone. (70) Participants who were employed and had higher baseline levels of self-efficacy were noted to have the greatest improvements.

Healthcare Utilization.

The economic burden of arthritis (medical expenditures and earning losses) in the US is considerable, totaling \$303.5 billion in 2013.(71) The average OA patient has a mean (SD) incremental total expenditure of \$2046 (\$862) and the average RA patient prescribed both traditional and biologic disease modifying agents has a mean (SD) annual expenditure of \$23,185 (\$1,295).(72,73)

For patients with rheumatic disease, pain reduction and improved physical function via strategies that strengthen self-efficacy and lessen mood disturbance may improve healthcare expenditures in high risk individuals. Generally, patients with chronic disease using self-management mobile/internet based interventions felt better cared for and perceived a more active role in their disease management compared to those treated traditionally.(74)

Improved emotion regulation and mindfulness of bodily symptoms may serve to decrease healthcare utilization over time (telephone encounters, EMR secured messaging, urgent appointments, and emergency room visits).(75) In a small study of patients with chronic disease (n=38; defined as chronic pain, chronic illness, or stress-related problems), an office-based mindfulness program was found to reduce primary care visits, specialty care visits, ER visits, and hospitalizations at one year.(75) Furthermore, a Canadian population based study of patients who had high healthcare utilization found that those treated with MBCT versus other non-mindfulness group-based therapies had reductions in utilization at one year.(76) Similar stress-reducing strategies, such as Relaxation Response and Resiliency training (3RP), have shown a large reduction in health care utilization.(77) The mechanism by which MBIs compared to other group based therapies facilitate the reduction in healthcare utilization is unknown.

Personal Insights and Observations.

We believe that using a multi-disciplinary approach that seeks to promote self-efficacy, mental health, and wellbeing are necessary to successfully treat the patient with arthritis. Mood disturbance is intimately tied to pain and patient perception of disease activity and if left untreated, may lead to worse clinical outcomes and increased healthcare expenditures. (6,78–80) Conversely, patients who are able to generate and maintain positive emotions tend to evidence improved pain-related outcomes.(81) Our clinical experience has shown that patients who lack psychological resources (e.g., self-efficacy) and demonstrate a hypervigilance to pain and disease activity often have poor long-term outcomes and may endure escalation of potent disease-modifying agents in RA and analgesics in OA in an effort to control symptoms. A similar, yet opposite clinical challenge may be found in the RA patient who avoids use of biologic disease-modifying agents due to fear of the immediate and long-term adverse effects.

While office-based interventions offer the opportunity for a patient-provider connection to contribute to more enduring and robust treatment effects, they come with the cost of increased patient burden in time and money. Internet-based platforms may improve program recruitment and adherence, and are typically low cost. Currently, there is a paucity of clinical trials evaluating whether treatment of anxiety and depression, especially early in the disease course, improves long-term clinical and patient-reported outcomes.(82) This is especially important in RA in which highly expensive pharmaceuticals are commonly utilized. Multi-disciplinary approaches addressing both the mental and physical needs of the patient should be considered in order to optimize clinical outcomes and reduce healthcare spending.

Summary.

Despite advances in the pharmacologic and interventional management of arthritis, there still remains a large role for adjunctive, behavioral strategies to improve self-efficacy, physical function, and HRQOL.(83) Self-management strategies, MBIs, and CBTs are evidenced-based treatments that may be implemented in those with arthritis and are offered in a variety of formats including office and internet-based settings.

Funding source:

Dr. DiRenzo is supported by the National Institute of Arthritis and Musculoskeletal and Skin Diseases of the National Institutes of Health under Award Number T32AR048522.

Dr. Finan is supported by the National Institute on Drug Abuse of the National Institutes of Health under Award Number K23DA035915.

References:

Recently published papers of particular interest have been highlighted as:

- Of importance
- Of major importance

1. Barbour KE, Helmick CG, Boring M, Brady TJ. Vital Signs: Prevalence of Doctor-Diagnosed Arthritis and Arthritis-Attributable Activity Limitation — United States, 2013–2015. *MMWR Morb Mortal Wkly Rep.* 2017 3 10;66(9):246–53. [PubMed: 28278145] **Most recent evaluation of arthritis-related disability in the US.
2. Martel-Pelletier J, Barr AJ, Cicuttini FM, Conaghan PG, Cooper C, Goldring MB, et al. Osteoarthritis. *Nat Rev Dis Prim.* 2016 10 13;2:16072. [PubMed: 27734845]
3. Singh JA, Saag KG, Louis Bridges SJ, Akl EA, Bannuru RR, Sullivan MC, et al. 2015 American College of Rheumatology Guideline for the Treatment of Rheumatoid Arthritis. *Arthritis Care Res (Hoboken).*
4. Hunter TM, Boytsov NN, Zhang X, Schroeder K, Michaud K, Araujo AB. Prevalence of rheumatoid arthritis in the United States adult population in healthcare claims databases, 2004–2014. *Rheumatol Int.* 2017 9 28;37(9):1551–7. [PubMed: 28455559]
5. Matcham F, Ali S, Irving K, Hotopf M, Chalder T. Are depression and anxiety associated with disease activity in rheumatoid arthritis? A prospective study. *BMC Musculoskelet Disord.* 2016 12 11;17(1):155. [PubMed: 27068100]
6. Sharma A, Kudesia P, Shi Q, Gandhi R. Anxiety and depression in patients with osteoarthritis: impact and management challenges. *Open Access Rheumatol Res Rev.* 2016 10;Volume 8:103–13.
7. Stubbs B, Aluko Y, Myint PK, Smith TO. Prevalence of depressive symptoms and anxiety in osteoarthritis: a systematic review and meta-analysis. *Age Ageing.* 2016 3;45(2):228–35. [PubMed: 26795974]
8. Covic T, Cumming SR, Pallant JF, Manolios N, Emery P, Conaghan PG, et al. Depression and anxiety in patients with rheumatoid arthritis: prevalence rates based on a comparison of the Depression, Anxiety and Stress Scale (DASS) and the hospital, Anxiety and Depression Scale (HADS). *BMC Psychiatry.* 2012 1 24;12:6. [PubMed: 22269280]
9. Smolen JS, Aletaha D, McInnes IB. Rheumatoid arthritis. *Lancet.* 2016 10 22;388(10055):2023–38. [PubMed: 27156434]
10. Zhang L, Xia Y, Zhang Q, Fu T, Yin R, Guo G, et al. The correlations of socioeconomic status, disease activity, quality of life, and depression/anxiety in Chinese patients with rheumatoid arthritis. *Psychol Health Med.* 2017 1 2;22(1):28–36. [PubMed: 27367831]
11. Dieppe P, Cushnaghan J, Tucker M, Browning S, Shepstone L. The Bristol ‘OA500 study’: progression and impact of the disease after 8 years. *Osteoarthr Cartil.* 2000 3;8(2):63–8. [PubMed: 10772234]
12. Kekow J, Moots R, Khandker R, Melin J, Freundlich B, Singh A. Improvements in patient-reported outcomes, symptoms of depression and anxiety, and their association with clinical remission among patients with moderate-to-severe active early rheumatoid arthritis. *Rheumatology.* 2011 2 1;50(2):401–9. [PubMed: 21059675]
13. Kronisch C, McLernon DJ, Dale J, Paterson C, Ralston SH, Reid DM, et al. Brief Report: Predicting Functional Disability: One-Year Results From the Scottish Early Rheumatoid Arthritis Inception Cohort. *Arthritis Rheumatol.* 2016 7;68(7):1596–602. [PubMed: 26866516] *Recent cohort study evaluating predictors of functional disability in RA
14. Scopaz KA, Piva SR, Wisniewski S, Fitzgerald GK. Relationships of Fear, Anxiety, and Depression With Physical Function in Patients With Knee Osteoarthritis. *Arch Phys Med Rehabil.* 2009 11;90(11):1866–73. [PubMed: 19887210]
15. Rathbun AM, Stuart EA, Shardell M, Yau MS, Baumgarten M, Hochberg MC. Dynamic Effects of Depressive Symptoms on Osteoarthritis Knee Pain. *Arthritis Care Res (Hoboken).* 2018 1;70(1): 80–8. [PubMed: 28320048]
16. Finney Rutten LJ, Hesse BW, St Sauver JL, Wilson P, Chawla N, Hartigan DB, et al. Health Self-Efficacy Among Populations with Multiple Chronic Conditions: the Value of Patient-Centered Communication. *Adv Ther.* 2016 8;33(8):1440–51. [PubMed: 27357639]
17. Meredith P, Strong J, Feeney JA. Adult attachment, anxiety, and pain self-efficacy as predictors of pain intensity and disability. *Pain.* 2006 7 1;123(1–2):146–54. [PubMed: 16644132]
18. Matcham F, Norton S, Scott DL, Steer S, Hotopf M. Symptoms of depression and anxiety predict treatment response and long-term physical health outcomes in rheumatoid arthritis: secondary analysis of a randomized controlled trial. *Rheumatology.* 2016 2;55(2):268–78. [PubMed: 26795974]

26350486] *Secondary analysis of RCT determining impact of mood disturbance on clinical outcomes

19. Treharne GJ, Lyons AC, Booth DA, Kitas GD. Psychological well-being across 1 year with rheumatoid arthritis: Coping resources as buffers of perceived stress. *Br J Health Psychol.* 2007 9;12(3):323–45. [PubMed: 17640450]
20. Liu L, Xu N, Wang L. Moderating role of self-efficacy on the associations of social support with depressive and anxiety symptoms in Chinese patients with rheumatoid arthritis. *Neuropsychiatr Dis Treat.* 2017;13:2141–50. [PubMed: 28860771]
21. Brekke M, Hjortdahl P, Kvien TK. Self-efficacy and health status in rheumatoid arthritis: a two-year longitudinal observational study. *Rheumatology.* 2001 4 1;40(4):387–92. [PubMed: 11312375]
22. Börso B, Gerdle B, Peolsson M. Impact of the interaction between self-efficacy, symptoms and catastrophising on disability, quality of life and health in with chronic pain patients. *Disabil Rehabil.* 2010 1;32(17):1387–96. [PubMed: 20513205]
23. Shelby RA, Somers TJ, Keefe FJ, Pells JJ, Dixon KE, Blumenthal JA. Domain specific self-efficacy mediates the impact of pain catastrophizing on pain and disability in overweight and obese osteoarthritis patients. *J Pain.* 2008 10;9(10):912–9. [PubMed: 18602871]
24. Somers TJ, Wren AA, Shelby RA. The Context of Pain in Arthritis: Self-efficacy for Managing Pain and Other Symptoms. *Curr Pain Headache Rep.* 2012 12 2;16(6):502–8. [PubMed: 23054977]
25. Somers TJ, Shelby RA, Keefe FJ, Godiwala N, Lumley MA, Mosley-Williams A, et al. Disease severity and domain-specific arthritis self-efficacy: Relationships to pain and functioning in patients with rheumatoid arthritis. *Arthritis Care Res (Hoboken).* 2010 6;62(6):848–56. [PubMed: 20535796]
26. Primdahl J, Wagner L, Hørslev-Petersen K. Self-Efficacy as an Outcome Measure and its Association with Physical Disease-Related Variables in Persons with Rheumatoid Arthritis: A Literature Review. *Musculoskeletal Care.* 2011 6;9(3):n/a–n/a.
27. Strahl C, Kleinknecht RA, Dinnel DL. The role of pain anxiety, coping, and pain self-efficacy in rheumatoid arthritis patient functioning. *Behav Res Ther.* 2000 9;38(9):863–73. [PubMed: 10957821]
28. Martinez-Calderon J, Meeus M, Struyf F, Luque-Suarez A. The role of self-efficacy in pain intensity, function, psychological factors, health behaviors, and quality of life in people with rheumatoid arthritis: A systematic review. *Physiother Theory Pract.* 2018 6 6;1–17. *Evaluation of the role of self-efficacy in RA
29. Lorig KR, Sobel DS, Ritter PL, Laurent D, Hobbs M. Effect of a self-management program on patients with chronic disease. *Eff Clin Pract.* 2001;4(6):256–62. [PubMed: 11769298]
30. Lorig K, Chastain RL, Ung E, Shoor S, Holman HR. Development and evaluation of a scale to measure perceived self-efficacy in people with arthritis. *Arthritis Rheum.* 1989 1;32(1):37–44. [PubMed: 2912463]
31. Lorig KR, Ritter PL, Laurent DD, Fries JF. Long-term randomized controlled trials of tailored-print and small-group arthritis self-management interventions. *Med Care.* 2004 4;42(4):346–54. [PubMed: 15076811]
32. Dishman RK, Motl RW, Sallis JF, Dunn AL, Birnbaum AS, Welk GJ, et al. Self-management strategies mediate self-efficacy and physical activity. *Am J Prev Med.* 2005 7;29(1):10–8. [PubMed: 15958246]
33. Sharpe PA, Wilcox S, Schoffman DE, Baruth M. “Participation, satisfaction, perceived benefits, and maintenance of behavioral self-management strategies in a self-directed exercise program for adults with arthritis.” *Eval Program Plann.* 2017 2 1;60:143–50. [PubMed: 27863325]
34. Marconcin P, Espanha M, Teles J, Bento P, Campos P, André R, et al. A randomized controlled trial of a combined self-management and exercise intervention for elderly people with osteoarthritis of the knee: the PLE² NO program. *Clin Rehabil.* 2018 2 17;32(2):223–32. [PubMed: 28714343]
**Recent RCT evaluating self-management in knee OA

35. Knittle K, De Gucht V, Hurkmans E, Peeters A, Ronday K, Maes S, et al. Targeting motivation and self-regulation to increase physical activity among patients with rheumatoid arthritis: a randomised controlled trial. *Clin Rheumatol*. 2015 2 9;34(2):231–8. [PubMed: 24213780]
36. Ndosi M, Johnson D, Young T, Hardware B, Hill J, Hale C, et al. Effects of needs-based patient education on self-efficacy and health outcomes in people with rheumatoid arthritis: a multicentre, single blind, randomised controlled trial. *Ann Rheum Dis*. 2016 6 1;75(6):1126–32. [PubMed: 26162769]
37. Azevedo ARP, de Sousa HML, Monteiro JAF, Lima ARNP. Future perspectives of Smartphone applications for rheumatic diseases self-management. *Rheumatol Int*. 2015 3 29;35(3):419–31. [PubMed: 25168866]
38. Rod K Finding Ways to Lift Barriers to Care for Chronic Pain Patients: Outcomes of Using Internet-Based Self-Management Activities to Reduce Pain and Improve Quality of Life. *Pain Res Manag*. 2016;2016:8714785. [PubMed: 27445632]
39. Revenäs Å, Opava CH, Ahlén H, Brusewitz M, Pettersson S, Åsenlöf P. Mobile internet service for self-management of physical activity in people with rheumatoid arthritis: evaluation of a test version. *RMD Open*. 2016 4 4;2(1):e000214. [PubMed: 27099777] *Preliminary evaluation of use of mobile devices
40. De Angelis G, Davies B, King J, Wells GA, Brosseau L. The use of social media by arthritis health professionals to disseminate a self-management program to patients: A feasibility study. *Digit Heal*. 2017 1 8;3:205520761770052.
41. Hofmann SG, Gómez AF. Mindfulness-Based Interventions for Anxiety and Depression. *Psychiatr Clin North Am*. 2017 12;40(4):739–49. [PubMed: 29080597]
42. Kabat-Zinn J An outpatient program in behavioral medicine for chronic pain patients based on the practice of mindfulness meditation: theoretical considerations and preliminary results. *Gen Hosp Psychiatry*. 1982 4;4(1):33–47. [PubMed: 7042457]
43. Sipe WEB, Stuart;, Eisendrath J. Mindfulness-Based Cognitive Therapy: Theory and Practice. *Can J PsychiatryTheCJP.ca CanJPsychiatry*. 2012;5757(632):63–9.
44. DiRenzo D, Crespo-Bosque M, Gould N, Finan P, Nanavati J, Bingham CO. Systematic Review and Meta-analysis: Mindfulness-Based Interventions for Rheumatoid Arthritis. *Curr Rheumatol Rep*. 2018 10 18;20(12):75. [PubMed: 30338418] *Systematic Review evaluating MBIs in RA
45. The Center For Mindfulness In Medicine, Health Care, And Society University Of Massachusetts Medical School Mindfulness-Based Stress Reduction (Mbsr): Standards Of Practice Mindfulness-Based Stress Reduction (MBSR) Standards of Practice Background and Overview. 2014;
46. Freudenthaler L, Turba JD, Tran US. Emotion Regulation Mediates the Associations of Mindfulness on Symptoms of Depression and Anxiety in the General Population. *Mindfulness (N Y)*. 2017 10 28;8(5):1339–44. [PubMed: 28989550]
47. Desrosiers A, Vine V, Klemanski DH, Nolen-Hoeksema S. Mindfulness And Emotion Regulation In Depression And Anxiety: Common And Distinct Mechanisms Of Action. *Depress Anxiety*. 2013 7;30(7):654–61. [PubMed: 23592556]
48. Garland EL, Farb NA, Goldin P, Fredrickson BL. Mindfulness Broadens Awareness and Builds Eudaimonic Meaning: A Process Model of Mindful Positive Emotion Regulation. *Psychol Inq*. 2015 10 1;26(4):293–314. [PubMed: 27087765]
49. Grossman P, Niemann L, Schmidt S, Walach H. Mindfulness-based stress reduction and health benefits: A meta-analysis. *J Psychosom Res*. 2004 7 1;57(1):35–43. [PubMed: 15256293]
50. Pradhan EK, Baumgarten M, Langenberg P, Handwerker B, Gilpin AK, Magyari T, et al. Effect of Mindfulness-Based stress reduction in rheumatoid arthritis patients. *Arthritis Rheum*. 2007 10 15;57(7):1134–42. [PubMed: 17907231]
51. Fogarty FA, Booth RJ, Gamble GD, Dalbeth N, Consedine NS. The effect of mindfulness-based stress reduction on disease activity in people with rheumatoid arthritis: a randomised controlled trial. *Ann Rheum Dis*. 2015 2 1;74(2):472–4. [PubMed: 25406303]
52. Zangi HA, Mowinckel P, Finset A, Eriksson LR, Høystad TØ, Lunde AK, et al. A mindfulness-based group intervention to reduce psychological distress and fatigue in patients with inflammatory rheumatic joint diseases: a randomised controlled trial. *Ann Rheum Dis*. 2012 6;71(6):911–7. [PubMed: 22186709]

53. Smith BW, Zautra AJ. The effects of anxiety and depression on weekly pain in women with arthritis. *Pain*. 2008 8 31;138(2):354–61. [PubMed: 18289792]
54. Lee AC, Harvey WF, Price LL, Han X, Driban JB, Wong JB, et al. Mindfulness Is Associated With Treatment Response From Nonpharmacologic Exercise Interventions in Knee Osteoarthritis. *Arch Phys Med Rehabil*. 2017 11;98(11):2265–2273.e1. [PubMed: 28506776] *Evaluation of mindfulness (trait) on treatment response in knee OA
55. Spijkerman MPJ, Pots WTM, Bohlmeijer ET. Effectiveness of online mindfulness-based interventions in improving mental health: A review and meta-analysis of randomised controlled trials. *Clin Psychol Rev*. 2016 4;45:102–14. [PubMed: 27111302]
56. van Emmerik AAP, Berings F, Lancee J. Efficacy of a Mindfulness-Based Mobile Application: a Randomized Waiting-List Controlled Trial. *Mindfulness (N Y)*. 2018 2 21;9(1):187–98. [PubMed: 29387266]
57. Firth J, Torous J, Nicholas J, Carney R, Prapat A, Rosenbaum S, et al. The efficacy of smartphone-based mental health interventions for depressive symptoms: a meta-analysis of randomized controlled trials. *World Psychiatry*. 2017 10;16(3):287–98. [PubMed: 28941113]
58. Ólason M, Andrason RH, Jónsdóttir IH, Kristbergisdóttir H, Jensen MP. Cognitive Behavioral Therapy for Depression and Anxiety in an Interdisciplinary Rehabilitation Program for Chronic Pain: a Randomized Controlled Trial with a 3-Year Follow-up. *Int J Behav Med*. 2018 2 1;25(1): 55–66. [PubMed: 29094283]
59. Nash VR, Ponto J, Townsend C, Nelson P, Bretz MN. Cognitive Behavioral Therapy, Self-Efficacy, and Depression in Persons with Chronic Pain. *Pain Manag Nurs*. 2013 12;14(4):e236–43. [PubMed: 23273826]
60. Turner JA, Anderson ML, Balderson BH, Cook AJ, Sherman KJ, Cherkin DC. Mindfulness-based stress reduction and cognitive behavioral therapy for chronic low back pain: similar effects on mindfulness, catastrophizing, self-efficacy, and acceptance in a randomized controlled trial. *Pain*. 2016 11;157(11):2434–44. [PubMed: 27257859]
61. Broderick JE, Keefe FJ, Schneider S, Junghaenel DU, Bruckenthal P, Schwartz JE, et al. Cognitive behavioral therapy for chronic pain is effective, but for whom? *Pain*. 2016 9;157(9):2115–23. [PubMed: 27227692]
62. Evers AWM, Kraaimaat FW, van Riel PLCM, de Jong AJL. Tailored cognitive-behavioral therapy in early rheumatoid arthritis for patients at risk: a randomized controlled trial. *Pain*. 2002 11;100(1–2):141–53. [PubMed: 12435467]
63. Khoo E-L, Small R, Cheng W, Hatchard T, Glynn B, Rice DB, et al. Comparative evaluation of group-based mindfulness-based stress reduction and cognitive behavioural therapy for the treatment and management of chronic pain: A systematic review and network meta-analysis. *Evid Based Ment Heal*. 2019 1 31;22(1):ebmental-2018–300062. **Systematic review comparing MBSR to CBT for management of chronic pain
64. Barsky AJ, Ahern DK, Orav EJ, Nestoriuc Y, Liang MH, Berman IT, et al. A randomized trial of three psychosocial treatments for the symptoms of rheumatoid arthritis. *Semin Arthritis Rheum*. 2010 12;40(3):222–32. [PubMed: 20621334]
65. Andersson G, Rozental A, Shafraan R, Carlbring P. Long-term effects of internet-supported cognitive behaviour therapy. *Expert Rev Neurother*. 2018 1 2;18(1):21–8. [PubMed: 29094622]
66. Ferwerda M, van Beugen S, van Middendorp H, Spillekom-van Koulil S, Donders ART, Visser H, et al. A tailored-guided internet-based cognitive-behavioral intervention for patients with rheumatoid arthritis as an adjunct to standard rheumatological care. *Pain*. 2017 5;158(5):868–78. [PubMed: 28106666] *Preliminary study evaluating use of internet-based CBT for RA
67. O'Moore KA, Newby JM, Andrews G, Hunter DJ, Bennell K, Smith J, et al. Internet Cognitive-Behavioral Therapy for Depression in Older Adults With Knee Osteoarthritis: A Randomized Controlled Trial. *Arthritis Care Res*. 2018;70(1):61–70.
68. Rini C, Porter LS, Somers TJ, McKee DC, DeVellis RF, Smith M, et al. Automated Internet-based pain coping skills training to manage osteoarthritis pain: a randomized controlled trial. *Pain*. 2015;156(5):837–48. [PubMed: 25734997]

69. Bennell KL, Nelligan RK, Rini C, Keefe FJ, Kasza J, French S, et al. Effects of internet-based pain coping skills training before home exercise for individuals with hip osteoarthritis (HOPE trial). *Pain*. 2018 5;159(9):1. [PubMed: 28885455]
70. Lawford BJ, Hinman RS, Kasza J, Nelligan R, Keefe F, Rini C, et al. Moderators of Effects of Internet-Delivered Exercise and Pain Coping Skills Training for People With Knee Osteoarthritis: Exploratory Analysis of the IMPACT Randomized Controlled Trial. *J Med Internet Res*. 2018 5 9;20(5):e10021. [PubMed: 29743149]
71. Murphy LB, Cisternas MG, Pasta DJ, Helmick CG, Yelin EH. Medical Expenditures and Earnings Losses Among US Adults With Arthritis in 2013. *Arthritis Care Res (Hoboken)*. 2018 6 1;70(6): 869–76. [PubMed: 28950426]
72. Park T Health care utilization and expenditures among adults with rheumatoid arthritis using specialty pharmaceuticals. *Res Soc Adm Pharm*. 2018 9 12;
73. Menon J, Mishra P. Health care resource use, health care expenditures and absenteeism costs associated with osteoarthritis in US healthcare system. *Osteoarthr Cartil*. 2018 4;26(4):480–4. [PubMed: 29269328]
74. Morton K, Dennison L, May C, Murray E, Little P, McManus RJ, et al. Using digital interventions for self-management of chronic physical health conditions: A meta-ethnography review of published studies. *Patient Educ Couns*. 2017 4 1;100(4):616–35. [PubMed: 28029572]
75. Mccubbin T, Dimidjian S, Kempe K, Glassey MS, Ross C, Beck A. Mindfulness-based stress reduction in an integrated care delivery system: one-year impacts on patient-centered outcomes and health care utilization. *Perm J*. 2014 11 3;18(4):4–9.
76. Kurdyak P, Newman A, Segal Z. Impact of mindfulness-based cognitive therapy on health care utilization: A population-based controlled comparison. *J Psychosom Res*. 2014 8 1;77(2):85–9. [PubMed: 25077847]
77. Stahl JE, Dossett ML, LaJoie AS, Denninger JW, Mehta DH, Goldman R, et al. Relaxation Response and Resiliency Training and Its Effect on Healthcare Resource Utilization. Dalal K, editor. *PLoS One*. 2015 10 13;10(10):e0140212. [PubMed: 26461184]
78. Khan NA, Spencer HJ, Abda E, Aggarwal A, Alten R, Ancuta C, et al. Determinants of discordance in patients' and physicians' rating of rheumatoid arthritis disease activity. *Arthritis Care Res (Hoboken)*. 2012 2 1;64(2):206–14. [PubMed: 22052672]
79. Agarwal P, Sambamoorthi U. Healthcare Expenditures Associated with Depression Among Individuals with Osteoarthritis: Post-Regression Linear Decomposition Approach. *J Gen Intern Med*. 2015 12 20;30(12):1803–11. [PubMed: 25990191]
80. Kronisch C, McLernon DJ, Dale J, Paterson C, Ralston SH, Reid DM, et al. Brief Report: Predicting Functional Disability: One-Year Results From the Scottish Early Rheumatoid Arthritis Inception Cohort. *Arthritis Rheumatol*. 2016 7 1;68(7):1596–602. [PubMed: 26866516]
81. Finan PH, Garland EL. The role of positive affect in pain and its treatment. *Clin J Pain*. 2015 2;31(2):177–87. [PubMed: 24751543]
82. Fiest KM, Hitchon CA, Bernstein CN, Peschken CA, Walker JR, Graff LA, et al. Systematic Review and Meta-analysis of Interventions for Depression and Anxiety in Persons With Rheumatoid Arthritis. *JCR J Clin Rheumatol*. 2017 12;23(8):425–34. [PubMed: 28221313]
83. Keefe FJ, Somers TJ, Martire LM. Psychologic interventions and lifestyle modifications for arthritis pain management. *Rheum Dis Clin North Am*. 2008 5;34(2):351–68. [PubMed: 18638681]

Table 1-

Current Management Strategies to Improve Self-Efficacy.

Management Category	General Description	Select Programs	Effect on Self-Efficacy
Self-Management	Group or online-based programs that encourage understanding of disease, centralize the role of the individual in managing symptoms, emotions, and medications, and promote healthy lifestyle behaviors such as diet and exercise.	Arthritis Self-Management Program, First Step to Active Health, PLE ² NO	Self-management strategies may mediate the association of self-efficacy with physical activity.
Mindfulness Based Interventions	Group-based therapies that are usually 8-weeks or longer in length designed to train individuals to cultivate mindfulness through present-moment awareness and without judgement. Programs typically include traditional seated meditations, whole-body scans, gentle yoga, and various exercises to cope with pain, bodily sensations, and emotions.	Mindfulness Based Stress Reduction, Mindfulness Based Cognitive Therapy (office), Vitality Training Programme, Internal Family Systems, Mindfulness Attention and Awareness Training	Mindfulness may improve self-efficacy via emotion regulation and positive reappraisal.
Cognitive Behavioral Therapy	Individualized psychotherapy that may be online or in-person; focuses on the relationship between thoughts, physical signs/symptoms, and behaviors.	(Traditional) Cognitive Behavioral Therapy, Pain Coping Skills Training	CBT may improve self-efficacy through restructuring of maladaptive thoughts and behaviors with a focus on symptom prediction and symptom control.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript