


4. Lee C, Adegunsoye AO, Chung J, Jablonski R, Pan I, Montner S, *et al.* Identification of predisposing environmental risk factors for interstitial lung disease [abstract]. *Am J Respir Crit Care Med* 2019;199:A1454.
5. Knoeller GE, Mazurek JM, Moorman JE. Work-related asthma among adults with current asthma in 33 states and DC: evidence from the Asthma Call-Back Survey, 2006-2007. *Public Health Rep* 2011;126:603-611.
6. De Matteis S, Jarvis D, Hutchings S, Darnton A, Fishwick D, Sadhra S, *et al.* Occupations associated with COPD risk in the large population-based UK Biobank cohort study. *Occup Environ Med* 2016;73:378-384.

Copyright © 2019 by the American Thoracic Society



Reply to Lee and Strek

From the Authors:

We welcome the comments by Dr. Lee and Dr. Strek calling for increased recognition, training, and data capture to address unrecognized occupational attributions in response to the recently published work entitled “The Occupational Burden of Nonmalignant Respiratory Diseases. An Official American Thoracic Society and European Respiratory Society Statement.” Indeed, those were implicit goals that led our international taskforce members to undertake this in-depth review (1). From the onset, we recognized that in focusing on selected nonmalignant respiratory conditions, there would be other occupational associations that we would not be able to address in depth. For example, Table E2 in the online supplement of Reference 1 describes a number of other pulmonary disorders associated with occupational exposures (with supporting citations in the main publication). These include acute eosinophilic pneumonia, bronchiolitis (obliterative, proliferative, and lymphocytic), cryptogenic organizing pneumonia, desquamative interstitial pneumonia, diffuse pulmonary hemorrhage, lipoid pneumonia, nonspecific interstitial pneumonia, and respiratory bronchiolitis interstitial lung disease (1). Many of these disorders fall into the interstitial lung disease group emphasized in the letter by Dr. Lee and Dr. Strek.

The authors also raise the important point that autoimmune conditions such as rheumatoid arthritis may not be appreciated as occupationally triggered by silica and other workplace exposures. In that context, it is important to note that concomitant lung disease may not be overt in such syndromes (2). It is our hope that further research and in-depth reviews will continue to shed light on underappreciated occupational contributors to disease and prioritize a reduction in the burden of these preventable conditions. ■

Author disclosures are available with the text of this letter at www.atsjournals.org.

Paul D. Blanc, M.D., M.S.P.H*
University of California, San Francisco
San Francisco, California

Carrie A. Redlich, M.D., M.P.H
Yale University School of Medicine
New Haven, Connecticut

On behalf of all the authors

ORCID IDs: 0000-0001-9043-175X (P.D.B.); 0000-0001-6486-6725 (C.A.R.).

*Corresponding author (e-mail: paul.blanc@ucsf.edu).

References

1. Blanc PD, Annesi-Maesano I, Balmes JR, Cummings KJ, Fishwick D, Miedinger D, *et al.* The occupational burden of nonmalignant respiratory diseases: an official American Thoracic Society and European Respiratory Society statement. *Am J Respir Crit Care Med* 2019;199:1312-1334.
2. Murphy D, Hutchinson D. Is male rheumatoid arthritis an occupational disease? A review. *Open Rheumatol J* 2017;11:88-105.

Copyright © 2019 by the American Thoracic Society



Expression of Concern: Inhaled Ethyl Nitrite Prevents Hyperoxia-impaired Postnatal Alveolar Development in Newborn Rats

The *Journal* is publishing this expression of concern about an article in the August 1, 2007, issue (1) because of potential problems with the reliability of its data. The authors have informed us that, although the data appear to be accurate, they have uncertainty about the validity of the data described in Figure 7 because of irregularities in the procedures of a lab that generated those data. ■

Reference

1. Auten RL, Mason SN, Whorton MH, Lampe WR, Foster WM, Goldberg RN, Li B, Stamler JS, Auten KM. Inhaled ethyl nitrite prevents hyperoxia-impaired postnatal alveolar development in newborn rats. *Am J Respir Crit Care Med* 2007;176:291-299.

Copyright © 2019 by the American Thoracic Society

Ⓓ This article is open access and distributed under the terms of the Creative Commons Attribution Non-Commercial No Derivatives License 4.0 (<http://creativecommons.org/licenses/by-nc-nd/4.0/>). For commercial usage and reprints, please contact Diane Gern (dgern@thoracic.org).

Originally Published in Press as DOI: 10.1164/rccm.201907-1379LE on July 25, 2019

Ⓓ This article is open access and distributed under the terms of the Creative Commons Attribution Non-Commercial No Derivatives License 4.0 (<http://creativecommons.org/licenses/by-nc-nd/4.0/>). For commercial usage and reprints, please contact Diane Gern (dgern@thoracic.org).