



# HHS Public Access

Author manuscript

*Ticks Tick Borne Dis.* Author manuscript; available in PMC 2021 January 01.

Published in final edited form as:

*Ticks Tick Borne Dis.* 2020 January ; 11(1): 101309. doi:10.1016/j.ttbdis.2019.101309.

## First report of *Candidatus Rickettsia mendelii* in *Ixodes brunneus* from the United States

Alexandra N. Cumbie, Eric L. Walters, Holly D. Gaff, Wayne L. Hynes\*

Department of Biological Sciences, Old Dominion University, Norfolk, VA 23529, USA

### Abstract

*Candidatus Rickettsia mendelii* is a novel rickettsial species recently identified in *Ixodes ricinus*. In this study, *Ixodes brunneus* collected from wild birds ( $n=77$  ticks) or vegetation ( $n=4$  ticks) in southeastern Virginia were surveyed for rickettsial agents. *Candidatus Rickettsia mendelii* was confirmed in *I. brunneus* through sequencing of the *gltA* and 16S rRNA genes. This is the first report of this rickettsial species in *Ixodes* ticks in North America.

### Keywords

*Ixodes brunneus*; *Candidatus Rickettsia mendelii*

### Introduction

*Candidatus Rickettsia mendelii* is a recently reported rickettsial species (Hajduskova et al., 2016) found in *Ixodes ricinus* and most closely related to *Rickettsia bellii*, a member of the rickettsial ancestral group (Philip et al., 1983). Classification of *Candidatus Rickettsia mendelii* was through sequencing of the citrate synthase gene (*gltA*) and 16S rRNA sequencing; these sequences were concatenated and used in constructing a phylogenetic tree (Hajduskova et al., 2016). In a survey for rickettsial agents conducted in the Czech Republic, 4524 *I. ricinus* were collected from various hosts / sources; most of the ticks were from wild birds, but also included ticks from canine hosts and vegetation. Seven of the ticks surveyed were positive for *Candidatus Rickettsia mendelii*: one was collected from vegetation, four from a dog, and two from wild birds (Hajduskova et al., 2016). Since its initial report, *Candidatus Rickettsia mendelii* has been documented in *I. ricinus* in other European countries including Slovakia (Minichova et al., 2017), Ukraine (Rogovskyy et al., 2018), and Poland (Stanczak et al., 2018).

\*Corresponding author at: Department of Biological Sciences, Old Dominion University, Norfolk, VA 23529, USA. Tel: +1 757-683-3613, whynes@odu.edu (W.L. Hynes).

**Publisher's Disclaimer:** This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

**Animal Care:** This study was conducted under Old Dominion University IACUC Protocol 12-006, Virginia Department of Game and Inland Fisheries permit 52070, and United States Department of the Interior permit 23803.

**Declaration of Interest:** None

*Ixodes brunneus* is not considered a primary vector of any human pathogens; as such, this tick species has not been studied in detail for human pathogenic agents. In the United States, *I. brunneus* is a host-specific black-legged tick that feeds exclusively on birds during all life stages (Bishopp and Trembley, 1945). In the southeastern United States, *I. brunneus* has been associated with avian paralysis, a phenomenon observed when female ticks release toxins from their salivary glands while feeding. This toxin can sometimes result in a fatal paralysis of the avian host (Luttrell et al., 1996). Previously, it has been reported that *I. brunneus* can harbor *R. rickettsii* (Clifford et al., 1969). In the study reported here, *I. brunneus* were collected from wild birds and vegetation in southeastern Virginia and tested for the presence of *Rickettsia* spp. This study was part of a larger effort to assess the role that birds play in the movement and maintenance of tick-borne bacteria in southeastern Virginia.

## Material and methods

### Tick Collection

Seventy-seven *I. brunneus* ticks, representing all three life stages, were from wild birds caught at various sites in southeastern Virginia through mist netting from 2012 to 2014. Four *I. brunneus* larvae were flagged from vegetation in 2016.

### Tick DNA Extraction

All adult ticks were cut bilaterally; one half of the tick was extracted using a DNeasy Blood and Tissue Extraction Kit (Qiagen, Valencia, CA) while the other half was stored at  $-80^{\circ}\text{C}$  for additional analysis, if required. All nymphal and larval ticks were extracted whole due to their small size. Samples were pulverized using 1.0-mm glass beads using a Mini-Beadbeater-1 (BioSpec Products, Inc., Bartlesville, OK, USA). Cells were lysed using lysis buffers; the DNA was column purified and eluted according to kit manufacturers' guidelines.

### Tick Identification

Each tick was identified using molecular methods, as morphological identification is often inconclusive with immature stages and engorged ticks (Anderson et al., 2004). *Ixodes brunneus* was identified using a PCR-amplified 454-bp segment of the 16S mitochondrial RNA gene (Macaluso et al., 2003; Wright et al., 2014). Each PCR reaction (final volume 25  $\mu\text{L}$ ) consisted of 12.5  $\mu\text{L}$  EconoTaq PLUS 2X Master Mix (Lucigen, Middleton, WI), 2  $\mu\text{L}$  of each primer (10  $\mu\text{M}$ ), 3.5  $\mu\text{L}$  of nuclease-free water, and 5  $\mu\text{L}$  DNA. The thermocycler protocol included a denaturing step at  $95^{\circ}\text{C}$  for 3 min, followed by 30 cycles of:  $95^{\circ}\text{C}$  for 30 s,  $52^{\circ}\text{C}$  for 45 s, and  $72^{\circ}\text{C}$  for 60 s; the reaction finished with an extension step at  $72^{\circ}\text{C}$  for 7 min. The 16S amplicons were sequenced using a BigDye Terminator v3.1 Cycle Sequencing Kit (ABI-ThermoFisher Scientific, Carlsbad, CA) with analysis on an ABI 3130x1 genetic analyzer. Ticks were identified to species by comparison with known tick 16S gene sequences using NCBI Blast (<http://blast.ncbi.nlm.nih.gov>). No template negative controls were included in PCR reactions.

### *Rickettsia* spp. Identification

Two real-time PCR assays were used: one amplifying a portion of the rickettsial 17-kDa gene to detect the presence of *Rickettsia* spp. in the ticks; the other amplified a portion of the

*ompB* gene specific to *R. parkeri* (Jiang et al., 2012). Each rickettsial 17-kDa gene real-time PCR reaction (final volume 25  $\mu$ L) consisted of 12.5  $\mu$ L EconoTaq PLUS 2X Master Mix (Lucigen, Middleton, WI), 1.25  $\mu$ L of each primer (10  $\mu$ M), 1  $\mu$ L of probe (10  $\mu$ M), 2  $\mu$ L  $MgCl_2$  (25 mM), 2  $\mu$ L of nuclease-free water, and 5  $\mu$ L DNA. Each *R. parkeri ompB* real-time PCR reaction (final volume 25  $\mu$ L) consisted of 12.5  $\mu$ L EconoTaq PLUS 2X Master Mix (Lucigen, Middleton, WI), 1.75  $\mu$ L of each primer (10  $\mu$ M), 1  $\mu$ L of probe (10  $\mu$ M), 2.5  $\mu$ L  $MgCl_2$  (25 mM), 0.5  $\mu$ L of nuclease-free water, and 5  $\mu$ L DNA. The thermocycler protocol included a pre-hold step of 50°C for 2 min, a denaturing step of 95°C for 2 min, followed by 45 cycles of: 95°C for 15 s and 60°C for 30 s. Using the rickettsial 17-kDa-specific assay (Jiang et al., 2012), ticks positive for *Rickettsia* spp. (Cq  $\leq$  35 cycles) were detected. Depending on availability of DNA, rickettsia-positive ticks were further examined by amplification and sequencing of one or more of *ompA*, *ompB*, *sca4*, *gltA*, and 16S rRNA genes (Table 1). No template negative controls were included in PCR reactions. Sequences were initially analyzed using Geneious software (Kearse et al., 2012) version R11 (<http://www.geneious.com>, Biomatters Inc., New Zealand) and compared using NCBI Blast (<http://blast.ncbi.nlm.nih.gov>) to identify the rickettsial species.

## Results

The sequence of the 16S mitochondrial RNA gene confirmed 81 ticks collected from wild birds and vegetation were *I. brunneus*. The rickettsial 17-kDa real-time PCR indicated 44 of 66 (67%) ticks tested positive for rickettsial DNA; we were not able to test the remaining 15 because the DNA was of poor quality due to DNA degradation, or the samples had already been used for other experiments. Five of the 66 ticks had *Candidatus Rickettsia mendelii* as determined by sequencing of the rickettsial *gltA* gene, based on a BLAST alignment to identify the species. A 790-bp partial *gltA* fragment (Accession number MH458574) was 99.9% (789 of 790-bp) identical to the sequence of *Candidatus Rickettsia mendelii* (Hajduskova et al., 2016). We amplified the rickettsial 16S rRNA gene from 3 of the 5 ticks. Some of the sequencing chromatograms contained multiple overlapping peaks, but BLAST analysis of clean single read (one direction) fragments of 627–745 bp were >99% identical to *Candidatus Rickettsia mendelii*. A 627-bp fragment of the 16S gene from one isolate, sequenced in both directions, was 99.2–99.7% identical to the three *Candidatus R. mendelii* 16S genes in GenBank (Accession numbers: [KJ882316](#), [KJ882317](#), [KJ882318](#)); the closest match to any *Rickettsia* sp. other than *Candidatus Rickettsia mendelii* was <98%. We were not able to amplify *ompA*, *ompB*, or *sca4* from any of the 5 tick samples that were positive for *Candidatus R. mendelii*; similar results were reported by Hajduskova et al. (2016). Using *R. parkeri* species-specific real-time PCR analysis (Jiang et al., 2012), *R. parkeri* was detected in 19 of the 81 (23.5%) ticks. Amplified portions of the *gltA* and *ompA* genes were sequenced, in one direction, from two ticks for each gene. The closest similarity for the sequence was to the respective *R. parkeri* genes. For the *gltA* gene, similarity was 100% over at least 225bp, while for *ompA* it was >98% over more than 475bp. The rickettsia present in the other 17-kDa positive ticks was not able to be determined due to lack of DNA or inability to obtain sequence data.

All ticks that were positive for *Candidatus R. mendelii* were *I. brunneus* juveniles: 3 nymphs and 2 larvae (Table 2). Two nymphs were collected from the same bird, a hermit thrush

(*Catharus guttatus*) in Zuni, one nymph from a tufted titmouse (*Baeolophus bicolor*) in Portsmouth, one larva from a hermit thrush in Chesapeake, and one larva flagged from vegetation in Cape Charles.

## Discussion

This is the first report of *Candidatus Rickettsia mendelii* in a tick population within the United States. In the Czech Republic study, *Candidatus Rickettsia mendelii* was determined to be phylogenetically unique based on the sequence of the *gltA* gene and 16S rRNA; other rickettsial genes: *ompA*, *ompB*, and *sca4* could not be detected (Hajduskova et al., 2016). Therefore, no sequence data are available for the *ompA*, *ompB*, and *sca4* genes of *Candidatus Rickettsia mendelii* for comparison. Likewise, we were unable to amplify or sequence these three genes from the *I. brunneus* infected with *Candidatus Rickettsia mendelii*. It is unknown why we were unable to amplify *ompA*, *ompB*, and *sca4* in *Candidatus Rickettsia mendelii*, but it could be attributed to the absence of these genes or a high degree of sequence heterogeneity in this *Rickettsia* sp. We were, however, able to amplify and sequence rickettsial genes (*ompA* and *gltA*) from other 17-kDa-positive and *R. parkeri*-positive *I. brunneus*. Results suggested the presence of other *Rickettsia* spp. in these ticks and, in some cases, coinfections (multiple peaks in sequencing chromatograms) with two different *Rickettsia* sp. were possible, although we could not determine this with certainty. *R. parkeri* species-specific real-time PCR amplified *R. parkeri* DNA in some ticks suggesting coinfections of *I. brunneus* with *Candidatus Rickettsia mendelii* and *R. parkeri*. Additionally, some of these ticks were from the same bird, potentially inflating the *R. parkeri* prevalence. Thus, the precise number of individual ticks within our sample set carrying *Candidatus R. mendelii* is unknown.

How and why *Candidatus Rickettsia mendelii* is present in two tick taxa from two isolated geographic regions, separated by the Atlantic Ocean, is unknown. Hajduskova et al. (2016) suggested a symbiotic rather than pathogenic relationship for this organism. It is possible that our identification of this organism in *I. brunneus* is a previously unidentified endosymbiont in ticks from the United States. *Candidatus Rickettsia mendelii* was found in an unfed *I. brunneus* larva flagged from vegetation, suggesting transovarial transmission from an infected adult female tick. A less likely possibility is that the infected nymphs and larvae acquired *Candidatus Rickettsia mendelii* from the bird host. Although birds serve as hosts for many tick species, information on systemic infection by *Rickettsia* spp. in bird blood is limited and with mixed results; some studies have detected *Rickettsia* spp. in bird blood (Berthova et al., 2016; Hornok et al., 2014), others have not (Cohen et al., 2015). Life histories of the hermit thrush and tufted titmouse reveal that the former does not normally migrate outside of North America (Dellinger et al., 2012), while the latter has never been recorded outside of North America (Ritchison et al., 2015). Thus, it would be extremely unlikely these birds traveled to Europe and played host to an infected *I. ricinus* to acquire the bacteria.

It is also possible that an infected *I. ricinus* travelled to North America and infected an *I. brunneus* via sharing of hosts or co-feeding. As far as is known, *I. ricinus* has not been found in North America where other *Ixodes* spp. are established (Rich et al., 1995). One means for

*I. ricinus* to travel to North America would be on a host. The two avian species that the infected *I. ricinus* were collected from in the Czech Republic were the common nightingale (*Luscinia megarhynchos*) and the European robin (*Erithacus rubecula*), neither of which migrate outside of Europe (Hajduskova et al., 2016), much less as far as North America.

One further explanation is that *I. ricinus* may have fed on livestock, which could have been transported to North America, although this is also unlikely as there have been no livestock exports historically from the Czech Republic to the United States (Mikulasova, 2016).

The most plausible explanation for our finding of *Candidatus Rickettsia mendelii* in *I. brunneus* is that it is already established in eastern North America and has merely gone undetected. With recent reports on foreign ticks (Rainey et al., 2018) potentially going undetected for many years (Buccino, 2018) it is possible that the presence of unusual bacteria in native ticks may indicate a past or contemporaneous introduction. Our study indicates a need for the identification of both endosymbiont and pathogenic rickettsial species. Understanding interactions between ticks and their microbes can provide insights into pathogen-vector-host activity. The work presented here emphasizes a need for more thorough studies on the movement and occurrence of different rickettsial species as they become established in different geographic regions. Although *Candidatus R. mendelii* is unlikely to be a human pathogen, continued active surveillance provides important information and early warning of potential infectious agents.

## Acknowledgements

We acknowledge Erin Heller and the many field assistants for their help with mist netting birds and Ashley Morris, Zach Bement, and Anna Phan for their support in molecular identification of the ticks. We also thank The Nature Conservancy, Kiptopeke State Park, and Blackwater Ecological Preserve for permission to use their land.

**Funding:** This work was funded in part by NIH grant 1R01AI136035 as part of the joint NIH-NSF-USDA Ecology and Evolution of Infectious Diseases program and by start-up funds from Old Dominion University to ELW and grants from the Virginia Academy of Science and the Virginia Society of Ornithology to Erin Heller.

## References

- Anderson JM, Ammerman NC, Norris DE, 2004 Molecular differentiation of metastriate tick immatures. *Vector Borne Zoonotic Dis.* 4, 334–342. [PubMed: 15682516]
- Bishopp FC, Trembley HL, 1945 Distribution and Hosts of Certain North American Ticks. *J. Parasit* 31, 1–54.
- Buccino N, 2018 Rutgers-Led “Tick Blitz” Finds Exotic Longhorned Ticks and Aggressive Lone Star Ticks in New Locations Across New Jersey. <https://news.rutgers.edu/rutgers-led-“tick-blitz”-finds-exotic-longhorned-ticks-and-aggressive-lone-star-ticks-new-locations/20180530#.WxpzgfZFwkG> (accessed 8 June 2018)
- Clifford CM, Sonenshine DE, Atwood EL, Robbins CS, Hughes LE, 1969 Tests on ticks from wild birds collected in the eastern United States for rickettsiae and viruses. *Am. J. Trop. Med. Hyg* 18, 1057–1061. [PubMed: 4982220]
- Cohen EB, Auckland LD, Marra PP, Hamer SA, 2015 Avian migrants facilitate invasions of neotropical ticks and tick-borne pathogens into the United States. *Appl. Environ. Microbiol* 81, 8366–8378. [PubMed: 26431964]
- Dellinger R, Wood PB, Jones PW, Donovan TM 2012 Hermit Thrush (*Catharus guttatus*), version 2.0 In Poole AF (Ed.), *The Birds of North America*. Ithaca, New York, USA: Cornell Lab or Ornithology Retrieved from <https://birdsna.org/Species-Account/bna/species/herthr/introduction>. doi:<https://doi.org/10.2173/bna.261>

- Hajduskova E, Literak I, Papousek I, Costa FB, Novakova M, Labruna MB, Zdrzilova-Dubska L, 2016 '*Candidatus Rickettsia mendelii*', a novel basal group rickettsia detected in *Ixodes ricinus* ticks in the Czech Republic. *Ticks Tick-Borne Dis.* 7, 482–486. [PubMed: 26873811]
- Jiang J, Blair PJ, Felices V, Moron C, Cespedes M, Anaya E, Schoeler GB, Sumner JW, Olson JG, Richards AL, 2005 Phylogenetic analysis of a novel molecular isolate of spotted fever group Rickettsiae from northern Peru: *Candidatus Rickettsia andeanae*. *Ann. N. Y. Acad. Sci* 1063, 337–342. [PubMed: 16481537]
- Jiang J, Stromdahl EY, Richards AL, 2012 Detection of *Rickettsia parkeri* and *Candidatus Rickettsia andeanae* in *Amblyomma maculatum* Gulf Coast ticks collected from humans in the United States. *Vector Borne Zoonotic Dis.* 12, 175–182. [PubMed: 22022815]
- Kearse M, Moir R, Wilson A, Stones-Havas S, Cheung M, Sturrock S, Buxton S, Cooper A, Markowitz S, Duran C, Thierer T, Ashton B, Meintjes P, Drummond A, 2012 Geneious Basic: an integrated and extendable desktop software platform for the organization and analysis of sequence data. *Bioinformatics* 28, 1647–1649. [PubMed: 22543367]
- Labruna MB, McBride JW, Bouyer DH, Camargo LM, Camargo EP, Walker DH, 2004 Molecular evidence for a spotted fever group *Rickettsia* species in the tick *Amblyomma longirostre* in Brazil. *J. Med. Entomol* 41, 533–537. [PubMed: 15185961]
- Luttrell MP, Creekmore LH, Mertins JW, 1996 Avian tick paralysis caused by *Ixodes brunneus* in the southeastern United States. *J. Wildl. Dis* 32, 133–136. [PubMed: 8627926]
- Macaluso KR, Mulenga A, Simser JA, Azad AF, 2003 Differential expression of genes in uninfected and rickettsia-infected *Dermacentor variabilis* ticks as assessed by differential-display PCR. *Infect. Immun* 71, 6165–6170. [PubMed: 14573632]
- Mikulasova J, 2016 Livestock and Products Annual 2016 Czech Republic GAIN Report Number: EZ1609. [https://gain.fas.usda.gov/Recent%20GAIN%20Publications/Livestock%20and%20Products%20Annual%202016\\_Prague\\_Czech%20Republic\\_8-19-2016.pdf](https://gain.fas.usda.gov/Recent%20GAIN%20Publications/Livestock%20and%20Products%20Annual%202016_Prague_Czech%20Republic_8-19-2016.pdf) (accessed 5/28/2018)
- Minichova L, Hamsikova Z, Mahrikova L, Slovak M, Kocianova E, Kazimirova M, Skultety L, Stefanidesova K, Spitalska E, 2017 Molecular evidence of *Rickettsia* spp. in ixodid ticks and rodents in suburban, natural and rural habitats in Slovakia. *Parasit. Vectors* 10.
- Nilsson K, Jaenson TG, Uhnöo I, Lindquist O, Pettersson B, Uhlen M, Friman G, Pahlson C, 1997 Characterization of a spotted fever group *Rickettsia* from *Ixodes ricinus* ticks in Sweden. *J. Clin. Microbiol* 35, 243–247. [PubMed: 8968916]
- Paddock CD, Sumner JW, Comer JA, Zaki SR, Goldsmith CS, Goddard J, McLellan SL, Tamminga CL, Ohl CA, 2004 *Rickettsia parkeri*: a newly recognized cause of spotted fever rickettsiosis in the United States. *Clin. Infect. Dis* 38, 805–811. [PubMed: 14999622]
- Philip RN, Casper EA, Anacker RL, Cory J, Hayes SF, Burgdorfer W, Yunker CE, 1983 *Rickettsia bellii* sp. nov.: a tick-borne Rickettsia, widely distributed in the United States, that is distinct from the spotted fever and typhus biogroups. *Int. J. Syst. Evol. Microbiol* 33, 94–106.
- Rainey T, Occi JL, Robbins RG, Egizi A, 2018 Discovery of *Haemaphysalis longicornis* (Ixodida: Ixodidae) parasitizing a sheep in New Jersey, United States. *J. Med. Entomol* 55, 757–759. [PubMed: 29471482]
- Rich SM, Caporale DA, Telford SR 3rd, Kocher TD, Hartl DL, Spielman A, 1995 Distribution of the *Ixodes ricinus*-like ticks of eastern North America. *Proc. Natl. Acad. Sci. U.S.A* 92, 6284–6288. [PubMed: 7603983]
- Ritchison G, Grubb TCJ, Pravosudov VV 2015 Tufted Titmouse (*Baeolophus bicolor*), version 2.0 In Rodewald PG (Ed.), *The Birds of North America* (Vol. version 2). Ithaca, New York, USA: Cornell Lab of Ornithology Retrieved from <https://birdsna.org/Species-Account/bna/species/tuftit/introduction>. doi:10.2173/bna.86
- Rogovskyy A, Batool M, Gillis DC, Holman PJ, Nebogatkin IV, Rogovska YV, Rogovskyy MS, 2018 Diversity of *Borrelia* spirochetes and other zoonotic agents in ticks from Kyiv, Ukraine. *Ticks Tick Borne Dis.* 9, 404–409. [PubMed: 29258801]
- Stanczak J, Biernat B, Racewicz M, Zalewska M, Matyjasek A, 2018 Prevalence of different *Rickettsia* spp. in *Ixodes ricinus* and *Dermacentor reticulatus* ticks (Acari: Ixodidae) in north-eastern Poland. *Ticks Tick Borne Dis.* 9, 427–434. [PubMed: 29290582]

Wright CL, Hynes WL, White BT, Marshall MN, Gaff HD, Gauthier DT, 2014 Single-tube real-time PCR assay for differentiation of *Ixodes affinis* and *Ixodes scapularis*. *Ticks Tick Borne Dis.* 5, 48–52. [PubMed: 24192510]

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

**Table 1.**

Primers used to amplify rickettsial genes. These primers were used to amplify outer membrane protein A (*ompA*), outer membrane protein B (*ompB*), gene D (*sca4*), citrate synthase (*gltA*), and 16S rRNA.

| Gene            | Fragment Size | Primers  | Reference              |
|-----------------|---------------|--|------------------------|
| <i>ompA</i>     | 540-bp        | 190-FN1 5'-AAGCAATACAACAAGGTC<br>190-RN1 5'-TGACAGTTATTATACCTC           | (Paddock et al., 2004) |
| <i>ompB</i>     | 1895-bp       | RompB11F 5'-ACCATAGTAGCMAGTTTGCAG<br>RompI902R 5'-CCGTCATTCCAATAACTAACTC | (Jiang et al., 2005)   |
| <i>sca4</i>     | 928-bp        | D1F 5'-ATGAGTAAAGACGGTAACCT<br>D928R 5'-AAGCTATTGCGTCATCTCCG             | (Jiang et al., 2005)   |
| <i>gltA</i>     | 830-bp        | CS239 5' GCTCTTCTCATCCTATGGCTATTAT<br>CS1069 5' CAGGGTCTTCGTGCATTCTT     | (Labruna et al., 2004) |
| <i>16S rRNA</i> | 757-bp        | Ric 5'- TCTAGAACGAACGCTATCGGTAT<br>Ric Rt 5'- TTTCATCGTTTAACGGCGTGGACT   | (Nilsson et al., 1997) |
| <i>16S rRNA</i> | 1385-bp       | Ric 5'- TCTAGAACGAACGCTATCGGT AT<br>Ric U8 5'- TGCGTTAGCTCACCACCTTCAGG   | (Nilsson et al., 1997) |



**Table 2.**

Collection locations of *Ixodes brunneus* ticks, infected with *Candidatus Rickettsia mendelii*, from various sites across southeastern Virginia.

| Tick Species           | Tick Life Stage | Location   | Source                                     |
|------------------------|-----------------|--|--|
| <i>Ixodes brunneus</i> | Nymph           | Blackwater Ecological Preserve in Zuni, Virginia       | Hermit thrush, <i>Catharus guttatus</i>    |
| <i>Ixodes brunneus</i> | Nymph           | Blackwater Ecological Preserve in Zuni, Virginia       | Hermit thrush, <i>Catharus guttatus</i>    |
| <i>Ixodes brunneus</i> | Nymph           | Residential area in Portsmouth, Virginia               | Tufted titmouse, <i>Baeolophus bicolor</i> |
| <i>Ixodes brunneus</i> | Larva           | Property of Nature Conservancy in Chesapeake, Virginia | Hermit thrush, <i>Catharus guttatus</i>    |
| <i>Ixodes brunneus</i> | Larva           | Kiptopeke State Park in Cape Charles, Virginia         | Vegetation                                 |