

Overview

A Review of Strain and Sex Differences in Response to Pain and Analgesia in Mice

Jennifer C Smith*

Pain and its alleviation are currently a highly studied issue in human health. Research on pain and response to analgesia has evolved to include the effects of genetics, heritability, and sex as important components in both humans and animals. The laboratory mouse is the major animal studied in the field of pain and analgesia. Studying the inbred mouse to understand how genetic heritable traits and/or sex influence pain and analgesia has added valuable information to the complex nature of pain as a human disease. In the context of biomedical research, identifying pain and ensuring its control through analgesia in research animals remains one of the hallmark responsibilities of the research community. Advancements in both human and mouse genomic research shed light not only on the need to understand how both strain and sex affect the mouse pain response but also on how these research achievements can be used to improve the humane use of all research animal species. A better understanding of how strain and sex affect the response to pain may allow researchers to improve study design and thereby the reproducibility of animal research studies. The need to use both sexes, along with an improved understanding of how genetic heritability affects nociception and analgesic sensitivity, remains a key priority for pain researchers working with mice. This review summarizes the current literature on how strain and sex alter the response to pain and analgesia in the modern research mouse, and highlights the importance of both strain and sex selection in pain research.

DOI: 10.30802/AALAS-CM-19-000066

Pain is a prevalent disease state and public health crisis in humans worldwide.^{70,97,105,124} In 2016, an estimated 20.4% of adults (50 million) in the United States suffered from chronic pain, with up to 8% (11 million) reporting pain lasting over 6 mo, resulting in substantial restriction of life activities, including employment.²⁶ Furthermore, pain has emerged as a primary symptom in both mental health and substance abuse treatment.⁶⁸

The science of pain has now become a unique research field, and animal models have played a significant role in this advancement.⁸⁴ A categorical and bibliometric meta-trend analysis of the number of articles submitted to the journal *Pain* during the years 1975 to 2007 further demonstrates the use of animal models in pain research.⁸⁹ Approximately one-third of all the published work in *Pain* during this time used animal subjects, the majority of which included rodents (rats and mice) which were conscious during the studies.^{84,89} During the past 2 decades of specific pain research, animal models have helped advance the hypothesis that genetics is a possible etiologic or confounding factor of pain.⁵⁸⁻⁶⁰ We now know that pain and analgesia traits are heritable in both humans and mice.^{59,75,81,91-93,102,108}

In humans, studies of the genetics of pain are somewhat limiting due to the large amount of resources needed and unique study design (for example, twin studies).^{1,79} Rodent models have advantages over their human counterparts in genetic studies due to their small size and ease of use in larger investigations.⁹² Rodent models are also advantageous due to their ability to provide better control over genotype, environment, and pain stimulus parameters.⁵⁹ Moreover, rodent models offer easier access

to aging populations¹¹³ and studies can be conducted over an animal's entire lifecycle.³⁸

The preferred animal model for the study of genetic heritability in pain research is the laboratory mouse.^{1,15,66,78,123} With its large number of inbred strains with known pedigrees, and the capacity to genetically modify the species, the laboratory mouse has become an important tool in the study of pain and analgesia.^{1,59} The purpose of this review was to identify, describe, and summarize the current trends in pain research, relating to the strain and sex differences of common inbred strains of mice.

Pain Terminology

Pain and nociception are 2 different terms. The International Association for the Study of Pain (IASP) defines pain as “an unpleasant sensory and emotional experience associated with actual or potential tissue damage or described in terms of such damage”.^{3,68} Pain typically involves a noxious stimulus that activates nociceptors in the body to send signals to the central nervous system, where they are processed to generate multiple responses,⁹⁸ including a complex biopsychosocial pain experience.^{68,71} The IASP defines nociception as “the neural process of encoding and processing noxious stimuli”.³ Nociception represents the nervous systems' processing of information, as generated by nociceptor activation. This information is processed at both spinal and supraspinal levels of the central nervous system, providing details about the noxious events.^{98,117} The study of pain in the laboratory also includes terms such as allodynia (pain in response to a stimulus that does not usually provoke pain), and hyperalgesia (increased pain sensitivity from a stimulus that usually provokes pain).^{3,44} To simplify the terms in this review, “pain” will be used to describe the overall process of

Received: 17 Jun 2019. Revision requested: 05 Aug 2019. Accepted: 16 Sept 2019.
Department of Bioresearches, Henry Ford Health System, Detroit, Michigan
*Corresponding author. Email: jsmith28@hfhs.org

nociception processing and response to noxious stimuli; while “nociception” will be referred to only in the discussion of specific laboratory methodology, which could include allodynia and/or hyperalgesia.

The Genetics of Pain

Background. Early research on pain documented significant variability in the response to pain in both humans and animals.^{78,91} This variability was hypothesized, and later confirmed, to have a genetic cause.⁷⁸ Much of this early pain research was developed in rodent models, with the laboratory mouse remaining the most studied animal species.^{21,66,74} The mouse models that were initially used were bred specifically for nociception testing, and include the CXBK recombinant inbred strain, and the High Analgesia/Low Analgesia (HA/LA), and High Analgesia Response/Low Analgesia Response (HAR/LAR) mouse lines.⁷⁸ Although advantageous, these mouse lines were soon replaced with more common inbred mouse strains (for example C57BL/6 and 129) as specific phenotypes for pain were characterized.^{85,87,91}

Pain at the Gene Level. Completion of the Human Genome Project (HGP) in 2003, and its subsequent mouse genome sequence, was a significant milestone in identifying, sequencing, and storing over 20,000 human genes.³⁷ Pain and analgesia traits are heritable in humans and mice.^{85,91} Trait heritability refers to the portion of the overall variability that is due to inherited genetic factors⁵⁹ and is an essential component to identifying “pain genes.” The heritability of a trait must be known before it can be identified at the gene level.⁵⁹ Several methods exist to determine if a trait is heritable, including selective breeding, genetic reference populations, and inbred strain differences, which will be the focus of this review. It is estimated that less than 50% of the total variability in a trait is due to genetic factors in humans and rodent pain models; however, this percentage can be increased with the appropriate mitigation of environmental factors.⁴⁸

Mapping and manipulating the genes specific to pain has greatly improved the tool kit for the modern researcher. More recently, the Pain Genes Database⁵⁷ has been established to provide an interactive web-based browser specific for mouse genes. Through the use of this new database, several hundred genes germane to pain or analgesia in the mouse have been identified and cataloged for use in research.^{57,78} The inclusion of functional genomics⁶⁷ and the ability to further exploit genetic information in “pain genes” has further advanced the study of pain and analgesia.

Similarly, the field of pain genetics has been advanced by the use of genetic linkage mapping to identify and establish the genomic position of the pain related trait.⁵⁹ By estimating the distance between the genomic loci, this technique can be used to map specific regions or quantitative trait loci (QTL) of the genome associated with a pain phenotype of interest. Using QTL to guide the homologous recombination breeding, the resulting genetic population provides an inexpensive path to the identification of potential pain gene candidates.^{59,108}

The Transgenic Knockout Mouse. Transgenesis, homologous recombination, and CRISPR gene editing have allowed new genetic models to be developed, and the “pain genes” that have been identified to be further evaluated.^{57,66} One review estimates that at least 60 publications per year are published demonstrating a significant behavioral pain phenotype resulting from the null mutation of at least one single gene.⁵⁷ Inclusion and summation of the myriad of knock-out strains specific to pain or analgesic response is beyond the scope of this review; however,

the relationship to common inbred strains will be addressed in the following section.

Common Inbred Mouse Strains. The inbred mouse remains the most studied species in heritability pain research.⁹² Common inbred mouse strains have been thoroughly characterized, making their similarities and differences well known.⁴ For instance, inbred strains can differ in behavior,^{15,22} serotonin levels,^{6,127} pigmentation,¹²⁵ immune system components,¹¹⁸ and responses to both chemotherapeutic agents¹³¹ and nicotine.⁴² These advanced strain characterizations have been loosely called “strain surveys” and have utility in pain modeling for disease states such as endometriosis,²⁸ arthritis,³³ fibromyalgia,¹¹ and psychiatric disorders.⁶¹

The earliest strain surveys were completed on common strains from different vendors.⁹⁶ More recently, common inbred strains have been compared in nociception assays.¹³ To date, several studies have been undertaken to survey the possible genetic contributions of strain to complex disorders, and these strain surveys are essential tools in providing researchers with a road map for selection of a strain.^{22,96} Caution is warranted in generalizing from a mouse strain survey to the variation in human responses; however, this connection between the mouse and human genome suggests the strain screening to be a useful tool in research.⁵²

Differences in mouse strain relating to pain and analgesia. Few reviews on mouse strains include a compilation of comparisons between strain and responses to nociception tests and/or analgesia.^{19,69,75,81} The seminal work on this topic by Mogil and colleagues demonstrated that inbred mice differ in their responses to common nociceptive challenges.⁹² Testing 11 inbred mouse strains (129/J, A/J, AKR/J, BALB/cJ, C3H/HeJ, C57BL/6J, C58J, CBA/J, DBA2/J, RIIS/J, and SM/J) against 12 common models of nociception (writhing, autotomy after hindlimb denervation, carrageenan hypersensitivity, formalin test, hot-plate test, Hargreaves’ test of thermal nociception, Chung peripheral nerve injury model, tail-withdrawal test, and von Frey filament test of mechanical sensitivity), this laboratory demonstrated that rodent populations display large and heritable differences in both nociception and analgesic sensitivity.⁹² Data from these common inbred mouse strains show that they react differently to nociception assays. Genetic variation between mouse strains and their response to nociception assays suggests that nociception in the mouse is heritable. Follow up strain screening studies^{60,93} continued to expand the number of nociception assays that were used, ultimately revealing 5 major “clusters” or “types” of nociception and pain processing, including baseline thermal nociception, spontaneous response to chemical stimuli, thermal hypersensitivity, mechanical hypersensitivity, and afferent input-dependent hypersensitivity.

To further highlight the importance of strain variation on pain research, we conducted a literature search to investigate the connection between mouse inbred strain differences and performance on nociceptive assays relating to pain and analgesia. Table 1 through Table 3 provides the reader with a compilation of this search, summarized by the strain(s) studied. Inclusion in this table was limited to published work that compared at least 2 strains (or substrains of one strain) in similar testing methodology. Published literature was identified by the key search terms “mouse”, “strain”, “pain”, “nociception”, “analgesia” (PubMed). In addition to the strain breakdown found in Table 1 through Table 3, the following mouse models of pain and nociception have been characterized and validated using the strain screening approach: neuropathic pain,^{39,110,134} inflammatory nociception,^{87,134} cutaneous thermal nociception,^{13,30,101,134} hot and

Table 1. Summary of literature (1998–2003) assessing common mouse strain differences relating to pain and analgesia.

	Mogil 1998 ⁸⁵	Mogil 1999 ⁸¹	Mogil 1999 ⁹²	Mogil 1999 ⁹³	Wan 2001 ¹²⁵	Kest 2002 ⁵²	Lariviere 2002 ⁶⁰	Bon 2003 ¹⁰	Chesler 2003 ¹⁹	Kamp 2003 ⁴⁶
129/J	x	x	x	x	x					
129P2/J										
129P3/J					x		x	x		
129Sv										
129S6										x
A/J	x	x	x	x	x		x	x		
A/HeJ										
AKR/J	x	x	x	x	x		x	x		
BALB/c										
BALB/cJ	x	x	x	x	x		x	x		
C3H/He										
C3H/HeJ	x	x	x	x	x		x	x		
C3H/HeN										
C57BL/6										x
C57BL/6J	x	x	x	x	x		x	x		
C57BL/6N										
C57BL/10J							x	x		
C58/J	x	x	x	x	x		x	x		
CBA										
CBA/CaCrI										
CBA/J	x	x	x	x	x			x		
CD1	x				x			x		
DBA/2										
DBA/2J	x	x	x	x	x		x	x		
FVB										
ICR	x				x					
LP						x				
RIIS/J	x	x	x	x			x	x		
SJL										
SM/J	x	x	x	x	x		x	x		
SW						x				

Table 2. Summary of literature (2003–2009) assessing common mouse strain differences relating to pain and analgesia.

	Levin-Arama 2003 ⁶⁴	Neilan 2003 ⁹⁹	Wilson 2003 ¹²⁸	Wilson 2003 ²⁹	Banik 2006 ²	Green 2006 ³⁶	Liang 2006 ⁶⁶	Mogil 2006 ⁸⁸	Rigaud 2008 ¹¹⁰	Dickinson 2009 ²⁷
129/J										
129P2/J										
129P3/J				x		x				
129Sv					x		x			
129S6										
A/J				x		x	x			
A/HeJ						x	x			
AKR/J				x		x	x			
BALB/c	x			x		x	x			
BALB/cJ				x		x	x			
C3H/He				x		x	x			
C3H/HeJ				x		x	x			
C3H/HeN										
C57BL/6	x				x					
C57BL/6J				x		x	x	x		
C57BL/6N										
C57BL/10J										
C58/J				x		x				
CBA										
CBA/CaCrI										x
CBA/J				x		x				
CD1				x						
DBA/2				x						
DBA/2J					x				x	
FVB										
ICR										
LP										
RIIIS/J				x		x				
SJL				x						
SM/J				x		x				
SW			x							

Table 3. Summary of literature (2010–2019) assessing common mouse strain differences relating to pain and analgesia.

	Wijnvoord 2010 ²⁷	Benedetti 2012 ⁶	Young 2014 ³⁴	Miller 2015 ⁷³	Moloney 2015 ⁷⁶	Ono 2015 ¹⁰¹	Tajerian 2015 ¹²¹	Blennerhassett 2017 ⁷	Cho 2018 ²⁰	Isami 2018 ⁴²	Bryant 2019 ¹³
129/J					x						
129P2/J											
129P3/J			x		x						
129Sv											
129S6	x										
A/J			x		x					x	
A/HeJ											
AKR/J			x								
BALB/c	x		x		x	x		x			
BALB/cJ		x									
C3H/He											
C3H/HeJ			x							x	
C3H/HeN					x						
C57BL/6						x					
C57BL/6J	x		x		x		x			x	x
C57BL/6N		x							x		x
C57BL/10J			x								
C58/J			x								
CBA				x							
CBA/CaCrI											
CBA/J			x		x						
CD1					x			x			
DBA/2				x							
DBA/2J	x		x		x					x	
FVB					x						
ICR											
LP											
RIIIS/J			x								
SJL											
SM/J			x		x						
SW					x						

cold nociception,^{29,81} visceral nociception,^{10,46,96} and scratching and itch behavior.^{36,134}

Similarly, strain surveys have documented variation in the sensitivity to analgesics.^{128,129} Given that strains of mice perform differently in various nociception assays,¹³⁴ one might assume that strains that are more sensitive to pain will be less responsive to analgesics. However, early strain surveys⁸⁶ demonstrated the opposite effect. Strains that were reactive to nociceptive assays were less responsive to morphine, and strains that were less responsive to nociceptive assays were more responsive to morphine.^{75,86} These results suggest the potential for mouse strains to differ in pain and/or analgesia sensitivity primarily due to genetic mechanisms.⁷⁵ Sensitivity to analgesics may also be influenced by a wide variety of other factors including age,³⁵ sex,⁸³ and laboratory environmental factors (for example diet, housing, stress).^{74,78,80} While the majority of strain surveys relating to analgesia centered around the evaluation of opioids (which will be covered in the “sex” section below), nonopioid analgesic responses also differed by strain, including electroacupuncture,¹²⁵ xylazine,⁶⁴ ketamine,⁶⁴ and over-the-counter medications (acetaminophen and NSAIDs).¹²⁸ Common mouse strains have also been surveyed for general anesthetic characterization and pain model development. For instance, heritable differences in respiratory drive and breathing pattern in C58BL/6, C3H/HeJ and B6C3F1/J strains of mice while under isoflurane anesthesia have been established.⁴⁰ Moreover, the effect of isoflurane anesthesia on the Mouse Grimace Scale in DBA and DBA/2 strains have been evaluated.⁷³

The work discussed in this section has the potential to positively impact the welfare of laboratory mice by providing targeted analgesia by strain and ensuring that the selection of an analgesic agent is appropriate for the strain and model being studied. A point of discussion in the review of these examples is the need for improved and consistent measures of pain in mice. Sources have demonstrated the variability in the assessment of rodent pain,¹⁶ especially at the strain level.^{128,129} The evaluation of contemporary methods of pain assessment has now been documented in laboratory animal literature.^{20,73,100,112} These adjuncts to typical nociception assays will enhance the body of literature and provide a more holistic and consistent approach to the assessment of rodent pain and analgesia.

C57BL/6 versus 129. The common use of knockout or transgenic mice in biomedical pain research requires the interpretation of the genetic background of the resulting strain. The genetically modified strain is almost always a mixture of alleles from 2 different strains, usually 129 or C57BL/6.⁶²

Of particular interest to pain researchers are the significant nociceptive and pain model differences identified between C57BL/6 and 129 inbred strains.^{13,15,58,69,85,92,93} (Table 1 through Table 3.) Significant differences between these 2 strains have also been demonstrated for sensitivity to analgesia.^{123,128,129} For example, early strain surveys demonstrated the C57BL/6 strain as one of the most sensitive inbred strains when tested on a battery of nociceptive assays.⁹² In the same survey, the C57BL/6 and 129 strains demonstrated significantly different responses to 8 of 12 nociceptive assays.⁹² Other laboratories have also established the nociceptive sensitivity of the C57BL/6 strain in inflammatory pain but not neuropathic or visceral pain conditions.⁶² Subsequent work on strain differences have highlighted the importance of substrain characterization, especially for the C57BL/6 and 129 strains.⁵⁶ For example, the C57BL/6J and C57BL/6N substrains vary in response to ethanol preference, conditioned fear, and pain sensitivity.^{15,56,69} This work is notable

because these strains are generally used as the default genetic background for null mutations.^{66,67}

Sex Differences in Response to Pain and Analgesia. Pain research has evolved from debating whether sex differences in the sensitivity to pain and analgesia exists to now recognizing the importance of these differences and how to understand their effects in human and animal models.^{38,95,103,126}

In humans, females are overrepresented in complex disease states such as autoimmune disorders and chronic pain.^{17,119} Differences between men and women regarding pain perception, tolerance and behavior, as well as the prevalence in seeking medical attention for the treatment of pain, are well known.¹⁰⁴ Pain studies in humans have also documented that variability exists in response to analgesia by sex, and that a portion of these variabilities are likely to have a heritable etiology.¹⁷ Historically, these variations in sex response to pain and analgesia and heritability have been studied independently;⁷⁷ however, contemporary pain research suggests that both play a fundamental role and should be studied concurrently.^{26,77,90}

In the mouse, sex differences have been demonstrated to significantly affect the response to nicotine,⁴³ Down’s Syndrome,⁸ neuroimmunity,^{114,115} obesity,¹¹³ diabetes,¹¹³ aging,¹¹³ cardiovascular health,⁷² liver disease,⁷² and cancer research.⁷²

Several anatomic/physiologic factors come into play when interpreting sex differences in regard to pain and analgesia. For example, adult male rodents have greater percentage of body fat than females³⁸ while the opposite is true for humans. This sexual dimorphism can affect the distribution of highly lipophilic drugs and influence analgesia potency, and duration of action. Similarly, metabolism, immune response, activity level, and response to analgesics all have a sex difference.^{2,8}

Sex differences to pain and analgesia in inbred mice are likely the result of genetic background, making variability by strain and sex important research considerations.^{9,76} However, the majority of animal subjects in pain research are overwhelmingly male,¹⁰⁹ which can lead to a bias in interpreting studies of animal pain.⁸²

In a recent review⁸³ of a 10-y history in the journal *Pain*, nearly 80% of publications included only male subjects. Conversely, 8% of the studies used only female subjects. Given that a greater percentage of clinical pain conditions affect women over men, and the growing supportive research on this topic, all pain research be performed in both sexes, and when only one can be used, it should be female.³⁸

The above point was further punctuated within the research community in 2014, when the United States National Institutes of Health (NIH) issued a requirement that grant applicants use both sexes of model organisms in their work.²¹ To date, strong justification from the scientific literature or preliminary data must be included in an application seeking to study only one sex²¹ to reduce the male bias of the study.¹⁰⁴

Examples of sex variation in common inbred mouse models of pain and analgesia include: morphine sensitivity,^{25,32,34,41,45,50,51,54,95} NMDA receptor antagonism,¹⁴ Kappa receptor sensitivity,¹⁸ response to nicotine,^{43,107,120} femoral cancer pain,³⁹ immune system,¹¹⁴ and sleep disruption.¹¹¹ In the evaluation of the sex variation studies in inbred mice described above, a portion of the research is conflicting, suggesting that in addition to sex variation, gonadal hormones should be evaluated.^{82,119}

Role of Gonadal Hormones. Animal studies have demonstrated variable results regarding the role of gonadal hormones on pain sensitivity and analgesic efficacy.^{24,104} Gonadal hormones are produced by the ovaries (estrogens and progesterins) and testes (androgens),²⁴ and pain researchers are primarily interested in the modulating effects of estradiol and testosterone.²⁴ The exact role

of gonadal hormones in pain are not well understood,⁴⁵ however they are suspected to affect pain and analgesia either activationally (in adulthood) or organizationally (in development).^{5,24}

A common method to evaluate gonadal hormones and their research effect is to include female subjects at different stages of the estrous cycle.³⁸ In humans, sex hormones have been shown to influence pain sensitivity, pain threshold, and pain tolerance and vary with stage of the menstrual cycle.¹²⁶ In mice, estrous cycle has been demonstrated to impact pain and symptom severity on a multiple sclerosis model;¹⁰⁶ however, the value of testing female mice at different stages of the estrous cycle for pain research is still debatable.^{38,119} While testing mice for pain during the entire estrous cycle is interesting; it can be prohibitive due to the large numbers of animals needed to adequately power the analysis.⁸³ In addition, the task of assigning the correct stage of estrus requires extra handling and vaginal cytology, which could confound the study.³⁸ To improve research reproducibility, it would be useful for the study design to include the evaluation of the stage of estrus. An advancement in this approach is the use of gonadectomy with or without hormone depletion and/or replacement.³⁸

Gonadectomy can be performed via surgery (most common approach) or via “chemical castration.” The surgical method, although useful in identifying the effects of hormones on pain and analgesia in animals, has drawbacks including alteration of pain thresholds and sensitivity to analgesic agents, and the disruption of normal hormonal feedback loop, potentially causing both males and females to have elevated or depressed circulating hormones.³⁸ The most common method of using gonadectomy in female mouse models of pain and analgesia is to provide hormone replacement (that is estradiol) after gonadectomy. While this is a popular approach, factors such as strain differences, inconsistent dose range, and age differences, can introduce variations in results.¹¹⁹

The Role of Sex-Based Factors Specific to Opioids. Similar to the concept of sex differences in nociception, studies have also shown a relationship between sex differences and sensitivity to analgesics.²³ The largest body of work on this subject involves opioid analgesia, specifically morphine, whose action is defined as a μ receptor agonist.⁵⁰⁻⁵⁴ Both humans and animals display variable responses to morphine sensitivity,⁵⁴ and some of this variability may be attributable to sex.⁵¹ Review of early pain research specific to mice documents that male and female mice differ in their sensitivity to morphine;^{50,76} however, the lack of consistent findings makes a broad statement of strain and morphine sensitivity problematic.⁵⁴ Rather, it is now generally believed that the variability in male and female mice to the effects of morphine are likely the result of several mechanisms, including strain differences.^{14,23,32,41,51,76}

Multiple studies have documented that sex affects opioid analgesia,²⁵ but the exact mechanism is unknown and is likely due to multiple factors. In mice, opioids can be more potent in male compared with female mice when given systemically,^{23,76} which may be due to the variable degree of morphine tolerance existing between male and females.

In addition to differences in sensitivity, analgesic responses to opioid classes also demonstrate a sex predilection (for example κ opioid analgesics).^{23,47} κ -Opioid analgesia has been further evaluated on the effects of detromethorphan (DEX) N-methyl-D-aspartate (NMDA) antagonist,^{47,77} and melanocortin-1-receptor⁹⁴ abilities to potentiate analgesia, especially in females.^{47,77}

Other factors

The nature of this broad literature review does not allow for detailed consideration of all of the factors proposed to influence

the effect of strain and/or sex on the response to pain and/or analgesia in the mouse. However, several potential factors have been discussed in this review. One potentially influential factor in the study of mouse pain could be age. In humans, gender differences in pain sensitivities emerge during adolescence, with females experiencing an increase during puberty while their male counterparts remain stable.⁶³ In rodents, the impact of advancing age on the biologic systems responsible for experiencing pain have not been fully established.^{132,133} Unfortunately, the majority of the studies reviewed for this manuscript did not include age as a factor in the study design. Another factor that could affect the response of mice to pain and/or analgesia is the source of the strain used. This review has identified and tabulated the strains surveyed based on the information and nomenclature included in the study design of each reference (Table 1 through Table 3). In many cases, the referenced manuscripts did not include the substrain or the source of the strain(s) of mice. To more clearly understand the effects that age or source of strain could have on mice in relation to pain research, future study designs must provide comprehensive detail on animal subjects and their background.

Conclusions

The use of mice in the study of pain and analgesia will likely continue, although should be enhanced to include updated modalities in human medicine (for example imaging, genetics).^{66,84} Studies examining nociception assays have established performance differences within common mouse strains,^{92,93,128,129} many of which can also be influenced by sex. Similarly, surveys of common mouse strain have been used to evaluate the differences of genes and sex on the sensitivity to analgesia.^{128,129} Collectively, this review establishes that over the past several decades, genetic heritability and sex are well documented to influence the experience of pain and response to analgesia in the laboratory mouse; however, the mechanisms of action and relationship to complex interactions are still under investigation.

Refinements Needed. Rice and colleagues¹⁰⁹ articulately describe key enhancements needed by the research community regarding animal models of pain and lists the choice of animal as one of the key factors for reducing experimental bias. These concepts, punctuated by the call to reduce the overwhelming male-dominated bias,¹⁰⁹ and to minimize the effects of laboratory environmental factors,⁸⁰ remain some of the hallmark goals for animal models of pain. Ensuring research transparency and minimizing method bias in the laboratory animal community¹⁶ will, in addition, improve the pain and analgesia related work.

To assess this concept in the laboratory animal literature, a survey of the last 5 y of literature published by AALAS journals¹²² (*JAALAS*, and *Comparative Medicine*) revealed that roughly half of the submissions relating to mouse anesthesia and/or analgesia included both male and female subjects.^{12,31,49,64,100,116,130} In only one case⁵⁵ was a justification for single-sex study based on published literature included in the Materials section. The deficiency in dual sex inclusion underscores the need to improve the study design of all laboratory animal studies, specifically those relating to pain and analgesic response.

Recent work in the laboratory animal community provides new tools to assess both pain and analgesia in mice. Evaluation of nest building behaviors,¹¹² species-specific cage side assessments,¹⁰⁰ and the Mouse Grimace Scale²⁰ are all examples of novel methods for assessing pain. These methods can be used to characterize strain and sex differences and should be considered for use in order to obtain accurate assessment of pain and analgesic agents in future studies.

Reproducibility of pain research relating to animals can only be achieved when these complex interactions are accurately included in the study design. This review was undertaken to solidify these important concepts and provide a workable summary for the laboratory animal community to reference.

References

- Allayee H, Andalibi A, Mehrabian M. 2006. Using inbred mouse strains to identify genes for complex diseases. *Front Biosci* **11**:1216–1226. <https://doi.org/10.2741/1874>.
- Banik RK, Woo YC, Park SS, Brennan TJ. 2006. Strain and sex influence on pain sensitivity after plantar incision in the mouse. *Anesthesiology* **105**:1246–1253. <https://doi.org/10.1097/00000542-200612000-00025>.
- Barrot M. 2012. Tests and models of nociception and pain in rodents. *Neuroscience* **211**:39–50. <https://doi.org/10.1016/j.neuroscience.2011.12.041>.
- Beck JA, Lloyd S, Hafezparast M, Lennon-Pierce M, Eppig JT, Festing MF, Fisher EM. 2000. Genealogies of mouse inbred strains. *Nat Genet* **24**:23–25. <https://doi.org/10.1038/71641>.
- Becker JB, Arnold AP, Berkley KJ, Blaustein JD, Eckel LA, Hampson E, Herman JP, Marts S, Sadee W, Steiner M, Taylor J, Young E. 2005. Strategies and methods for research on sex differences in brain and behavior. *Endocrinology* **146**:1650–1673. <https://doi.org/10.1210/en.2004-1142>.
- Benedetti M, Merino R, Kusuda R, Ravanelli MI, Cadetti F, dos Santos P, Zanon S, Lucas G. 2011. Plasma corticosterone levels in mouse models of pain. *Eur J Pain* **16**:803–815. <https://doi.org/10.1002/j.1532-2149.2011.00066.x>.
- Blennerhassett MG, Lourens SR, Parlow LRG, Ghasemlou N, Winterborn AN. 2017. Analgesia and mouse strain influence neuromuscular plasticity in inflamed intestine. *Neurogastroenterol Motil* **29**:1–12. <https://doi.org/10.1111/nmo.13097>.
- Block A, Ahmed MM, Dhanasekaran AR, Tong S, Gardiner KJ. 2015. Sex differences in protein expression in the mouse brain and their perturbations in a model of Down syndrome. *Biol Sex Differ* **6**:1–18. <https://doi.org/10.1186/s13293-015-0043-9>.
- Boerner KE, Chambers CT, Gahagan J, Keogh E, Fillingim RB, Mogil JS. 2018. Conceptual complexity of gender and its relevance to pain. *Pain* **159**:2137–2141. <https://doi.org/10.1097/j.pain.0000000000001275>.
- Bon K, Lichtensteiger CA, Wilson SG, Mogil J. 2003. Characterization of cyclophosphamide cystitis, a model of visceral and referred pain, in the mouse: species and strain differences. *J Urol* **170**:1008–1012. <https://doi.org/10.1097/01.ju.0000079766.49550.94>.
- Bonaterra GA, Then H, Oezel L, Schwarzbach H, Ocker M, Thieme K, Di Fazio P, Kinscherf R. 2016. Morphological alterations in gastrocnemius and soleus muscles in male and female mice in a fibromyalgia model. *PLoS One* **11**:1–18. <https://doi.org/10.1371/journal.pone.0151116>.
- Braden GC, Brice AK, Hankenson FC. 2015. Adverse effects of vapocoolant and topical anesthesia for tail biopsy of preweaning mice. *J Am Assoc Lab Anim Sci* **54**:291–298.
- Bryant CD, Bagdas D, Goldberg LR, Khalefa T, Reed ER, Kirkpatrick SL, Kelliher JC, Chen MM, Johnson WE, Mulligan MK, Imad Damaj M. 2019. C57BL/6 substrain differences in inflammatory and neuropathic nociception and genetic mapping of a major quantitative trait locus underlying acute thermal nociception. *Mol Pain* **15**:1–15. <https://doi.org/10.1177/1744806918825046>.
- Bryant CD, Eitan S, Sinchak K, Fanselow MS, Evans CJ. 2006. NMDA receptor antagonism disrupts the development of morphine analgesic tolerance in male, but not female C57BL/6J mice. *Am J Physiol Regul Integr Comp Physiol* **291**:R315–R326. <https://doi.org/10.1152/ajpregu.00831.2005>.
- Bryant CD, Zhang NN, Sokoloff G, Fanselow MS, Ennes HS, Palmer AA, McRoberts JA. 2008. Behavioral differences among C57BL/6 substrains: implications for transgenic and knockout studies. *J Neurogenet* **22**:315–331. <https://doi.org/10.1080/01677060802357388>.
- Carbone L, Austin J. 2016. Pain and laboratory animals: publication practices for better data reproducibility and better animal welfare. *PLoS One* **11**:1–24. <https://doi.org/10.1371/journal.pone.0155001>.
- Chanda ML, Mogil JS. 2006. Sex differences in the effects of amiloride on formalin test nociception in mice. *Am J Physiol Regul Integr Comp Physiol* **291**:R335–R342. <https://doi.org/10.1152/ajpregu.00902.2005>.
- Chavkin C, Cohen JH, Land BB. 2019. Repeated administration of norbinaltorphimine produces cumulative k opioid receptor inactivation. *Front Pharmacol* **10**:1–5. <https://doi.org/10.3389/fphar.2019.00088>.
- Chesler EJ, Ritchie J, Kokayeff A, Lariviere WR, Wilson SG, Mogil JS. 2003. Genotype-dependence of gabapentin and pregabalin sensitivity: the pharmacogenetic mediation of analgesia is specific to the type of pain being inhibited. *Pain* **106**:325–335. [https://doi.org/10.1016/S0304-3959\(03\)00330-0](https://doi.org/10.1016/S0304-3959(03)00330-0).
- Cho C, Michalidis V, Lecker I, Collymore C, Hanwell D, Loka M, Danesh M, Pham C, Urban P, Bonin RP, Martin LJ. 2019. Evaluating analgesic efficacy and administration route following craniotomy in mice using the grimace scale. *Sci Rep* **9**:1–9. <https://doi.org/10.1038/s41598-018-36897-w>.
- Collins FS, Tabak LA. 2014. Policy: NIH plans to enhance reproducibility. *Nature* **505**:612–613. <https://doi.org/10.1038/505612a>.
- Crabbe JC, Wahlsten D, Dudek BC. 1999. Genetics of mouse behavior: interactions with laboratory environment. *Science* **284**:1670–1672. <https://doi.org/10.1126/science.284.5420.1670>.
- Craft RM. 2003. Sex differences in opioid analgesia: “from mouse to man. *Clin J Pain* **19**:175–186. <https://doi.org/10.1097/00002508-200305000-00005>.
- Craft RM, Mogil JS, Aloisi AM. 2004. Sex differences in pain and analgesia: the role of gonadal hormones. *Eur J Pain* **8**:397–411. <https://doi.org/10.1016/j.ejpain.2004.01.003>.
- Dahan A, Kest B, Waxman AR, Sarton E. 2008. Sex-specific responses to opiates: animal and human studies. *Anesth Analg* **107**:83–95. <https://doi.org/10.1213/ane.0b013e31816a66a4>.
- Dahlhamer J, Lucas J, Zelaya C, Nahin R, Mackey S, DeBar L, Kerns R, Von Korff M, Porter L, Helmick C. 2018. Prevalence of chronic pain and high-impact chronic pain among adults—United States, 2016. *MMWR Morb Mortal Wkly Rep* **67**:1001–1006. <https://doi.org/10.15585/mmwr.mm6736a2>.
- Dickinson AL, Leach MC, Flecknell PA. 2009. The analgesic effects of oral paracetamol in 2 strains of mice undergoing vasectomy. *Lab Anim* **43**:357–361. <https://doi.org/10.1258/la.2009.009005>.
- Dodds KN, Beckett EAH, Evans SF, Hutchinson MR. 2017. Lesion development is modulated by the natural estrous cycle and mouse strain in a minimally invasive model of endometriosis. *Biol Reprod* **97**:810–821. <https://doi.org/10.1093/biolre/iox132>.
- Duman EN, Kesim M, Kadioglu M, Ulku C, Kalyoncu NI, Yaris E. 2006. Effect of gender on antinociceptive effect of paroxetine in hot plate test in mice. *Prog Neuropsychopharmacol Biol Psychiatry* **30**:292–296. <https://doi.org/10.1016/j.pnpbp.2005.10.012>.
- Elhabazi K, Ayachi S, Ilien B, Simonin F. 2014. Assessment of morphine-induced hyperalgesia and analgesic tolerance in mice using thermal and mechanical nociceptive modalities. *J Vis Exp* **89**:1–10.
- Erickson RL, Blevins CE, Souza Dyer C, Marx JO. 2019. Alfa-xalone-Xylazine anesthesia in laboratory mice (*Mus musculus*). *J Am Assoc Lab Anim Sci* **58**:30–39. <https://doi.org/10.30802/AALAS-JAALAS-18-000010>.
- Fillingim RB, Gear RW. 2004. Sex differences in opioid analgesia: clinical and experimental findings. *Eur J Pain* **8**:413–425. <https://doi.org/10.1016/j.ejpain.2004.01.007>.
- Fischer BD, Adeyemo A, O’Leary ME, Bottaro A. 2017. Animal models of rheumatoid pain: experimental systems and insights. *Arthritis Res Ther* **19**:1–9. <https://doi.org/10.1186/s13075-017-1361-6>.
- Gioiosa L, Chen X, Watkins R, Klanfer N, Bryant CD, Evans CJ, Arnold AP. 2008. Sex chromosome complement affects nociception in tests of acute and chronic exposure to morphine in mice. *Horm Behav* **53**:124–130. <https://doi.org/10.1016/j.yhbeh.2007.09.003>.
- Gioiosa L, Chen X, Watkins R, Umeda EA, Arnold AP. 2008. Sex chromosome complement affects nociception and analgesia

- in newborn mice. *J Pain* 9:962–969. <https://doi.org/10.1016/j.jpain.2008.06.001>.
36. **Green AD, Young KK, Lehto SG, Smith SB, Mogil JS.** 2006. Influence of genotype, dose and sex on pruritogen-induced scratching behavior in the mouse. *Pain* 124:50–58. <https://doi.org/10.1016/j.pain.2006.03.023>.
37. **Green ED, Watson JD, Collins FS.** 2015. Human genome project: twenty-five years of big biology. *Nature* 526:29–31. <https://doi.org/10.1038/526029a>.
38. **Greenspan JD, Craft RM, LeResche L, Arendt-Nielsen L, Berkley KJ, Fillingim RB, Gold MS, Holdcroft A, Lautenbacher S, Mayer EA, Mogil JS, Murphy AZ, Traub RJ: Consensus Working Group of the Sex, Gender, and Pain SIG of the IASP.** 2007. Studying sex and gender differences in pain and analgesia: a consensus report. *Pain* 132 Suppl 1:S26–S45. <https://doi.org/10.1016/j.pain.2007.10.014>.
39. **Gregory NS, Gibson-Corley K, Frey-Law L, Sluka KA.** 2013. Fatigue-enhanced hyperalgesia in response to muscle insult: induction and development occur in a sex-dependent manner. *Pain* 154:2668–2676. <https://doi.org/10.1016/j.pain.2013.07.047>.
40. **Groeben H, Meier S, Tankersley CG, Mitzner W, Brown RH.** 2003. Heritable differences in respiratory drive and breathing pattern in mice during anaesthesia and emergence. *Br J Anaesth* 91:541–545. <https://doi.org/10.1093/bja/aeg222>.
41. **Hopkins E, Rossi G, Kest B.** 2004. Sex differences in systemic morphine analgesic tolerance following intrathecal morphine injections. *Brain Res* 1014:244–246. <https://doi.org/10.1016/j.brainres.2004.03.056>.
42. **Isami K, Imai S, Sukeishi A, Nagayasu K, Shirakawa H, Nakagawa T, Kaneko S.** 2018. The impact of mouse strain-specific spatial and temporal immune responses on the progression of neuropathic pain. *Brain Behav Immun* 74:121–132. <https://doi.org/10.1016/j.bbi.2018.08.013>.
43. **Isiegas C, Mague SD, Blendy JA.** 2009. Sex differences in response to nicotine in C57Bl/6:129SvEv mice. *Nicotine Tob Res* 11:851–858. <https://doi.org/10.1093/ntr/ntp076>.
44. **Jensen TS, Finnerup NB.** 2014. Allodynia and hyperalgesia in neuropathic pain: clinical manifestations and mechanisms. *Lancet Neurol* 13:924–935. [https://doi.org/10.1016/S1474-4422\(14\)70102-4](https://doi.org/10.1016/S1474-4422(14)70102-4).
45. **Juni A, Klein G, Kowalczyk B, Ragnauth A, Kest B.** 2008. Sex differences in hyperalgesia during morphine infusion: effect of gonadectomy and estrogen treatment. *Neuropharmacology* 54:1264–1270. <https://doi.org/10.1016/j.neuropharm.2008.04.004>.
46. **Kamp EH, Jones RC 3rd, Tillman SR, Gebhart GF.** 2003. Quantitative assessment and characterization of visceral nociception and hyperalgesia in mice. *Am J Physiol Gastrointest Liver Physiol* 284:G434–G444. <https://doi.org/10.1152/ajpgi.00324.2002>.
47. **Kavaliers M, Colwell DD.** 1991. Sex differences in opioid and non-opioid mediated predator-induced analgesia in mice. *Brain Res* 568:173–177. [https://doi.org/10.1016/0006-8993\(91\)91394-G](https://doi.org/10.1016/0006-8993(91)91394-G).
48. **Kc M, Steer CJ.** 2019. A new era of gene editing for the treatment of human diseases. *Swiss Med Wkly* 149:w20021.
49. **Kennedy LH, Hwang H, Wolfe AM, Hauptman J, Nemzek-Hamlin JA.** 2014. Effects of buprenorphine and estrous cycle in a murine model of cecal ligation and puncture. *Comp Med* 64:270–282.
50. **Kest B, Hopkins E.** 2001. Morphine tolerance after chronic intracerebroventricular injection in male and female mice. *Brain Res* 892:208–210. [https://doi.org/10.1016/S0006-8993\(00\)03301-1](https://doi.org/10.1016/S0006-8993(00)03301-1).
51. **Kest B, Palmese C, Hopkins E.** 2000. A comparison of morphine analgesic tolerance in male and female mice. *Brain Res* 879:17–22. [https://doi.org/10.1016/S0006-8993\(00\)02685-8](https://doi.org/10.1016/S0006-8993(00)02685-8).
52. **Kest B, Palmese CA, Hopkins E, Adler M, Juni A, Mogil JS.** 2002. Naloxone-precipitated withdrawal jumping in 11 inbred mouse strains: evidence for common genetic mechanisms in acute and chronic morphine physical dependence. *Neuroscience* 115:463–469. [https://doi.org/10.1016/S0306-4522\(02\)00458-X](https://doi.org/10.1016/S0306-4522(02)00458-X).
53. **Kest B, Sarton E, Dahan A.** 2000. Gender differences in opioid-mediated analgesia: animal and human studies. *Anesthesiology* 93:539–547. <https://doi.org/10.1097/00000542-200008000-00034>.
54. **Kest B, Wilson SG, Mogil JS.** 1999. Sex differences in supraspinal morphine analgesia are dependent on genotype. *J Pharmacol Exp Ther* 289:1370–1375.
55. **Kick BL, Shu P, Wen B, Sun D, Taylor DK.** 2017. Pharmacokinetic profiles of Nalbuphine after intraperitoneal and subcutaneous administration to C57BL/6 mice. *J Am Assoc Lab Anim Sci* 56:534–538.
56. **Kiselycznyk C, Holmes A.** 2011. All (C57BL/6) mice are not created equal. *Front Neurosci* 5:1–3. <https://doi.org/10.3389/fnins.2011.00010>.
57. **Lacroix-Fralish ML, Ledoux JB, Mogil JS.** 2007. The pain genes database: an interactive web browser of pain-related transgenic knockout studies. *Pain* 131:3.e1–3.e4. <https://doi.org/10.1016/j.pain.2007.04.041>.
58. **Lariviere WR, Chesler EJ, Mogil JS.** 2001. Transgenic studies of pain and analgesia: mutation or background genotype? *J Pharmacol Exp Ther* 297:467–473.
59. **Lariviere WR, Mogil JS.** 2010. The genetics of pain and analgesia in laboratory animals. *Methods Mol Biol* 617:261–278. https://doi.org/10.1007/978-1-60327-323-7_20.
60. **Lariviere WR, Wilson SG, Laughlin TM, Kokayeff A, West EE, Adhikari SM, Wan Y, Mogil JS.** 2002. Heritability of nociception. III. Genetic relationships among commonly used assays of nociception and hypersensitivity. *Pain* 97:75–86. [https://doi.org/10.1016/S0304-3959\(01\)00492-4](https://doi.org/10.1016/S0304-3959(01)00492-4).
61. **Leite-Almeida H, Pinto-Ribeiro F, Almeida A.** 2015. Animal models for the study of comorbid pain and psychiatric disorders. *Mod Trends Pharmacopsychiatry* 30:1–21. <https://doi.org/10.1159/000435929>.
62. **Leo S, Straetemans R, D’Hooge R, Meert T.** 2008. Differences in nociceptive behavioral performance between C57BL/6J, 129S6/SvEv, B6 129 F1 and NMRI mice. *Behav Brain Res* 190:233–242. <https://doi.org/10.1016/j.bbr.2008.03.001>.
63. **LeResche L.** 2011. Defining gender disparities in pain management. *Clin Orthop Relat Res* 469:1871–1877. <https://doi.org/10.1007/s11999-010-1759-9>.
64. **Levin-Arama M, Abraham L, Waner T, Harmelin A, Steinberg DM, Lahav T, Harlev M.** 2016. Subcutaneous compared with intraperitoneal KetamineXylazine for anesthesia of mice. *J Am Assoc Lab Anim Sci* 55:794–800.
65. **Liang DY, Liao G, Wang J, Usuka J, Guo Y, Peltz G, Clark JD.** 2006. A genetic analysis of opioid-induced hyperalgesia in mice. *Anesthesiology* 104:1054–1062. <https://doi.org/10.1097/00000542-200605000-00023>.
66. **Liu ET, Bolcun-Filas E, Grass DS, Lutz C, Murray S, Shultz L, Rosenthal N.** 2017. Of mice and CRISPR: The post-CRISPR future of the mouse as a model system for the human condition. *EMBO Rep* 18:187–193. <https://doi.org/10.15252/embr.201643717>.
67. **Lötsch J, Doebering A, Mogil JS, Arndt T, Geisslinger G, Ultsch A.** 2013. Functional genomics of pain in analgesic drug development and therapy. *Pharmacol Ther* 139:60–70. <https://doi.org/10.1016/j.pharmthera.2013.04.004>.
68. **Lumley MA, Cohen JL, Borszcz GS, Cano A, Radcliffe AM, Porter LS, Schubiner H, Keefe FJ.** 2011. Pain and emotion: a biopsychosocial review of recent research. *J Clin Psychol* 67:942–968. <https://doi.org/10.1002/jclp.20816>.
69. **Matsuo N, Takao K, Nakanishi K, Yamasaki N, Tanda K, Miyakawa T.** 2010. Behavioral profiles of three C57BL/6 substrains. *Front Behav Neurosci* 4:1–12.
70. **Megat S, Shiers S, Moy JK, Barragan-Iglesias P, Pradhan G, Seal RP, Dussor G, Price TJ.** 2017. A critical role for Dopamine D5 receptors in pain chronicity in male mice. *J Neurosci* 38:379–397. <https://doi.org/10.1523/JNEUROSCI.2110-17.2017>.
71. **Meints SM, Edwards RR.** 2018. Evaluating psychosocial contributions to chronic pain outcomes. *Prog Neuropsychopharmacol Biol Psychiatry* 87:168–182. <https://doi.org/10.1016/j.pnpbp.2018.01.017>.
72. **Miaskowski CG, Levine JD.** 2000. Sex-related differences in analgesic responses, p 209–230. In: Fillingim RB, editor. *Progress in pain research and management*, vol 17. Seattle (WA): IASP Press.
73. **Miller A, Kitson G, Skalkoyannis B, Leach M.** 2015. The effect of isoflurane anaesthesia and buprenorphine on the mouse grimace scale and behaviour in CBA and DBA/2 mice. *Appl Anim Behav Sci* 172:58–62. <https://doi.org/10.1016/j.applanim.2015.08.038>.

74. Minett MS, Eijkelkamp N, Wood JN. 2014. Significant determinants of mouse pain behaviour. *PLoS One* **9**:1–6. <https://doi.org/10.1371/journal.pone.0104458>.
75. Mogil JS. 1999. The genetic mediation of individual differences in sensitivity to pain and its inhibition. *Proc Natl Acad Sci USA* **96**:7744–7751. <https://doi.org/10.1073/pnas.96.14.7744>.
76. Mogil JS. 2012. Interactions between sex and genotype in the mediation and modulation of nociception in rodents, p 25–40 In: Fillingim RB, editor. *Progress in pain research and management*, vol 17. Seattle (WA): IASP Press.
77. Mogil JS. 2003. Interaction between sex and genotype in the mediation of pain and pain inhibition. *Seminars in Pain Medicine* **1**:197–205. [https://doi.org/10.1016/S1537-5897\(03\)00028-4](https://doi.org/10.1016/S1537-5897(03)00028-4).
78. Mogil JS. 2012. Pain genetics: past, present and future. *Trends Genet* **28**:258–266. <https://doi.org/10.1016/j.tig.2012.02.004>.
79. Mogil JS. 2012. Sex differences in pain and pain inhibition: multiple explanations of a controversial phenomenon. *Nat Rev Neurosci* **13**:859–866. <https://doi.org/10.1038/nrn3360>.
80. Mogil JS. 2017. Laboratory environmental factors and pain behavior: the relevance of unknown unknowns to reproducibility and translation. *Lab Anim (NY)* **46**:136–141. <https://doi.org/10.1038/labani.1223>.
81. Mogil JS, Adhikari SM. 1999. Hot and cold nociception are genetically correlated. *J Neurosci* **19**:RC25. <https://doi.org/10.1523/JNEUROSCI.19-18-j0003.1999>.
82. Mogil JS, Bailey AL. 2010. Sex and gender differences in pain and analgesia. *Prog Brain Res* **186**:141–157.
83. Mogil JS, Chanda ML. 2005. The case for the inclusion of female subjects in basic science studies of pain. *Pain* **117**:1–5. <https://doi.org/10.1016/j.pain.2005.06.020>.
84. Mogil JS, Davis KD, Derbyshire SW. 2010. The necessity of animal models in pain research. *Pain* **151**:12–17. <https://doi.org/10.1016/j.pain.2010.07.015>.
85. Mogil JS, Grisel JE. 1998. Transgenic studies of pain. *Pain* **77**:107–128. [https://doi.org/10.1016/S0304-3959\(98\)00093-1](https://doi.org/10.1016/S0304-3959(98)00093-1).
86. Mogil JS, Kest B, Sadowski B, Belknap JK. 1996. Differential genetic mediation of sensitivity to morphine in genetic models of opiate antinociception: influence of nociceptive assay. *J Pharmacol Exp Ther* **276**:532–544.
87. Mogil JS, Lichtensteiger CA, Wilson SG. 1998. The effect of genotype on sensitivity to inflammatory nociception: characterization of resistant (A/J) and sensitive (C57BL/6J) inbred mouse strains. *Pain* **76**:115–125. [https://doi.org/10.1016/S0304-3959\(98\)00032-3](https://doi.org/10.1016/S0304-3959(98)00032-3).
88. Mogil JS, Ritchie J, Sotocinal SG, Smith SB, Croteau S, Levitin DJ, Naumova AK. 2006. Screening for pain phenotypes: analysis of 3 congenic mouse strains on a battery of nine nociceptive assays. *Pain* **126**:24–34. <https://doi.org/10.1016/j.pain.2006.06.004>.
89. Mogil JS, Simmonds K, Simmonds MJ. 2009. Pain research from 1975 to 2007: a categorical and bibliometric meta-trend analysis of every Research Paper published in the journal, *Pain*. *Pain* **142**:48–58. <https://doi.org/10.1016/j.pain.2008.11.012>.
90. Mogil JS, Sorge RE, LaCroix-Fralish ML, Smith SB, Fortin A, Sotocinal SG, Ritchie J, Austin JS, Schorscher-Petcu A, Melmed K, Czereminski J, Bittong RA, Mokris JB, Neubert JK, Campbell CM, Edwards RR, Campbell JN, Crawley JN, Lariviere WR, Wallace MR, Sternberg WF, Balaban CD, Belfer I, Fillingim RB. 2011. Pain sensitivity and vasopressin analgesia are mediated by a gene-sex-environment interaction. *Nat Neurosci* **14**:1569–1573. <https://doi.org/10.1038/nn.2941>.
91. Mogil JS, Sternberg WF, Marek P, Sadowski B, Belknap JK, Liebeskind JC. 1996. The genetics of pain and pain inhibition. *Proc Natl Acad Sci USA* **93**:3048–3055. <https://doi.org/10.1073/pnas.93.7.3048>.
92. Mogil JS, Wilson SG, Bon K, Lee SE, Chung K, Raber P, Pieper JO, Hain HS, Belknap JK, Hubert L, Elmer GI, Chung JM, Devor M. 1999. Heritability of nociception I: responses of 11 inbred mouse strains on 12 measures of nociception. *Pain* **80**:67–82. [https://doi.org/10.1016/S0304-3959\(98\)00197-3](https://doi.org/10.1016/S0304-3959(98)00197-3).
93. Mogil JS, Wilson SG, Bon K, Lee SE, Chung K, Raber P, Pieper JO, Hain HS, Belknap JK, Hubert L, Elmer GI, Chung JM, Devor M. 1999. Heritability of nociception II. 'Types' of nociception revealed by genetic correlation analysis. *Pain* **80**:83–93. [https://doi.org/10.1016/S0304-3959\(98\)00196-1](https://doi.org/10.1016/S0304-3959(98)00196-1).
94. Mogil JS, Wilson SG, Chesler EJ, Rankin AL, Nemmani KVS, Lariviere WR, Groce MK, Wallace MR, Kaplan L, Staud R, Ness TJ, Glover TL, Stankova M, Mayorov A, Hrubby VJ, Grisel JE, Fillingim RB. 2003. The melanocortin-1 receptor gene mediates female-specific mechanisms of analgesia in mice and humans. *Proc Natl Acad Sci USA* **100**:4867–4872. <https://doi.org/10.1073/pnas.0730053100>.
95. Mogil JS, Kest B. 1999. Sex differences in opioid analgesia: of mice and women. *Pain Forum* **8**:48–50.
96. Moloney RD, Dinan TG, Cryan JF. 2015. Strain-dependent variations in visceral sensitivity: relationship to stress, anxiety and spinal glutamate transporter expression. *Genes Brain Behav* **14**:319–329. <https://doi.org/10.1111/gbb.12216>.
97. Nahin RL. 2015. Estimates of pain prevalence and severity in adults: United States, 2012. *J Pain* **16**:769–780. <https://doi.org/10.1016/j.jpain.2015.05.002>.
98. National Research Council Committee on Recognition and Alleviation of Pain in Laboratory Animals. 2009. *Recognition and alleviation of pain in laboratory animals*. Washington (DC): National Academies Press.
99. Neilan CL, King MA, Rossi G, Ansonoff M, Pintar JE, Schiller PW, Pasternak GW. 2003. Differential sensitivities of mouse strains to morphine and [Dmt1]DALDA analgesia. *Brain Res* **974**:254–257. [https://doi.org/10.1016/S0006-8993\(03\)02590-3](https://doi.org/10.1016/S0006-8993(03)02590-3).
100. Oliver VL, Thurston SE, Lofgren JL. 2018. Using cageside measures to evaluate analgesic efficacy in mice (*Mus musculus*) after surgery. *J Am Assoc Lab Anim Sci* **57**:186–201.
101. Ono K, Ye Y, Viet CT, Dang D, Schmidt BL. 2015. TRPV1 expression level in isolectin B(4)-positive neurons contributes to mouse strain difference in cutaneous thermal nociceptive sensitivity. *J Neurophysiol* **113**:3345–3355. <https://doi.org/10.1152/jn.00973.2014>.
102. Packiasabapathy S, Sadhasivam S. 2018. Gender, genetics, and analgesia: understanding the differences in response to pain relief. *J Pain Res* **11**:2729–2739. <https://doi.org/10.2147/JPR.S94650>.
103. Paller CJ, Campbell CM, Edwards RR, Dobs AS. 2009. Sex-based differences in pain perception and treatment. *Pain Med* **10**:289–299. <https://doi.org/10.1111/j.1526-4637.2008.00558.x>.
104. Palmeira CC, Ashmawi HA, Posso Ide P. 2011. [Sex and pain perception and analgesia.] *Rev Bras Anestesiol* **61**:814–828. [Article in English, multiple languages]. [https://doi.org/10.1016/S0034-7094\(11\)70091-5](https://doi.org/10.1016/S0034-7094(11)70091-5).
105. Pitcher MH, Von Korff M, Bushnell MC, Porter L. 2019. Prevalence and profile of high-impact chronic pain in the United States. *J Pain* **20**:146–160. <https://doi.org/10.1016/j.jpain.2018.07.006>.
106. Rahn EJ, Iannitti T, Donahue RR, Taylor BK. 2014. Sex differences in a mouse model of multiple sclerosis: neuropathic pain behavior in females but not males and protection from neurological deficits during proestrus. *Biol Sex Differ* **5**:4. <https://doi.org/10.1186/2042-6410-5-4>.
107. Rasakham K, Liu-Chen LY. 2011. Sex differences in k opioid pharmacology. *Life Sci* **88**:2–16. <https://doi.org/10.1016/j.lfs.2010.10.007>.
108. Recla JM, Robledo RF, Gatti DM, Bult CJ, Churchill GA, Chesler EJ. 2014. Precise genetic mapping and integrative bioinformatics in Diversity Outbred mice reveals Hydin as a novel pain gene. *Mamm Genome* **25**:211–222. <https://doi.org/10.1007/s00335-014-9508-0>.
109. Rice AS, Cimino-Brown D, Eisenach JC, Kontinen VK, Lacroix-Fralish ML, Machin I, Preclinical Pain C, Mogil JS, Stöhr T. 2008. Animal models and the prediction of efficacy in clinical trials of analgesic drugs: a critical appraisal and call for uniform reporting standards. *Pain* **139**:243–247. <https://doi.org/10.1016/j.pain.2008.08.017>.
110. Rigaud M, Gemes G, Barabas ME, Chernoff DI, Abram SE, Stucky CL, Hogan QH. 2008. Species and strain differences in rodent sciatic nerve anatomy: implications for studies of neuropathic pain. *Pain* **136**:188–201. <https://doi.org/10.1016/j.pain.2008.01.016>.
111. Robinson-Junker A, O'Hara B, Durkes A, Gaskill B. 2019. Sleeping through anything: The effects of unpredictable disruptions on mouse sleep, healing, and affect. *PLoS One* **14**:1–17. <https://doi.org/10.1371/journal.pone.0210620>.

112. **Rock ML, Karas AZ, Rodriguez KB, Gallo MS, Pritchett-Corning K, Karas RH, Aronovitz M, Gaskill BN.** 2014. The time-to-integrate-to-nest test as an indicator of wellbeing in laboratory mice. *J Am Assoc Lab Anim Sci* 53:24–28.
113. **Rodgers HM, Liban S, Wilson LM.** 2014. Attenuated pain response of obese mice (B6.Cg-lep(ob)) is affected by aging and leptin but not sex. *Physiol Behav* 123:80–85. <https://doi.org/10.1016/j.physbeh.2013.10.007>.
114. **Rosen S, Ham B, Mogil JS.** 2017. Sex differences in neuroimmunity and pain. *J Neurosci Res* 95:500–508. <https://doi.org/10.1002/jnr.23831>.
115. **Rosen SF, Ham B, Haichin M, Walters IC, Tohyama S, Sotocinal SG, Mogil JS.** 2019. Increased pain sensitivity and decreased opioid analgesia in T-cell-deficient mice and implications for sex differences. *Pain* 160:358–366. <https://doi.org/10.1097/j.pain.0000000000001420>.
116. **Siriarchavatana P, Ayers JD, Kendall LV.** 2016. Anesthetic activity of alfaxalone compared with ketamine in mice. *J Am Assoc Lab Anim Sci* 55:426–430.
117. **Sneddon LU.** 2018. Comparative physiology of nociception and pain. *Physiology (Bethesda)* 33:63–73.
118. **Sorge RE, Mapplebeck JC, Rosen S, Beggs S, Taves S, Alexander JK, Martin LJ, Austin JS, Sotocinal SG, Chen D, Yang M, Shi XQ, Huang H, Pillon NJ, Bilan PJ, Tu Y, Klip A, Ji RR, Zhang J, Salter MW, Mogil JS.** 2015. Different immune cells mediate mechanical pain hypersensitivity in male and female mice. *Nat Neurosci* 18:1081–1083. <https://doi.org/10.1038/nn.4053>.
119. **Sorge RE, Totsch SK.** 2017. Sex differences in pain. *J Neurosci Res* 95:1271–1281. <https://doi.org/10.1002/jnr.23841>.
120. **Sternberg WF, Ritchie J, Mogil JS.** 2004. Qualitative sex differences in k-opioid analgesia in mice are dependent on age. *Neurosci Lett* 363:178–181. <https://doi.org/10.1016/j.neulet.2004.04.004>.
121. **Tajerian M, Sahbaie P, Sun Y, Leu D, Yang HY, Li W, Huang TT, Kingery W, David Clark J.** 2015. Sex differences in a murine model of complex regional pain syndrome. *Neurobiol Learn Mem* 123:100–109. <https://doi.org/10.1016/j.nlm.2015.06.004>.
122. **Toth LA, Compton S, Tolwani R, Farrar J.** 2019. The AALAS Journals: 2018 in Review. *Comp Med* 69:80–85.
123. **Tuttle AH, Philip VM, Chesler EJ, Mogil JS.** 2018. Comparing phenotypic variation between inbred and outbred mice. *Nat Methods* 15:994–996. <https://doi.org/10.1038/s41592-018-0224-7>. Author correction: comparing phenotypic variation between inbred and outbred mice. [*Nat Methods*. 2019]
124. **Von Korff M, Scher AI, Helmick C, Carter-Pokras O, Dodick DW, Goulet J, Hamill-Ruth R, LeResche L, Porter L, Tait R, Terman G, Veasley C, Mackey S.** 2016. United States national pain strategy for population research: concepts, definitions, and pilot data. *J Pain* 17:1068–1080. <https://doi.org/10.1016/j.jpain.2016.06.009>.
125. **Wan Y, Wilson SG, Han J, Mogil JS.** 2001. The effect of genotype on sensitivity to electroacupuncture analgesia. *Pain* 91:5–13. [https://doi.org/10.1016/S0304-3959\(00\)00416-4](https://doi.org/10.1016/S0304-3959(00)00416-4).
126. **Wiesenfeld-Hallin Z.** 2005. Sex differences in pain perception. *Gend Med* 2:137–145. [https://doi.org/10.1016/S1550-8579\(05\)80042-7](https://doi.org/10.1016/S1550-8579(05)80042-7).
127. **Wijnvoord N, Albuquerque B, Haussler A, Myrczek T, Popp L, Tegeder I.** 2010. Inter-strain differences of serotonergic inhibitory pain control in inbred mice. *Mol Pain* 6:1–11. <https://doi.org/10.1186/1744-8069-6-70>.
128. **Wilson SG, Bryant CD, Lariviere WR, Olsen MS, Giles BE, Chesler EJ, Mogil JS.** 2003. The heritability of antinociception II: pharmacogenetic mediation of three over-the-counter analgesics in mice. *J Pharmacol Exp Ther* 305:755–764. <https://doi.org/10.1124/jpet.102.047902>.
129. **Wilson SG, Smith SB, Chesler EJ, Melton KA, Haas JJ, Mitton B, Strasburg K, Hubert L, Rodriguez-Zas SL, Mogil JS.** 2003. The heritability of antinociception: common pharmacogenetic mediation of 5 neurochemically distinct analgesics. *J Pharmacol Exp Ther* 304:547–559. <https://doi.org/10.1124/jpet.102.041889>.
130. **Wolfe AM, Kennedy LH, Na JJ, Nemzek-Hamlin JA.** 2015. Efficacy of tramadol as a sole analgesic for postoperative pain in male and female mice. *J Am Assoc Lab Anim Sci* 54:411–419.
131. **Woller SA, Corr M, Yaksh TL.** 2015. Differences in cisplatin-induced mechanical allodynia in male and female mice. *Eur J Pain* 19:1476–1485. <https://doi.org/10.1002/ejp.679>.
132. **Yeziarski RP.** 2012. The effects of age on pain sensitivity: preclinical studies. *Pain Med* 13 Suppl 2:S27–S36. <https://doi.org/10.1111/j.1526-4637.2011.01311.x>.
133. **Yeziarski RP, King CD, Morgan D, Carter CS, Vierck CJ.** 2010. Effects of age on thermal sensitivity in the rat. *J Gerontol A Biol Sci Med Sci* 65A:353–362. <https://doi.org/10.1093/gerona/glq024>.
134. **Young EE, Costigan M, Herbert TA, Lariviere WR.** 2014. Heritability of nociception IV: neuropathic pain assays are genetically distinct across methods of peripheral nerve injury. *Pain* 155:868–880. <https://doi.org/10.1016/j.jpain.2013.09.018>.