



HHS Public Access

Author manuscript

Resuscitation. Author manuscript; available in PMC 2020 February 01.

Published in final edited form as:

Resuscitation. 2019 February ; 135: 1–5. doi:10.1016/j.resuscitation.2018.12.006.

Variations in the Application of Exception from Informed Consent in a Multicenter Clinical Trial

Jestin Carlson

Emergency Department, Saint Vincent Health Center, 232 West 25th St, 16544,
jcarlson@svhs.org, 14128643830

Abstract

Background—Exception from informed consent (EFIC) is allowed using federal regulations 21 CFR 50.24 and facilitates research on patients with critical conditions such as cardiac arrest. Little is known regarding the differences in the application of EFIC requirements such as community consultation (CC), public disclosure (PD) and patient notification. We sought to characterize variations in the fulfillment of EFIC requirements in a national multicenter clinical trial in the United States.

Methods—We determined the strategies for fulfillment of EFIC requirements at five regional coordinating centers of the Pragmatic Airway Resuscitation Trial (PART), a cluster-crossover randomized trial comparing airway devices in out-of-hospital cardiac arrest. We collected information from the including site demographics, how CC and PD were implemented, methods undertaken by the site investigative team to meet the local IRB's interpretation, and patient notification timing (post-enrollment). We analyzed the data using descriptive statistics.

Results—Sites had multiple approaches to CC, including social media advertising, random digit dialing surveys, working with city officials, and websites with embedded surveys. All sites used more than one approach for conducting CC. Public Disclosure activities included press releases through various means, website documentation, and letters to community members and local officials. Time from CC to study approval ranged from 42 days to 253 days.

Conclusion—EFIC implementation varies across sites and highlight community and regional variation. Different EFIC approaches may be needed to effectively accomplish the goals of community consultation, public disclosure, and patient notification.

Introduction

Prospective research on acutely ill and injured patients is challenging because obtaining informed consent, due to the nature of the critical illness, is often impossible or impracticable. To address this, the US Department of Health and Human Services (DHHS) and the Food and Drug Administration (FDA) developed regulations allowing such research

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

to occur, Federal Regulation 21 CFR 50.24, called exception from informed consent (EFIC) under emergency circumstances.¹

The additional requirements of 21 CFR 50.24 include community consultation (CC) and public disclosure (PD). According to the FDA Guidance for Institutional Review Boards (IRBs), Clinical Investigators, and Sponsors, community consultation “means providing the opportunity for discussions with, and soliciting opinions from, the community in which the study will take place” with the goals of “showing respect for persons by informing the community about the study in advance”, “provide a means for affected communities to provide meaningful input to the IRB before its decision to approve, require modifications to, or disapprove the study”, “identify potential community-level concerns and effects of the research”, and “obtain input from a group that is expected to be similar to the eventual study subjects”.² Public disclosure has been interpreted by the FDA to mean dissemination of information by the sponsor (i.e., one-way communication) to the community, the public, and researchers about emergency research. It occurs before initiation of the clinical trial, after the trial has been completed, and whenever the IRB determines that new disclosures are appropriate.

An additional requirement of 21 CFR 50.24 is that “IRBs must ensure there are procedures in place to provide information at the earliest feasible opportunity about the subject’s inclusion in the clinical investigation, the details of the investigation and other information contained in the informed consent document”.²

Because of local cultural, population demographics, and IRB differences, optimizing these required processes of federal regulation 21 CFR 50.24 may vary in each community. Currently, there is limited understanding of the variation that occurs with any given multicenter EFIC study. For these reasons, we utilized an established clinical trials network (Resuscitation Outcomes Consortium; ROC) to describe the variation in implementation of CC and PD by all participating study sites in the Pragmatic Airway Resuscitation Trial (PART), a multicenter randomized clinical trial using exception from informed consent under emergency circumstances comparing the effectiveness of endotracheal versus laryngeal tube airway insertion on outcomes after out-of-hospital cardiac arrest.³

Methods

We report the CC and PD performed at each site involved in PART, a multi-center, prospective, randomized cluster with crossover clinical trial comparing two airway devices in out-of-hospital cardiac arrest (OHCA). Details of the study methodology have been described previously.³ Briefly, PART compared 72-hour survival from OHCA in adult patients initially managed with either endotracheal intubation or a supraglottic airway.⁴

Because of the nature of the condition of cardiac arrest, 21 CFR 50.24, exception from informed consent under emergency circumstances was used for this study. PART was performed in 27 EMS agencies surrounding five US cities: Birmingham, Alabama; Dallas-Fort Worth, Texas; Milwaukee, Wisconsin; Pittsburgh, Pennsylvania; and Portland, Oregon.

The IRB for each site approved the parent study and, when required, the IRBs of other local hospitals also reviewed and approved the study.

Data were collected during regular discussions between the site principal investigators and study authors and verified by the lead study coordinators for the parent study at the sites. We report CC and PD requirements regarding local demographics (number of EMS agencies, number of local IRBs that reviewed the protocol), IRB requirements for CC, PD, and patient notification, how the local investigators completed CC and PD, and timeliness of patient notification after study enrollment for each of the five sites. We analyzed the data using descriptive techniques.

Results

Sites had multiple approaches to CC, including social media advertising, random digit dialing surveys, working with city officials, and websites with embedded surveys (Table 1). All sites used more than one approach for conducting CC. The time from initiation of CC to local IRB approval ranged from 42 to 253 days. Public Disclosure activities included press releases through various means, website documentation, and letters to community members and local officials. The median interval (interquartile range [IQR]; % unable to be notified) across sites for notifying patients who had been enrolled was 21 days (IRQ: 15–23 days; 6.8%) for patients who did not survive to be transported, 24 days (IRQ 18–42 days; 3.3%) for patients who died within 72 hours of ED arrival, and 8 days (IRQ: 3–25 days; 3.2%) for those who survived for 72 hours or more post-arrival (Table 2).

Discussion

Within the context of a national multicenter clinical trial, we observed variations in the strategies used to fulfill the EFIC requirements. Sites reported different EFIC approaches, suggesting that the EFIC process may be impacted by regional demographics, epidemiology and outcomes of the studied event, local or regional course of care, variations in interpretation of the FDA guidance and cultural differences. While it is unknown why these differences occur, it is important to recognize this variability and future work will be needed to understand why the approach varies across different sites for the same study.

Our sites have completed multiple prospective, randomized clinical trials^{4–10} over the last 10 years incorporating EFIC and have likely refined the CC, PD and notification practices over time to what has proved most successful. They have also integrating newer technologies (e.g. social media) when available, suggesting an iterative and adaptive methodology focused on a site or region's specific makeup.¹¹ Even with this experience, there remains great variability in the time from CC initiation to IRB approval to enroll patients and all sites required IRB review of proposed CC activities before conducting CC. These time intervals (ranging from 42 to 253 days) can represent significant challenges in terms of resources; specifically investigator time and cost. While we did not examine the financial or individual time resources (i.e. number of hours required to completed these tasks) associated with these activities, future work may be needed to help develop efficient strategies across sites to help reduce time and labor while ensuring the ethical treatment of human subjects. However, our

work helps to better describe the techniques undertaken to achieve CC and PD in an EFIC trial while also reporting the successes with notification of enrollment in a mature research network.

While EFIC guidelines require patient notification as soon as feasible post-enrollment to allow an opportunity for the patient or a legally authorized representative (LAR) to opt-out of continued participation, there is a paucity of published guidance for these activities. In the PART study, ongoing participation was only in relationship to the collection of hospital-based data. Given the high mortality of OHCA, it was important to consider notification to the next of kin for patients who did not survive their cardiac arrest. Most sites adopted a strategy to notify and consent patients surviving to 72 hours post-admission in-person, while sending letters to known next of kin for patients who died prior to EMS transport for ongoing treatment, or those who died in the ED or within the first 72 hours of hospitalization. Given this timing, sites must have the ability to quickly know of and monitor patient status post-enrollment to identify availability of patients or LARs for notification and consenting for ongoing participation, and to consider the clinical status in the timing.

As more studies have been completed using EFIC, greater focus has been spent understanding stakeholder impact of CC and PD, including EMS providers¹² and IRBs^{13, 14}, and examinations of ethical implications of this research.^{15, 16} Previous work has described EFIC application in multi-site trials.^{3, 17–20} The methods described in these previous trials include newspaper, radio, or television advertisements, direct letters to community groups or individuals, presentations at community meetings or health fairs, random digit dialing surveys, and, more recently, the inclusion of technology and social media through internet advertising, web-based surveying and comment collection, and specifically-targeted Facebook advertising. These and other specific methods have been increasingly shared^{21–26}, and most note the variation in approaches given community needs and responsiveness.

These observations may have implications for future EFIC trials. As researchers and institutions at all study sites of NIH-funded multi-center clinical trials will soon be required to use a single, centralized IRB for approval, such as proposed for the Strategies to Innovate EmeRgENcy Care Clinical Trials Network (SIREN), stakeholders need to better understand and consider how this will affect execution of CC and PD. Recognizing and honoring local variation in implementing CC and PD may be crucial to optimize these processes locally. A 2018 study systematically reviewed FDA-reported CC and PD activities for 34 trials²⁷, indicating similar variability as our study, and concluded that there is a need to identify best practices and less uncertainty as to the implementation of CC and PD. With the current local variation in implementing CC and PD processes, there may be misunderstandings of what is required for CC and PD between local and centralized IRBs.

Uniform, standard approaches to CC and PD, while potentially helpful to centralized IRBs, may fail to address the nuances of local CC and PD processes necessary to address local cultural and demographic differences. Future work will be needed to detail the ideal approach to local CC and PD with centralized IRBs.

Limitations

There are several limitations to our work. We report the CC and PD process of several sites in one trial. Our results were derived from five study sites. A larger sample size involving a greater number of sites over multiple trials may result in different findings. However, these sites have participated in multiple previous trials through the ROC network. As such, we believe they have a refined approach to CC and PD that may be beneficial for other sites that have less experience conducting CC and PD for EFIC trials. The EFIC approach in sites without extensive experience may be different. There was variability in both the approach to EFIC and the time from CC to IRB approval. Understanding why these differences occur, will require future study. Individuals who died in the prehospital setting or who were admitted to the ED and survived <72 hours had longer time to notification than those who were admitted to the ED and survived ≥ 72 hours. We believe this is related to the challenges in notification the LAR when an individual dies while recognizing the balance between these challenges and respecting the grieving period of families however, the ideal methodology and time interval for identifying and notifying LARs in these cases is unknown. All sites were relatively large metropolitan centers, and how these findings extrapolate to smaller communities and rural areas is unknown. The parent study (PART) focused on airway management in OHCA. As a result, these findings may not be generalizable outside clinical trials studying pre-hospital interventions for OHCA. The current EFIC study involved only adult study subjects and our findings do not necessarily apply to the pediatric study population.

Conclusion

Significant variability exists in implementation of CC and PD as well as the methods utilized to achieve these but less so with notification. The drivers of this variability are likely multifactorial but may be related to community factors such as demographics and cultural needs; regulatory factors such as interpretation of the FDA guidance variation; and other considerations including feasibility, convenience and study specific issues. Further study will be needed to determine the driving factors behind local variability. Recognizing and honoring local variation in implementing EFIC may be crucial to optimize these processes locally if centralized IRBs are utilized for future multi-centered EFIC trials. Future work will be needed to establish the ideal methods for recognizing local variability within the context of centralized IRBs.

Acknowledgments

Financial Support:

Jestin N Carlson: ID #118243 from the American Heart Association and UH2-HL125163 from the National Heart, Lung and Blood Institute (NHLBI)

Dana Zive: UH2-HL125163 from the National Heart, Lung and Blood Institute (NHLBI)

Denise Griffiths: UH2-HL125163 from the National Heart, Lung and Blood Institute (NHLBI)

Karen N. Brown: None

Robert H Schmicker: UH2-HL125163 from the National Heart, Lung and Blood Institute (NHLBI)

Heather Herren: UH2-HL125163 from the National Heart, Lung and Blood Institute (NHLBI)

George Sopko: None

Sara DiFiore: None

Dixie Climer: None

Caroline Herdeman: None

Ahamed Idris: UH2-HL125163 from the National Heart, Lung and Blood Institute (NHLBI)

Graham Nichol: UH2-HL125163 from the National Heart, Lung and Blood Institute (NHLBI)

Henry E Wang: ID #118243 from the American Heart Association and UH2-HL125163 from the National Heart, Lung and Blood Institute (NHLBI)

References

1. 21 cfr 50.24 exception from informed consent requirements for emergency research. 1996;5024
2. U.S. Department of Health and Human Services FaDA. Guidance for institutional review boards, clinical investigators, and sponsors: Exception from informed consent requirements for emergency research 2013
3. Wang HE, Prince DK, Stephens SW, Herren H, Daya M, Richmond N, Carlson J, Warden C, Colella MR, Brienza A, Aufderheide TP, Idris AH, Schmicker R, May S, Nichol G. Design and implementation of the resuscitation outcomes consortium pragmatic airway resuscitation trial (part). Resuscitation. 2016;101:57–64 [PubMed: 26851059]
4. Wang HE, Schmicker RH, Daya MR, Stephens SW, Idris AH, Carlson JN, Colella MR, Herren H, Hansen M, Richmond NJ, Puyana JCJ, Aufderheide TP, Gray RE, Gray PC, Verkest M, Owens PC, Brienza AM, Sternig KJ, May SJ, Sopko GR, Weisfeldt ML, Nichol G. Effect of a strategy of initial laryngeal tube insertion vs endotracheal intubation on 72-hour survival in adults with out-of-hospital cardiac arrest: A randomized clinical trial. JAMA. 2018;320:769–778 [PubMed: 30167699]
5. Aufderheide TP, Nichol G, Rea TD, Brown SP, Leroux BG, Pepe PE, Kudenchuk PJ, Christenson J, Daya MR, Dorian P, Callaway CW, Idris AH, Andrusiek D, Stephens SW, Hostler D, Davis DP, Dunford JV, Pirrallo RG, Stiell IG, Clement CM, Craig A, Van Ottingham L, Schmidt TA, Wang HE, Weisfeldt ML, Ornato JP, Sopko G, Resuscitation Outcomes Consortium I. A trial of an impedance threshold device in out-of-hospital cardiac arrest. N Engl J Med. 2011;365:798–806 [PubMed: 21879897]
6. Stiell IG, Nichol G, Leroux BG, Rea TD, Ornato JP, Powell J, Christenson J, Callaway CW, Kudenchuk PJ, Aufderheide TP, Idris AH, Daya MR, Wang HE, Morrison LJ, Davis D, Andrusiek D, Stephens S, Cheskes S, Schmicker RH, Fowler R, Vaillancourt C, Hostler D, Zive D, Pirrallo RG, Vilke GM, Sopko G, Weisfeldt M, Investigators ROC. Early versus later rhythm analysis in patients with out-of-hospital cardiac arrest. N Engl J Med. 2011;365:787–797 [PubMed: 21879896]
7. Bulger EM, Jurkovich GJ, Nathens AB, Copass MK, Hanson S, Cooper C, Liu PY, Neff M, Awan AB, Warner K, Maier RV. Hypertonic resuscitation of hypovolemic shock after blunt trauma: A randomized controlled trial. Archives of surgery. 2008;143:139–148; [PubMed: 18283138]
8. Kudenchuk PJ, Brown SP, Daya M, Rea T, Nichol G, Morrison LJ, Leroux B, Vaillancourt C, Wittwer L, Callaway CW, Christenson J, Egan D, Ornato JP, Weisfeldt ML, Stiell IG, Idris AH, Aufderheide TP, Dunford JV, Colella MR, Vilke GM, Brienza AM, Desvigne-Nickens P, Gray PC, Gray R, Seals N, Straight R, Dorian P, Resuscitation Outcomes Consortium I. Amiodarone, lidocaine, or placebo in out-of-hospital cardiac arrest. N Engl J Med. 2016;374:1711–1722 [PubMed: 27043165]
9. Nichol G, Leroux B, Wang H, Callaway CW, Sopko G, Weisfeldt M, Stiell I, Morrison LJ, Aufderheide TP, Cheskes S, Christenson J, Kudenchuk P, Vaillancourt C, Rea TD, Idris AH, Colella R, Isaacs M, Straight R, Stephens S, Richardson J, Conde J, Schmicker RH, Egan D, May S, Ornato JP, Investigators ROC. Trial of continuous or interrupted chest compressions during cpr. N Engl J Med. 2015;373:2203–2214 [PubMed: 26550795]

10. Holcomb JB, Tilley BC, Baraniuk S, Fox EE, Wade CE, Podbielski JM, del Junco DJ, Brasel KJ, Bulger EM, Callcut RA, Cohen MJ, Cotton BA, Fabian TC, Inaba K, Kerby JD, Muskat P, O’Keeffe T, Rizoli S, Robinson BR, Scalea TM, Schreiber MA, Stein DM, Weinberg JA, Callum JL, Hess JR, Matijevic N, Miller CN, Pittet JF, Hoyt DB, Pearson GD, Leroux B, van Belle G, Group PS. Transfusion of plasma, platelets, and red blood cells in a 1:1:1 vs a 1:1:2 ratio and mortality in patients with severe trauma: The proppr randomized clinical trial. *JAMA*. 2015;313:471–482 [PubMed: 25647203]
11. Stephens SW, Williams C, Gray R, Kerby JD, Wang HE. Preliminary experience with social media for community consultation and public disclosure in exception from informed consent trials. *Circulation*. 2013;128:267–270 [PubMed: 23857233]
12. Ripley E, Ramsey C, Prorock-Ernest A, Foco R, Lockett S Jr., Ornato JP. Ems providers and exception from informed consent research: Benefits, ethics, and community consultation. *Prehospital emergency care: official journal of the National Association of EMS Physicians and the National Association of State EMS Directors*. 2012;16:425–433
13. Deiorio NM, McClure KB, Nelson M, McConnell KJ, Schmidt TA. Ethics committee experience with emergency exception from informed consent protocols. *J Empir Res Hum Res Ethics*. 2007;2:23–30 [PubMed: 19385848]
14. McClure KB, Delorio NM, Schmidt TA, Chiodo G, Gorman P. A qualitative study of institutional review board members’ experience reviewing research proposals using emergency exception from informed consent. *J Med Ethics*. 2007;33:289–293 [PubMed: 17470507]
15. Biros MH, Dickert NW, Wright DW, Scicluna VM, Harney D, Silbergleit R, Denninghoff K, Pentz RD. Balancing ethical goals in challenging individual participant scenarios occurring in a trial conducted with exception from informed consent. *Academic emergency medicine: official journal of the Society for Academic Emergency Medicine*. 2015;22:340–346 [PubMed: 25716051]
16. Govindarajan P, Dickert NW, Meeker M, De Souza N, Harney D, Hemphill CJ, Pentz R. Emergency research: Using exception from informed consent, evaluation of community consultations. *Academic emergency medicine: official journal of the Society for Academic Emergency Medicine*. 2013;20:98–103 [PubMed: 23570483]
17. Brasel KJ, Bulger E, Cook AJ, Morrison LJ, Newgard CD, Tisherman SA, Kerby JD, Coimbra R, Hata JS, Hoyt DB. Hypertonic resuscitation: Design and implementation of a prehospital intervention trial. *J Am Coll Surg*. 2008;206:220–232 [PubMed: 18222373]
18. Mosesso VN Jr., Brown LH, Greene HL, Schmidt TA, Aufderheide TP, Sayre MR, Stephens SW, Travers A, Craven RA, Weisfeldt ML. Conducting research using the emergency exception from informed consent: The public access defibrillation (pad) trial experience. *Resuscitation*. 2004;61:29–36 [PubMed: 15081178]
19. Dickert NW, Mah VA, Baren JM, Biros MH, Govindarajan P, Pancioli A, Silbergleit R, Wright DW, Pentz RD. Enrollment in research under exception from informed consent: The patients’ experiences in emergency research (peer) study. *Resuscitation*. 2013;84:1416–1421 [PubMed: 23603291]
20. Eubank L, Lee KS, Seder DB, Strout T, Darrow M, MacDonald C, May T, Riker RR, Kern KB. Approaches to community consultation in exception from informed consent: Analysis of scope, efficiency, and cost at two centers. *Resuscitation*. 2018;130:81–87 [PubMed: 29964145]
21. Stephens SW, Williams C, Gray R, Kerby JD, Wang HE, Bosarge PL. Using social media for community consultation and public disclosure in exception from informed consent trials. *The journal of trauma and acute care surgery*. 2016;80:1005–1009 [PubMed: 26998781]
22. Holsti M, Zemek R, Baren J, Stanley RM, Mahajan P, Vance C, Brown KM, Gonzalez V, King D, Jacobsen K, Shreve K, van de Bruinhorst K, Jones AM, Chamberlain JM. Variation of community consultation and public disclosure for a pediatric multi-centered “exception from informed consent” trial. *Clinical trials*. 2015;12:67–76 [PubMed: 25369796]
23. Bulger EM, Schmidt TA, Cook AJ, Brasel KJ, Griffiths DE, Kudenchuk PJ, Davis D, Bardarson B, Idris AH, Aufderheide TP. The random dialing survey as a tool for community consultation for research involving the emergency medicine exception from informed consent. *Annals of emergency medicine*. 2009;53:341–350, 350 e341–342 [PubMed: 18824274]
24. Nelson M, Schmidt TA, DeIorio NM, McConnell KJ, Griffiths DE, McClure KB. Community consultation methods in a study using exception to informed consent. *Prehospital emergency care :*

official journal of the National Association of EMS Physicians and the National Association of State EMS Directors. 2008;12:417–425

25. Salzman JG, Frascone RJ, Godding BK, Provo TA, Gertner E. Implementing emergency research requiring exception from informed consent, community consultation, and public disclosure. *Annals of emergency medicine*. 2007;50:448–455, 455 e441–444 [PubMed: 17222939]
26. Richardson LD, Quest TE, Birnbaum S. Communicating with communities about emergency research. *Acad Emerg Med*. 2005;12:1064–1070 [PubMed: 16264076]
27. Haggins AN, Harney D, Scott S, Silbergleit R. A systematic review of federal drug administration docket for community consultation and public disclosure in exception from informed consent trials. *Clinical trials*. 2018;15:29–35 [PubMed: 29039683]

Table 1.

Variation in EFIC processes

	Total Sites (n=5)
Satellite IRBs	Yes (n=5, 16 total)
IRB Required activities for CC	Not defined (n=4) One 'town hall' style meeting in each county (n=1)
Methods for conducting CC	-Facebook Ads (n=4) -Random Digit Dialing surveys (n=3) -Contact Local City Officials (n=2) -Website with survey (n=3) -Booth at Festivals/Health Fairs (n=1) -Town hall' style meeting with a survey (n=1)
IRB Required activities for PD	Not defined (n=4) Press release announcing study start date (n=1)
Methods for conducting PD	-Press release via any method(n=4) -Press release via newspaper (n=3) -Press release via radio (n=2) -Press release via TV (n=1) - Press release via bus ads (n=1) - Press release via web ads (n=1) -Website (n=4) -Letters to local Officials and community members (n=1)
IRB Required sequence of CC/PD Activities	CC and PD performed simultaneously (n=3) CC then PD (n=2)
Time from initiation of CC to study approval in days	Site A - 42 Site B - 164 Site C - 141 Site D - 253 Site E - 209

IRB - Institutional Review board

CC - Community consultation

PD - Public disclosure

TV - Television

Table 2.

Time from enrollment to notification in days

	Died in PH ¹			Admitted to ED Died <72 hours ²			Admitted to ED Survived 72 hours ³		
	25 th %ile	50 th %ile	75 th %ile	25 th %ile	50 th %ile	75 th %ile	25 th %ile	50 th %ile	75 th %ile
Site 1	12	14	18	11	14	18	7	15	31
Site 2	19	22	28	4	15	25	2	6	11
Site 3	21	21	22	23	39	57	3	7	36
Site 4	12	16	21	10	15	19	3	13	24
Site 5	21	24	27	22	24	27	3	11	25
All Sites	15	21	23	18	24	42	3	8	25

PH – Prehospital setting

ED – Emergency Department

¹ – 6.8 % of these patients were unable to be notified² – 3.3% of these patients were unable to be notified³ – 3.2% of these patients were unable to be notified

Note – Sites A-E (Table 1) are different from Sites 1–5 (Table 2) to maintain site confidentiality.