## CORRECTION




## Correction to: Vascular endothelial PDPK1 plays a pivotal role in the maintenance of pancreatic beta cell mass and function in adult male mice


Atsushi Obata<sup>1</sup> • Tomohiko Kimura<sup>1</sup> • Yoshiyuki Obata<sup>1</sup> • Masashi Shimoda<sup>1</sup> • Tomoe Kinoshita<sup>1</sup> • Kenji Kohara<sup>1</sup> • Seizo Okauchi<sup>1</sup> • Hidenori Hirukawa<sup>1</sup> • Shinji Kamei<sup>1</sup> • Shuhei Nakanishi<sup>1</sup> • Tomoatsu Mune<sup>1</sup> • Kohei Kaku<sup>1</sup> • Hideaki Kaneto<sup>1</sup>

Published online: 11 October 2019 © Springer-Verlag GmbH Germany, part of Springer Nature 2019

## Correction to: Diabetologia https://doi.org/10.1007/s00125-019-4878-1

It has been brought to our attention that Fig. 5a showing the vasculature in islets of control *flox* mice is not in fact an endocrine cell but, rather, exocrine tissue. In the new Fig. 5a presented here we show the vasculature adjacent to pancreatic beta cells of control *flox* mice. In addition, the  $\times$  5000 image in the left-hand panel of Fig. 5b is replaced by a lower-resolution image ( $\times$  3000) to provide a broader cellular context.





**Fig. 5** Vascular structure of islets from VE-PDPK1-KO mice and control *flox* mice. Electron microscopic images showing capillaries surrounded by pancreatic beta cells in (**a**) control *flox* mice and (**b**) VE-PDPK1-KO mice. EC, endothelial cells. Images are representative of four mice. Magnification  $\times$  5000 (**a**, left) or  $\times$  3000 (**b**, left) scale bars, 1 µm; or  $\times$  10,000 (right), scale bars, 500 nm

The online version of the original article can be found at https://doi.org/10.1007/s00125-019-4878-1

Atsushi Obata obata-tky@med.kawasaki-m.ac.jp

<sup>&</sup>lt;sup>1</sup> Department of Diabetes, Endocrinology and Metabolism, Kawasaki Medical School, 577 Matsushima, Kurashiki 701-0192, Japan