

Mutant Muddle: Some Arabidopsis *eds5* Mutant Lines Have a Previously Unnoticed Second-Site Mutation in *FAH1*^[OPEN]

Sravani Ram Veeragoni,^a Birgit Lange,^a Mario Serrano,^b Christiane Nawrath,^c Sibylle Bauer,^a Anton Rudolf Schöffner,^a Hans Thordal-Christensen,^d Jörg Durner,^{a,e} and Frank Gaupels^{a,1,2}

^aInstitute of Biochemical Plant Pathology, Helmholtz Zentrum München, D-85764 Neuherberg, Germany

^bCentro de Ciencias Genómicas, Universidad Nacional Autónoma de México, 62209 Cuernavaca, Morelos, México

^cDepartment of Plant Molecular Biology, University of Lausanne, 1015 Lausanne, Switzerland

^dDepartment of Plant and Environmental Sciences, Copenhagen Plant Science Centre, University of Copenhagen, DK-1871 Frederiksberg, Denmark

^eBiochemical Plant Pathology, Technische Universität München, D-85354 Freising, Germany

ORCID IDs: 0000-0002-1510-756X (M.S.); 0000-0003-1470-8459 (C.N.); 0000-0001-5501-7481 (S.B.); 0000-0002-9424-5548 (A.R.S.); 0000-0002-6079-6800 (F.G.).

Salicylic acid (SA) is produced by the enzyme isochorismate synthase (ICS) within the chloroplast and is subsequently exported to the cytosol by the multi-drug and toxic compound extrusion transporter ENHANCED DISEASE SUSCEPTIBILITY5 (*EDS5*; Nawrath et al., 2002; Serrano et al., 2013). The ICS pathway is the major source of SA during plant responses to various fungal and bacterial pathogens (Nawrath and Métraux, 1999; Wildermuth et al., 2001). Accordingly, Arabidopsis (*Arabidopsis thaliana*) mutants defective in *EDS5* are impaired in the SA-dependent establishment of resistance against *Pseudomonas syringae* pv. *tomato* (Nawrath and Métraux, 1999). Several SA-deficient *eds5* loss-of-function alleles have been established. The mutant alleles *eds5-1* and *eds5-3* (Glazebrook et al., 1996; Nawrath and Métraux, 1999) are widely used in plant pathology research, whereas *eds5-2* (Volko et al., 1998) is less frequently used. We discovered that the *eds5-3*, *syp121-1 syp122-1 eds5-3*, and *syp121-1 syp122-1 eds5-3 sid2-1* mutants all carried an unnoticed second-site mutation in *FERULIC ACID 5-HYDROXYLASE1* (*FAH1*). The *eds5-1* mutant lines might harbor a similar mutation

in *FAH1*. Therefore, it is strongly recommended to test all stocks of *eds5-3* and *eds5-1* for the *FAH1* background mutation before use.

eds5 mutants do not have an obvious growth phenotype. However, under UV-A illumination, leaves of *eds5-3* exhibit red chlorophyll fluorescence rather than the blue fluorescence observed in wild-type (Col-0) plants (Fig. 1A). UV-excited blue fluorescence is mainly emitted by sinapoylmalate, which is a major phenylpropanoid in vacuoles of the leaf upper epidermis, where it serves as a protective barrier against harmful UV irradiation (Fraser and Chapple, 2011). *eds5-1* mutant seedlings grown from two different seed stocks (the Mario Serrano laboratory stock and the Arabidopsis Biological Resource Center stock CS3735, donated by the Frederick Ausubel laboratory) had similar levels of blue fluorescence as the wild type (Fig. 1A), suggesting that the defect in *EDS5* expression did not cause the lack of sinapoylmalate in *eds5-3*. NahG *sid2-1* plants cannot accumulate SA due to a mutation in the ICS coding gene *ICS1/SID2* and transgenic expression of the SA hydroxylase gene NahG from *Pseudomonas putida* (Delaney et al., 1994; Supplemental Materials and Methods S1). Under UV, NahG *sid2-1* was indistinguishable from wild-type plants, underpinning again that SA deficiency does not affect sinapoylmalate biosynthesis (Fig. 1A).

The *eds5-3* mutant had a similar UV phenotype as reported in the previously identified *reduced epidermal fluorescence* mutants, including the *fah1-2* line (Fraser and Chapple, 2011; Fig. 1A). *fah1-2* has a defect in *FAH1*, which encodes ferulate-5-hydroxylase, an essential enzyme for the conversion of ferulate into sinapates such as sinapoylmalate (Chapple et al., 1992; Fraser and Chapple, 2011). Ultra-performance liquid chromatography coupled to mass spectrometry (Tzin et al., 2012) was employed for molecular phenotyping of the mutants (Supplemental Materials

¹Author for contact: frank.gaupels@gmx.de.

²Senior author.

The author responsible for distribution of materials integral to the findings presented in this article in accordance with the policy described in the Instructions for Authors (www.plantphysiol.org) is: Frank Gaupels (frank.gaupels@gmx.de).

S.R.V., F.G., and J.D. planned and designed the research; S.R.V. and B.L. performed the UV imaging, UPLC-MS, and data analysis; M.S., C.N., S.B., A.R.S., and H.T.-C. generated, characterized, and maintained the Arabidopsis mutant lines; F.G. wrote the article with contributions from all other authors; F.G. agrees to serve as the author responsible for contact and ensure communication.

^[OPEN]Articles can be viewed without a subscription.

www.plantphysiol.org/cgi/doi/10.1104/pp.19.01125




Figure 1. *eds5-3* carries a *FAH1* mutation. A, Detection of blue fluorescent sinapoylmalate under UV-A illumination (365-nm wavelength). Mutant lines lacking sinapoylmalate emit red fluorescence under UV. B, Semiquantification of SA and sinapoylmalate (SinM) by ultra-performance liquid chromatography coupled to mass spectrometry. Plants were infected with *P. syringae* pv. *tomato* *AvrRpm1*, and leaf extracts were sampled at 4- and 24-h post infection (hpi). Error bars indicate the SD ($n = 4-5$). C, The *syp121-1 syp122-1 eds5-3* and *syp121-1 syp122-1 eds5-3 sid2-1* mutants carry the *fah1-2* background mutation as evidenced by their red fluorescence phenotype. *snc1-1 eds5-3* showed blue fluorescence under UV illumination, suggesting that the *fah1-2* mutation had been crossed out. Refer to Supplemental Materials and Methods S1 for a detailed description of the mutant lines and methods.

and Methods S1). Pathogen infection with avirulent *P. syringae* pv. *tomato* *AvrRpm1* triggered the accumulation of SA at 4 h and more pronouncedly at 24 h after infection in wild-type and *fah1-2* plants (Fig. 1B). By contrast, the SA peak was reduced in *eds5-3* and

eds5-1, and absent in NahG *sid2-1*. Sinapoylmalate levels were high and not strongly regulated upon pathogen infection in wild type, *eds5-1*, and NahG *sid2-1*. However, sinapoylmalate was not detectable in *eds5-3* and *fah1-2* (Fig. 1B). Collectively, these results point to sinapoylmalate biosynthesis being disrupted in *eds5-3* independently of the *EDS5* mutation.

This prompted us to consider whether an unknown second-site mutation caused the unexpected *eds5-3* phenotype. In more recent reports, Col-0 was given as the background of *eds5-3* (Nawrath et al., 2002; Serrano et al., 2013). Only the initial publications, describing the generation, identification, and characterization of *eds5* mutants, provided an explanation for the sinapoylmalate deficiency of *eds5-3*. In these publications, it was mentioned that *eds* mutants were generated by EMS mutagenesis of *fah1-2* rather than Col-0 seeds (Glazebrook et al., 1996; Rogers and Ausubel, 1997). The *fah1-2* mutant (in the Col-0 background) was chosen because the red fluorescence phenotype is a useful marker in genetic crosses (Glazebrook et al., 1996). *eds5-1* and *eds5-3* both originated from the same mutant population. The *fah1-2* and *eds5* mutations are only 10 centimorgans apart (Reuber et al., 1998), which rendered removal of the *fah1-2* mutation by backcrossing difficult due to genetic linkage. *eds5-1* was successfully backcrossed with Col-0 before submission to the Arabidopsis Biological Resource Center (Fig. 1; J. Glazebrook, personal communication), whereas *eds5-3* still carries the *fah1-2* background mutation (Fig. 1; Nawrath and Métraux, 1999). *eds5-2* originated from an independent screen, not involving *fah1-2* (Volko et al., 1998).

eds5 lines were crossed with various mutants to study the interaction of SA signaling with other defense mechanisms (Zhang et al., 2008; Venugopal et al., 2009; Dong et al., 2016). Inspecting multiple mutants under UV revealed that *syp121-1 syp122-1 eds5-3* and *syp121-1 syp122-1 eds5-3 sid2-1* (mutants from H.T.-C.'s laboratory; Zhang et al., 2008) displayed the red fluorescence phenotype, whereas in *snc1-1 eds5-3* (mutant from Xin Li's laboratory; Dong et al., 2016) the *fah1-2* mutation was seemingly outcrossed (Fig. 1C). Thus, due to genetic linkage, *fah1-2* can be an unnoticed background mutation in multiple mutants containing *eds5-3*. Some laboratories could have stocks of the original *eds5-1* line with the *fah1-2* background. However, we have yet to test multiple mutants containing *eds5-1*, such as *acd11 eds5-1* and *ssi2 eds1-2 eds5-1* (Brodersen et al., 2005; Venugopal et al., 2009).

Previous studies provided evidence that the lack of sinapates and syringyl lignin in *fah1-2* increased susceptibility to the fungal pathogens *Botrytis cinerea* and *Verticillium longisporum* (Lloyd et al., 2011; Demkura and Ballaré, 2012; König et al., 2014). Hence, the unnoticed *fah1-2* background mutation could influence pathogen resistance of *eds5-3* and possibly *eds5-1* plants in a SA-independent manner, thereby leading to false conclusions on the role of EDS5 and SA in plant-pathogen interactions. For this reason, it is strongly

recommended to check all lab stocks of *eds5-3*, *eds5-1*, and multiple mutants containing one of these mutations under UV-A illumination for the characteristic red fluorescence phenotype of *fah1-2*, followed by molecular confirmation of the genotype by PCR with *FAH1*-specific primers as described in Weng et al. (2010).

Supplemental Data

The following supplemental materials are available.

Supplemental Materials and Methods S1. Description of the mutant lines, materials, and methods used.

ACKNOWLEDGMENTS

We thank Clint Chapple and Xin Li for donating the mutant lines *fah1-2* and *snc1 eds5-3*, respectively.

Received September 18, 2019; accepted October 24, 2019; published November 4, 2019.

LITERATURE CITED

- Brodersen P, Malinovsky FG, Hématy K, Newman MA, Mundy J** (2005) The role of salicylic acid in the induction of cell death in *Arabidopsis acd11*. *Plant Physiol* **138**: 1037–1045
- Chapple CC, Vogt T, Ellis BE, Somerville CR** (1992) An *Arabidopsis* mutant defective in the general phenylpropanoid pathway. *Plant Cell* **4**: 1413–1424
- Delaney TP, Uknes S, Vernooij B, Friedrich L, Weymann K, Negrotto D, Gaffney T, Gut-Rella M, Kessmann H, Ward E, et al** (1994) A central role of salicylic acid in plant disease resistance. *Science* **266**: 1247–1250
- Demkura PV, Ballaré CL** (2012) UVR8 mediates UV-B-induced *Arabidopsis* defense responses against *Botrytis cinerea* by controlling sinapate accumulation. *Mol Plant* **5**: 642–652
- Dong OX, Tong M, Bonardi V, El Kasmi F, Woloshen V, Wunsch LK, Dangl JL, Li X** (2016) TNL-mediated immunity in *Arabidopsis* requires complex regulation of the redundant *ADR1* gene family. *New Phytol* **210**: 960–973
- Fraser CM, Chapple C** (2011) The phenylpropanoid pathway in *Arabidopsis*. *Arabidopsis Book* **9**: e0152
- Glazebrook J, Rogers EE, Ausubel FM** (1996) Isolation of *Arabidopsis* mutants with enhanced disease susceptibility by direct screening. *Genetics* **143**: 973–982
- König S, Feussner K, Kaever A, Landesfeind M, Thurow C, Karlovsky P, Gatz C, Polle A, Feussner I** (2014) Soluble phenylpropanoids are involved in the defense response of *Arabidopsis* against *Verticillium longisporum*. *New Phytol* **202**: 823–837
- Lloyd AJ, William Allwood J, Winder CL, Dunn WB, Heald JK, Cristescu SM, Sivakumaran A, Harren FJM, Mulema J, Denby K, et al** (2011) Metabolomic approaches reveal that cell wall modifications play a major role in ethylene-mediated resistance against *Botrytis cinerea*. *Plant J* **67**: 852–868
- Nawrath C, Heck S, Parinthewong N, Métraux JP** (2002) EDS5, an essential component of salicylic acid-dependent signaling for disease resistance in *Arabidopsis*, is a member of the MATE transporter family. *Plant Cell* **14**: 275–286
- Nawrath C, Métraux JP** (1999) Salicylic acid induction-deficient mutants of *Arabidopsis* express *PR-2* and *PR-5* and accumulate high levels of camalexin after pathogen inoculation. *Plant Cell* **11**: 1393–1404
- Reuber TL, Plotnikova JM, Dewdney J, Rogers EE, Wood W, Ausubel FM** (1998) Correlation of defense gene induction defects with powdery mildew susceptibility in *Arabidopsis* enhanced disease susceptibility mutants. *Plant J* **16**: 473–485
- Rogers EE, Ausubel FM** (1997) *Arabidopsis* enhanced disease susceptibility mutants exhibit enhanced susceptibility to several bacterial pathogens and alterations in *PR-1* gene expression. *Plant Cell* **9**: 305–316
- Serrano M, Wang B, Aryal B, Garcion C, Abou-Mansour E, Heck S, Geisler M, Mauch F, Nawrath C, Métraux J-P** (2013) Export of salicylic acid from the chloroplast requires the multidrug and toxin extrusion-like transporter EDS5. *Plant Physiol* **162**: 1815–1821
- Tzin V, Malitsky S, Ben Zvi MM, Bedair M, Sumner L, Aharoni A, Galili G** (2012) Expression of a bacterial feedback-insensitive 3-deoxy-D-arabino-heptulosonate 7-phosphate synthase of the shikimate pathway in *Arabidopsis* elucidates potential metabolic bottlenecks between primary and secondary metabolism. *New Phytol* **194**: 430–439
- Venugopal SC, Jeong RD, Mandal MK, Zhu S, Chandra-Shekara AC, Xia Y, Hersh M, Stromberg AJ, Navarre D, Kachroo A, et al** (2009) Enhanced disease susceptibility 1 and salicylic acid act redundantly to regulate resistance gene-mediated signaling. *PLoS Genet* **5**: e1000545
- Volko SM, Boller T, Ausubel FM** (1998) Isolation of new *Arabidopsis* mutants with enhanced disease susceptibility to *Pseudomonas syringae* by direct screening. *Genetics* **149**: 537–548
- Weng JK, Akiyama T, Bonawitz ND, Li X, Ralph J, Chapple C** (2010) Convergent evolution of syringyl lignin biosynthesis via distinct pathways in the lycophyte *Selaginella* and flowering plants. *Plant Cell* **22**: 1033–1045
- Wildermuth MC, Dewdney J, Wu G, Ausubel FM** (2001) Isochorismate synthase is required to synthesize salicylic acid for plant defence. *Nature* **414**: 562–565
- Zhang Z, Lenk A, Andersson MX, Gjetting T, Pedersen C, Nielsen ME, Newman MA, Hou BH, Somerville SC, Thordal-Christensen H** (2008) A lesion-mimic syntaxin double mutant in *Arabidopsis* reveals novel complexity of pathogen defense signaling. *Mol Plant* **1**: 510–527