



Published in final edited form as:

J Am Board Fam Med. 2019 ; 32(4): 607–613. doi:10.3122/jabfm.2019.04.180326.

Physician and Family Discussions about Driving Safety: Findings from the LongROAD Study

Marian E. Betz, MD, MPH¹, Leon Villavicencio, MPH², Deepika Kandasamy, MPH¹, Tara Kelley-Baker, PhD², Woon Kim, PhD², Carolyn DiGuseppi, MD, PhD³, Thelma J. Mielenz, PT, PhD^{4,5}, David W. Eby, PhD⁶, Lisa J. Molnar, PhD⁶, Linda Hill, MD, MPH⁷, David Strogatz, PhD⁸, David B. Carr, MD⁹, Guohua Li, MD, DrPH^{4,5,10} LongROAD Research Team^a

¹Department of Emergency Medicine, University of Colorado School of Medicine, Aurora, Colorado, USA

²AAA Foundation for Traffic Safety, Washington, DC, USA

³Department of Epidemiology, Colorado School of Public Health, University of Colorado Anschutz Medical Campus, Aurora, Colorado, USA

⁴Department of Epidemiology, Mailman School of Public Health, Columbia University, New York, NY, USA

⁵Center for Injury Epidemiology and Prevention, Columbia University Medical Center, New York, NY, USA

⁶University of Michigan Transportation Research Institute and Center for Advancing Transportation Leadership and Safety, Ann Arbor, MI, USA

⁷Department of Family Medicine and Public Health, University of California, San Diego, CA, USA

⁸Bassett Research Institute, Bassett Healthcare Network, Cooperstown, NY, USA

⁹Departments of Medicine and Neurology, Washington University School of Medicine, St. Louis, MO, United States

¹⁰Department of Anesthesiology, Columbia University College of Physicians and Surgeons, New York, NY, USA

Abstract

Background: Older adult drivers may experience decreases in driving safety with age or health status change. Discussing driving safety may help them plan for driving restriction and eventual cessation. Here, we sought to examine conversations between older adults and their family members and physicians.

Corresponding Author: Marian E. Betz, MD, MPH, University of Colorado School of Medicine; 12401 E. 17th Ave B-215; Aurora, CO 80045, Telephone: (720) 848-6770; Fax: (720) 848-7374; marian.betz@ucdenver.edu, **Twitter handles:** @EmmyBetz, @AAAFTS.

^aalso includes Howard Andrews, Vanya Jones, Robert Santos, David LeBlanc, and Lindsay Ryan

Conflicts of Interest: None of the authors have any conflicts of interest to disclose.

Note to NIH: Please add that the published version of this paper is available for free at: <https://www.jabfm.org/content/32/4/607>

Methods: In this multi-site cross-sectional analysis of baseline data from the AAA Longitudinal Research on Aging Drivers (AAA LongROAD) cohort study, we measured the prevalence and characteristics of family and physician driving discussions. We examined associations between having driving discussions and participant characteristics using multivariate logistic regression.

Results: Of 2990 current drivers aged 65–79 years (53% female, 85.5% White), only 14.2% reported discussing driving safety with family and 5.5% had discussions with physicians. Men (adjusted OR: 1.32, 95% CI 1.05–1.66) and those with Master’s degrees or higher (adjusted OR: 1.65, 95% CI, 1.27–2.13) more often had family discussions. Those with at least a Master’s degree were also more likely to speak with their physician (adjusted OR: 1.77, 95% CI 1.17–2.68).

Conclusion: Few older adults had driving safety conversations with their family or physicians. Practical and effective interventions are needed to engage family and physicians in assisting older adults with risk assessment and driving cessation planning to maintain mobility and well-being.

Keywords

Older drivers; driving safety; driving cessation; family; healthcare provider; communication

INTRODUCTION

Medical conditions, medications, and physiologic changes from aging can affect driving ability.¹ Driving safety, self-regulation, and cessation are issues many older adults eventually face.² Family and physicians may influence driving decisions, depending on their availability and involvement and the older adult’s counseling preferences.^{3–5}

Prior research showed that drivers aged 50 years preferred discussing driving with family versus non-family.⁶ In another driving survey, 64% of community-dwelling older adults were open to family discussion but fewer to family (32%) or physicians (44%) making driving cessation decisions.⁷ How much interaction older adults have with family often determines their comfort with family involvement in driving decisions.⁸

Yet healthcare providers also play key roles in driving safety and cessation.¹ One study found that married older adults preferred hearing first from their spouse and adult children but also desired conversations with their physician.⁶ Those indicating physicians as first choice believed the physician could judge their driving capabilities. However, the study’s physician participants were reluctant to partake in driving decisions. Many physicians report assessing driving fitness or discussing driving with their older patients, but only with *some* patients and not frequently with *each* of them.^{1,9} A retrospective medical record review documented that physicians discussed driving with only 8–22% of patients within a 12-month period.¹⁰

The LongROAD study, a multi-site longitudinal cohort study of older drivers, offers opportunities to examine the prevalence of driving discussions with family and providers in a larger sample than prior work. Understanding the frequency of such discussions and the demographic and driving characteristics associated with them may enhance communication strategy development and refinement.

METHODS

Design and Participants

This cross-sectional study utilized baseline data from participants enrolled at the five LongROAD sites in California, Colorado, Michigan, Maryland, and New York; LongROAD was designed to examine driving behaviors and outcomes (and their associations with health and functional variables) in a large cohort over time, and the study is described in detail elsewhere.¹¹ Research assistants (RAs) identified potentially eligible participants from primary care clinic rosters and assessed eligibility and interest by phone. Eligible individuals: were 65–79 years; possessed valid driver's licenses; drove at least once weekly on average; drove one car (1996 model or newer) primarily (>80% of the time); and had no significant cognitive impairment (verified through medical records at some sites and the Six-Item Screener at all sites).¹² RAs obtained written informed consent at the in-person enrollment visit. Institutional Review Boards at each site approved the study.

Measures

This analysis used baseline data from all LongROAD participants for variables surrounding driving discussions (Have you ever [spoken to your family doctor or any other doctor] / [had a discussion with a family member] about your driving safety?) and driving limitation recommendations. Additional variables included demographic characteristics, self-reported health condition-related driving reduction in the past year and driving experiences (Table 1). Drivers were asked to rate their average driving ability and comfort driving on a separate scale of 1 (poor ability/not at all comfortable) to 7 (excellent ability/completely comfortable). They reported driving lapses in attention, errors and violations on separate scales from 1 (never) to 6 (nearly all the time). Strategic self-regulation was measured as the sum of 13 binary variables on whether they indicated avoiding behaviors due to concerns regarding their cognitive, physical, perceptual performance as it relates to driving. Tactical self-regulation was measured as the sum of 7 binary variables indicating behavior avoidance due to concerns regarding their cognitive, physical, or perceptual performance as it relates to driving.

Analysis

We described family and physician driving discussion prevalence with 95% confidence intervals (95%CI). We used Chi-square or Fisher's exact test for categorical variables and one-way ANOVA for continuous variables, in comparing demographic characteristics and family and physician discussions. We utilized multivariable logistic regression, identifying variables associated with discussions using a $p < 0.20$ cut-off for initial model inclusion and backwards elimination for model optimization.

RESULTS

Of 2,990 total LongROAD participants, approximately half were female and 65–69 years old; most were White and non-Hispanic. A majority were married and had household incomes \$50,000 yearly and Bachelor's degrees or higher (Table 1). Overall, 337

participants (11.3%) reported driving reduction due to a health problem in the past year. Only 2.2% reported that, in the past year, someone had recommended limiting their driving.

Overall, 17.3% of participants reported ever having driving discussions with family or physicians. They were more likely to have spoken with family (14.2%; Table 1) than physicians (5.5%). Seventy-four participants (2.5%) had conversations with both. Similar patterns manifested across gender and age. Men and those aged 75–79 years (versus younger age groups) were significantly more likely to speak with family (Table 1). Most family conversations were initiated by family (60.6%), while most physicians conversations were initiated by older drivers (55.0%). When asked what triggered a family discussion, most said driving safety concerns (64.8%), followed by health issue (22.3%), driving infraction (8.7%), and crash (8.7%). Patterns were similar across gender and age (data not shown).

Common outcomes of physician discussions were medical or physical assessment (41.8%), education or advice (38.2%), follow-up visits to monitor health concerns (32.2%), and medical treatment (29.0%; Figure 1). Few reported referrals for driving assessment to a licensing bureau (3.6%) or occupational therapist (2.4%). Nearly one-quarter (22.5%) said no action resulted from these conversations.

Driving discussions with family were significantly associated with demographic characteristics (Table 2). Men, participants with Master's degrees and higher (versus less than Bachelor's degrees), and those who reduced driving for self-regulatory reasons in the past year were more likely to have driving conversations with family as well as those with lower self-rated driving ability, higher strategic self-regulation (i.e., pre-trip decisions attributed to self-regulation), and more driving errors or violations. Similar characteristics were associated with having driving discussions with physicians (Table 2).

DISCUSSION

In the large LongROAD cohort of older drivers, most had not discussed driving with their family or physician. Conversations that did occur were more common with family than with physicians, perhaps reflecting the relatively good health of participants. Few older drivers having physician discussions were referred for driving assessment.

Older drivers who anticipate and prepare for driving cessation experience better health outcomes, emphasizing the benefits of advance planning.³ Yet few participants had discussed driving safety with family or physicians, which is consistent with past research and may reflect avoidance of this sensitive subject¹³ or unawareness that many older adults are unable to drive for the last years of life.² Physicians in particular may believe that conversations about driving safety adversely affects patient-physician relationships, especially when they include recommendations for driving cessation.^{14–16} Physicians may also feel they lack training or resources regarding driving assessments, licensing laws, or linkages to older driver testing or education programs.^{17,18} Yet physicians do have obligations to protect patient and public safety,¹⁹ and available resources can help them with these difficult situations.^{1,20}

Participants were twice as likely to discuss driving safety with family as with physicians. While some resources exist to help physicians address this topic in clinical settings,¹ there are fewer available for families, who may also benefit from public awareness campaigns and education.^{13,21} Also, physicians should pay special attention to older adults who are widowed, single, or without family, since this population may not have social support in driving matters.

Driving discussions with family were more likely to occur when participants were male and more educated. Gender-related findings may reflect that, similar to past studies, men may want to maintain a traditional provider role and continue driving later in life (hence prompting family discussions), while women are willing to relinquish driving earlier.²² Gender was not associated with physician conversations, but those with higher education and who self-restricted their driving often discussed driving safety with physicians. Generally, previous research shows physicians and older drivers tend not to initiate driving conversations until specific “red flags,” (e.g., crashes) manifest.⁸ Circumstances of conversation initiation with physicians were not considered in the LongROAD study and should be examined in future research.

Very few participants had driving discussions resulting in referral for driving assessment. Such evaluations are often unavailable, though they are often recommended and useful.²³ Creating and testing triage tools to identify older adults who may benefit from driving assessments is vital, as reliable tests or combinations of tests have not yet been identified.²⁴ Physicians desire simple protocols to screen older drivers so developing such resources may help them to properly address older adult driving in clinical settings.^{17,25}

Limitations

Study limitations include an inability to determine when participants had conversations with family or physicians. The LongROAD study’s longitudinal follow-up may help us better address the temporal relationship between discussions and subsequent driving behaviors, including attention to patterns by age or gender. Since the LongROAD study recruited active primary care patients, the prevalence of physician conversations may be higher than in the general population. Additionally, findings were based on a sample of mainly well-educated, White, non-Hispanic drivers aged 65 to 79 and may not be generalizable to other populations. Findings also are not generalizable to older adults with cognitive impairment, who were excluded from the parent LongROAD study but are a population at risk of adverse driving outcomes.

CONCLUSION

In this study, most older adults did not report past driving discussions with family or physicians. Family and physicians may benefit from resources that address driving safety in the older population. Future interventions should utilize health education techniques targeting both families and physicians to raise awareness and provide information about this impending issue so they might knowledgeably participate in maintaining and improving older adults’ quality of life.

Funding Sources:

This research was supported by the AAA Foundation for Traffic Safety (Washington, DC) through the Longitudinal Research on Aging Drivers (LongROAD) Project and by a Paul Beeson Career Development Award Program [The National Institute on Aging; AFAR; The John A. Hartford Foundation; and The Atlantic Philanthropies; grant number-K23AG043123]. Dr. Guohua Li and Dr. Thelma J. Mielenz were also supported by Grant 1 R49 [Centers of Disease Control and Prevention; grant number-CE002096]. The contents of the manuscript are solely the responsibility of the authors and do not necessarily reflect the official views of the funding agencies.

REFERENCES

1. Pomidor A Clinician's guide to assessing and counseling older drivers (3rd edition). American Geriatrics Society and National Highway Traffic Safety Administration; 2016.
2. Foley DJ, Heimovitz HK, Guralnik JM, Brock DB. Driving life expectancy of persons aged 70 years and older in the United States. *Am. J. Public Health* 2002;92(8):1284–1289. [PubMed: 12144985]
3. Betz ME, Scott K, Jones J, DiGuseppi C. "Are you still driving?" Meta-synthesis of older adults' preferences for communication with healthcare providers about driving. *Traffic Inj Prev.* 2016;17(4):367–373. [PubMed: 26507251]
4. Liddle J, Gustafsson L, Mitchell G, Pachana NA. A difficult journey: Reflections on driving and driving cessation from a team of clinical researchers. *Gerontologist.* 2017;57(1):82–88. [PubMed: 27102058]
5. Meuser TM, Carr DB, Unger EA, Ulfarsson GF. Family reports of medically impaired drivers in Missouri: cognitive concerns and licensing outcomes. *Accid. Anal. Prev* 2015;74:17–23. [PubMed: 25463940]
6. D'Ambrosio L, Coughlin J, Mohyde M, Gilbert J, Reimer B. Family Matter: Older Drivers and the Driving Decision. *Transportation Research Record: Journal of the Transportation Research Board.* 2007;2009:23–39.
7. Lum HD, Brown JB, Juarez-Colunga E, Betz ME. Physician involvement in life transition planning: a survey of community-dwelling older adults. *BMC Fam. Pract* 2015;16(1):92. [PubMed: 26219548]
8. Betz ME, Jones J, Petroff E, Schwartz R. "I Wish We Could Normalize Driving Health:" A Qualitative Study of Clinician Discussions with Older Drivers. *J. Gen. Intern. Med* 2013;28 (12): 1573–1580. [PubMed: 23715688]
9. Huseth-Zosel AL, Sanders G, O'Connor M. Predictors of health care provider anticipatory guidance provision for older drivers. *Traffic Inj Prev.* 2016;17(8):815–820. [PubMed: 26940031]
10. Betz ME, Kanani H, Juarez-Colunga E, Schwartz R. Discussions about driving between older adults and primary care providers. *J Am Geriatr Soc.* 2016;64(6):1318–1323. [PubMed: 27321612]
11. Li G, Eby DW, Santos R, et al. Longitudinal Research on Aging Drivers (LongROAD): Study design and methods. *Injury Epidemiology* 2017;4(1):22. [PubMed: 28736796]
12. Callahan CM, Unverzagt FW, Hui SL, Perkins AJ, Hendrie HC. Six-item screener to identify cognitive impairment among potential subjects for clinical research. *Med. Care* 2002;40(9):771–781. [PubMed: 12218768]
13. Eby DW, Molnar LJ, Shope JT, Vivoda JM, Fordyce TA. Improving older driver knowledge and self-awareness through self-assessment: the driving decisions workbook. *J Safety Res.* 2003;34(4): 371–381. [PubMed: 14636659]
14. Dugan E *The Driving Dilemma: The Complete Resource Guide for Older Drivers and Their Families.* 2006.
15. Jang RW, Man-Son-Hing M, Molnar FJ, et al. Family physicians' attitudes and practices regarding assessments of medical fitness to drive in older persons. *J. Gen. Intern. Med* 2007;22(4):531–543. [PubMed: 17372806]
16. Huseth-Zosel AL, Sanders G, O'Connor M, Fuller-Iglesias H, Langley L. Health Care Provider Mobility Counseling Provision to Older Adults: A Rural/Urban Comparison. *J. Community Health* 2016;41(1):1–10. [PubMed: 26070871]

17. Leinberger RL, Janz NK, Musch DC, Niziol LM, Gillespie BW. Discussing driving concerns with older patients: I. vision care providers' attitudes and behaviors. *JAMA Ophthalmology*. 2013;131(2):205–212. [PubMed: 23544207]
18. Keskinen E Education for older drivers in the future. *IATSS Research*. 2014;38(1):14–21.
19. Code of Medical Ethics Opinion 8.2: Impaired Drivers & Their Physicians. <https://www.ama-assn.org/delivering-care/ethics/impaired-drivers-their-physicians>. Accessed January 8, 2019.
20. Senior Driving: State Laws. 2019; <https://seniordriving.aaa.com/states/>. Accessed January 8, 2019.
21. Adler G Driving decision-making in older adults with dementia. *Dementia*. 2010;9(1):45–60.
22. Choi M, Adams KB, Kahana E. Self-regulatory driving behaviors: gender and transportation support effects. *J. Women Aging* 2013;25(2):104–118. [PubMed: 23488647]
23. Betz ME, Dickerson A, Coolman T, Schold Davis E, Jones J, Schwartz R. Driving rehabilitation programs for older drivers in the United States. *Occup Ther Health Care*. 2014;28(3):306–317. [PubMed: 24971897]
24. Dickerson AE. Screening and assessment tools for determining fitness to drive: a review of the literature for the pathways project. *Occup Ther Health Care*. 2014;28(2):82–121. [PubMed: 24754758]
25. Bogner HR, Straton JB, Gallo JJ, Rebok GW, Keyl PM. The role of physicians in assessing older drivers: barriers, opportunities, and strategies. *J. Am. Board Fam. Pract* 2004;17(1):38–43. [PubMed: 15014051]




Figure 1. Reported outcomes of discussions with physicians, by age and gender (n=165).

Table 1.

Participant characteristics by reported discussions with family members or physicians (n=2990)

Characteristic	n	With Family (n=426)			With Physician (n=165)	
		%	95% CI	%	95% CI	
Total	2990	14.2	13.0–15.5	5.5	4.7–6.3	
Age (years)^a						
65–69	1243	13.8	11.9–15.8	5.6	4.3–6.8	
70–74	1037	12.5	10.5–14.6	5.5	4.1–6.9	
75–79	710	17.5	14.7–20.3	5.5	3.8–7.2	
Gender^a						
Male	1404	15.9	14.0–17.8	5.6	4.4–6.8	
Female	1586	12.8	11.2–14.4	5.4	4.3–6.5	
Race						
White	2557	14.6	13.2–16	5.5	4.6–6.4	
Black	212	11.3	7.1–15.6	5.7	2.5–8.8	
Asian	64	14.1	5.5–22.6	7.8	1.2–14.4	
Other	157	12.7	7.5–18.0	4.5	1.2–7.7	
Hispanic						
Yes	83	13.3	6.0–20.6	1.2	0.0–3.6	
No	2794	14.5	13.2–15.8	5.8	4.9–6.6	
Education^b						
Some College	1062	12.0	10.0–13.9	5.2	3.8–6.5	
Bachelors	698	12.0	9.6–14.4	4.3	2.8–5.8	
Masters	1221	17.5	15.4–19.7	6.6	5.2–7.9	
Income						
\$49,999	775	12.6	10.3–15	6.2	4.5–7.9	
\$50,000–\$79,999	719	14.3	11.8–16.9	7.0	5.1–8.8	
\$80,000–\$99,999	431	14.8	11.5–18.2	5.8	3.6–8.0	
\$100,000	959	16.2	13.8–18.5	4.1	2.8–5.3	
Paid employment						
Yes	904	13.4	11.3–15.7	4.2	2.9–5.5	
No	2084	14.5	13.0–16.1	6.1	5.1–7.1	
Marital status^a						
Married/Living with Partner	1974	15.3	13.7–16.9	5.5	4.5–6.5	
Separated/Divorced	608	11.3	8.8–13.9	6.1	4.2–8.0	
Widowed	378	13.0	9.6–16.4	4.5	2.4–6.6	
Traffic event^e						
Yes	703	16.1	13.4–18.8	5.1	3.5–6.8	
No	2286	13.7	12.3–15.1	5.6	4.7–6.5	
Driving reduction^{b,d}						
No reduction	2442	13.4	12.0–14.7	5.0	4.1–5.9	
For self-regulation	173	26.6	20.0–33.2	13.9	8.7–19.0	
For other reason(s)	355	13.5	10.0–17.1	5.4	3.0–7.7	
Other people...						
...depend on you for rides						
Yes	762	15.9	13.3–18.5	5.2	3.7–6.8	
No	2222	13.7	12.3–15.1	5.6	4.6–6.5	
...can give you rides						
Yes	2826	14.2	12.9–15.5	5.7	4.9–6.6	
No	148	13.5	8.0–19.0	2.0	0–4.3	
Driving scales^f			Mean (SD)		Mean (SD)	
Self-rated average ability ^{b,d}			5.65 (0.73)		5.67 (0.75)	
Errors ^b			1.54 (0.34)		1.47 (0.35)	

Characteristic	n	With Family (n=426)		With Physician (n=165)	
		%	95% CI	%	95% CI
Violations ^b		1.7	(0.41)	1.61	(0.37)
Strategic Self Regulation ^{b,d}		2.96	(2.42)	3.22	(2.56)
Tactical Self Regulation ^{b,e}		3.18	(1.7)	3.26	(1.68)

Family discussion prevalence differs by characteristic at $p < 0.05^a$ or $p < 0.001^b$ via Chi Square tests; Physician discussion prevalence differs at $p < 0.05^c$ or $p < 0.001^d$ via Chi Square tests or Fisher's exact tests

^e“Traffic event” was 1 motor vehicle crash, police stop, or traffic ticket within the past year

^fSee Methods for full description of scales

Table 2.

Factors associated with older adults reporting driving discussions with family members or physicians

Characteristic		Family member AOR (95%CI)	Physician AOR (95%CI)
Age (years)	65–69	1.00 (Ref)	--
	70–74	0.86 (0.6–1.12)	--
	75–79	1.16 (0.88–1.53)	--
Gender	Male	1.32 (1.05–1.66) ^a	--
	Female	1.00 (Ref)	--
Hispanic	Yes	--	0.24 (0.03–1.75)
	No	--	1.00 (Ref)
Education	Some College	1.00 (Ref)	1.00 (Ref)
	Bachelors	0.92 (0.67–1.27)	1.03 (0.64–1.68)
	Masters	1.65 (1.27–2.13) ^b	1.77 (1.17–2.68) ^b
Income	\$49,999	--	1.00 (Ref)
	\$50,000–\$79,999	--	1.13 (0.73–1.75)
	\$80,000–\$99,999	--	0.91 (0.53–1.57)
	\$100,000	--	0.69 (0.42–1.13)
Employed (paid)	No	--	1.00 (Ref)
	Yes	--	0.75 (0.51–1.11)
Traffic event^d	No	1.00 (Ref)	--
	Yes	1.27 (0.99–1.63)	--
Driving reduction	None	1.00 (Ref)	1.00 (Ref)
	Self-regulation	1.68 (1.12–2.53) ^b	2.10 (1.25–3.50) ^b
	Other reason(s)	0.94 (0.66–1.33)	1.14 (0.69–1.90)
Others...	...dependent for rides		
	No	1.00 (Ref)	--
	Yes	1.23 (0.96–1.58)	--
...can give rides	No	--	1.00 (Ref)
	Yes	--	3.13 (0.98–10.04)
Driving scales^e	Self-Rated Driving Ability	0.67 (0.56–0.79) ^c	0.75 (0.59–0.95) ^a
	Errors	1.65 (1.13–2.39) ^a	--
	Violations	1.45 (1.06–1.99) ^a	--
	Strategic Self-regulation	1.09 (1.04–1.15) ^b	1.10 (1.03–1.19) ^b
	Tactical Self-regulation	--	--

^a $P < 0.05$;^b $P < 0.01$;^c $P < 0.001$

AOR: Adjusted odds ratio (adjusted for all other characteristics in AOR model)

^d“Traffic event” was 1 crash, police stop, or traffic ticket within the past year

^eSee Methods for full description of scales

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript