



Published in final edited form as:

Bio Protoc. 2015 March 5; 5(5): . doi:10.21769/BioProtoc.1408.

Differentiation of Naturally Produced Extracellular Membrane Vesicles from Lipid Aggregation by Glucuronoxylomannan Immunogold Transmission Electron Microscopy in *Bacillus subtilis*

Lisa Brown^{1,*}, Geoff Perumal², Arturo Casadevall^{1,3}

¹Department of Microbiology and Immunology, Albert Einstein College of Medicine, New York, USA;

²Analytical Imaging Facility, Albert Einstein College of Medicine, New York, USA;

³Department of Medicine, Albert Einstein College of Medicine, New York, USA

Abstract

Recently, membrane vesicle (MV) production was described in Gram-positive bacteria, which harbor a variety of components such as toxins, antibiotic resistance proteins, proteases, DNA, and immune modulators. Free lipids have the ability to form micelles, thus it is important to rule out spontaneous association of lipids into vesicle-like structures and rather, that MVs are produced naturally by a metabolically active cell. Here, we describe a protocol utilizing the polysaccharide, glucuronoxylomannan (GXM) from *Cryptococcus neoformans* (*C. neoformans*) as a marker to differentiate naturally produced MVs from vesicles that form spontaneously in the Gram-positive model organism, *Bacillus subtilis* (*B. subtilis*). MVs are purified from bacterial cultures grown in the presence of GXM; MVs naturally produced by cells would not contain GXM in the lumen whereas vesicular structures forming in the media could encapsulate GXM and this can be visualized via immunogold transmission electron microscopy.

Materials and Reagents

1. Purified glucuronoxylomannan (GXM) (protocol included)
2. YPD broth (Difco, catalog number: 242820)
3. *C. neoformans* H99 fungal strain
4. 100 kDa, 50 kDa, 10 kDa, 1 kDa cut-off EMD Millipore Ultrafiltration Membranes (Millipore, catalog number: 14442AM)
5. *B. subtilis* 168 bacterial strain
6. BHI broth (Difco, catalog number: 299070)
7. 8% glutaraldehyde (Polysciences, catalog number: 111-30-8)

*For correspondence: Lisa.Brown@phd.einstein.yu.edu.

8. 16% paraformaldehyde (Electron Microscopy Sciences, catalog number: 15700)
9. 0.5 M sodium cacodylate (pH 7.4) (Electron Microscopy Sciences, catalog number: 11650)
10. 100% ethanol
11. Lowicryl HM-20 monostep resin (Electron Microscopy Sciences, catalog number: 14340)
12. Gelatin (Thermo Fisher Scientific)
13. Aurion Donkey Block (Electron Microscopy Sciences, catalog number: 25599)
14. α -GXM monoclonal antibody 18B7 (mouse monoclonal IgG1) (Casadevall lab-generated)
15. BSA-c (Electron Microscopy Sciences, catalog number: 25557)
16. 10 nm conjugated secondary Ab (donkey α -mouse) (Electron Microscopy Sciences, catalog number: 25814)
17. 4% uranyl acetate (aq) (SPI Supplies, catalog number: 615-44-0)
18. 1 \times phosphate buffered saline (PBS) (see Recipes)

Equipment

1. Express PLUS Membrane Filters - Pore (0.22 μ m) (Millipore, catalog number: SCGVU01RE)
2. Ultracentrifugation tubes (thickwall, polyallomer/polypropylene, 3.5 ml, 13 \times 51 mm) (Beckman Coulter, catalog number: 349623)
3. 200 mesh nickel grids (Polysciences, catalog number: 24916)
4. Centrifuge capable of 15,000 \times g
5. TLA 100.3 rotor (Beckman Coulter, catalog number: 349490)
6. Amicon ultrafiltration system (Millipore, catalog number: 5124)
7. Optima TL ultracentrifuge (Beckman Coulter, discontinued comparable to B11229)
8. Sonicator (sonic dismembrator) (Thermo Fisher Scientific, model: 100 cpn-214-161)
9. Freeze substitution system (RMC, model: FS-7500)
10. Reichert Ultracut UCT Ultramicrotome
11. JEOL 100CXII or JEOL 1200EX Electron Microscopes

Procedure

A. Purification of GXM

1. Grow 1 L of YPD broth inoculated with *C. neoformans* strain H99 overnight at 37 °C with shaking (~18 h 200 rpm).
2. Spin out cells by centrifuging 15,000 *x g* for 20 min at 4 °C.
3. Filter cell-free supernatant with 0.22 µm Express PLUS Membrane Filters to remove debris.
4. Sequentially concentrate the cell-free supernatant with 100 kDa, 50 kDa, 10 kDa, and 1 kDa cutoff EMD Millipore Ultrafiltration Membranes. The jelly that forms on the filters is GXM (save jelly that is <10 kDa and >1 kDa) (see Figure 1 for concentrator set-up).

B. Purification of gram-positive extracellular vesicles

1. Inoculate 100 ml of BHI broth with fractionated GXM <10 kDa and *B. subtilis* strain 168.
2. Grow overnight at 37 °C with shaking (~18 h 200 rpm).
3. Centrifuge culture to remove cells at 15,000 *x g* for 20 m at 4 °C.
4. Filter cell-free supernatant with 0.22 µm Express PLUS Membrane Filters to remove debris.
5. Concentrate cell-free supernatant to small volumes (approximately 6 ml) with ultrafiltration system and 100 kDa cut-off EMD Millipore Ultrafiltration Membranes (Figure 1).
6. Ultracentrifuge concentrated supernatant at 100,000 *x g* for 1 h at 4 °C in thickwall, polyallomer, 3.5 ml, 13 × 51 mm tubes to pellet vesicles (Figure 2).
7. Wash/resuspend vesicle pellet with 500 µl PBS and repeat spin (repeat wash twice).
8. Remove supernatant from sample without disturbing pellet.
 - a. For sonication (production of disrupted/unnatural vesicles-positive control) resuspend pellet in PBS and sonicate (20 sec sonication and 30 sec incubation at 4 °C on ice, repeated 4 times at power 2).
 - b. Ultracentrifuge again to pellet vesicles and carefully remove supernatant.
9. Carefully add 4% paraformaldehyde, 0.1% glutaraldehyde in 0.1 M sodium cacodylate without disturbing pellet.
10. Incubate for 1 h at room temperature (RT).

C. Sample preparation for immunogold labeling

1. After fixing for 1 h at RT, wash with 0.1 M sodium cacodylate 3 times.

2. Enrobe samples in 5% gelatin.
3. Dehydrate through a graded ethanol series in a RMC FS 7500; progressively lowering the temperature 5 °C /h from 4 °C to –50 °C.
4. Use Lowicryl HM-20 monostep resin to embed sample.
5. Polymerize embedded sample with UV light for 48 h at –50 °C.
6. Use Reichert Ultracut UCT to cut 80 nm ultrathin sections onto 200 mesh nickel grids.

D. Immunogold labeling of GXM in vesicle preparations

1. Block with Aurion Donkey Block for 1 h.
2. Incubate samples with the primary α -GXM mAb antibody, 18B7, in 0.1% BSA-c/PBS incubation buffer for ~18 h at 4 °C.
3. Wash samples with 0.1% BSA-c/PBS.
4. Incubate samples with the secondary donkey α -mouse IgG 10 nm immunogold-conjugated antibody for 2 h at RT.
5. Wash samples with BSA-C.
6. Post-fix samples for 5 m at 25 °C with 2% glutaraldehyde/PBS to stabilize the gold particles.
7. Counterstain samples for 15 min 25 °C with 4% uranyl acetate (aq).
8. Also make control samples with an irrelevant IgG antibodies and without the 1^o antibody (18B7).

Representative data

Vesicles are not incredibly stable so it is suggested to process them immediately after isolation. Only Beckman polyallomer/propylene ultracentrifugation tubes can be used for IEM preparation. Polycarbonate tubes cannot be used for this procedure but can be used when purifying vesicles for other experiments. The sonicated (unnatural) vesicles serve as a positive control. It is expected that vesicles from the sonicated vesicle preparation will contain significantly more gold particles than the naturally produced vesicles. One may also expect that sonicated vesicles have a smaller and less variable diameter than that of natural vesicles.

Notes

B. subtilis strain 168 concentrates very quickly but other strains of bacteria may not. *B. subtilis* strain 168 produces a large quantity of recoverable vesicles and only requires 100 ml of culture whereas other bacteria may require volumes larger than 1 L. Concentrating to small volumes allows for less ultracentrifugation spins and cuts down on the use of the ultracentrifuge tubes. Ultracentrifuge tubes can be washed and re-used but be aware there may be some contamination.

Recipes

1. 1 L 1× phosphate buffered saline (PBS) (pH 7.4)
137 mM NaCl
2.7 mM KCl
10 mM Na₂HPO₄
1.8 mM KH₂PO₄
Bring to pH 7.4 with HCl
Dissolve in H₂O up to 1 L

Acknowledgements

Funding from NIH Grant Numbers: HL059842, AI033774, AI033142, AI052733 and Center for AIDS Research at Albert Einstein College of Medicine.

Protocol is adapted from Brown *et al.* (2014) and vesicle purification is adapted from Prados-Rosales *et al.* (2014).

References

1. Brown L, Kessler A, Cabezas-Sanchez P, Luque-Garcia JL and Casadevall A (2014). Extracellular vesicles produced by the Gram-positive bacterium *Bacillus subtilis* are disrupted by the lipopeptide surfactin. *Mol Microbiol* 93(1): 183–198. [PubMed: 24826903]
2. Prados-Rosales R, Brown L, Casadevall A, Montalvo-Quirós S and Luque-Garcia JL (2014). Isolation and identification of membrane vesicle-associated proteins in Gram-positive bacteria and *mycobacteria*. *MethodsX* (1): 124–129. [PubMed: 26150943]



Figure 1. Amicon ultrafiltration setup.

The ultrafiltration cells are connected to compressed nitrogen gas. The gas applies pressure, forcing the supernatant through the 100 kDa filter membrane. Vesicles stay in the concentrated supernatant while smaller proteins and liquid supernatant flow into waste container.



Figure 2. Beckman ultracentrifuge tubes.
Ultracentrifugation tubes utilized to pellet vesicles.




Figure 3. Transmission electron micrograph of immunogold labeling of GXM in vesicles from *Bacillus subtilis*. Micrograph shows vesicles and gold particles. Gold particles indicate the presence of GXM. Notice all of the gold is outside of vesicles. White arrows indicate vesicles and black arrows indicate gold particles. Scale bar = 100 nm