

Robust CD4⁺ T-cell recovery in adults transplanted with cord blood and no antithymocyte globulin

Ioannis Politikos,^{1,2} Jessica A. Lavery,³ Patrick Hilden,³ Christina Cho,^{1,2} Taylor Borrill,¹ Molly A. Maloy,¹ Sergio A. Giralt,^{1,2} Marcel R. M. van den Brink,^{1,2} Miguel-Angel Perales,^{1,2,*} and Juliet N. Barker^{1,2,*}

¹Adult Bone Marrow Transplantation Service, Department of Medicine, Memorial Sloan Kettering Cancer Center, New York, NY; ²Department of Medicine, Weill Cornell Medical College, New York, NY; and ³Department of Epidemiology and Biostatistics, Memorial Sloan Kettering Cancer Center, New York, NY

Key Points

- ATG-free CBT in adults is associated with rapid CD4⁺-biased, thymus-independent, T-cell reconstitution.
- Recovery of CD4⁺ T cells and T-cell function is associated with improved survival in adult CBT recipients.

Quality of immune reconstitution after cord blood transplantation (CBT) without antithymocyte globulin (ATG) in adults is not established. We analyzed immune recovery in 106 engrafted adult CBT recipients (median age 50 years [range 22-70]) transplanted for hematologic malignancies with cyclosporine/mycophenolate mofetil immunoprophylaxis and no ATG. Patients were treated predominantly for acute leukemia (66%), and almost all (96%) underwent myeloablation. Recovery of CD4⁺ T cells was faster than CD8⁺ T cells with median CD4⁺ T-cell counts exceeding 200/mm³ at 4 months. Early post-CBT, effector memory (EM), and central memory cells were the most common CD4⁺ subsets, whereas effector and EM were the most common CD8⁺ T-cell subsets. Naive T-cell subsets increased gradually after 6 to 9 months post-CBT. A higher engrafting CB unit infused viable CD3⁺ cell dose was associated with improved CD4⁺ and CD4⁺CD45RA⁺ T-cell recovery. Cytomegalovirus reactivation by day 60 was associated with an expansion of total, EM, and effector CD8⁺ T cells, but lower CD4⁺ T-cell counts. Acute graft-versus-host disease (aGVHD) did not significantly compromise T-cell reconstitution. In serial landmark analyses, higher CD4⁺ T-cell counts and phytohemagglutinin responses were associated with reduced overall mortality. In contrast, CD8⁺ T-cell counts were not significant. Recovery of natural killer and B cells was prompt, reaching medians of 252/mm³ and 150/mm³ by 4 months, respectively, although B-cell recovery was delayed by aGVHD. Neither subset was significantly associated with mortality. ATG-free adult CBT is associated with robust thymus-independent CD4⁺ T-cell recovery, and CD4⁺ recovery reduced mortality risk.

Introduction

Cord blood (CB) is a valuable alternative hematopoietic stem cell (HSC) source for patients who lack suitable adult donors, especially racial and ethnic minorities.^{1,2} Double-unit CB grafts have successfully extended cord blood transplantation (CBT) to larger children and adults,³ and both single- and double-unit CBT has been associated with potent graft-versus-leukemia (GVL) effects,^{4,5} low rates of chronic graft-versus-host disease (GVHD),⁶⁻⁸ and high rates of disease-free survival in patients with hematologic malignancies.^{4-6,8,9} CBT, however, has also been associated with delayed immune reconstitution compared with T-cell replete HLA-matched adult donor allografts with multiple reports of high infection rates early posttransplant.¹⁰⁻¹³

CB grafts contain low numbers of progenitor stem and immune cells compared with adult donor HSC grafts.¹⁴ In addition, CB-derived lymphocyte populations have unique phenotypic and immunological

Submitted 14 August 2019; accepted 18 November 2019; published online 14 January 2020. DOI 10.1182/bloodadvances.2019000836.

*M.-A.P. and J.N.B. contributed equally to this study.

To request original data, contact corresponding author, Ioannis Politikos (politiki@mskcc.org).

The full-text version of this article contains a data supplement.

© 2020 by The American Society of Hematology

properties, including almost exclusively naive T cells that do not transfer immune memory.^{15,16} Although these CB graft attributes could contribute to delayed immune reconstitution, many previous CBT series have included antithymocyte globulin (ATG), a platform that has detrimental effects on both immune reconstitution and survival after CBT.¹⁷⁻²² Notably, low ATG exposure or omission of ATG has been associated with rapid thymus-independent T-cell expansion and robust immune reconstitution in pediatric CBT recipients.^{19,22-25} In contrast to children, however, relatively little is known about immune reconstitution after ATG-free CBT in adults.^{12,26-30}

Herein, we report the kinetics of immune reconstitution in a large cohort of adult CBT recipients transplanted for hematologic malignancies at a single center without ATG. We also analyzed the impact of patient, graft, and early posttransplant factors on immune recovery, as well as the immune variables associated with improved survival. Our hypothesis was that, similar to pediatric series, ATG-free adult CBT is associated with prompt immune reconstitution and that early T-cell recovery improves survival post-CBT.

Methods

Patient and transplant characteristics

All consecutive adult patients ≤ 70 years old who underwent first allogeneic transplantation using single- or double-unit CB grafts for the treatment of hematologic malignancies at Memorial Sloan Kettering Cancer Center (MSKCC) between April 2012 and May 2016 were eligible for analysis ($n = 114$). Those who did not achieve CB-derived engraftment ($n = 4$) or had no immune reconstitution assays performed due to development of fatal early posttransplant complications before day 30 ($n = 4$) were excluded. Of the 106 evaluable patients, 93 were treated on Institutional Review Board (IRB)-approved protocols (#NCT00739141, #NCT01682226, and #NCT00387959). The remaining 13 patients were treated off protocol due to either protocol ineligibility ($n = 8$) or insurance denial for clinical trials in otherwise eligible patients ($n = 5$).

CB units were at least 4/6 HLA-A, -B antigen, -DRB1 allele matched to the recipient, and each unit had a cryopreserved total nucleated cell (TNC) dose $\geq 1.5 \times 10^7/\text{kg}$. High-resolution HLA typing, CD34⁺ cell dose, CB quality, and bank of origin were also considered in unit selection as previously detailed.³¹ Some patients also received mobilized peripheral blood-derived haploidentical CD34⁺ cells as a myeloid bridge prior to CB engraftment (#NCT01682226).

Most patients received myeloablative conditioning (outlined in Table 1). GVHD prophylaxis was with cyclosporine-A/mycophenolate mofetil, and no patient received ATG. Granulocyte colony-stimulating factor was given starting day +7. Cytomegalovirus (CMV)-seropositive patients were monitored by quantitative polymerase chain reaction as of day +14 posttransplant. Preemptive antiviral therapy was given upon detection of any level of CMV viremia.

This study was performed with MSKCC IRB approval.

Immune reconstitution monitoring

Immune reconstitution monitoring was per standard MSKCC clinical practice and not influenced by protocol enrollment. Flow cytometric

Table 1. Patient and graft characteristics (n = 106)

Variable	Value
Median age (range), y	50 (22-70)
Male, n (%)	55 (52)
Median weight (range), kg	80 (36-138)
Recipient CMV ⁺ , n (%)	59 (56)
Diagnosis, n (%)	
Acute leukemia (AML/ALL/other)	70 (66)
MDS/MPN	14 (13)
Lymphoma (NHL/HD)	22 (21)
Conditioning, n (%)*	
High intensity	1 (1)
Intermediate intensity	101 (95)
Nonmyeloablative	4 (4)
Donor-recipient 8-allele HLA match,† median (range)	5 (3 to 7)
Infused TNC dose $\times 10^7/\text{kg}$,‡ median (range)	2.35 (1.23-5.31)
Infused viable CD34 ⁺ dose $\times 10^5/\text{kg}$,‡ median (range)	1.18 (0.18-4.08)
Infused viable CD3 ⁺ dose $\times 10^6/\text{kg}$,‡ median (range)	3.34 (0.45-10.61)
Graft composition, n (%)‡	
dCB	45 (42)
dCB-haploCD34 ⁺	59 (56)
sCB-haploCD34 ⁺	2 (2)

ALL, acute lymphoblastic leukemia; AML, acute myeloid leukemia; HD, Hodgkin disease; MDS, myelodysplastic syndrome; MPN, myeloproliferative neoplasm; NHL, non-Hodgkin lymphoma.

*High-intensity myeloablative conditioning was with cyclophosphamide 120 mg/kg, fludarabine 75 mg/m², and total body irradiation (TBI) 1320 cGy; intermediate-intensity myeloablative was with cyclophosphamide 50 mg/kg, fludarabine 150 mg/m², thiotepa 5 to 10 mg/kg, and TBI 400 cGy³³; nonmyeloablative included cyclophosphamide 50 mg/kg, fludarabine 150 mg/m², and TBI 200 cGy ($n = 3$), or fludarabine 150 mg/m² and TBI 400 cGy ($n = 1$).

†Engrafting CB unit.

‡Fifty-nine double-unit CB (dCB) grafts and 2 single-unit CB (sCB) grafts were supplemented with haploidentical CD34⁺ cells to provide a myeloid bridge prior to CB engraftment.

immunophenotyping for the monitoring of absolute lymphocyte count (ALC) and lymphocyte subset recovery was performed prospectively on fresh whole blood samples at the MSKCC Clinical Immunology Laboratory using BD FACS Canto II and BD FACS Canto 10 color flow cytometers. Until April 2014, analyzed lymphocyte subsets included total T cells (CD3⁺), total CD4⁺, CD4⁺45RA⁺, and CD8⁺ T cells, natural killer (NK; CD3⁻CD56⁺CD16⁺), and B cells (CD19⁺). Since May 2014, subset analysis of CD4⁺ and CD8⁺ naive (CCR7⁺CD45RA⁺), central memory (CM; CCR7⁺CD45RA⁻), effector memory (EM; CCR7⁻CD45RA⁻), effector (CCR7⁻CD45RA⁺), and activated (CD38⁺HLADR⁺) T cells, and naive (CD27⁻IgD⁺), nonswitched memory (CD27⁺IgD⁺), switched memory (CD27⁺IgD⁻), transitional (CD24⁺CD38⁺), and plasmablast (CD27⁺CD20⁻CD38⁺) B cells was performed. T-cell proliferative responses to mitogen phytohemagglutinin (PHA) were assessed by measuring 3H-thymidine incorporation. Assessments were usually performed at days 30, 60, 120, 180, 270, 1 year, 18 months, and 2 years post-CBT. Immune recovery data were censored at the time of second allogeneic transplantation ($n = 1$) or after infusion of viral-specific cytotoxic T cells ($n = 5$; 4 CMV specific, 1 Epstein-Barr virus specific).

Statistical analysis

Acute graft-versus-host disease (aGVHD) was diagnosed clinically with histologic confirmation when possible and was graded according to International Bone Marrow Transplant Registry criteria.³² CMV reactivation was defined as detection of CMV DNA by quantitative polymerase chain reaction regardless of the viremia level. Transplant-related mortality (TRM) was defined as death from any cause other than disease relapse or progression. The cumulative incidence of aGVHD, CMV reactivation, relapse, and TRM was calculated in the competing risks framework considering relapse/death without developing aGVHD, death without CMV reactivation, TRM, and relapse as the competing events, respectively. Overall survival (OS) and progression-free survival estimates were calculated using Kaplan-Meier methodology.

Loess-smoothed curves were estimated to visually inspect immune recovery trends over time. The association between baseline or early posttransplant factors and immune variables was investigated with linear mixed effects models. Immune recovery data were natural log transformed before the analysis. The mixed effects models contained linear and quadratic terms for time, as well as random intercepts for each patient and fixed effects for variables of interest. From the mixed effects models, the slopes and the 95% confidence intervals (CIs) of the baseline and early posttransplant factors were estimated. Models for day 60 CMV reactivation and day 100 aGVHD considered only immune recovery data beyond 60 and 100 days, respectively.

Landmark analyses at 2, 4, and 6 months were performed to investigate the association between each of the immune variables and OS. For the 2-, 4-, and 6-month landmark analyses, the most recent immune parameter values prior to day 65, 130, and 195 were used, respectively. Univariable OS analyses were first performed for each immune variable, baseline factors, and early posttransplant factors. Exploratory analyses were also performed to potentially identify an optimal CD4⁺ T-cell count based on its univariable association with OS at each of the landmark time points. Multivariable Cox regression models were built for each immune variable that was statistically significant controlling for significant baseline and/or posttransplant covariates. Immune variables were highly correlated with each other and therefore could not be included in the same models (correlations were assessed using Spearman's rank correlation coefficient).

Tests with $P < .01$ were considered statistically significant, to account for multiple comparisons. All statistical analyses were performed in R version 3.5.3 (The R Foundation for Statistical Computing).

Results

Patient and graft characteristics

The characteristics of the 106 analyzed patients (median 50 years, range 22 to 70) and their grafts are summarized in Table 1. The most common diagnosis was acute leukemia (66%), followed by lymphoma (21%), or myelodysplasia/myeloproliferative disease (13%). The majority of patients (95%) received intermediate-intensity myeloablative conditioning.³³

Nearly all patients ($n = 104$) received double-unit CB grafts. The median infused TNC and infused viable CD34⁺ cell doses of the engrafting unit were 2.35 (range 1.23-5.31) $\times 10^7$ /kg and 1.18

(range 0.18-4.08) $\times 10^5$ /kg, respectively. The median 8-allele engrafting unit-recipient HLA match was 5/8 (range 3/8-7/8).

Transplant outcomes

In all of the 106 patients (including all 61 recipients of CB grafts supplemented with haploidentical CD34⁺ cells), long-term hematopoiesis was mediated by 1 CB unit (termed the engrafting unit), with no contribution from the host. Of the 59 CMV-seropositive patients, 44 reactivated CMV by day 60 for a cumulative incidence of 75% (95% CI, 61-84). The cumulative incidence of grade II to IV and III to IV aGVHD by day 100 was 77% (95% CI, 68-84) and 17% (95% CI, 11-25), respectively. Of the 64 patients with grade II day 100 aGVHD, 22 (34%) were treated with systemic corticosteroids by day 100; remaining patients were treated with nonabsorbable oral or topical corticosteroids.

With a median survivor follow-up of 2 years (range 7 months to 4.7 years), the 2-year estimates of OS and progression-free survival for the 106 patients evaluable for immune recovery (8 nonevaluable patients with graft failure or early death excluded) were 73% (95% CI, 62-81) and 69% (95% CI, 58-78), respectively. Two-year TRM was 19% (95% CI, 12-28) and 2-year relapse was 12% (95% CI, 6-20).

T-, NK-, and B-cell immune reconstitution

The median ALC surpassed 500/mm³ by 2 months. The speed of recovery varied markedly between lymphocyte subsets. NK cells were the most common lymphocyte population early posttransplant, followed by CD3⁺ T cells, and B cells. However, CD3⁺ T cells comprised the largest lymphocyte subset beyond 4 months (Table 2; Figure 1).

Of CD3⁺ T-cell subsets, CD4⁺ lymphocytes recovered faster than CD8⁺ lymphocytes (Table 2; Figure 2A). The median CD4⁺ T-cell count surpassed 200/mm³ by 4 months. CD4⁺CD45RA⁺ T cells, considered to represent naive CD4⁺ T cells, remained low in the first 6 to 9 months post-CBT (Figure 2A). CD8⁺ T cells increased steadily until 15 to 18 months post-CBT, followed by a relative contraction of the CD8⁺ T-cell compartment (Figure 2A). T-cell function, as assessed by PHA responses, increased rapidly and remained relatively stable beyond 6 months (Figure 2B).

Eighty-eight patients had 1 or more time points assayed with the extended lymphocyte subset immunophenotyping panel. EM and CM cells were the most prevalent CD4⁺ subsets post-CBT, followed by naive and effector subsets (Figure 2C). EM and CM CD4⁺ T-cell numbers increased rapidly in the first 6 months and remained relatively stable between 9 and 24 months. Naive CD4⁺ T cells increased steadily beyond 6 months. In contrast to CD4⁺ T-cell subsets, effector CD8⁺ T cells comprised the majority of CD8⁺ T cells post-CBT, followed by the EM CD8⁺ subset (Figure 2D). Following their marked initial increase, effector CD8⁺ T cells decreased beyond 18 months posttransplant. Recovery of naive CD8⁺ T cells paralleled that of naive CD4⁺ T cells beyond 6 months post-CBT (Figure 2D).

NK cells increased rapidly in the first 2 months post-CBT and remained stable thereafter (Table 2; Figure 1). B cells also rose rapidly with the median B-cell count reaching normal range by 4 months posttransplant (Table 2; Figure 1). This was explained by a marked increase of naive B cells in the first 6 months post-CBT. Naive B cells continued to be the most prevalent B-cell subset

Table 2. Recovery of lymphocyte subsets and PHA responses after CBT

Immune variable, median (range)	Posttransplant time point			
	2 mo	4 mo	6 mo	1 y
ALC, cells/mm ³ (NR: 500-5300)	608 (85-7200)	745 (145-2025)	933 (173-6287)	1554 (384-4774)
CD3 ⁺ T cells, cells/mm ³ (NR: 822-1825)	171 (7-3024)	252 (8-1112)	380 (38-1572)	669 (148-3246)
CD4 ⁺ T cells, cells/mm ³ (NR: 429-1131)	120 (5-1584)	203 (8-912)	294 (30-1069)	420 (22-891)
CD8 ⁺ T cells, cells/mm ³ (NR: 209-768)	22 (0-1440)	25 (0-761)	47 (0-909)	204 (11-2574)
B cells, cells/mm ³ (NR: 87-441)	22 (0-3024)	150 (0-1357)	201 (0-3960)	324 (0-1968)
NK cells, cells/mm ³ (NR: 78-424)	222 (41-1333)	252 (29-717)	221 (65-1519)	220 (70-896)
PHA, CPM (NR: 109 576-256 486)	66 816 (231-156 128)	76 319 (667-162 912)	86 944 (14 590-161 140)	86 091 (6724-164 435)

CPM, counts per minute; NR, normal range.

through 2 years post-CBT (Figure 3). Transitional B cells were the next most common B-cell subset, whereas nonswitched memory, switched memory, and plasmablast B-cell subsets remained low (Figure 3).

Association of patient, graft, and posttransplant variables with immune reconstitution

The association of baseline and early posttransplant variables with recovery of lymphocyte subsets and PHA responses is shown in Tables 3 and 4. No patient, graft (including the addition of haploidentical CD34⁺ cells), or early posttransplant variables were significantly associated with ALC or CD3⁺ T-cell recovery (Table 3).

Patient characteristics, including age, sex, or diagnosis, were not significantly associated with CD4⁺ or CD8⁺ T-cell recovery. A

higher engrafting unit infused viable CD3⁺ cell dose was associated with higher CD4⁺ and CD4⁺CD45RA⁺ T-cell counts after CBT, although these associations were not significant at the 0.01 level (Table 4). Conversely, CMV reactivation was associated with lower CD4⁺ T cells (Figure 4A) and PHA responses post-CBT (Table 4). In contrast, CMV reactivation was the only factor significantly associated with higher CD8⁺ T-cell counts posttransplant (Table 4; Figure 4B).

The association of transplant variables with the recovery of CD4⁺ T-cell subsets is shown in supplemental Table 1. A higher engrafting unit CD34⁺ cell dose was associated with higher EM CD4⁺ T-cell counts, whereas a higher engrafting unit CD3⁺ dose was associated with higher naive CD4⁺ T-cell counts post-CBT, although these associations did not reach significance at the 0.01 level. CMV reactivation was associated with lower CM and, to a lesser extent, naive CD4⁺ T-cell counts (supplemental Figure 1). No transplant factors were associated with effector CD4⁺ T-cell reconstitution. Increased numbers of activated CD4⁺ T cells were observed in association with a higher engrafting unit CD34⁺ cell dose, whereas lower activated CD4⁺ T-cell counts were seen in patients with aGVHD, especially corticosteroid-requiring aGVHD (data not shown). These associations, however, did not reach the prespecified level of statistical significance.

The association of transplant variables with the recovery of CD8⁺ T-cell subsets is shown in supplemental Table 2. A higher engrafting unit CD34⁺ cell dose was associated with higher EM and, to a lesser extent, effector CD8⁺ T-cell counts after CBT. In addition, a marked expansion of EM and effector CD8⁺ T cells was observed in patients with CMV reactivation (supplemental Figure 2). No transplant factors were significantly associated with CM or naive CD8⁺ T-cell reconstitution. A higher number of activated CD8⁺ T cells were observed in association with higher CD34⁺ dose of the engrafting CB unit and CMV reactivation.

Older recipient age and higher CD34⁺ cell dose of the engrafting CB unit were associated with NK-cell recovery early posttransplant (Table 3). B-cell reconstitution was adversely affected by the development of grade II to IV aGVHD (Table 3), especially aGVHD requiring systemic corticosteroids (data not shown). The negative




Figure 1. ALC and lymphocyte subset recovery in adult CBT recipients. ALC, CD3⁺ cells, NK cells, B cells. Curves are Loess-smoothed averages.




Figure 2. Recovery of T-cell subsets and T-cell function in adult CBT recipients. (A) CD3⁺, CD4⁺, CD8⁺, CD4⁺CD45RA⁺ T cells. (B) PHA responses. (C) CD4⁺ and CD4⁺ T-cell subsets (naïve, CM, EM, effector). (D) CD8⁺ and CD8⁺ T-cell subsets (naïve, CM, EM, effector). Curves are Loess-smoothed averages. In panels C and D, Loess-smoothed curves for total CD4⁺ and total CD8⁺ T cells, respectively, are based on values from the time points of which the extended immunophenotyping panel was also performed.

effect of aGVHD was also observed in naïve, transitional, nonswitched memory, and to a lesser extent, switched memory and plasmablast B-cell subset recovery (data not shown). No other transplant factors were associated with B-cell recovery.

Association of immune variables with OS

Next, we investigated the association of immune reconstitution with OS in serial landmark analyses. At 2 months, recipient age was the only baseline factor associated with OS (Table 5). A higher PHA response was the only immune variable associated with improved OS in univariable and multivariable analysis controlling for age. This was due to decreased TRM in patients with higher PHA responses, whereas relapse risk was unaffected (data not shown).

In the 4-month landmark analysis (Table 6), recipient age was again the only baseline factor impacting OS. In addition, there was no association of day 100 aGVHD grade with OS. In univariable analysis, 4-month ALC, total CD3⁺ and CD4⁺ T cells, and PHA responses were significantly associated with OS beyond 4 months. Although an association of higher B-cell counts with improved OS was also observed, it was not significant at the 0.01 level. In multivariable analyses controlling for age, each of the variables of ALC, CD3⁺ and CD4⁺ T cells, and PHA remained significant. As expected, CD4⁺ T-cell counts and PHA responses at 4 months were highly correlated ($r = 0.35$). Therefore, they could not be concurrently included in a multivariable model.

At 6 months (Table 7), the association of recipient age with OS was no longer significant at the 0.01 level ($P = .025$), but was retained in




Figure 3. Recovery of B cells and subsets in adult CBT recipients. Curves are Loess-smoothed averages. Loess-smoothed curves for total B-cell counts are based on values from the time points of which the extended immunophenotyping panel was also performed.

multivariable modeling because it is a clinically important factor. Of the immune variables, higher ALC, total CD3⁺ and CD4⁺ T cells, and PHA responses were significantly associated with subsequent improved OS in univariable and age-adjusted analyses. As with the 4-month landmark, CD4⁺ T-cell counts and PHA responses at 6 months were highly correlated ($r = 0.39$); therefore, their independent prognostic significance could not be examined.

CD4⁺CD45RA⁺ T-cell, CD8⁺ T-cell, or NK-cell counts were not associated with OS at any time point post-CBT (Tables 5-7). In

addition, due to the limited number of events in assayed patients, the prognostic value of specific CD4⁺ and CD8⁺ T-cell subsets could not be evaluated. Finally, in exploratory analyses, no optimal CD4⁺ T-cell count cutoff for OS could be identified at any time point post-CBT.

Discussion

This study represents the largest analysis of immune reconstitution after ATG-free CBT in adults to date. In contrast to the protracted T-cell lymphopenia reported after ATG-based CBT, we show rapid early thymus-independent CD4⁺ T-cell recovery consistent with that reported in pediatric ATG-free CBT.^{23,24} In this adult population with a median age of 50 years, the median CD4⁺ T-cell count already exceeded 200/mm³ by 4 months and increased steadily through 2 years posttransplant. EM and CM T cells were the most prevalent CD4⁺ subsets early post-CBT,^{11,34} whereas naive CD4⁺ T-cell counts gradually increased beyond 6 to 9 months post-CBT.

Recovery of CD8⁺ T cells was initially slow, but marked CD8⁺ T-cell expansion was noted beyond 6 months post-CBT.²⁸ Effector and EM were the most common CD8⁺ T-cell subsets early post-CBT.^{11,34} As with naive CD4⁺ T cells, naive CD8⁺ T-cell counts increased steadily beyond 6 to 9 months, suggesting thymus-dependent recovery from this time point, even despite the relatively advanced age of this patient cohort.¹⁵ Recovery of T-cell function, as assayed by PHA responses, was also prompt.

Of factors associated with T-cell reconstitution, a notable finding was the lack of an adverse effect of recipient age on total, CD4⁺, CD8⁺, and subset T-cell recovery. It is possible that the relatively older age of this patient cohort did not allow the detection of such

Table 3. Association of transplant variables with the recovery of ALC and total T, NK, and B cells

Variable	ALC		CD3 ⁺ T cells		NK cells		B cells	
	Slope (95% CI)	P	Slope (95% CI)	P	Slope (95% CI)	P	Slope (95% CI)	P
Age (10 y)	0.01 (−0.08, 0.09)	.897	−0.06 (−0.16, 0.04)	.260	0.15 (0.07, 0.23)	.001	−0.20 (−0.46, 0.07)	.146
8-allele HLA-match*		.135		.733		.327		.465
<5/8	reference		reference		reference		reference	
≥5/8	0.17 (−0.05, 0.39)		0.05 (−0.22, 0.31)		0.12 (−0.12, 0.35)		0.26 (−0.44, 0.96)	
Infused viable CD34⁺ cell dose*		.099		.840		.018		.245
<1.18 × 10 ⁶ /kg	reference		reference		reference		reference	
≥1.18 × 10 ⁶ /kg	0.17 (−0.03, 0.37)		0.03 (−0.22, 0.27)		0.25 (0.05, 0.46)		0.38 (−0.25, 1.01)	
Infused viable CD3⁺ cell dose*		.727		.185		.093		.612
<3.34 × 10 ⁶ /kg	reference		reference		reference		reference	
≥3.34 × 10 ⁶ /kg	0.04 (−0.16, 0.24)		0.16 (−0.08, 0.40)		−0.18 (−0.39, 0.03)		0.164 (−0.47, 0.80)	
CMV reactivation by day 60		.783		.391		.577		.488
No CMV reactivation	reference		reference		reference		reference	
CMV reactivation	−0.03 (−0.23, 0.17)		0.12 (−0.15, 0.38)		0.06 (−0.15, 0.27)		−0.26 (−1.01, 0.48)	
Day 100 aGVHD grade		.313		.960		.708		<.001
0-I	reference		reference		reference		reference	
II	−0.16 (−0.43, 0.11)		−0.05 (−0.41, 0.31)		0.09 (−0.19, 0.38)		−1.15 (−2.04, −0.26)	
III-IV	−0.26 (−0.60, 0.08)		−0.02 (−0.47, 0.43)		−0.01 (−0.37, 0.35)		−2.38 (−3.50, −1.25)	

There was no association with the variables of recipient sex, diagnosis (acute leukemia vs MDS/MPD vs lymphoma), or the addition of haploidentical CD34⁺ cells (data not shown). P values significant at the .01 level are indicated in bold.

*Engrafting CB unit.

Table 4. Association of transplant variables with the recovery of T-cell subsets and function

Variable	CD4 ⁺ T cells		CD4 ⁺ CD45RA ⁺ T cells		CD8 ⁺ T cells		PHA	
	Slope (95% CI)	P	Slope (95% CI)	P	Slope (95% CI)	P	Slope (95% CI)	P
Age (10 y)	-0.09 (-0.20, 0.01)	.087	-0.05 (-0.22, 0.12)	.539	0.01 (-0.15, 0.16)	.924	-0.08 (-0.21, 0.04)	.205
8-allele HLA-match*		.200		.545		.105		.565
<5/8	reference		reference		reference		reference	
≥5/8	0.19 (-0.10, 0.47)		0.14 (-0.30, 0.58)		-0.33 (-0.73, 0.07)		0.09 (-0.22, 0.41)	
Infused viable CD34⁺ cell dose*		.646		.333		.245		.574
<1.18 × 10 ⁶ /kg	reference		reference		reference		reference	
≥1.18 × 10 ⁶ /kg	-0.06 (-0.32, 0.20)		-0.20 (-0.60, 0.20)		0.22 (-0.15, 0.58)		0.08 (-0.21, 0.37)	
Infused viable CD3⁺ cell dose*		.045		.020		.893		.065
<3.34 × 10 ⁶ /kg	reference		reference		reference		reference	
≥3.34 × 10 ⁶ /kg	0.26 (0.01, 0.52)		0.47 (0.08, 0.87)		0.03 (-0.34, 0.39)		0.27 (-0.01, 0.56)	
CMV reactivation by day 60		.005		.171		<.001		<.001
No CMV reactivation	reference		Reference		reference		reference	
CMV reactivation	-0.39 (-0.66, -0.13)		-0.31 (-0.75, 0.13)		1.15 (0.77, 1.52)		-0.58 (-0.86, -0.29)	
Day 100 aGVHD grade		.651		.386		.630		.435
0-I	reference		reference		reference		reference	
II	-0.16 (-0.52, 0.20)		-0.05 (-0.62, 0.51)		0.29 (-0.32, 0.90)		-0.20 (-0.61, 0.21)	
III-IV	-0.19 (-0.65, 0.27)		0.37 (0.36, 1.10)		0.30 (-0.48, 1.07)		-0.33 (-0.83, 0.17)	

*Engrafting CB unit.

There was no association with the variables of recipient sex, diagnosis (acute leukemia vs MDS/MPD vs lymphoma), or the addition of haploidentical CD34⁺ cells (data not shown).

P values significant at the .01 level are indicated in bold.

an effect. Alternatively, early thymus-independent T-cell recovery after ATG-free CBT may not be significantly influenced by advanced recipient age, in contrast to the thymus-dependent pathway of T-cell reconstitution.¹⁵ Data concerning the potential impact of CB graft characteristics on T-cell recovery after CBT are limited.^{23,35,36} We observed a notable trend for a higher engrafting CB unit infused viable CD3⁺ cell dose being associated with higher total and naive CD4⁺ T cells post-CBT. In addition, higher engrafting CB unit infused viable CD34⁺ cell dose was associated with higher EM and activated CD8⁺ T-cell counts, and a similar trend was observed for the same CD4⁺ T-cell subsets. Interestingly, despite high levels of engrafting CB unit-recipient HLA-mismatch, such mismatch had no discernible effect on total T-cell and subset recovery. In addition, despite concerns that aGVHD may hamper immune

reconstitution,³⁶⁻³⁸ overall grade II to IV aGVHD did not significantly affect T-cell recovery in our series, with the exception of activated CD4⁺ T-cell counts.

CMV reactivation significantly impacts quantitative and qualitative T-cell recovery after HSC transplantation (HSCT).^{39,40} CMV reactivation after HSCT has been associated with a rapid increase of CD8⁺ T cells due to EM and effector CD8⁺ subset expansion.³⁹⁻⁴⁵ The impact on total CD4⁺ T-cell and subset recovery, however, is less clear,³⁹⁻⁴³ with a previous report suggesting that CMV reactivation may compromise thymopoiesis.³⁹ These effects lead to contraction of the naive T-cell compartment and TCR repertoire diversity.^{39,43,46} In our analysis, a marked expansion of total CD8⁺, EM and effector CD8⁺ T-cell subsets, and

Figure 4. Association of CMV reactivation with CD4⁺ and CD8⁺ T-cell recovery in adult CBT recipients. (A) CD4⁺ T-cell recovery according to CMV reactivation by day 60 post-CBT. (B) CD8⁺ T-cell recovery according to CMV reactivation by day 60 post-CBT. Individual patient time points are color coded according to CMV reactivation. Curves are Loess-smoothed averages.




Table 5. Two-month landmark analysis for OS

Immune variable*	n	HR (95% CI)	P
ALC	90	0.67 (0.34-1.31)	.246
CD3 ⁺ T cells	90	0.73 (0.45-1.18)	.219
CD4 ⁺ T cells	90	0.69 (0.43-1.12)	.150
CD4 ⁺ CD45RA ⁺ T cells	90	1.11 (0.75-1.64)	.587
CD8 ⁺ T cells	90	1.00 (0.66-1.49)	.984
NK cells	78	1.34 (0.63-2.84)	.439
B cells	90	0.85 (0.68-1.07)	.163
PHA	88	0.56 (0.40-0.78)†	.009

P values significant at the .01 level are indicated in bold.

*In univariable analysis of patient and graft variables, only recipient age was significantly associated with OS (hazard ratio [HR], 2.11 [95% CI, 1.30-3.43] per decade, *P* = .001). Patient sex, diagnosis, CMV seropositivity, engrafting CB unit–recipient HLA-match, engrafting CB unit infused viable CD34⁺ and CD3⁺ cell doses, addition of haploidentical CD34⁺ cells were not significant.

†HR adjusted for age.

activated CD8⁺ T cells⁴⁴ was observed in patients with CMV reactivation post-CBT. Because CB T cells are exclusively naive,⁴⁷ this CMV-driven CD8⁺ T-cell expansion reflects the emergence of CMV effectors from naive CB-derived T cells. Although T-cell specificity was not examined, our findings are supported by reports that CMV effectors with memory phenotype are generated from naive CB-derived T cells early post-CBT⁴⁸⁻⁵⁰ and after CMV infection in neonates.⁵¹ We also observed that CMV reactivation was associated with lower total and CM CD4⁺ T-cell counts. Although naive CD4⁺ T cells were also numerically lower, this difference was not significant at the prespecified significance threshold. Therefore, although we demonstrate a negative impact of CMV reactivation on CD4⁺ T-cell recovery after CBT, whether thymopoiesis is impaired is uncertain.³⁹

A critical finding of this analysis is that early T-cell recovery, especially CD4⁺ counts and T-cell function, was associated with improved survival. Although an association of CD4⁺ counts with OS was not observed at 2 months, >75% of assayed patients already had CD4⁺ T cells >50/mm³ by that time, a threshold previously associated with protection against infections and improved survival in pediatric CBT.^{19,22,52,53} Importantly, we observed a continuous reduction in overall mortality with increasing CD4⁺ T-cell counts at 4 and 6 months post-CBT. Because CD4⁺ T cells were the most prevalent CD3⁺ subset at these time points, this lymphocyte population likely also accounted for the prognostic significance of CD3⁺ T-cell counts and ALC on survival. In contrast, CD8⁺ T-cell counts were not associated with survival, as previously reported after adult donor HSCT.^{38,54} Early recovery of T-cell function as assessed by PHA responses was also associated with improved survival, as has previously been reported in CD34⁺ selected HSCT recipients,⁵⁵ and was highly correlated with CD4⁺ T-cell counts.

Notably, in contrast to previous reports supporting an important role of thymopoiesis on CBT outcomes,^{15,30,48} we did not observe an association of CD4⁺CD45RA⁺ T-cell count recovery with improved survival. A possible explanation is that ATG omission may decrease reliance on thymopoiesis for CD4⁺ T-cell reconstitution due to early thymus-independent T-cell recovery.

Previous CBT studies have reported prompt posttransplant NK- and B-cell recovery, regardless of ATG use.^{12,28,29,56} Similarly, we observed early NK-cell recovery in our cohort. Unexpectedly,

Table 6. Four-month landmark analysis for OS

Immune variable*	n	HR (95% CI)	P
ALC	93	0.20 (0.08-0.51)†	.004
CD3 ⁺ T cells	93	0.41 (0.24-0.71)†	.006
CD4 ⁺ T cells	93	0.43 (0.25-0.73)†	.006
CD4 ⁺ CD45RA ⁺ T cells	93	1.03 (0.70-1.51)	.887
CD8 ⁺ T cells	93	0.86 (0.60-1.24)	.434
NK cells	91	0.71 (0.32-1.60)	.420
B cells	93	0.80 (0.66-0.97)	.029
PHA	89	0.43 (0.26-0.71)†	.008

P values significant at the .01 level are indicated in bold.

*In univariable analysis of patient, graft, and posttransplant variables, only recipient age was significantly associated with OS (HR, 2.21 [95% CI, 1.31-3.75] per decade, *P* = .001). Patient sex, diagnosis, CMV seropositivity, engrafting CB unit–recipient HLA match, engrafted CB unit infused viable CD34⁺ and CD3⁺ cell doses, addition of haploidentical CD34⁺ cells, and day 100 aGVHD grade were not significant.

†HR adjusted for age.

advanced recipient age was associated with higher NK-cell counts, as has previously been suggested in some HSCT studies,^{36,55} and may represent a compensatory mechanism as recovery of adaptive immunity may be delayed in older HSCT recipients. We also observed an association of a higher engrafting unit infused viable CD34⁺ cell dose with higher NK-cell counts post-CBT. In contrast, NK-cell counts tended to be lower in recipients of a higher engrafting unit infused viable CD3⁺ cell dose, a finding possibly explained by NK- and T-cell competition for homeostatic cytokines.⁵⁷ Similar findings have been reported in adult donor HSCT.^{58,59} Importantly, in contrast to multiple HSCT series,^{54,58,60-62} we found no association between NK-cell counts and CBT outcomes.

B-cell counts normalized by 4 months and were primarily characterized by naive immunophenotype. Development of grade II to IV aGVHD by day 100, and especially corticosteroid-requiring GVHD, was the only factor that adversely affected B-cell and subset recovery. Whether B-cell recovery has any prognostic value for CBT outcomes remains unclear, as the prespecified statistical significance threshold in our analysis was not achieved.

A striking finding of our study is that T-cell recovery after ATG-free CBT in adults is robust, thymus independent, and CD4⁺ biased. These results add to the growing literature supporting an underappreciated prompt T-cell recovery after T-replete CBT.^{19,22-24,28} They suggest that the reconstituting neonatal immune system in adult CBT recipients has unique biology as previously demonstrated in children.²⁴ The observed CD4⁺ bias is in contrast to the inversion of the CD4⁺/CD8⁺ T-cell ratio after adult donor HSCT. Moreover, our study demonstrates that early CD4⁺ T-cell recovery after adult CBT, as well as T-cell function, is associated with reduction in overall mortality, corroborating previous HSCT and primarily pediatric CBT studies.^{22,28,38,53-55,63,64}

We acknowledge that not all patients had early T-cell recovery. Therefore, as with adult donor HSCT, strategies to further augment immune reconstitution are needed. Notably, we found that increased cell dose improved T-cell recovery, whereas HLA mismatch was not detrimental. If these findings are confirmed, they would support prioritizing cell dose over HLA match in CB unit selection as a potentially modifiable factor to promote both myeloid engraftment and T-cell recovery. In addition, although grade II

Table 7. Six-month landmark analysis for OS

Immune variable*	n	HR (95% CI)	P
ALC	92	0.20 (0.07-0.54)†	.003
CD3 ⁺ T cells	93	0.34 (0.19-0.61)†	.001
CD4 ⁺ T cells	93	0.30 (0.16-0.56)†	<.001
CD4 ⁺ CD45RA ⁺ T cells	93	0.90 (0.59-1.39)	.651
CD8 ⁺ T cells	93	0.74 (0.45-1.21)	.221
NK cells	93	0.85 (0.31-2.34)	.757
B cells	93	0.80 (0.64-0.99)	.054
PHA	89	0.34 (0.17-0.68)†	.004

P values significant at the .01 level are indicated in bold.

*In univariable analysis of patient, graft, and posttransplant variables, only recipient age was associated with OS, although not at the 0.01 significance level (HR, 1.78 [95% CI, 1.04-3.03] per decade, *P* = .025). Patient sex, diagnosis, CMV seropositivity, engrafting CB unit–recipient HLA-match, engrafting CB unit infused viable CD34⁺ and CD3⁺ cell doses, addition of haploidentical CD34⁺ cells, and day 100 aGVHD grade were not significant.

†HR adjusted for age.

to IV aGVHD did not significantly affect overall T-cell recovery or mortality, severe aGVHD is often associated with significant morbidity and opportunistic infections. Moreover, mitigation of aGVHD burden may further speed thymus-dependent T-cell recovery due to the deleterious effect of aGVHD on thymopoiesis.^{37,65} Therefore, based on the promising results reported in adult donor HSCT,^{66,67} our center is now investigating enhanced aGVHD prophylaxis with the addition of tocilizumab (#NCT03434730) as a nonlymphodepleting strategy. Finally, although CMV-specific immunity ultimately develops, most seropositive CBT recipients reactivate CMV early posttransplant.⁶⁸ Therefore, our center has adopted letermovir prophylaxis starting day +7 for all adult CMV-seropositive CBT recipients.⁶⁹ It is possible that reduction in CMV reactivation could in turn improve CD4⁺ T-cell recovery in these patients.

Our study could have benefited from an even larger patient sample size and more frequent lymphocyte subset immunophenotyping assessments. In addition, detailed evaluation of thymopoiesis, including T-cell receptor excision circles¹⁵ and T-cell repertoire diversity,^{43,46} virus-specific T-cell responses,^{29,39,48-50} and analysis of phenotypic or functional NK-cell reconstitution,^{62,70,71} were not routinely performed. These should be investigated in the future to further elucidate the biology of immune recovery after CBT. Future correlation of quantitative and qualitative T-cell reconstitution with functional immune competence, as evidenced by protection against viral infections,^{52,72} will also be critically important. Such studies could identify immune milestones that may inform the duration of needed viral monitoring and the safe cessation of antiviral prophylaxis, including letermovir. Correlation of B-cell recovery with vaccine responses⁷³ will also be of great interest. Finally, CBT has been associated with robust GVL effects,^{4,5,74} and early CD4⁺ T-cell recovery has been associated with reduction in relapse risk in patients

with myeloid malignancies.^{22,53} Thus, further research should focus on the impact of T-cell recovery on TRM and relapse risk separately, as well as the unique immune biology of CB-mediated GVL.

Acknowledgments

The authors thank Theodore and Laura Hromadka for their generous support.

This research was supported in part by the National Institutes of Health, National Cancer Institute grants P01 CA23766 and P30 CA008748.

Authorship

Contribution: I.P., M.-A.P., and J.N.B. designed the study and assembled and analyzed the data; I.P. and J.N.B. wrote the manuscript; J.A.L. and P.H. performed the statistical analysis; T.B. and M.A.M. maintained the patient database and procured data for the study; I.P., J.A.L., P.H., C.C., S.A.G., M.R.M.v.d.B., M.-A.P., and J.N.B. interpreted the data and reviewed and edited the manuscript; and all authors have approved the final version of the manuscript.

Conflict-of-interest disclosure: I.P. serves on a data and safety monitoring board for ExcellThera. S.A.G. has served as a consultant for Amgen, Actinium, Celgene, Johnson & Johnson, Jazz Pharmaceutical, Takeda, Novartis, Kite, and Spectrum Pharma and has received research funding from Amgen, Actinium, Celgene, Johnson & Johnson, Miltenyi, and Takeda. M.R.M.v.d.B. has received research support from Seres Therapeutics; has consulted, received honorarium from or participated in advisory boards for Seres Therapeutics, Flagship Ventures, Novartis, Evelo, Jazz Pharmaceuticals, Therakos, Amgen, Magenta Therapeutics, Merck & Co, Inc, Acute Leukemia Forum, and DKMS Medical Council (Board); has IP Licensing with Seres Therapeutics, Juno Therapeutics, and has stock options from Smart Immune. M.-A.P. has received honoraria from AbbVie, Bellicum, Bristol-Myers Squibb, Incyte, Merck, Novartis, Nektar Therapeutics, and Takeda; serves on data and safety monitoring boards for Servier and Medigene and the scientific advisory boards of MolMed and NexImmune; and has received research support for clinical trials from Incyte, Kite (Gilead), and Miltenyi Biotec. J.N.B. has received clinical trial funding from Angiocrine Bioscience and unrestricted educational grants from Gamida Cell and Merck. The remaining authors declare no competing financial interests.

The current affiliation for P.H. is RWJ Barnabas Health, St. Barnabas Medical Center, Livingston, NJ.

ORCID profiles: J.A.L., 0000-0002-2746-5647; S.A.G., 0000-0003-1944-5053.

Correspondence: Ioannis Politiki, Adult Bone Marrow Transplantation Service, Department of Medicine, Memorial Sloan Kettering Cancer Center, 1275 York Ave, Box 259, New York, NY 10065; e-mail: politiki@mskcc.org.

References

- Barker JN, Byam CE, Kernan NA, et al. Availability of cord blood extends allogeneic hematopoietic stem cell transplant access to racial and ethnic minorities. *Biol Blood Marrow Transplant*. 2010;16(11):1541-1548.
- Barker JN, Boughan K, Dahi PB, et al. Racial disparities in access to HLA-matched unrelated donor transplants: a prospective 1312-patient analysis. *Blood Adv*. 2019;3(7):939-944.

3. Scaradavou A, Brunstein CG, Eapen M, et al. Double unit grafts successfully extend the application of umbilical cord blood transplantation in adults with acute leukemia. *Blood*. 2013;121(5):752-758.
4. Milano F, Gooley T, Wood B, et al. Cord-blood transplantation in patients with minimal residual disease. *N Engl J Med*. 2016;375(10):944-953.
5. Ponce DM, Hilden P, Devlin SM, et al. High disease-free survival with enhanced protection against relapse after double-unit cord blood transplantation when compared with T cell-depleted unrelated donor transplantation in patients with acute leukemia and chronic myelogenous leukemia. *Biol Blood Marrow Transplant*. 2015;21(11):1985-1993.
6. Eapen M, Rocha V, Sanz G, et al; National Cord Blood Program of the New York Blood Center. Effect of graft source on unrelated donor haemopoietic stem-cell transplantation in adults with acute leukaemia: a retrospective analysis. *Lancet Oncol*. 2010;11(7):653-660.
7. Gutman JA, Ross K, Smith C, et al. Chronic graft versus host disease burden and late transplant complications are lower following adult double cord blood versus matched unrelated donor peripheral blood transplantation. *Bone Marrow Transplant*. 2016;51(12):1588-1593.
8. Brunstein CG, Gutman JA, Weisdorf DJ, et al. Allogeneic hematopoietic cell transplantation for hematologic malignancy: relative risks and benefits of double umbilical cord blood. *Blood*. 2010;116(22):4693-4699.
9. Eapen M, Rubinstein P, Zhang MJ, et al. Outcomes of transplantation of unrelated donor umbilical cord blood and bone marrow in children with acute leukaemia: a comparison study. *Lancet*. 2007;369(9577):1947-1954.
10. Parody R, Martino R, Rovira M, et al; Infectious/Non-infectious Complications Subcommittee of the Grupo Español de Trasplante Hematopoyético (GETH). Severe infections after unrelated donor allogeneic hematopoietic stem cell transplantation in adults: comparison of cord blood transplantation with peripheral blood and bone marrow transplantation. *Biol Blood Marrow Transplant*. 2006;12(7):734-748.
11. Komanduri KV, St John LS, de Lima M, et al. Delayed immune reconstitution after cord blood transplantation is characterized by impaired thymopoiesis and late memory T-cell skewing. *Blood*. 2007;110(13):4543-4551.
12. Jacobson CA, Turki AT, McDonough SM, et al. Immune reconstitution after double umbilical cord blood stem cell transplantation: comparison with unrelated peripheral blood stem cell transplantation. *Biol Blood Marrow Transplant*. 2012;18(4):565-574.
13. Ballen K, Woo Ahn K, Chen M, et al. Infection rates among acute leukemia patients receiving alternative donor hematopoietic cell transplantation. *Biol Blood Marrow Transplant*. 2016;22(9):1636-1645.
14. Storek J. Immunological reconstitution after hematopoietic cell transplantation—its relation to the contents of the graft. *Expert Opin Biol Ther*. 2008;8(5):583-597.
15. Politikos I, Boussiotis VA. The role of the thymus in T-cell immune reconstitution after umbilical cord blood transplantation. *Blood*. 2014;124(22):3201-3211.
16. Lucchini G, Perales MA, Veys P. Immune reconstitution after cord blood transplantation: peculiarities, clinical implications and management strategies. *Cytotherapy*. 2015;17(6):711-722.
17. Castillo N, García-Cadenas I, Barba P, et al. Early and long-term impaired T lymphocyte immune reconstitution after cord blood transplantation with antithymocyte globulin. *Biol Blood Marrow Transplant*. 2017;23(3):491-497.
18. Pascal L, Tucunduva L, Ruggeri A, et al; Eurocord and the European Society for Blood and Marrow Transplantation. Impact of ATG-containing reduced-intensity conditioning after single- or double-unit allogeneic cord blood transplantation. *Blood*. 2015;126(8):1027-1032.
19. Lindemans CA, Chiesa R, Amrolia PJ, et al. Impact of thymoglobulin prior to pediatric unrelated umbilical cord blood transplantation on immune reconstitution and clinical outcome. *Blood*. 2014;123(1):126-132.
20. Pascal L, Mohty M, Ruggeri A, et al. Impact of rabbit ATG-containing myeloablative conditioning regimens on the outcome of patients undergoing unrelated single-unit cord blood transplantation for hematological malignancies. *Bone Marrow Transplant*. 2015;50(1):45-50.
21. Brunstein CG, Weisdorf DJ, DeFor T, et al. Marked increased risk of Epstein-Barr virus-related complications with the addition of antithymocyte globulin to a nonmyeloablative conditioning prior to unrelated umbilical cord blood transplantation. *Blood*. 2006;108(8):2874-2880.
22. Admiraal R, Lindemans CA, van Kesteren C, et al. Excellent T-cell reconstitution and survival depend on low ATG exposure after pediatric cord blood transplantation. *Blood*. 2016;128(23):2734-2741.
23. Chiesa R, Gilmour K, Qasim W, et al. Omission of in vivo T-cell depletion promotes rapid expansion of naïve CD4⁺ cord blood lymphocytes and restores adaptive immunity within 2 months after unrelated cord blood transplant. *Br J Haematol*. 2012;156(5):656-666.
24. Hiwarkar P, Hubank M, Qasim W, et al. Cord blood transplantation recapitulates fetal ontogeny with a distinct molecular signature that supports CD4⁺ T-cell reconstitution. *Blood Adv*. 2017;1(24):2206-2216.
25. de Koning C, Gabelich JA, Langenhorst J, et al. Filgrastim enhances T-cell clearance by antithymocyte globulin exposure after unrelated cord blood transplantation. *Blood Adv*. 2018;2(5):565-574.
26. Kanda J, Chiou LW, Szabolcs P, et al. Immune recovery in adult patients after myeloablative dual umbilical cord blood, matched sibling, and matched unrelated donor hematopoietic cell transplantation. *Biol Blood Marrow Transplant*. 2012;18(11):1664-1676.e1661.
27. Sauter C, Abboud M, Jia X, et al. Serious infection risk and immune recovery after double-unit cord blood transplantation without antithymocyte globulin. *Biol Blood Marrow Transplant*. 2011;17(10):1460-1471.
28. Bejanyan N, Brunstein CG, Cao Q, et al. Delayed immune reconstitution after allogeneic transplantation increases the risks of mortality and chronic GVHD. *Blood Adv*. 2018;2(8):909-922.
29. Saliba RM, Rezvani K, Leen A, et al. General and virus-specific immune cell reconstitution after double cord blood transplantation. *Biol Blood Marrow Transplant*. 2015;21(7):1284-1290.
30. Politikos I, Kim HT, Nikiforow S, et al. IL-7 and SCF levels inversely correlate with T cell reconstitution and clinical outcomes after cord blood transplantation in adults. *PLoS One*. 2015;10(7):e0132564.

31. Purtill D, Smith K, Devlin S, et al. Dominant unit CD34+ cell dose predicts engraftment after double-unit cord blood transplantation and is influenced by bank practice. *Blood*. 2014;124(19):2905-2912.
32. Rowlings PA, Przepiorka D, Klein JP, et al. IBMTR Severity Index for grading acute graft-versus-host disease: retrospective comparison with Glucksberg grade. *Br J Haematol*. 1997;97(4):855-864.
33. Ponce DM, Sauter C, Devlin S, et al. A novel reduced-intensity conditioning regimen induces a high incidence of sustained donor-derived neutrophil and platelet engraftment after double-unit cord blood transplantation. *Biol Blood Marrow Transplant*. 2013;19(5):799-803.
34. Ruggeri A, Peffault de Latour R, Carmagnat M, et al. Outcomes, infections, and immune reconstitution after double cord blood transplantation in patients with high-risk hematological diseases. *Transpl Infect Dis*. 2011;13(5):456-465.
35. Castillo N, García-Cadenas I, Díaz-Heredia C, et al. Cord blood units with high CD3(+) cell counts predict early lymphocyte recovery after in vivo T cell-depleted single cord blood transplantation. *Biol Blood Marrow Transplant*. 2016;22(6):1073-1079.
36. Niehues T, Rocha V, Filipovich AH, et al. Factors affecting lymphocyte subset reconstitution after either related or unrelated cord blood transplantation in children—a Eurocord analysis. *Br J Haematol*. 2001;114(1):42-48.
37. Clave E, Busson M, Douay C, et al. Acute graft-versus-host disease transiently impairs thymic output in young patients after allogeneic hematopoietic stem cell transplantation. *Blood*. 2009;113(25):6477-6484.
38. Berger M, Figari O, Bruno B, et al. Lymphocyte subsets recovery following allogeneic bone marrow transplantation (BMT): CD4+ cell count and transplant-related mortality. *Bone Marrow Transplant*. 2008;41(1):55-62.
39. Suessmuth Y, Mukherjee R, Watkins B, et al. CMV reactivation drives posttransplant T-cell reconstitution and results in defects in the underlying TCR β repertoire. *Blood*. 2015;125(25):3835-3850.
40. Lugthart G, van Ostaijen-Ten Dam MM, Jol-van der Zijde CM, et al. Early cytomegalovirus reactivation leaves a specific and dynamic imprint on the reconstituting T cell compartment long-term after hematopoietic stem cell transplantation. *Biol Blood Marrow Transplant*. 2014;20(5):655-661.
41. Kook H, Goldman F, Padley D, et al. Reconstruction of the immune system after unrelated or partially matched T-cell-depleted bone marrow transplantation in children: immunophenotypic analysis and factors affecting the speed of recovery. *Blood*. 1996;88(3):1089-1097.
42. Drylewicz J, Schellens IM, Gaiser R, et al. Rapid reconstitution of CD4 T cells and NK cells protects against CMV-reactivation after allogeneic stem cell transplantation. *J Transl Med*. 2016;14(1):230.
43. Kanakry CG, Coffey DG, Towler AM, et al. Origin and evolution of the T cell repertoire after posttransplantation cyclophosphamide. *JCI Insight*. 2016;1(5):e86252.
44. Itzykson R, Robin M, Moins-Teisserenc H, et al. Cytomegalovirus shapes long-term immune reconstitution after allogeneic stem cell transplantation. *Haematologica*. 2015;100(1):114-123.
45. Jeljeli M, Guérin-El Khourouj V, Porcher R, et al. Relationship between cytomegalovirus (CMV) reactivation, CMV-driven immunity, overall immune recovery and graft-versus-leukaemia effect in children. *Br J Haematol*. 2014;166(2):229-239.
46. van Heijst JW, Ceberio I, Lipuma LB, et al. Quantitative assessment of T cell repertoire recovery after hematopoietic stem cell transplantation. *Nat Med*. 2013;19(3):372-377.
47. Garderet L, Dulphy N, Douay C, et al. The umbilical cord blood alphabeta T-cell repertoire: characteristics of a polyclonal and naive but completely formed repertoire. *Blood*. 1998;91(1):340-346.
48. Brown JA, Stevenson K, Kim HT, et al. Clearance of CMV viremia and survival after double umbilical cord blood transplantation in adults depends on reconstitution of thymopoiesis. *Blood*. 2010;115(20):4111-4119.
49. McGoldrick SM, Bleakley ME, Guerrero A, et al. Cytomegalovirus-specific T cells are primed early after cord blood transplant but fail to control virus in vivo. *Blood*. 2013;121(14):2796-2803.
50. Cohen G, Carter SL, Weinberg KI, et al. Antigen-specific T-lymphocyte function after cord blood transplantation. *Biol Blood Marrow Transplant*. 2006;12(12):1335-1342.
51. Marchant A, Appay V, Van Der Sande M, et al. Mature CD8(+) T lymphocyte response to viral infection during fetal life. *J Clin Invest*. 2003;111(11):1747-1755.
52. Admiraal R, de Koning CCH, Lindemans CA, et al. Viral reactivations and associated outcomes in the context of immune reconstitution after pediatric hematopoietic cell transplantation. *J Allergy Clin Immunol*. 2017;140(6):1643-1650.e1649.
53. Admiraal R, van Kesteren C, Jol-van der Zijde CM, et al. Association between anti-thymocyte globulin exposure and CD4+ immune reconstitution in paediatric haemopoietic cell transplantation: a multicentre, retrospective pharmacodynamic cohort analysis. *Lancet Haematol*. 2015;2(5):e194-e203.
54. Bühlmann L, Buser AS, Cantoni N, et al. Lymphocyte subset recovery and outcome after T-cell replete allogeneic hematopoietic SCT. *Bone Marrow Transplant*. 2011;46(10):1357-1362.
55. Goldberg JD, Zheng J, Ratan R, et al. Early recovery of T-cell function predicts improved survival after T-cell depleted allogeneic transplant. *Leuk Lymphoma*. 2017;58(8):1859-1871.
56. Nakatani K, Imai K, Shigeno M, et al. Cord blood transplantation is associated with rapid B-cell neogenesis compared with BM transplantation. *Bone Marrow Transplant*. 2014;49(9):1155-1161.
57. Bunting MD, Varelias A, Souza-Fonseca-Guimaraes F, et al. GVHD prevents NK-cell-dependent leukemia and virus-specific innate immunity. *Blood*. 2017;129(5):630-642.
58. Savani BN, Mielke S, Adams S, et al. Rapid natural killer cell recovery determines outcome after T-cell-depleted HLA-identical stem cell transplantation in patients with myeloid leukemias but not with acute lymphoblastic leukemia. *Leukemia*. 2007;21(10):2145-2152.

59. Chang YJ, Zhao XY, Huang XJ. Effects of the NK cell recovery on outcomes of unmanipulated haploidentical blood and marrow transplantation for patients with hematologic malignancies. *Biol Blood Marrow Transplant.* 2008;14(3):323-334.
60. Minculescu L, Marquart HV, Friis LS, et al. Early natural killer cell reconstitution predicts overall survival in T cell-replete allogeneic hematopoietic stem cell transplantation. *Biol Blood Marrow Transplant.* 2016;22(12):2187-2193.
61. Dunbar EM, Buzzeo MP, Levine JB, Schold JD, Meier-Kriesche HU, Reddy V. The relationship between circulating natural killer cells after reduced intensity conditioning hematopoietic stem cell transplantation and relapse-free survival and graft-versus-host disease. *Haematologica.* 2008;93(12):1852-1858.
62. Bergerson RJ, Williams R, Wang H, et al. Fewer circulating natural killer cells 28 days after double cord blood transplantation predicts inferior survival and IL-15 response. *Blood Adv.* 2016;1(3):208-218.
63. Kim DH, Sohn SK, Won DI, Lee NY, Suh JS, Lee KB. Rapid helper T-cell recovery above 200 x 10⁶/l at 3 months correlates to successful transplant outcomes after allogeneic stem cell transplantation. *Bone Marrow Transplant.* 2006;37(12):1119-1128.
64. Bartelink IH, Belitser SV, Knibbe CA, et al. Immune reconstitution kinetics as an early predictor for mortality using various hematopoietic stem cell sources in children. *Biol Blood Marrow Transplant.* 2013;19(2):305-313.
65. Toubert A, Glauzy S, Douay C, Clave E. Thymus and immune reconstitution after allogeneic hematopoietic stem cell transplantation in humans: never say never again. *Tissue Antigens.* 2012;79(2):83-89.
66. Kennedy GA, Varelias A, Vuckovic S, et al. Addition of interleukin-6 inhibition with tocilizumab to standard graft-versus-host disease prophylaxis after allogeneic stem-cell transplantation: a phase 1/2 trial. *Lancet Oncol.* 2014;15(13):1451-1459.
67. Drobyski WR, Szabo A, Zhu F, et al. Tocilizumab, tacrolimus and methotrexate for the prevention of acute graft-versus-host disease: low incidence of lower gastrointestinal tract disease. *Haematologica.* 2018;103(4):717-727.
68. Dahi PB, Perales MA, Devlin SM, et al. Incidence, nature and mortality of cytomegalovirus infection after double-unit cord blood transplant. *Leuk Lymphoma.* 2015;56(6):1799-1805.
69. Marty FM, Ljungman P, Chemaly RF, et al. Letermovir prophylaxis for cytomegalovirus in hematopoietic-cell transplantation. *N Engl J Med.* 2017;377(25):2433-2444.
70. Pical-Izard C, Crocchiolo R, Granjeaud S, et al. Reconstitution of natural killer cells in HLA-matched HSCT after reduced-intensity conditioning: impact on clinical outcome. *Biol Blood Marrow Transplant.* 2015;21(3):429-439.
71. Nguyen S, Achour A, Souchet L, et al. Clinical impact of NK-cell reconstitution after reduced intensity conditioned unrelated cord blood transplantation in patients with acute myeloid leukemia: analysis of a prospective phase II multicenter trial on behalf of the Société Française de Greffe de Moelle Osseuse et Thérapie Cellulaire and Eurocord. *Bone Marrow Transplant.* 2017;52(10):1428-1435.
72. Camargo JF, Wieder ED, Kimble E, et al. Deep functional immunophenotyping predicts risk of cytomegalovirus reactivation after hematopoietic cell transplantation. *Blood.* 2019;133(8):867-877.
73. Shah GL, Shune L, Purtil D, et al. Robust vaccine responses in adult and pediatric cord blood transplantation recipients treated for hematologic malignancies. *Biol Blood Marrow Transplant.* 2015;21(12):2160-2166.
74. Hiwarkar P, Qasim W, Ricciardelli I, et al. Cord blood T cells mediate enhanced antitumor effects compared with adult peripheral blood T cells. *Blood.* 2015;126(26):2882-2891.