



REVIEW ARTICLE

Continuous vital sign analysis for predicting and preventing neonatal diseases in the twenty-first century: big data to the forefront

Navin Kumar¹, Gangaram Akangire², Brynne Sullivan³, Karen Fairchild³ and Venkatesh Sampath²

In the neonatal intensive care unit (NICU), heart rate, respiratory rate, and oxygen saturation are vital signs (VS) that are continuously monitored in infants, while blood pressure is often monitored continuously immediately after birth, or during critical illness. Although changes in VS can reflect infant physiology or circadian rhythms, persistent deviations in absolute values or complex changes in variability can indicate acute or chronic pathology. Recent studies demonstrate that analysis of continuous VS trends can predict sepsis, necrotizing enterocolitis, brain injury, bronchopulmonary dysplasia, cardiorespiratory decompensation, and mortality. Subtle changes in continuous VS patterns may not be discerned even by experienced clinicians reviewing spot VS data or VS trends captured in the monitor. In contrast, objective analysis of continuous VS data can improve neonatal outcomes by allowing heightened vigilance or preemptive interventions. In this review, we provide an overview of the studies that have used continuous analysis of single or multiple VS, their interactions, and combined VS and clinical analytic tools, to predict or detect neonatal pathophysiology. We make the case that big-data analytics are promising, and with continued improvements, can become a powerful tool to mitigate neonatal diseases in the twenty-first century.

Pediatric Research (2020) 87:210–220; <https://doi.org/10.1038/s41390-019-0527-0>

INTRODUCTION

Continuous assessment of cardiorespiratory stability is a critical component of neonatal intensive care in preterm infants. Vital signs including heart rate (HR), respiratory rate (RR), and pulse oximetry (SpO₂) are continuously monitored throughout hospital stay, while blood pressure (BP) may be continuously monitored in the first days after birth and during acute critical illness. Temperature is another important VS for assessing illness,¹ but is not considered here, as its measurement is confounded by use of external heat sources. VS are under complex physiological control, and while patterns of variation often reflect normal physiology such as circadian rhythms, they may also represent the earliest signs of decompensation triggered by pathological states.^{2–6} While clinical care is strongly influenced by interval assessments of VS displayed continuously on bedside monitors and periodic nursing assessments, the majority of continuous VS data is not utilized (Fig. 1). Furthermore, periodic reporting of VS lends itself to bias as well as under- or overreporting of abnormal values, which might adversely impact clinical decisions.⁷ Recent studies in neonates suggest that cumulative analysis of monitor-recorded continuous vital signs trends can predict impending clinical deterioration or disease such as sepsis, and may also predict long-term neurological or respiratory outcomes.^{2–4,8–12} In this review, we highlight studies that have evaluated the utility of continuous HR, RR, SpO₂, and BP data analytics to predict disease, and thereby potentially prevent adverse outcomes. Incorporating continuous VS analytics into the clinical record may improve

outcomes for preterm infants in the twenty-first century neonatal intensive care unit (NICU).

METHODS

We searched MEDLINE, CENTRAL, and CINAHL from 1990 to March 2019. Select studies were included in which continuous VS data on heart rate, respiratory rate, pulse oximetry, and/or blood pressure were captured and analyzed in neonates during their stay in the NICU, for any period of time. Outcomes of interest were analytics of VS physiologic variability, comparisons between continuous and interval VS measurements, and associations with short-term or long-term clinical outcomes.

CONTINUOUS HEART RATE MONITORING AND ANALYTICS

Physiologic determinants of heart rate monitoring HR is continuously monitored from birth until discharge for all NICU infants. The electrocardiogram (ECG) signal is obtained from three electrodes, two on the chest and one on the abdomen, and the bedside monitor displays one lead of ECG. HR can be calculated from the inter-beat interval (R to R wave of the QRS complexes), and in most NICU monitors the numerical HR displayed represents a moving average of several QRS complexes. Pulse rate (PR) calculated from the pulse oximetry signal is also continuously displayed. A significant discrepancy between the pulse oximetry-derived PR and the ECG-derived HR often indicates

¹Division of Neonatology, Hurley Medical Center, Flint, MI, USA; ²Division of Neonatology, Department of Pediatrics, Children's Mercy Hospital, Kansas City, MO, USA and ³Division of Neonatology, Department of Pediatrics, University of Virginia, Charlottesville, VA, USA
Correspondence: Venkatesh Sampath (vsampath@cmh.edu)

Received: 19 April 2019 Revised: 20 July 2019 Accepted: 25 July 2019
Published online: 4 August 2019




Fig. 1 Physiologic and pathologic factors regulating vital signs. Diagram depicting how HR, BP, RR, and SpO₂ are physiologically regulated in the preterm neonate, and patterns of vital signs changes that can be associated with pathophysiological states. HR heart rate, BP blood pressure, RR respiratory rate, SpO₂ pulse oximetry

poor pulse oximetry signal quality, and therefore, the HR is generally more reliable than the PR.¹³ Normative data on HR in preterm NICU patients have recently been published.¹⁴

Using HRV patterns to detect disease

HR variability (HRV), or normal beat to beat fluctuations in HR, is regulated by the sympathetic and parasympathetic nervous systems.¹⁵ In healthy preterm infants, HR accelerations and decelerations reflect high variability and are indicative of a functioning, adaptable autonomic nervous system, while decreased HRV suggests a stressed or pathological state. More than a century ago, obstetricians recognized that low HR variability punctuated by HR decelerations was a sign of fetal compromise,¹⁶ often associated with acidemia, and sometimes with chorioamnionitis and a fetal inflammatory response.^{17,18} A similar pattern of low variability with superimposed decelerations was observed in preterm neonates with sepsis and replicated in a preclinical model.^{19–21} Data from mice suggest that low HRV upon exposure to pathogens is mediated in part by cytokines released during sepsis. HR decelerations, in the same model, were mediated both by direct and indirect pathogen effects on the vagus nerve, as part of the cholinergic anti-inflammatory response.²²

In preterm infants, the most significant cause of abnormal heart rate characteristics (HRC) is sepsis. Late-onset sepsis (LOS, >3 days of age) and necrotizing enterocolitis (NEC) can be life-threatening, especially if not diagnosed until the infant is displaying VS and clinical instability. The use of continuous HR analytics for earlier detection and treatment could prevent mortality and long-term morbidity^{23–25} by alerting clinicians, and prompting a more careful clinical examination to support decisions about testing and treatment (Table 1). The first commercially available device that

serves as an early warning system for sepsis in the NICU is the HRC index monitor (HeRO monitor). A mathematical algorithm was developed by Moorman et al. to measure the observed phenomenon of decreased HRV punctuated by HR decelerations during systemic inflammation, and thereby detect an early signature of sepsis in premature infants.²⁶ The algorithm incorporates measures of HRV, and measures of sample asymmetry and entropy (lack of predictability) to detect pathologic decelerations.^{27,28} The HRC index, which is the fold-increased risk of sepsis diagnosis within 24 h, is continuously calculated from the previous 12 h of monitored ECG signal and an hourly updated value between zero and 7 is displayed, together with the HR values and the 5-day HRC index trend.

HRC in sepsis and infection-related outcomes

The impact of the HRC index monitor was tested in a multicenter, randomized controlled trial of 3003 VLBW infants. The overall incidence of LOS was not significantly different among infants with or without HRC display. However, infants whose HRC index was displayed to clinicians had a 22% relative reduction in all-cause mortality in the 120 days after randomization, and a 40% reduction in death within 30 days of septicemia compared to infants whose HRC index was not made available to clinicians.^{2,29} Although the HRC index was designed as a risk score for sepsis, subsequent analyses have demonstrated its utility in predicting other outcomes, including necrotizing enterocolitis.³⁰ In another small study, decreased high-frequency HRV in the first week after birth, suggesting reduced parasympathetic nervous system activity, was associated with later NEC diagnosis.³¹ Such early risk stratification could be useful for heightened vigilance or preventative bundles aimed at reducing incidence or severity of common preterm morbidities.

Table 1. Continuous heart rate monitoring analytics and outcome studies

Investigators	Study N Site(s)	Inclusion criteria	Results/conclusions
Griffin et al. ¹⁹	633 Two centers	27 0/7–37 weeks	Abnormal HRC (reduced variability, decelerations) is a valid tool for early diagnosis of impending LOS or sepsis-like illness.
Griffin et al. ⁴¹	337 Single center	27 0/7–35 weeks	HRC along with clinical signs of illness can add independent information in the diagnosis of LOS.
Moorman et al. ²⁹	3003 Multicenter	<1500 g	Display of HRC index is associated with 22% decreased mortality and trend towards increased ventilator-free days in comparison to control subjects with recording of HRC index without display.
Fairchild et al. ²	2989 Multicenter	<1500 g	40% decreased mortality within 30 days of septicemia with HRC index display due to earlier diagnosis.
Stone et al. ³⁰	97 Multicenter	<1500 g	Abnormal HRC occurs prior to diagnosis of medical and surgical NEC and can help with early diagnosis and treatment.
Addison et al. ³²	65 Single center	<1500 g	Cumulative frequency of abnormal HRC score is associated with increased risk of cerebral palsy and delayed early cognitive development.
Fairchild et al. ⁴⁶	1065 Two centers	<1500 g	Best metric for predicting LOS/NEC was cross correlation(X-Corr)-HR-SpO2 and a 3-variable vital sign model.
Fairchild et al. ⁴⁵	629 Single center	<1500 g	High cross correlation (X-Corr)-HR-SpO2 is associated with apnea and adverse events including NEC and LOS.
Sullivan et al. ³³	566 Single center	<1500 g	HRC index on day of life 1 and 7 compares favorably to established risk scores to predict death and morbidities (LOS, BPD, severe IVH, NEC and severe ROP).
Doheny et al. ³¹	70 Single center	28 0/7–34 6/ 7 weeks	HF-HRV can be a potential noninvasive biomarker for NEC diagnosis.
Fairchild et al. ³⁷	384 Single center	<1000 g	Increased HRC index is associated with abnormal brain imaging and can predict adverse neurologic outcomes.
Vergales et al. ³⁸	37–64 Single center	Term gestation	Low HRV (and corresponding high HRC index) is associated with worse EEG, MRI and 2-year Bayley neurodevelopmental outcomes.
Goel et al. ³⁶	102 Single center	All ventilated infants	HRC index is higher in infants with extubation failures compared to infants with successful extubation. Infants with failed extubation had lower gestational age, extubated at lower corrected gestational age, longer duration of ventilation prior to extubation and culture positive sepsis.
Sullivan et al. ³⁴	274 Single center	<1500 g	Sepsis, respiratory deterioration, surgical procedures and other infectious or inflammatory conditions are associated with large increase in HRC index.

HRC heart rate characteristics LOS late onset sepsis, NEC necrotizing enterocolitis, X-Corr-HR-SpO2 cross correlation of heart rate and oxygen saturation, BPD bronchopulmonary dysplasia, ROP retinopathy of prematurity, IVH intraventricular hemorrhage, HUS head ultrasound, MRI magnetic resonance imaging

HRC in respiratory and neurologic morbidities

A limitation of HRC monitoring for LOS and NEC is that infants with acute or chronic lung or brain injury often have abnormal HRC.^{32,33} An analysis of large increases, or “spikes” in the HRC index for infants in the control arm of the RCT in one NICU showed that about half of the spikes were associated with suspected or proven infection and about a third were associated with a significant acute respiratory deterioration in absence of infection.³⁴ In the setting of acute (or acute on chronic) respiratory distress, a high or rising HRC index may reflect repetitive apnea (with associated HR deceleration), lung inflammation, or effects of hypoxemia or acidosis on HRV. Another study found that, of multiple cardiorespiratory signs tested, the most useful analytic predicting respiratory deterioration requiring urgent intubation was the HRC index.³⁵ A subsequent study found that the HRC index assisted in determining extubation readiness in preterm infants.³⁶ This suggests that continuous display of an HR or cardiorespiratory index might alert clinicians to preterm infants’ need for either more or less respiratory support.

Abnormal heart rate patterns are associated with acute brain injury presumably due to autonomic dysfunction and perhaps reflecting a systemic inflammatory response associated with neuroinflammation.^{33–38} Severe IVH has been shown to lead to a chronically high HRC index for the first month after birth, followed by normalization.³⁷ Another study found that among neonates undergoing therapeutic hypothermia for hypoxic-ischemic encephalopathy, low HRV was associated with worse

brain MRI changes and neurological outcomes.³⁸ Use of HR analytics for IVH prediction has also been explored using such methods as detrended fluctuation analysis³⁹ and sample entropy.⁴⁰

HRC analytics combined with clinical assessment tools and other VS trends

Even with advanced analytics, abnormal HR patterns must be interpreted in the context of clinical variables and other VS trends. Griffin et al. developed a clinical risk score to use in conjunction with HRC monitoring for early detection of LOS.⁴¹ Adding respiratory analysis to HRC analysis may improve diagnostic accuracy of HRC analysis. This is especially true for sepsis-related complications as increased apneas and periodic breathing frequently accompany abnormal HR patterns, in part related to endogenous prostaglandins release.^{42–44} Apnea and periodic breathing analysis for sepsis detection is discussed in the respiratory section.^{45,46} The “PhysiScore” incorporating HRV and other continuously monitored VS in the first hours after birth was shown in preterm infants to predict later sepsis and other adverse outcomes.⁴⁷ The NeoNEEDS score that incorporates clinical evaluation of behavior and abdominal examination along with HR, blood pressure, FiO2 changes, and respiratory rate was shown to decrease the rate of NEC in a small single-center study.⁴⁵ The RALIS (Integralis-medical.com) system allows clinicians to input 6 monitor VS every 2 h to generate an alarm for possible onset of sepsis in subsequent days.⁴⁸

In summary, although display of the HRC index or other HR-based early warning systems for sepsis has the potential to improve outcomes, clinicians in most NICUs continue to rely solely on observation of ECG HR trends on bedside monitors to make determinations about clinical status, and the need for escalation of care. Ongoing research aims to further improve sepsis early-warning systems by incorporating other VS, clinical signs, and biomarkers to develop decision support tools to reduce sepsis and NEC-associated outcomes.

CONTINUOUS PULSE OXIMETRY MONITORING AND ANALYTICS

Physiological determinants and importance of oxygen saturation monitoring

Pulse oximetry allows for transcutaneous estimation of oxygen saturation in the blood using principles of spectrophotometry.⁴⁹ The pulse oximetry probe contains two light emitting diodes positioned on one side of the tissue (generally the hand or foot), and a photodetector on the opposing side. The probe measures the differential light absorption of deoxyhemoglobin and oxyhemoglobin at 600 nm (red) and 940 nm (infrared) wavelengths, to determine their respective concentrations in the blood passing through tissue, and calculate the percentage of oxyhemoglobin (SpO₂). The final SpO₂ determination is made by comparing the absorbance during pulsatile and nonpulsatile flow to isolate the light absorption of blood.^{50,51}

While fetal growth occurs in a relatively hypoxic intrauterine environment (PaO₂ 20–28 torr), premature infants are exposed to a hyperoxic environment ex-utero. This poses the distinct challenge of maintaining optimal oxygenation to ensure adequate growth and development, while limiting free-radical damage in an oxygen-rich milieu.^{52,53} SpO₂ monitoring allows continuous assessment of both hypoxemia and hyperoxemia.^{52,53} The importance of SpO₂ targeting is established by the meta-analysis of large clinical trials (SUPPORT, BOOST, and COT) that enrolled over 5000 infants <1250 g birth weight.^{54–57} Pooled data from these trials demonstrated that targeting SpO₂ between 85 and 89% was associated with a higher risk of mortality and necrotizing enterocolitis, whereas targeting higher SpO₂ between 91 and 95% was associated with higher risk of retinopathy of prematurity.

Maintenance of SpO₂ in the intended target range

Most NICUs have guidelines to keep preterm infants' SpO₂ within certain target limits based on gestational and chronologic age, and disease state. SpO₂ analytics have been used to evaluate the precision of oxygen targeting. In a study involving premature infants receiving supplemental oxygen at 14 centers, SpO₂ was maintained in the intended range only 48% of the time, being above and below the range 36% and 16% of the time, respectively.⁵⁸ Similar findings were observed in another study for infants on continuous positive airway pressure, where SpO₂ was maintained in the targeted range only 31% of the time.⁵⁹ Recent studies also suggest that bedside recognition of both hypoxemia and hyperoxemia plays a role in achieving targeted SpO₂ range.^{3,4,12}

Pulse oximetry analytics in prediction of clinical outcomes

Analysis of continuous SpO₂ trends, either alone or in combination with other VS parameters has been shown to predict short-term clinical deterioration as well as long-term outcomes such as BPD (Table 2). In one study involving 150 babies born with a gestational age (GA) <34 weeks, the mean, baseline and residual SpO₂ variability was combined with HR and RR patterns in the first 3 h of life, to develop a probability score (PhysiScore) for predicting adverse outcomes.⁴⁷ PhysiScore was more accurate in predicting overall morbidity with 86% sensitivity and 96% specificity compared to other scoring systems such as APGAR,⁶⁰

SNAP-II,⁶¹ SNAPPE-II,⁶¹ and CRIB.⁶² Using Bayesian modeling, the investigators demonstrated that overall morbidity was significantly associated with mean SpO₂ <92% and prolonged time spent (>5% of time) at SpO₂ less than 85%. PhysiScore also achieved almost 100% specificity for two major combined outcome categories: infection (NEC, sepsis, UTI and pneumonia; with 90% sensitivity) and cardiopulmonary complication (including BPD, hemodynamic instability, pulmonary hypertension and pulmonary hemorrhage with 96% sensitivity). In another study, the mean, standard deviation, kurtosis and skewness of SpO₂ data was combined with continuous HR trends in the first 12 h, and first 7 days of life to develop a pulse oximetry predictive score (POPS) in 800 VLBW infants.⁶³ POPS is a combination of pulse oximetry data score and a clinical predictive score. The predictive performance of POPS was most significant for BPD (AUC: 0.935) and mortality (AUC: 0.864), but was also significant for severe IVH, NEC, treated ROP, and length of stay. Associations between a lower mean SpO₂ in the first 12 h and mortality, and lower SpO₂ in the first 7 days and BPD were also found. In another study of 137 extremely preterm infants, BPD was associated with increasing intermittent hypoxemias in the first 4 weeks of life.⁶⁴ A recent study of 645 VLBW infants at three NICUs found that early severe hypoxemia burden (time spent with SpO₂ <70%) was significantly higher in infants with severe grade 3–4 IVH.⁶⁵

Retinopathy of prematurity (ROP) is known to be associated with both hypoxemia and hyperoxemia, and a study of 56 ELBW infants found that an increase in weekly mean SpO₂ fluctuations over the first 28 days of life was associated with moderate to severe ROP.⁸ In another study of 63 infants, severe ROP was associated with early intermittent hypoxemia events.⁶⁶ Overall, data from these studies reveal that analyzing trends in continuous SpO₂ is a useful tool to predict several preterm diseases.

Prediction of imminent clinical deterioration using continuous SpO₂ data

In a two-center study of 1065 VLBW infants, mean, SD, and cross correlation of SpO₂, HR and RR, were analyzed for 1.15 million hours to detect correlations with imminent diagnosis LOS and NEC.⁴⁶ In one center, a small but statistically significant decrease in mean SpO₂ preceded LOS or NEC, whereas a decrease in SD of SpO₂ preceded LOS or NEC in another center. Importantly, the cross correlation of HR and SpO₂, measuring co-trending of the two VS, was the single best predictor of illness at both centers. The best combined model (mean SpO₂, SD HR, and cross correlation of HR-SpO₂) provided additive value to the HRC index for illness prediction (net improvement 0.25, 95% CI 0.113, 0.328). SpO₂ data have also been used in predicting short-term respiratory outcomes. In a study of 31 premature infants, failure to wean from nasal positive pressure support to nasal cannula was correlated with the proportion of time spent with saturations <86% pre-wean.⁶⁷ In another study of 100 premature infants, daily mean, and SD of SpO₂ data, along with cumulative time of hypoxemia was analyzed to detect impending respiratory deterioration.⁴ Analysis of >3 million data points from relatively stable infants on noninvasive respiratory support or room air showed a 3.7-fold increase in odds of respiratory support escalation within the next 3 days if hypoxemia was present for 5–10% of total time. This study also showed a significant correlation between the hypoxemia time and decreased Fenton weight z-scores.⁴ Additionally, nursing documentation of hypoxemia did not correlate with short-term respiratory deterioration. Interestingly, similar associations between hypoxemia and impaired growth outcomes has also been shown in animal models.⁶⁸ These studies highlight the potential for utilizing the continuous pulse oximetry data in prediction of adverse outcomes such as LOS, respiratory deterioration and growth failure.

In summary, analysis of continuous SpO₂ data can improve clinicians' ability to maintain preterm neonates in a target range, and provide high resolution data to supplement bedside

Table 2. Continuous pulse oximetry monitoring analytics and outcome studies

Investigators	Study N Site(s)	Inclusion criteria	Results/conclusions
Di Fiore et al. ⁵⁷	1316 Multicenter	24 0/7–27 6/7 weeks	SGA infants in lower target SpO ₂ (85–89%) range achieved lowest oxygen saturation and higher incidence of intermittent hypoxemia in comparison with higher target SpO ₂ (91–95%) during first 3 days of life. Lowest quartile for saturation (≤92%) in the first 3 days of life is associated with increased 90-day mortality in AGA and SGA infants.
Poet et al. ⁵⁶	1019 Multicenter	23 0/7–27 6/7 weeks	Prolonged hypoxemic episodes in first 2–3 months of life increased risk of death or cognitive/motor impairment at 18 months of age. Bradycardia did not alter the prognostic value of hypoxemia.
Sullivan et al. ⁶³	778 Two centers	<1500 g	Pulse oximetry predictive score (POPS) performed better for prediction of death, IVH and BPD compared to HRC alone.
Raffay et al. ⁶⁴	137 Single center	<28 weeks	More frequent, longer and elevated intermittent Hypoxemia (IH) nadirs with increased oxygen and pressure exposure in first 28 days of life increases the risk of BPD. Early IH patterns may contribute to the development of BPD and may help identify infants at risk.
Di Fiore et al. ⁶⁶	63 Single center	24 0/7–27 6/7 weeks	Increased variability, longer duration, and lower nadir of hypoxemia is associated with increased risk of severe ROP. Identification of spectral component of SpO ₂ waveform may aid in identification of infants at risk for severe ROP.
Fairchild et al. ⁴⁶	1065 Two centers	<1500 g	HR-SpO ₂ cross correlation model performed better than HR or SpO ₂ for preclinical detection of NEC and sepsis.
Warburton et al. ⁴	94 Single center	<36 weeks gestation	% of time with SpO ₂ <90% is associated with risk of respiratory support escalation and decreased weight gain.
Das et al. ⁸	56 Single center	<1000 g	Increased SpO ₂ fluctuation during first 4 weeks of life was higher in infants with severe ROP. No association was found if the time spent was in the target range of SpO ₂ .
Mascoll-Robertson et al. ⁶⁷	31 Single center	24 0/7–32 6/7 weeks	Pulse oximetry histogram may be helpful in determining readiness of weaning from CPAP/HFNC to LFNC/oxyhood/RA transition.
Vesoulis et al. ⁶⁵	645 Multicenter	<32 weeks or <1500 g	Greater hypoxemia burden in first week of life was significantly associated with grade III/IV IVH.
Bizzarro et al. ⁶⁹	700 Single center	<1500 g	Significant reduction in severe ROP and ROP requiring surgery after staff education and implementation of signal extraction technology to quantify success with achieved target oxygen saturations.

SGA small for gestational age, AGA appropriate for gestational age, SpO₂ oxygen saturation, POPS pulse oximetry predictive scores, IVH intraventricular hemorrhage, BPD bronchopulmonary dysplasia, HRC heart rate characteristics, ROP retinopathy of prematurity, NEC necrotizing enterocolitis, CPAP continuous positive airway pressure, HFNC high flow nasal cannula, LFNC low flow nasal cannula, RA room air, HUS head ultrasound

assessments in predicting clinical deterioration. Many newer pulse oximeters and bedside monitors have the option to display cumulative SpO₂ histograms or other data displays to inform caregivers about percentage of time spent in lower and higher SpO₂ ranges. In one quality improvement study, educating caregivers about the risks of hyperoxemia and the importance of reviewing monitor data and improving oximetry targeting, resulted in significant decrease in severe ROP.⁶⁹

In light of the inherent difficulty for ICU nurses and respiratory therapists to titrate supplemental oxygen to keep infants in safe target ranges, efforts have been directed over the past several decades toward automated closed-loop FiO₂ titration systems. Several studies have shown these sophisticated systems to be more effective than manual control of oxygen delivery to achieve optimal oxygen targeting in preterm infants.^{70–76} Since NICU nurses have many responsibilities in addition to oxygen titration, the real-world NICU is likely to use automated FiO₂ titration in the twenty-first century. However, well-designed trials are needed to establish the efficacy of these approaches to improve clinical outcomes.

CONTINUOUS RESPIRATORY RATE MONITORING AND ANALYTICS

Physiological determinants and methods of respiratory rate (RR) monitoring

In most NICU monitors, RR rate is measured by detection of alterations in thoracic impedance between two electrodes placed

on the chest and abdomen. Due to the differential impedance of air and tissue, the electrodes detect a change in impedance as the air-to-tissue ratio increases during inspiration, thereby allowing display of an RR or absence of respiratory effort during central apnea.^{77,78} Thoracic impedance measurement of respiratory rate, despite being universally used in NICUs, has a number of drawbacks. Obstructive apnea is not detected since chest wall impedance changes during airway obstruction. Obstructive apnea can be monitored using a nasal cannula thermistor detecting warm exhaled breath or capnography detecting exhaled carbon dioxide, but these are not performed in routine clinical care. Another drawback of chest impedance monitoring is that it can provide an inaccurate RR due to the tendency of infants to have irregular and sometimes shallow respirations and due to motion and cardiac artifact.⁷⁹ Respiratory inductance plethysmography measures RR more reliably than impedance monitoring but requires placement of multiple electrodes circling the chest, and as such is not practicable for monitoring infants in the NICU.⁸⁰ Other promising methodologies, including devices based on monitoring breath sounds in the neck,⁸¹ and using ECG or photoplethysmography waveform data to determine RR⁸² have not been evaluated in neonates.

Importance of apnea and periodic breathing (PB) and improved methods for their quantitation
Respiratory physiology in premature infants has unique characteristics when compared to older children, and assessment of

Table 3. Continuous respiratory rate, effort and blood pressure data analysis and outcome studies

Investigators (respiratory rate)	Study N Site(s)	Inclusion criteria	Results/conclusions
Hofstetter et al. ⁹⁶	33 Single center	23 0/7–27 6/ 7 weeks	Apnea/hypopnea, bradycardia and hypoxemia episodes decreased with age, but continued at term equivalent, and even after hospital discharge. Infection increased apnea/hypopnea and hypoxemia events.
Fairchild et al. ⁴²	1211 Single center	<35 weeks	Number and duration of apnea events decreased with increasing gestational age. ABD events has higher frequency in <31 weeks infants but not increased in infants with severe ROP, BPD and severe IVH after adjusting for GA. ABD events increased before the diagnosis of LOS and NEC.
Tabacaru et al. ⁹⁹	302 Single center	<32 weeks	Intermittent caffeine boluses and discontinuation at 33 weeks PMA were associated with small changes in ABD events.
Patel et al. ⁴³	1211 Single center	<35 weeks	Periodic breathing (PB) increases with gestational age and the highest amount was between 30–33 weeks and 2 weeks chronological age. Extreme PB is associated with infection, NEC, caffeine discontinuation and immunizations.
Warburton et al. ⁴	94 Single center	<36 weeks	Tachypnea (RR > 70) is associated with poor growth and respiratory support escalation. >30% tachypnea/day is associated with increased respiratory support in subsequent 3 days.
Mohr et al. ⁹⁵	70 Single center	All infants in the NICU	For 32 weeks gestation infants, PB peaked 7–14 days after birth (6.5%). Infant with death (SIDS) had 40% PB each day and her twin had 15% PB each day.
Investigators (blood pressure)	Study N Site(s)	Inclusion criteria	Results/conclusions
Goldstein et al. ¹¹⁷	191 Single center	<1500 g	Metabolic acidosis and respiratory acidosis are related to adverse cognitive, motor and neurologic outcome at 6 months of age, while only metabolic component is related to adverse outcomes at 24 months of age.
Miall-Allen et al. ¹¹⁹	33 Single center	<31 weeks	Hypotension (<30 mmHg) for over an hour was associated with IVH, ischemic cerebral lesions and death (within 48 h).
Miall-Allen et al. ¹³⁹	22 Single center	<31 weeks	No association was found between blood pressure fluctuation and IVH in first 36 h of life.
Low et al. ¹²²	98 Single center	<34 weeks	Combination of hypotension and hypoxemia in first 96 h of life significantly increased the risk of brain damage and poor outcomes.
Cunningham et al. ¹⁰⁰	232 Single center	<1500 g	IVH was associated with low or variable BP. PVL and ROP were not associated with BP. BP variability was associated with death.
Bada et al. ¹²⁰	100 Single center	<1500 g	Infants with periventricular IVH had a greater minute to minute BP variability compared to infants with no periventricular IVH.
Perlman et al. ¹⁴⁰	50 Single center	<1500 g	Fluctuating cerebral blood-flow velocity in infants with RDS increases the risk of IVH.
Soul et al. ¹⁴¹	90 Two centers	<1500 g	Cerebral pressure passivity is associated with gestational age and low birth weight, systemic hypotension and maternal hemodynamic factors.
Dacosta et al. ¹⁴²	44 Single center	23 0/7–26 6/ 7 weeks	Defining the MAP with strongest cerebrovascular activity is feasible and deviations in that increased the risk of IVH and death.
Semenova et al. ¹⁴³	25 Single center	<32 weeks	Normal well-being is associated with nonlinear association between EEG and BP. Presence of weak association with distinctive directionality of information flow is associated with increased mortality.
Hoffman et al. ¹³⁶	61 Single center	23 0/7–28 6/ 7 weeks	More time with impaired cerebral autoregulation and less time with cerebral reactivity was associated with grade 3-4 IVH.

ABD apnea bradycardia desaturation, *BPD* bronchopulmonary dysplasia, *ROP* retinopathy of prematurity, *IVH* intraventricular hemorrhage, *LOS* late onset sepsis, *NEC* necrotizing enterocolitis, *GA* gestational age, *PMA* post menstrual age, *MAP* mean arterial pressure, *EEG* electroencephalogram, *RDS* respiratory distress syndrome, *PVL* periventricular leukomalacia, *SIDS* sudden infant death syndrome

respiratory function in the NICU has been largely limited to nursing documentation of oxygen saturation, respiratory rate, and apnea events. Prolonged apneic pauses may be associated with decreases in heart rate and/or SpO₂.^{75,76,83,84} Although apnea and periodic breathing predominantly reflect normal physiologic immaturity of respiratory control,^{85–88} pathological states such as LOS and NEC can also predispose preterm infants to increased number or severity of apnea spells (Table 3).^{89,90} It is well known that interval monitoring and documentation of apnea/periodic breathing in the medical record do not accurately capture trends in breathing patterns or apnea.^{4,9,10,91,92} To address this challenge, an automated apnea detection system for preterm NICU patients was developed. This

algorithm analyzes the waveform chest impedance signal, after removing motion and cardiac artifact, for episodes of very low variance indicative of central apneic pauses.⁷⁹ Multiple studies have been published using this algorithm, quantitating and describing “ABD” events, central apnea with associated decline in HR and SpO₂.^{91,93,94} An algorithm to quantitate periodic breathing (PB), which if prolonged, can be pathological has also been developed. This algorithm which analyzes chest impedance apneic pauses was useful for defining normal PB, and identifying NICU patients with severely exaggerated PB who later died of SIDS or sepsis-like illness.⁹⁵ Algorithms for continuous evaluation of central ABD and PB are not yet available for clinical use.

Continuous RR monitoring and clinical outcomes

Limited studies suggest that continuously monitored RR data can be used for prediction or prognostication in preterm infants. The PhysiScore study previously cited in the HRV section also analyzed RR mean and variability in a small cohort of preterm infants in the first 3 h of life, and showed that an average RR of less than 35 or greater than 75 breaths per minute was associated with higher mortality. In contrast to low HRV being associated with adverse outcomes, for RR the effect was bi-modal, with both high and low variability associated with higher risk of morbidity.⁴⁷ In another study of 100 preterm infants, analysis of continuous RR data was able to predict impending worsening of respiratory status. Among infants on noninvasive respiratory support or room air, infants who were tachypneic (RR > 70) for >30% of time had 2.8-fold increased odds of requiring escalation of respiratory support in the subsequent 24–72 h window.⁴ In the same study, a significant correlation between time spent per day with RR > 70 and decreased weight gain was reported in infants on RA or noninvasive respiratory support. In a prospective study of 33 ELBW infants, continuous chest impedance, electrocardiographic

waveforms and SpO2 data were collected for apnea, bradycardia and desaturations.⁹⁶ Blood stream infection was noted in 42% of the study participants, and a significant increase in apnea events was noted prior to the onset of infection. In a large retrospective study of 1211 premature infants (GA < 35 weeks), a twofold increase in apnea, bradycardia and desaturations (ABD) events was noted in about one-third cases of NEC and almost half of the cases with LOS 24 h prior to diagnosis.⁴² In addition to apneas, periodic breathing (PB), commonly defined as short repetitive cycles of respiratory pauses may also be a harbinger of disease.^{97,98} The association between extreme PB, defined, in this study, as PB for >10% of time and higher than 6 standard deviations above the mean for GA, and clinical outcomes were evaluated retrospectively in the above cohort.⁴³ PB was found in 76 of 1211 infants, and was associated 45% of the time with infection or NEC, immunizations, or caffeine discontinuation. Analysis of continuous cardiorespiratory data has also been able to shed some light on the impact of caffeine for treatment of apnea of prematurity. In another retrospective study of 300 preterm infants (GA < 32 weeks), caffeine loading was associated




Fig. 2 Twenty-first century analytics of continuous vital sign data to prevent diseases in neonates. This illustration depicts a three-step approach combining: **a** single or multi-tier analytics of vital signs, **b** careful clinical assessment, and **c** auxiliary blood and other imaging studies to prevent and decrease morbidity from life-threatening illnesses. HeRo heart rate characteristics, HRC index heart rate characteristics index, POPS pulse oximetry predictive score, BP blood pressure, CBC complete blood count, CRP C-reactive protein, USG ultrasonography, ABG arterial blood gas, TCM transcutaneous carbon dioxide monitoring, NIRS near-infrared spectroscopy, EEG electroencephalogram, HUS head ultrasound, NEC necrotizing enterocolitis, BPD bronchopulmonary dysplasia, ROP retinopathy of prematurity, IVH intraventricular hemorrhage, MCA Doppler middle cerebral artery Doppler

with a significant decrease in ABDs events.⁹⁹ While these early studies suggest that variation in RR or apnea/PB events can portend adverse events, the sensitivity and specificity for predicting outcomes appear modest. Moreover, unlike HRC-based algorithms, standardized criteria to delineate normal from abnormal ranges have not been tested in randomized clinical trials. Whether addition of continuous RR and apnea monitoring into complex algorithms including other VS would increase sensitivity or specificity for predicting clinical outcomes remains to be answered (Fig. 2).

CONTINUOUS BLOOD PRESSURE (BP) MONITORING

Physiological determinants and monitoring of blood pressure BP is determined by the product of cardiac output and systemic vascular resistance and increases with birth weight, higher GA, and with postnatal adaptation to extra-uterine life.^{100,101} BP in preterm infants is influenced by factors such as restricted myocardial compliance and contractility, poor tolerance to high SVR, antenatal steroids, chorioamnionitis, relative adrenal insufficiency, PDA, and mechanical ventilation.^{102–109} While BP is used as a surrogate to monitor systemic blood flow and tissue perfusion, it is impacted by the presence of ductal and atrial shunts, and thus may not accurately reflect cardiac output or tissue perfusion.^{110,111} BP is measured noninvasively by cuffs that estimate blood pressure by quantifying oscillometric changes in the arterial wall generated by pulsatile blood flow.¹¹² Although easy to use, this method can overestimate pressures, especially in sick, hypotensive preterm infants.^{113–116} Invasive arterial BP measurement allows continuous direct monitoring of the BP generated by every heartbeat. In the immediate postnatal period, an umbilical artery catheter may be inserted into the abdominal or thoracic aorta and connected to a transducer to allow continuous measurement of BP.¹¹² Later in the NICU stay, arterial BP may be continuously monitored through a peripheral arterial catheter during times of critical illness.

In preterm neonates, low mean arterial blood pressure (MBP) in the first week of life has been associated with higher risk of mortality, and worse neurodevelopmental outcomes.^{117,118} Continuous BP studies have primarily focused on the first week of life because of a putative “window” of the enhanced vulnerability of preterm brain to perfusion-related brain injury and ready availability of umbilical arterial catheter data. Studies utilizing continuous BP trends have been used to glean normative blood pressure data in preterm infants, assess cerebral perfusion, and study patterns of brain injury and neurological outcomes (Table 3).^{119–122}

Defining normative BP

A study of 35 infants with mean GA of 25 weeks analyzed >11 million BP values generated from continuous sampling every 2 s in the first 72 h of life.¹⁰⁹ Optimum MBP estimated from these data was higher with a narrow range than previously published, at around 33 ± 3 mmHg. In a retrospective study, where mean BP was averaged over 1 min in 232 VLBW over the first 7 days of life, similar values were generated.¹⁰⁰ The discrepancies between normative data generated by continuous BP sampling from published studies may be due to the use of interval data collection with different sampling periods, and inconsistent methodology such as utilizing both noninvasive oscillometer and intra-arterial manometric measurements.^{123–125}

Association of early postnatal hypotension with neurological outcomes

Computerized analysis of continuous BP data suggests that early low MBP can predict adverse events in preterm infants. In a study of infants with GA <31 weeks, MBP <30 mmHg for over an hour

was associated with severe IVH, ischemic cerebral lesions, and mortality within 48 h.¹¹⁹ A similar relationship between low BP and grade II + IVH was observed in 100 VLBW infants.¹²⁰ In this study IVH was also associated with wider swings in MBP. In a study of 98 infants with GA <34 weeks, continuous HR, BP, and transcutaneous oxygen tension (PaO₂) data from the first 4 days of life were analyzed.¹²² Among infants with hypotension and hypoxemia, 53% had either postmortem evidence of brain injury or abnormal neurological outcome at 1 year of age, compared to an 8% incidence without hypotension or hypoxemia. In contrast to the findings in these studies, several studies^{126–133} including the large ELGAN study^{134,135} did not find an association between early hypotension and adverse neurological outcomes. These different findings may be attributable to analysis of intermittent data either from monitors or medical records, and mixed use of cuff-based and invasive BP measurements.

Immature autoregulation of cerebral blood flow in preterm infants leads to a pressure passive state, a known risk factor for cerebrovascular injury.^{136–138} In past studies, BP variability alone has produced conflicting results related to IVH prediction,^{100,139} but showed a strong association when combined with changes in cerebral blood-flow velocity in the anterior cerebral artery.¹⁴⁰ The unreliability of isolated MBP monitoring is exemplified by a study in 90 VLBW infants where both continuous MBP and cerebral perfusion using near-infrared spectroscopy (NIRS) in the first 5 days was measured.¹⁴¹ Although this study did find a strong correlation between pressure passivity and hypotension, MBP and NIRS were not invariably associated with each other. Another study of 44 infants with median GA of 25 weeks suggested that identifying individualized optimum MBP based on the correlation of cerebrovascular reactivity (estimated from HR and NIRS tissue oxygenation index) can better predict IVH.¹⁴² In a study of 25 preterm infants continuous MBP and electrical cortical activity by spectral analysis using continuous multichannel EEG recordings was measured.¹⁴³ This study found increased mortality if there was a stronger directionality of interaction between MBP and aEEG leading to decreasing spectral power of EEG with decreasing MBP. Although mean BP alone was predominantly used in these prior studies, recently BP parameters such as systolic (SBP), diastolic (DBP), and pulse pressure (PP) are also being explored. In a study of 485 neonates with mean GA of 28 weeks, left ventricular output was positively associated with SBP and PP, and not MBP and DBP.¹⁴⁴

In summary, continuous BP data, rather than intermittent values, have the potential to predict short- and long-term adverse neurological outcomes. However, rapid hemodynamic changes occurring in the first week after birth decrease sensitivity of BP as a tool to monitor cardiac output and tissue perfusion. Ideally, BP analysis should be combined with other measurements such as functional echocardiography, NIRS, aEEG, and Doppler studies of regional blood flow to predict and potentially prevent adverse outcomes.^{110,111,145–148}

CONCLUSION

It is increasingly evident that intermittent assessment of VS from review of standard bedside monitor data or medical record documentation fails to capture trends that portend impending pathophysiology, and that analysis of continuous VS data can improve outcomes of infants in the NICU. Display to clinicians of a score based on continuous analysis of HR characteristics has been shown in a large randomized clinical trial to decrease sepsis-associated mortality in preterm VLBW infants. While various analytics of continuous SpO₂, RR, apnea, and BP data either alone or in combination show promise for disease prevention or mitigation, they require more validation in randomized clinical

trials. The prospect of computerized, artificial intelligence-based multi-tier analysis (Fig. 2) incorporating several vital signs will likely offer a plethora of actionable data to guide bedside management. A three-tiered approach incorporating VS analytics, careful consideration of clinical variables, complimented by laboratory testing and imaging represents a “trifecta” for using readily accessible continuous VS data to prevent neonatal diseases in the twenty-first century.

ACKNOWLEDGEMENTS

This study is supported by Children’s Mercy Research Institute (V.S.); NIH HD072071 (K.F.), Translational Health Research Institute of Virginia Scholars Award (B.S.).

AUTHOR CONTRIBUTIONS

Conception and design: N.K., K.F., and V.S.; data acquisition, drafting manuscript/figures, revising article: N.K., G.A., B.S., K.F., V.S.

ADDITIONAL INFORMATION

Competing interests: The authors declare no competing interests.

Publisher’s note: Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

REFERENCES

- Leante-Castellanos, J. L. et al. Central-peripheral temperature monitoring as a marker for diagnosing late-onset neonatal sepsis. *Pediatr. Infect. Dis. J.* **36**, e297 (2017).
- Fairchild, K. D. et al. Septicemia mortality reduction in neonates in a heart rate characteristics monitoring trial. *Pediatr. Res.* **74**, 570–575 (2013).
- Ruiz, T. L., Trzaski, J. M., Sink, D. W. & Hagadorn, J. I. Transcribed oxygen saturation vs oximeter recordings in very low birth weight infants. *J. Perinatol.* **34**, 130–135 (2014).
- Warburton, A., Monga, R., Sampath, V. & Kumar, N. Continuous pulse oximetry and respiratory rate trends predict short-term respiratory and growth outcomes in premature infants. *Pediatr. Res.* **85**, 494–501 (2019).
- Cardona-Morrell, M. et al. Effectiveness of continuous or intermittent vital signs monitoring in preventing adverse events on general wards: a systematic review and meta-analysis. *Int. J. Clin. Pract.* **70**, 806–824 (2016).
- Chapman, S. M. & Maconochie, I. K. Early warning scores in paediatrics: an overview. *Arch. Dis. Child.* **104**, 395–399 (2018).
- Nghiem, T. et al. Nurse opinions and pulse oximeter saturation target limits for preterm infants. *Pediatrics* **121**, e1046 (2008).
- Das, A. et al. Effect of fluctuation of oxygenation and time spent in the target range on retinopathy of prematurity in extremely low birth weight infants. *J. Neonatal-Perinat. Med.* **11**, 257–263 (2018).
- Amin, S. & Burnell, E. Monitoring apnea of prematurity: validity of nursing documentation and bedside cardiorespiratory monitor. *Am. J. Perinatol.* **30**, 643–648 (2013).
- Muttitt, S. C., Finer, N. N., Tierney, A. J. & Rossmann, J. Neonatal apnea: diagnosis by nurse versus computer. *Pediatrics* **82**, 713 (1988).
- Brockmann, P. E. et al. Under-recognition of alarms in a neonatal intensive care unit. *Arch. Dis. Child. Fetal Neonatal Ed.* **98**, F527 (2013).
- Das, A. et al. A comparison of manual versus automated saturation of peripheral oxygenation in the neonatal intensive care unit. *J. Matern.-Fetal Neonatal Med.* **29**, 1631–1635 (2016).
- Hay, W. W. et al. Reliability of conventional and new pulse oximetry in neonatal patients. *J. Perinatol.* **22**, 360–366 (2002).
- Alonzo, C. J. et al. Heart rate ranges in premature neonates using high resolution physiologic data. *J. Perinatol.* **38**, 1242–1245 (2018).
- Shaffer, F. & Ginsberg, J. P. An overview of heart rate variability metrics and norms. *Front. public health* **5**, 258 (2017).
- Hon, E. H. & Lee, S. T. Electronic evaluation of the fetal heart rate. VIII. Patterns preceding fetal death, further observations. *Am. J. Obstet. Gynecol.* **87**, 814 (1963).
- Herry, C. L. et al. Temporal patterns in sheep fetal heart rate variability correlate to systemic cytokine inflammatory response: a methodological exploration of monitoring potential using complex signals bioinformatics. *PLoS ONE* **11**, e0153515 (2016).
- Durosier, L. D. et al. Does heart rate variability reflect the systemic inflammatory response in a fetal sheep model of lipopolysaccharide-induced sepsis? *Physiol. Meas.* **36**, 2089–2102 (2015).
- Griffin, M. P. et al. Abnormal heart rate characteristics preceding neonatal sepsis and sepsis-like illness. *Pediatr. Res.* **53**, 920–926 (2003).
- Fairchild, K. D., Srinivasan, V., Moorman, J. R., Gaykema, R. P. A. & Goehler, L. E. Pathogen-induced heart rate changes associated with cholinergic nervous system activation. *Am. J. Physiol.* **300**, 330–339 (2011).
- Fairchild, K. D. et al. Endotoxin depresses heart rate variability in mice: cytokine and steroid effects. *Am. J. Physiol.* **297**, 1019–1027 (2009).
- Huston, J. M. & Tracey, K. J. The pulse of inflammation: heart rate variability, the cholinergic anti-inflammatory pathway and implications for therapy. *J. Intern. Med.* **269**, 45–53 (2011).
- Greenberg, R. G. et al. Late-onset sepsis in extremely premature infants: 2000–2011. *Pediatr. Infect. Dis. J.* **36**, 774–779 (2017).
- Yee, W. H. et al. Incidence and timing of presentation of necrotizing enterocolitis in preterm infants. *Pediatrics* **129**, e304 (2012).
- Adams-Chapman, I. & Stoll, B. Neonatal infection and long-term neurodevelopmental outcome in the preterm infant. *Curr. Opin. Infect. Dis.* **19**, 290–297 (2006).
- Griffin, M. P. et al. Heart rate characteristics: novel physiologic markers to predict neonatal infection and death. *Pediatrics* **116**, 1070–1074 (2005).
- Lake, D. E., Richman, J. S., Griffin, M. P. & Moorman, J. R. Sample entropy analysis of neonatal heart rate variability. *Am. J. Physiol.* **283**, 789–797 (2002).
- Kovatchev, B. P. et al. Sample asymmetry analysis of heart rate characteristics with application to neonatal sepsis and systemic inflammatory response syndrome. *Pediatr. Res.* **54**, 892–898 (2003).
- Moorman, J. R. et al. Mortality reduction by heart rate characteristic monitoring in very low birth weight neonates: a randomized trial. *J. Pediatr.*, **159**, 906.e1 (2011).
- Stone, M. L. et al. Abnormal heart rate characteristics before clinical diagnosis of necrotizing enterocolitis. *J. Perinatol.* **33**, 847–850 (2013).
- Doheny, K. K. et al. Diminished vagal tone is a predictive biomarker of necrotizing enterocolitis-risk in preterm infants. *Neurogastroenterol. Motil.* **26**, 832–840 (2014).
- Addison, K., Griffin, M. P., Moorman, J. R., Lake, D. E. & O’Shea, T. M. Heart rate characteristics and neurodevelopmental outcome in very low birth weight infants. *J. Perinatol.* **29**, 750–756 (2009).
- Sullivan, B. A. et al. Early heart rate characteristics predict death and morbidities in preterm infants. *J. Pediatr.* **174**, 57–62 (2016).
- Sullivan, B. A., Grice, S. M., Lake, D. E., Moorman, J. R. & Fairchild, K. D. Infection and other clinical correlates of abnormal heart rate characteristics in preterm infants. *J. Pediatr.* **164**, 775–780 (2014).
- Clark, M. T. et al. Predictive monitoring for respiratory decompensation leading to urgent unplanned intubation in the neonatal intensive care unit. *Pediatr. Res.* **73**, 104–110 (2013).
- Goel, N., Chakraborty, M., Watkins, W. J. & Banerjee, S. Predicting extubation outcomes—a model incorporating heart rate characteristics index. *J. Pediatr.* **195**, 58.e1 (2018).
- Fairchild, K. D. et al. Abnormal heart rate characteristics are associated with abnormal neuroimaging and outcomes in extremely low birth weight infants. *J. Perinatol.* **34**, 375–379 (2014).
- Vergales, B. et al. Depressed heart rate variability is associated with abnormal EEG, MRI, and death in neonates with hypoxic ischemic encephalopathy. *Am. J. Perinatol.* **31**, 855–862 (2014).
- Tuzcu, V., Nas, S., Ulusar, U., Ugur, A. & Kaiser, J. R. Altered heart rhythm dynamics in very low birth weight infants with impending intraventricular hemorrhage. *Pediatrics* **123**, 810–815 (2009).
- Sortica da Costa, C. et al. Complexity of brain signals is associated with outcome in preterm infants. *J. Cereb. Blood Flow. Metab.* **37**, 3368–3379 (2017).
- Griffin, M. P., Lake, D. E., O’Shea, T. M. & Moorman, J. R. Heart rate characteristics and clinical signs in neonatal sepsis. *Pediatr. Res.* **61**, 222–227 (2007).
- Fairchild, K. et al. Clinical associations of immature breathing in preterm infants: part 1—central apnea. *Pediatr. Res.* **80**, 21–27 (2016).
- Patel, M. et al. Clinical associations with immature breathing in preterm infants: part 2—periodic breathing. *Pediatr. Res.* **80**, 28–34 (2016).
- Siljehav, V., Hofstetter, A. M., Leifsdottir, K. & Herlenius, E. Prostaglandin E2 mediates cardiorespiratory disturbances during infection in neonates. *J. Pediatr.* **167**, 1213.e3 (2015).
- Fairchild, K. D. & Lake, D. E. Cross-correlation of heart rate and oxygen saturation in very low birthweight infants: association with apnea and adverse events. *Am. J. Perinatol.* **35**, 463–469 (2018).
- Fairchild, K. D. et al. Vital signs and their cross-correlation in sepsis and NEC: a study of 1,065 very-low-birth-weight infants in two NICUs. *Pediatr. Res.* **81**, 315–321 (2017).

47. Saria, S., Rajani, A. K., Gould, J., Koller, D. & Penn, A. A. Integration of early physiological responses predicts later illness severity in preterm infants. *Sci. Transl. Med.* **2**, 48ra65 (2010).
48. Gur, I. et al. Pilot study of a new mathematical algorithm for early detection of late-onset sepsis in very low-birth-weight infants. *Am. J. Perinatol.* **32**, 321–330 (2015).
49. Wukitsch, M. W., Petterson, M. T., Tobler, D. R. & Pologe, J. A. Pulse oximetry: analysis of theory, technology, and practice. *J. Clin. Monit.* **4**, 290–301 (1988).
50. Avidan, A. & Levin, P. D. Pulse oximetry. *New Engl. J. Med.* **365**, 184 (2011). author reply 185.
51. Jubran, A. Pulse oximetry. *Crit. Care* **19**, 272 (2015).
52. Buonocore, G., Perrone, S. & Tataranno, M. L. Oxygen toxicity: chemistry and biology of reactive oxygen species. *Semin. Fetal Neonatal Med.* **15**, 186–190 (2010).
53. Davis, J. M. & Auten, R. L. Maturation of the antioxidant system and the effects on preterm birth. *Semin. Fetal Neonatal Med.* **15**, 191–195 (2010).
54. Saugstad, O. D. & Aune, D. Optimal oxygenation of extremely low birth weight infants: a meta-analysis and systematic review of the oxygen saturation target studies. *Neonatology* **105**, 55–63 (2013).
55. Askie, L. M. et al. Association between oxygen saturation targeting and death or disability in extremely preterm infants in the neonatal oxygenation prospective meta-analysis collaboration. *JAMA* **319**, 2190 (2018).
56. Poets, C. F. et al. Association between intermittent hypoxemia or bradycardia and late death or disability in extremely preterm infants. *JAMA* **314**, 595–603 (2015).
57. Di Fiore, J. M. et al. Patterns of oxygenation, mortality, and growth status in the surfactant positive pressure and oxygen trial cohort. *J. Pediatr.* **186**, 56.e1 (2017).
58. Hagadorn, J. I. et al. Achieved versus intended pulse oximeter saturation in infants born less than 28 weeks' gestation: the AVIOx Study. *Pediatrics* **118**, 1574–1582 (2006).
59. Lim, K. et al. Oxygen saturation targeting in preterm infants receiving continuous positive airway pressure. *J. Pediatr.* **164**, 736.e1 (2014).
60. APGAR, V. A proposal for a new method of evaluation of the newborn infant. *Curr. Res. Anesth. Analg.* **32**, 260 (1953).
61. Richardson, D. K., Corcoran, J. D., Escobar, G. J. & Lee, S. K. SNAP-II and SNAPPE-II: simplified newborn illness severity and mortality risk scores. *J. Pediatr.* **138**, 92–100 (2001).
62. Parry, G., Tucker, J. & Tarnow-Mordi, W. CRIB II: an update of the clinical risk index for babies score. *Lancet* **361**, 1789–1791 (2003).
63. Sullivan, B. A. et al. Early pulse oximetry data improves prediction of death and adverse outcomes in a two-center cohort of very low birth weight infants. *Am. J. Perinatol.* **35**, 1331–1338 (2018).
64. Raffay, T. M. et al. Neonatal intermittent hypoxemia events are associated with diagnosis of bronchopulmonary dysplasia at 36 weeks postmenstrual age. *Pediatr. Res.* **85**, 318–323 (2019).
65. Vesoulis, Z. A. et al. Early hypoxemia burden is strongly associated with severe intracranial hemorrhage in preterm infants. *J. Perinatol.* **39**, 48–53 (2019).
66. Di Fiore, J. M. et al. The relationship between patterns of intermittent hypoxia and retinopathy of prematurity in preterm infants. *Pediatr. Res.* **72**, 606–612 (2012).
67. Mascoll-Robertson, K. K., Viscardi, R. M. & Woo, H. C. The objective use of pulse oximetry to predict respiratory support transition in preterm infants: an observational pilot study. *Respir. Care* **61**, 416–422 (2016).
68. Pozo, M. E. et al. Effect of postnatal intermittent hypoxia on growth and cardiovascular regulation of rat pups. *Neonatology* **102**, 107–113 (2012).
69. Bizzarro, M. J. et al. Temporal quantification of oxygen saturation ranges: an effort to reduce hyperoxia in the neonatal intensive care unit. *J. Perinatol.* **34**, 33–38 (2014).
70. Claire, N. et al. Multicenter crossover study of automated control of inspired oxygen in ventilated preterm infants. *Pediatrics* **127**, e83 (2011).
71. van Kaam, A. H. et al. Automated versus manual oxygen control with different saturation targets and modes of respiratory support in preterm infants. *J. Pediatr.* **167**, 550.e2 (2015).
72. Zapata, J., Gómez, J. J., Araque Campo, R., Matiz Rubio, A. & Sola, A. A randomized controlled trial of an automated oxygen delivery algorithm for preterm neonates receiving supplemental oxygen without mechanical ventilation. *Acta Paediatr.* **103**, 928–933 (2014).
73. Saihi, K. et al. Feasibility and reliability of an automated controller of inspired oxygen concentration during mechanical ventilation. *Crit. Care* **18**, R35 (2014).
74. Hallenberger, A., Poets, C. F., Horn, W., Seyfang, A. & Urschitz, M. S. Closed-loop automatic oxygen control (CLAC) in preterm infants: a randomized controlled trial. *Pediatrics* **133**, e385 (2014).
75. Eduardo, C. N. Closed-loop control of inspired oxygen in premature infants. *Semin. Fetal Neonatal Med.* **20**, 198–204 (2015).
76. Claire, N. & Bancalari, E. Automated closed loop control of inspired oxygen concentration. *Respir. Care* **58**, 151–161 (2013).
77. Di Fiore, J. M. Neonatal cardiorespiratory monitoring techniques. *Semin. Neonatol.* **9**, 195–203 (2004).
78. Folke, M., Cernerud, L., Ekström, M. & Hök, B. Critical review of non-invasive respiratory monitoring in medical care. *Med Biol. Eng. Comput.* **41**, 377–383 (2003).
79. Lee, H. et al. A new algorithm for detecting central apnea in neonates. *Physiol. Meas.* **33**, 1–17 (2012).
80. Weese-Mayer, D. E. et al. Comparison of apnea identified by respiratory inductance plethysmography with that detected by end-tidal CO₂ or thermistor. The CHIME Study Group. *Am. J. Respir. Crit. Care Med.* **162**(2 Pt 1), 471–480 (2000).
81. Massaroni, C. et al. Contact-based methods for measuring respiratory rate. *Sensors* **19**, 908 (2019).
82. Charlton, P. H. et al. Breathing rate estimation from the electrocardiogram and photoplethysmogram: a review. *RBME* **11**, 2–20 (2018).
83. Gerhardt, T. & Bancalari, E. Apnea of prematurity: I. Lung function and regulation of breathing. *Pediatrics* **74**, 58 (1984).
84. Martin, R. J. et al. Vulnerability of respiratory control in healthy preterm infants placed supine. *J. Pediatr.* **127**, 609–614 (1995).
85. Katz-Salamon, M. Delayed chemoreceptor responses in infants with apnoea. *Arch. Dis. Child.* **89**, 261–266 (2004).
86. Gauda, E. B., McLemore, G. L., Tolosa, J., Marston-Nelson, J. & Kwak, D. Maturation of peripheral arterial chemoreceptors in relation to neonatal apnoea. *Semin. Neonatol.* **9**, 181–194 (2004).
87. Abu-Shaweeh, J. M. & Martin, R. J. Neonatal apnea: what's new? *Pediatr. Pulmonol.* **43**, 937–944 (2008).
88. Darnall, R. A. The role of CO₂ and central chemoreception in the control of breathing in the fetus and the neonate. *Respir. Physiol. Neurobiol.* **173**, 201–212 (2010).
89. Balan, K. V. et al. Vagal afferents modulate cytokine-mediated respiratory control at the neonatal medulla oblongata. *Respir. Physiol. Neurobiol.* **178**, 458–464 (2011).
90. Balan, K. V. et al. Intrapulmonary lipopolysaccharide exposure upregulates cytokine expression in the neonatal brainstem. *Acta Paediatr.* **101**, 466–471 (2012).
91. Vergales, B. et al. Accurate automated apnea analysis in preterm infants. *Am. J. Perinatol.* **31**, 157–162 (2014).
92. Southall, D. P. et al. Undetected episodes of prolonged apnea and severe bradycardia in preterm infants. *Pediatrics* **72**, 541 (1983).
93. Mohr, M. A. et al. Very long apnea events in preterm infants. *J. Appl. Physiol.* **118**, 558–568 (2015).
94. Zagol, K. et al. Anemia, apnea of prematurity, and blood transfusions. *J. Pediatr.* **161**, 421.e1 (2012).
95. Mohr, M. A. et al. Quantification of periodic breathing in premature infants. *Physiol. Meas.* **36**, 1415–1427 (2015).
96. Hofstetter, A. O., Legnevall, L., Herlenius, E. & Katz-Salamon, M. Cardiorespiratory development in extremely preterm infants: vulnerability to infection and persistence of events beyond term-equivalent age. *Acta Paediatr.* **97**, 285–292 (2008).
97. Shannon, D. C., Carley, D. W. & Kelly, D. H. Periodic breathing: quantitative analysis and clinical description. *Pediatr. Pulmonol.* **4**, 98–102 (1988).
98. Barrington, K. J. & Finer, N. N. Periodic breathing and apnea in preterm infants. *Pediatr. Res.* **27**, 118–121 (1990).
99. Tabacaru, C. R. et al. Impact of caffeine boluses and caffeine discontinuation on apnea and hypoxemia in preterm infants. *J. Caffeine Res.* **7**, 13–110 (2017).
100. Cunningham, S., Symon, A. G., Elton, R. A., Zhu, C. & McIntosh, N. Intra-arterial blood pressure reference ranges, death and morbidity in very low birthweight infants during the first seven days of life. *Early Hum. Dev.* **56**, 151–165 (1999).
101. Jones, J. E. & Jose, P. A. Neonatal blood pressure regulation. *Semin. Perinatol.* **28**, 141–148 (2004).
102. Toyono, M., Harada, K., Takahashi, Y. & Takada, G. Maturation changes in left ventricular contractile state. *Int. J. Cardiol.* **64**, 247–252 (1998).
103. Takahashi, Y. et al. Postnatal left ventricular contractility in very low birth weight infants. *Pedia. Cardiol.* **18**, 112–117 (1997).
104. Alan, F. A. Physiology of the fetal and transitional circulation. *Semin. Fetal Neonatal Med.* **20**, 210–216 (2015).
105. Ng, P. C. et al. Transient adrenocortical insufficiency of prematurity and systemic hypotension in very low birthweight infants. *Arch. Dis. Child. Fetal Neonatal Ed.* **89**, 119 (2004).
106. Laughon, M. M., Simmons, M. A. & Bose, C. L. Patency of the ductus arteriosus in the premature infant: is it pathologic? Should it be treated? *Curr. Opin. Pediatr.* **16**, 146–151 (2004).

107. Kluckow, M. Low systemic blood flow and pathophysiology of the preterm transitional circulation. *Early Hum. Dev.* **81**, 429–437 (2005).
108. Fernandez, E. F. & Watterberg, K. L. Relative adrenal insufficiency in the preterm and term infant. *J. Perinatol.* **29**(S2), S49 (2009).
109. Vesoulis, Z. A., El Ters, N. M., Wallendorf, M. & Mathur, A. M. Empirical estimation of the normative blood pressure in infants. *J. Perinatol.* **36**, 291–295 (2016).
110. Pladys, P., Wodey, E., Beuchée, A., Branger, B. & Bétrémieux, P. Left ventricle output and mean arterial blood pressure in preterm infants during the 1st day of life. *Eur. J. Pediatr.* **158**, 817–824 (1999).
111. Kluckow, M. & Evans, N. Relationship between blood pressure and cardiac output in preterm infants requiring mechanical ventilation. *J. Pediatr.* **129**, 506–512 (1996).
112. Weindling, A. M. Blood pressure monitoring in the newborn. *Arch. Dis. Child.* **64** (4 Spec No), 444–447 (1989).
113. Gevers, M., van Genderingen, H. R., Lafeber, H. N. & Hack, W. W. Accuracy of oscillometric blood pressure measurement in critically ill neonates with reference to the arterial pressure wave shape. *Intensive Care Med.* **22**, 242–248 (1996).
114. Takci, S., Yigit, S., Korkmaz, A. & Yurdakök, M. Comparison between oscillometric and invasive blood pressure measurements in critically ill premature infants. *Acta Paediatr.* **101**, 132–135 (2012).
115. Dannevig, I., Dale, H. C., Liestol, K. & Lindemann, R. Blood pressure in the neonate: three non-invasive oscillometric pressure monitors compared with invasively measured blood pressure. *Acta Paediatr.* **94**, 191–196 (2005).
116. Werther, T. et al. Discrepancy between invasive and non-invasive blood pressure readings in extremely preterm infants in the first four weeks of life. *PLoS ONE* **13**, e0209831 (2018).
117. Goldstein, R. F., Thompson, R. J. J., Oehler, J. M. & Brazy, J. E. Influence of acidosis, hypoxemia, and hypotension on neurodevelopmental outcome in very low birth weight infants. *Pediatrics* **95**, 238 (1995).
118. Fanaroff, J. M., Wilson-Costello, D. E., Newman, N. S., Montpetite, M. M. & Fanaroff, A. A. Treated hypotension is associated with neonatal morbidity and hearing loss in extremely low birth weight infants. *Pediatrics* **117**, 1131–1135 (2006).
119. Miall-Allen, V. M., de Vries, L. S. & Whitelaw, A. G. Mean arterial blood pressure and neonatal cerebral lesions. *Arch. Dis. Child.* **62**, 1068–1069 (1987).
120. Bada, H. S. et al. Mean arterial blood pressure changes in premature infants and those at risk for intraventricular hemorrhage. *J. Pediatr.* **117**, 607–614 (1990).
121. Fok, T. F., Davies, D. P. & Ng, H. K. A study of periventricular haemorrhage, post-haemorrhagic ventricular dilatation and periventricular leucomalacia in Chinese preterm infants. *J. Paediatr. Child Health* **26**, 271–275 (1990).
122. Low, J. A. et al. The association between preterm newborn hypotension and hypoxemia and outcome during the first year. *Acta Paediatr.* **82**, 433–437 (1993).
123. Versmold, H. T., Kitterman, J. A., Phibbs, R. H., Gregory, G. A. & Tooley, W. H. Aortic blood pressure during the first 12 h of life in infants with birth weight 610 to 4,220 grams. *Pediatrics* **67**, 607 (1981).
124. Lee, J., Rajadurai, V. S. & Tan, K. W. Blood pressure standards for very low birthweight infants during the first day of life. *Arch. Dis. Child. Fetal Neonatal Ed.* **81**, F170 (1999).
125. Batton, B. et al. Evolving blood pressure dynamics for extremely preterm infants. *J. Perinatol.* **34**, 301–305 (2014).
126. Trounce, J. Q., Shaw, D. E., Levene, M. I. & Rutter, N. Clinical risk factors and periventricular leucomalacia. *Arch. Dis. Child.* **63**, 17–22 (1988).
127. Grönlund, J. U., Korvenranta, H., Kero, P., Jalonen, J. & Välimäki, I. A. Elevated arterial blood pressure is associated with peri-intraventricular haemorrhage. *Eur. J. Pediatr.* **153**, 836–841 (1994).
128. D'Souza, S. W. et al. Blood pressure, heart rate, and skin temperature in preterm infants: associations with periventricular haemorrhage. *Arch. Dis. Child. Fetal Neonatal Ed.* **72**, F167 (1995).
129. Perlman, J. M., Risser, R. & Broyles, R. S. Bilateral cystic periventricular leukomalacia in the premature infant: associated risk factors. *Pediatrics* **97**, 822 (1996).
130. Wiswell, T. E., Graziani, L. J., Kornhuaser, M. S. & Stanley, C. Effects of hypocarbia on the development of cystic periventricular leukomalacia in premature infants treated with high-frequency jet ventilation. *Pediatrics* **98**, 918 (1996).
131. Dammann, O. et al. Systemic hypotension and white-matter damage in preterm infants. *Dev. Med. Child Neurol.* **44**, 82–90 (2002).
132. Limperopoulos, C. et al. Current definitions of hypotension do not predict abnormal cranial ultrasound findings in preterm infants. *Pediatrics* **120**, 966–977 (2007).
133. Batton, B. et al. Blood pressure, anti-hypotensive therapy, and neurodevelopment in extremely preterm infants. *J. Pediatr.* **154**, 357.e1 (2009).
134. Logan, J. W. et al. Early postnatal hypotension and developmental delay at 24 months of age among extremely low gestational age newborns. *Arch. Dis. Child. Fetal Neonatal Ed.* **96**, F328 (2011).
135. Logan, J. W. et al. Early postnatal hypotension is not associated with indicators of white matter damage or cerebral palsy in extremely low gestational age newborns. *J. Perinatol.* **31**, 524–534 (2011).
136. Hoffman, S. B., Cheng, Y., Magder, L. S., Shet, N. & Viscardi, R. M. Cerebral autoregulation in premature infants during the first 96 h of life and relationship to adverse outcomes. *Archives of disease in childhood. Arch. Dis. Child. Fetal Neonatal Ed.* **0**, F1–F7 (2018).
137. Lou, H. C., Lassen, N. A. & Friis-Hansen, B. Impaired autoregulation of cerebral blood flow in the distressed newborn infant. *J. Pediatr.* **94**, 118–121 (1979).
138. Tsuji, M. et al. Cerebral intravascular oxygenation correlates with mean arterial pressure in critically ill premature infants. *Pediatrics* **106**, 625–632 (2000).
139. Miall-Allen, V. M., de Vries, L. S., Dubowitz, L. M. & Whitelaw, A. G. Blood pressure fluctuation and intraventricular hemorrhage in the preterm infant of less than 31 weeks' gestation. *Pediatrics* **83**, 657 (1989).
140. Perlman, J. M., McMenam, J. B. & Volpe, J. J. Fluctuating cerebral blood-flow velocity in respiratory-distress syndrome. *New Engl. J. Med.* **309**, 204–209 (1983).
141. Soul, J. S. et al. Fluctuating pressure-passivity is common in the cerebral circulation of sick premature infants. *Pediatr. Res.* **61**, 467–473 (2007).
142. da Costa, C. S., Czosnyka, M., Smielewski, P. & Austin, T. Optimal mean arterial blood pressure in extremely preterm infants within the first 24 h of life. *J. Pediatr.* **203**, 242–248 (2018).
143. Semenova, O. et al. Coupling between mean blood pressure and EEG in preterm neonates is associated with reduced illness severity scores. *PLoS ONE* **13**, e0199587 (2018).
144. Kharrat, A. et al. The relationship between blood pressure parameters and left ventricular output in neonates. *J. Perinatol.* **39**, 619–625 (2019).
145. Groves, A. M., Kuschel, C. A., Knight, D. B. & Skinner, J. R. Relationship between blood pressure and blood flow in newborn preterm infants. *Arch. Dis. Child. Fetal Neonatal Ed.* **93**, F32 (2008).
146. Miletin, J. & Dempsey, E. M. Low superior vena cava flow on day 1 and adverse outcome in the very low birthweight infant. *Arch. Dis. Child. Fetal Neonatal Ed.* **93**, F371 (2008).
147. Miletin, J., Pichova, K. & Dempsey, E. Bedside detection of low systemic flow in the very low birth weight infant on day 1 of life. *Eur. J. Pediatr.* **168**, 809–813 (2009).
148. Kissack, C. M., Garr, R., Wardle, S. P. & Weindling, A. M. Cerebral fractional oxygen extraction in very low birth weight infants is high when there is low left ventricular output and hypocarbia but is unaffected by hypotension. *Pediatr. Res.* **55**, 400–405 (2004).