

Sipha maydis sensitivity to defences of *Lolium multiflorum* and its endophytic fungus *Epichloë occultans*

Daniel A. Bastías^{1,2}, Maria Alejandra Martínez-Ghersa¹, Jonathan A. Newman³, Stuart D. Card², Wade J. Mace² and Pedro E. Gundel¹

¹ IFEVA, Consejo Nacional de Investigaciones Científicas y Técnicas, Universidad de Buenos Aires, Buenos Aires, Argentina

² Forage Science, AgResearch Limited, Grasslands Research Centre, Palmerston North, New Zealand

³ Department of Biology, Wilfrid Laurier University, Waterloo, Ontario, Canada

ABSTRACT

Background. Plants possess a sophisticated immune system to defend from herbivores. These defence responses are regulated by plant hormones including salicylic acid (SA) and jasmonic acid (JA). Sometimes, plant defences can be complemented by the presence of symbiotic microorganisms. A remarkable example of this are grasses establishing symbiotic associations with *Epichloë* fungal endophytes. We studied the level of resistance provided by the grass' defence hormones, and that provided by *Epichloë* fungal endophytes, against an introduced herbivore aphid. These fungi protect their hosts against herbivores by producing bioactive alkaloids. We hypothesized that either the presence of fungal endophytes or the induction of the plant salicylic acid (SA) defence pathway would enhance the level of resistance of the grass to the aphid.

Methods. *Lolium multiflorum* plants, with and without the fungal endophyte *Epichloë occultans*, were subjected to an exogenous application of SA followed by a challenge with the aphid, *Sipha maydis*.

Results. Our results indicate that neither the presence of *E. occultans* nor the induction of the plant's SA pathway regulate *S. maydis* populations. However, endophyte-symbiotic plants may have been more tolerant to the aphid feeding because these plants produced more aboveground biomass. We suggest that this insect insensitivity could be explained by a combination between the ineffectiveness of the specific alkaloids produced by *E. occultans* in controlling *S. maydis* aphids and the capacity of this herbivore to deal with hormone-dependent defences of *L. multiflorum*.

Subjects Agricultural Science, Ecology, Entomology

Keywords Alkaloids, Beneficial microorganisms, Endophyte symbiosis, *Epichloë* fungal endophytes, Plant defences, Salicylic acid, Plant-herbivore interaction

INTRODUCTION

To defend from herbivore attacks, plants harbour a sophisticated immune system in which hormone pathways including salicylic acid (SA) and jasmonic acid (JA) mediate defence responses. These hormone pathways are known to be differentially involved in the response to distinct natural enemies. While the plant SA-dependent defence pathway

Submitted 12 August 2019
Accepted 20 November 2019
Published 18 December 2019

Corresponding author
Daniel A. Bastías,
daniel.bastias@agresearch.co.nz

Academic editor
Dezene Huber

Additional Information and
Declarations can be found on
page 14

DOI 10.7717/peerj.8257

© Copyright
2019 Bastías et al.

Distributed under
Creative Commons CC-BY 4.0

OPEN ACCESS

is generally induced by sap-sucking insect herbivores and biotrophic pathogens, the JA-dependent defence pathway acts in response to chewing insect herbivores and necrotrophic pathogens (Thaler, Humphrey & Whiteman, 2012; Ballaré, 2014). In some cases, the plant immune system can be complemented by hormone-independent mechanisms of defences. A particularly striking example of this is provided by grasses establishing symbiotic associations with *Epichloë* fungal endophytes (Clay, 1988; Bastías et al., 2017a; Bastías et al., 2017b). These fungi protect their host grasses against herbivores by the production of bioactive alkaloids (Panaccione, Beaulieu & Cook, 2014).

Some grasses establish symbiotic associations with asexual *Epichloë* fungal species that are strictly vertically-transmitted. These grass-*Epichloë* endophyte associations are usually mutualistic, since plants provide the fungus a place to live (the fungus is an obligate symbiont) and the fungus provides the grass with beneficial traits such as anti-herbivore defences (Gundel, Rudgers & Ghersa, 2011; Saikkonen, Saari & Helander, 2010; Schardl et al., 2013a; Schardl et al., 2013b). *Epichloë* fungal endophyte species can collectively synthesize a vast number of secondary metabolites and this diversity of bioactive compounds provides their host grasses defences against a range of herbivore species (Saikkonen, Saari & Helander, 2010; Schardl et al., 2013a; Schardl et al., 2013b). Four classes of *Epichloë*-derived alkaloids have been well-studied: pyrrolizidines (e.g., lolines), peramine, indole-diterpenes (e.g., lolitrem B, terpendoles), and ergot alkaloids (e.g., ergovaline) (Panaccione, Beaulieu & Cook, 2014; Saikkonen, Gundel & Helander, 2013; Schardl et al., 2013a; Schardl et al., 2013b). The alkaloid profile produced depends on the fungal species/strain, with some species/strains producing only one type of alkaloid (Schardl et al., 2013a; Schardl et al., 2013b). Moreover, the effectiveness of a given alkaloid type depends on its concentration and the herbivore species (Fuchs et al., 2017b; Bastías et al., 2017a; Bastías et al., 2017b; Bultman et al., 2018). Alkaloid production is dependent on the fungal biomass, herbivory level, plant ontogenic stage and nutritional status, plant tissue type, and some abiotic conditions (e.g., temperature, CO₂ levels) (Justus, Witte & Hartmann, 1997; Hunt et al., 2005; Rasmussen et al., 2007; Ryan et al., 2014; Fuchs et al., 2017a; Fuchs et al., 2017b; Gundel et al., 2018; Bultman et al., 2018).

The presence of biotrophic pathogens within plants is usually controlled by SA-dependent responses (Thaler, Humphrey & Whiteman, 2012). Recent research shows that the same pathway can also regulate the interaction with beneficial symbionts. For instance, the plant induction of the SA pathway inhibits the development and establishment of rhizobacterial and mycorrhizal symbionts in plant tissues (Cao et al., 2017; Bedini et al., 2018). Moreover, plant SA immune responses can also affect the activities performed by plant beneficial symbionts. For example, alkaloid production and nitrogen fixation performed by beneficial symbionts is reduced by the induction of the SA pathway in grasses, legumes, and ferns (Hayat et al., 2010; Bastías et al., 2018a; De Vries et al., 2018). Independently of the control exerted by plants on their beneficial symbionts, these symbionts can, in turn, modulate the SA pathway (Johnson et al., 2003; Stacey et al., 2006; Navarro-Meléndez & Heil, 2014; Dupont et al., 2015; Moreira, Abdala-Roberts & Castagneyrol, 2018; Ramos et al., 2018). For instance, genes encoding proteins pertaining to SA biosynthesis and signalling were downregulated by the presence of the endophyte

E. festucae F11 in *Lolium perenne* plants (Dupont et al., 2015). The symbiont's suppression of SA-dependent responses might be a mechanism used by these microorganisms to facilitate growth into plant tissues (Jung et al., 2012; Bastías et al., 2018a).

Here, we studied the level of resistance mediated by plant hormones and provided by an *Epichloë* endophyte fungus against an introduced aphid species in grasses. We hypothesized that either the presence of fungal endophytes or the induction of the plant SA pathway would enhance the level of resistance of the grass against the aphid. We subjected Italian ryegrass plants (*Lolium multiflorum*), symbiotic and non-symbiotic with the endophyte *Epichloë occultans* (Moon et al., 2000), to an exogenous SA application followed by a challenge with the hedgehog grain aphid (*Sipha maydis*). This aphid species is native to Eurasia, and individuals feed on cereals and other grasses (Skvarla et al., 2017). In Argentina, where this work was conducted, *S. maydis* was first discovered in 2002 and has since spread throughout temperate grasslands and cropping regions of the country (Corrales et al., 2007). Corrales et al. (2007) reported for this aphid species, average densities from 18 to 98 individuals per plant depending of the season and the geographic location. In temperate grasslands, the first populations were found in 2003 in the south-east of the Buenos Aires province (34°55'S, 57°57'W) (Corrales et al., 2007). In these grasslands, *Sipha maydis* is commonly found feeding on *L. multiflorum* plants with populations reaching sizes of around 250 individuals per plant (Chaneton & Omacini, 2007). The ryegrass *L. multiflorum* is a naturalised and abundant species in the Argentinian temperate grasslands. This species is native to the European Mediterranean region and was introduced in Argentina more than a century ago (Uchitel, Omacini & Chaneton, 2011). While many different alkaloids are produced across the genus *Epichloë*, *E. occultans* produces only loline alkaloids (i.e., N-formyllooline (NFL) and N-acetylnorloline (NANL)) (Sugawara et al., 2006; Moore et al., 2015; Bastías et al., 2017a; Bastías et al., 2017b). It has been shown that both the presence of *E. occultans* (Omacini et al., 2001; Miranda, Marina & Chaneton, 2011; Gundel et al., 2012; Ueno et al., 2015; Bastías et al., 2017a; Bastías et al., 2017b), and the loline alkaloids produced by this and other endophyte species provide the plant with protection from aphids (Johnson et al., 1985; Eichenseer, Dahlman & Bush, 1991; Wilkinson et al., 2000; Panaccione, Beaulieu & Cook, 2014).

We predicted that the presence of *Epichloë* fungal endophytes within their host plants would enhance the level of plant resistance against the aphid and consequently reduce the aphid's performance (i.e., aphid individual metabolic rates and population sizes). In addition, the stimulation of the SA-dependent defence response, by the exogenous application of the hormone, would increase the resistance in non-symbiotic plants, affecting negatively the aphid performance. However, we expected that the SA treatment would affect the endophyte-conferred plant resistance against aphids. Specifically, since *Epichloë* fungal endophytes are biotrophic microorganisms, the exogenously applied SA would impair the *Epichloë*, subsequently reducing its alkaloid production, and thus decreasing the resistance level and consequently, increasing the metabolic rates and population sizes of the aphids.

MATERIAL AND METHODS

Plant and aphid stocks

We worked with the annual grass plants of *Lolium multiflorum* both symbiotic (E+) and non-symbiotic (E-) with its common fungal endophyte *E. occultans* (Moon *et al.*, 2000). More than a decade ago, seeds of *L. multiflorum* with high percentages of endophyte infection were hand-collected from a naturalised population in the Pampean grassland (Argentina) (36°00'S, 61°5'W) (Gundel *et al.*, 2009). Immediately after collection, a proportion of these seeds were treated with the systemic fungicide Triadimenol (150 g kg⁻¹; Baytan®) in order to obtain endophyte-free individuals. Since then, plants of these two biotypes (i.e., E+ and E-) have been cultivated annually in a common garden (recall that *E. occultans* is strictly vertically transmitted), multiplying fresh seeds for experimentation [IFEVA - CONICET, Universidad de Buenos Aires, Argentina (34°35'S, 58°28'W)]. Genetic segregation between plant biotypes has been prevented by allowing individual plants to freely exchange pollen during flowering (Gundel *et al.*, 2012). Each late spring–early summer, ripe seeds produced by each plant biotype are harvested and evaluated for endophyte presence; after confirming the level of endophytes in each biotype, the seeds are stored in a 4 °C freezer. The endophyte detection is carried out by looking for fungal hypha in stained individual seeds following the “seed squash technique” (Bacon & White, 1994; Card *et al.*, 2011). For this, 100 seeds from each seed lot (E+ and E-), are incubated in NaOH (5%), stained with rose bengal and examined under a light microscope at 40X power. The seeds produced in 2014 were examined for endophyte infection frequency (E+: 99%, and E-: 1%; $n = 100$ each) and stored in cold and dry conditions until use (2015).

In early-spring 2015, individual aphids *S. maydis* (Passerini) were collected from the local extant vegetation dominated by cereals and wild grasses. Starting with approximately 150 apterous adults, an aphid population was established within a growth chamber [21 °C (± 1) constant, radiation 150 $\mu\text{mol m}^{-2} \text{s}^{-1}$, and photoperiod L16:D8 h] on wheat plants (Var. Cronox; Don Mario). Wheat plants were replaced periodically to provide fresh material for the aphid population. After 6 weeks, the aphid population was large enough to provide the required number of individual adult aphids for the experiment (see next section).

Experimental description

In 2015 during the normal growing season for *L. multiflorum* (autumn-winter-spring), 50 E+ and 50 E- plants were grown in 1.5 L pots, filled with a mix of soil, sand, and peat in equal proportions. Plants were watered to field capacity, as needed, to avoid water deficits. In early-spring, 28 E+ and 28 E- healthy plants were selected and transferred to a growth chamber with the same environmental conditions as described earlier. At that time, the plants averaged 48 tillers (range: 25–73) and were starting the reproductive stage (spike appearance). After careful examination to ensure there were no invertebrates present on the *L. multiflorum* plants, each plant was individually enclosed within a white cotton voile fabric bag supported by a tubular plastic net. Before the application of the hormone treatment (see below), the plants were acclimated to the growth chamber conditions for one week.

The experiment comprised a 2×2 full factorial design, with endophyte (E+, E-) and salicylic acid (SA+, SA-) as the experimental treatments. Fourteen plants from each biotype were sprayed with 10 ml of SA solution (0.5 mM; Biopack®, Argentina). The same procedure was done on the other 14 E+ and 14 E- plants but sprayed with 10 ml of distilled water. Three days later, each plant was challenged with 5 apterous adult aphids. This number of aphids as starting population was chosen based on field observations, where the colonization of *L. multiflorum* plants by *S. maydis* aphids is usually carried out by only few individuals (Chaneton & Omacini, 2007). The 3-days period between the application of SA and the aphid challenge was previously identified as enough time for the plants to develop a defence response prior to contact with aphids (Bastías et al., 2018a).

The aphid populations on the *L. multiflorum* plants developed over the next 24 days. We counted the number of insects on each plant (aphid population size) at days 7 and 14 (corresponding to days 10 and 17 since the SA application, respectively). At day 24 (27th since SA application), a group of approximately 35 randomly chosen aphids were sampled from a subset of 5 individual plants per treatment to measure the mass-specific standard metabolic rate (SMR) by means of open-flow respirometry.

Two serial samples of plant tissues were taken to measure the physiological concentrations of plant defence hormones and fungal alkaloids. The tissues were sampled from a subset of 8 plants per treatment, selected at random. The first sample was taken on day 3, just prior to the introduction of the aphids. Two leaf-blades were excised from one tiller per plant just before the aphid challenge (day 3 post SA application) to evaluate the concentrations of SA and JA hormones. Since the recognized role of JA-signalling pathway responses in plant defences (Thaler, Humphrey & Whiteman, 2012), JA concentration levels were also measured in response to the endophyte presence and the SA treatment. The second harvest comprised the removal of one tiller per E+ plant 7 days after the aphid challenge (10 days post SA application). We used the pseudostem from this tiller to evaluate the concentration of loline alkaloids [note that the fungal endophyte *E. occultans* only produces this type of alkaloid (Bastías et al., 2017a; Bastías et al., 2017b)]. We selected tillers in visibly good conditions but with symptoms of aphids feeding activities. Even though E- plants are incapable of producing loline alkaloids, E- were subjected to the same sampling procedure as E+ plants to avoid any manipulation-dependent effects (Cahill Jr, Castelli & Casper, 2002). We estimated that the total tissue removed for hormone and alkaloid assessments represented around 1% of the aboveground plant biomass. At day 27 post SA application, the total aboveground biomass of the plants was harvested, and immediately dried in an oven (2 d at 60 °C) to evaluate the individual plant dry weight (Analytical scale, ± 0.01 g, Mettler Toledo).

Quantification of SA and JA hormones

Starting from freeze-dried and ground leaf material, subsamples of 50–100 mg were extracted with 100% Acetonitrile containing 100 ng of d6-SA and d5-JA as internal standards. The extracts were dried, derivatized with N-Methyl-N-(trimethylsilyl)trifluoroacetamide, and injected into an Agilent DB-5MS column (30 m, 0.25 mm inner diameter, 0.25 μ m film thickness with a 10 m guard column). The column effluent was added into the ion source

of a Scion TQ GC-MS/MS (Bruker Daltonics Inc.). The mass spectrometer was operated in positive ionization mode with multiple reactions monitoring (MRM) as previously described in [Bastías et al. \(2018a\)](#). SA and JA hormones were quantified relative to the peak area of their corresponding internal standards.

Quantification of loline alkaloids

Lolines were extracted from 50 mg of freeze-dried, ground plant samples using a solution of 40% methanol/5% ammonia and 1,2-dichloroethane containing 54.8 ng mL⁻¹ 4-phenylmorpholine as internal standard. Plant extracts were centrifugated, and supernatants transferred to glass GC vials via a 20 µm filter for analysis. The analysis was conducted using a GC flame ionization detector (GC2010Plus, Shimadzu Corporation, Japan) and separation was achieved on a ZB-5 capillary column (30 m × 0.32 mm × 0.25 µm film). More information can be found in [Bastías et al. \(2018a\)](#). The detection limit was 25 µg g⁻¹ DW.

Measurements of standard metabolic rate (SMR)

The SMR quantifies the energy budget required for insects to maintain homeostasis, thus this variable represents a measure of the general physiological status of insects ([Nespolo, Roff & Fairbairn, 2008](#)). Aphid maximum annual growth rates (r_m) are negatively correlated with SMR. Aphids with high metabolic rates (i.e., high levels of maintenance costs) might have less energy available for reproduction thus negatively impacting their population sizes ([Castañeda, Figueroa & Nespolo, 2010](#)). The SMR was measured by the production of CO₂ (VCO₂) of aphid groups placed within of an open-flow respirometry system (LI-6400; Li-Cor, Lincoln, USA). We estimated the mass-specific SMR that is, the amount of CO₂ produced per mass unit of aphid per hour (i.e., µL of VCO₂ per mg of aphid per hour). Aphids were obtained from a subset of 5 randomly chosen plants per treatment. From each of these plants, 35 adult and non-winged aphids were carefully removed and placed in Eppendorf tubes. We considered each group of aphids from one individual plant to be a replicate (i.e., 5 replicates per treatment). All the aphid groups were weighted (± 0.01 mg, analytical balance, Mettler Toledo), and kept for one hour without food before the metabolic measurements. The VCO₂ of each aphid group was registered every second during a period of 10 min at 24 °C (± 0.5). For this, the insects were placed in 10 mL-metabolic chamber that received CO₂-scrubbed air at a constant rate of 70 mL min⁻¹ and connected to a sensor of CO₂ (LI-6400; Li-Cor, Lincoln, USA). The mass-specific SMR was obtained averaging the 2-mins continuous and most stable VCO₂ values (from the 10-mins register) and dividing this averaged value by the insect group weight. Aphids were discarded after the SMR measurements.

Statistical analyses

The effects of the plant symbiotic status and SA application on the concentrations of SA and JA hormones, and on the plant above-ground biomass were analysed separately with linear effects models, using the function `gls` from the `nlme` package in R software ([R Core Team, 2013](#)), and assuming independent, identically distributed normal random errors ([Pinheiro et al., 2009](#)). The models included the plant's symbiotic status (E+, E-) and SA

treatment (SA+, SA-) as categorical factors. VarIdent variance structures were used on the SA treatments to accommodate deviations in the variance homogeneity in the SA and JA concentrations response variables (Zuur *et al.*, 2009). After this procedure, all the ANOVA assumptions were met.

Similarly, the effects of the SA treatment on the concentrations of alkaloids (total lolines, NFL, and NANL) were analysed separately with lineal effect models, using the same software package mentioned earlier, and assuming independent, identically distributed normal random errors. The models included the SA treatment (SA+, SA-) as a categorical factor. When required, we used the function VarIdent on the SA treatment to accommodate deviations in the variance homogeneity (Zuur *et al.*, 2009). ANOVA assumptions were then met.

The effects of plant symbiotic status and SA treatment on the population size of aphids (number of individuals) were analysed with linear mixed-effects models using the package glmmADMB in R software, and assuming that random errors were distributed independently and following a negative binomial distribution (Fournier *et al.*, 2012). The model included plant symbiotic status (E+, E-), SA treatment (SA+, SA-), and time (7d and 14d since the aphid challenge) as categorical factors, and the random effect included the time nested in pot. Temporal autocorrelation between the repeated measurements was not observed.

The effects of the plant symbiotic status and the SA treatment on mass-specific SMR of aphids were analysed with linear effects models using the same R package and assuming the same random error distribution that for hormone concentration variables. The model included the plant symbiotic status (E+, E-) and SA treatment (SA+, SA-) as categorical factors. All the ANOVA assumptions were met. We performed post-hoc analyses between treatments when significant interactions were detected using the package *lsmeans* in R (Lenth, 2016). All the presented values in the result section are means \pm standard errors (S.E.M). All data obtained in the present study are available in Table S1.

RESULTS

Effects of plant endophyte presence and SA on hormone levels and plant growth

The concentration of the hormones SA and JA within the plants responded differentially to the endophyte presence and the exogenous application of the SA. The plant SA concentration was independently affected by the plant endophyte status and the hormonal treatment (Table 1). The presence of the endophyte in the plant reduced the SA by *ca.* 10% [E-: 1114.00 ± 264.50 , and E+: 1014.00 ± 248.00 (ng SA g⁻¹ DW)]. In addition, both plant biotypes (i.e., endophyte-symbiotic and non-symbiotic) showed increased SA concentrations of about 19-fold 3 days after the exogenous application of the hormone (Fig. 1A). Although the interaction effect between the treatments was not significant (Table 1), the reduction in SA concentration due to the endophyte presence was much more evident in plants not exposed to SA (the mean SA concentration difference between E+ and E- plants in the SA- treatment was around 22%) (Fig. 1A). The concentration of JA

Table 1 Effects of plant symbiotic status (E+, E−) and the exposure to the hormone salicylic acid (SA+, SA−) on different response variables of *Lolium multiflorum* plants symbiotic with the endophyte fungus *Epichloë occultans*. Note that for lolines, only the effect of SA is evaluated since endophyte-free plants do not contain these alkaloids. NFL and NANL mean N-formyllooline and N-acetylnorlooline alkaloids, respectively. Statistically significant effects are highlighted in bold. Mean values, S.E.M, and post hoc statistical differences are shown in Figures 1 and 2.

Response variable	Treatment	df	F	P-value	
Salicylic acid (ng g ⁻¹ DW) (n = 8)	Symbiosis	1,28	4.70	0.038	
	SA	1,28	316.87	<0.001	
	Symbiosis × SA	1,28	0.45	0.506	
Jasmonic acid (ng g ⁻¹ DW) (n = 8)	Symbiosis	1,28	2.57	0.120	
	SA	1,28	0.49	0.487	
	Symbiosis × SA	1,28	0.25	0.614	
Above-ground plant biomass (g) (n = 14)	Symbiosis	1,108	4.41	0.038	
	SA	1,108	1.08	0.300	
	Symbiosis × SA	1,108	3.86	0.052	
Lolines (μg g ⁻¹ DW) (n = 8)	Total	SA	1,14	0.25	0.626
	NFL	SA	1,14	0.36	0.555
	NANL	SA	1,14	0.06	0.809

was not significantly modified by either the plant endophyte status or by the SA treatment (Fig. 1B) (Table 1).

The above-ground plant tissues at the end of the aphid challenge (day 27 from the application of SA) was 11% higher in endophyte-symbiotic than in non-symbiotic plants (E-: 4.42 ± 0.28 g, and E+: 4.99 ± 0.39 g). This effect was, however, not affected by the treatment with SA (Table 1).

Effects of SA on fungal loline concentrations

Ten days after plant exposure to the SA hormone, the concentration of loline alkaloids (total and the derivatives NFL and NANL) in endophyte-symbiotic plants did not vary among SA-treated and SA-untreated plants (Fig. 2) (Table 1).

Effects of plant endophyte presence and SA on *S. maydis* populations

The aphid population size on *L. multiflorum* plants increased over time but this increase was independent of both the endophyte presence or the plant exposure to salicylic acid (Table 2 and Fig. 3). On average, the aphid population size increased 2.60 fold in 7 days (from days 10 to 17 since plants exposure to SA, 5.96 ± 0.63 and 15.91 ± 1.74, respectively) (Fig. 3).




Figure 1 Physiological concentration of salicylic acid (panel A) and jasmonic acid (panel B) of *Lolium multiflorum* plants symbiotic with the endophyte fungus *Epichloë occultans*. Concentrations were measured three days after the salicylic acid application [treated: SA+ (shaded bars), and untreated: SA- (unshaded bars)] on *L. multiflorum* plants with (E+) and without (E-) the endophyte fungus. Different letters indicate significant differences at $P < 0.05$. Bars represent mean values \pm S.E.M. ($n = 8$).

Full-size DOI: [10.7717/peerj.8257/fig-1](https://doi.org/10.7717/peerj.8257/fig-1)

Effects of plant endophyte presence and SA on *S. maydis* metabolic rate

The aphid mass-specific SMR, evaluated at day 24 since the insects were placed on the plants, was unaffected by the endophytic fungus, the SA hormone, or the interaction between them (Table 2).

DISCUSSION

Since *Epichloë* fungi produce anti-herbivore alkaloids (Schardl et al., 2013a; Schardl et al., 2013b), we expected that endophytes would provide protection to host plants against the




Figure 2 Concentrations of loline alkaloids produced by the fungal endophyte *Epichloë occulta* in *Lolium multiflorum* plants. Loline alkaloids were measured ten days after plants were exposed to salicylic acid (treated: SA+, untreated: SA-). Total lolines (black bars) are the sum of N-formylloline (NFL, grey bars) and N-acetyl norloline (NANL, white bars) derivatives. Non-symbiotic plants do not produce loline alkaloids. Each loline compound was analysed separately (see ‘Material and Method’ section). n.s. means non-significant differences between treatments. The bars represent mean values \pm S.E.M. ($n = 8$).

Full-size [DOI: 10.7717/peerj.8257/fig-2](https://doi.org/10.7717/peerj.8257/fig-2)

Table 2 Effects of plant symbiotic status (E+, E-) and the exposure salicylic acid (SA+, SA-) on aphids number and standard metabolic rate of *Sipha maydis* aphids grown on *Lolium multiflorum* plants with the endophyte fungus *Epichloë occulta*. Aphids number were measured at days 10 and 17 post salicylic acid application. Specific statistical differences for ‘aphids number’ response variable are shown in Figure 3. The volume of CO₂ produced by aphids is abbreviated as ‘VCO₂’. No significant differences were observed in standard metabolic rate (SMR) values. Replicate numbers are indicated in parenthesis. Values are mean \pm S.E.M.

Response variable	Treatment	df	χ^2 or F	P-value	SA-		SA+	
					E-	E+	E-	E+
Aphids number ($n = 14$)								
	Symbiosis	1,52	2.30	0.127	-	-	-	-
	SA	1,52	0.01	0.922	-	-	-	-
	Time	1,52	7.26	0.007	-	-	-	-
	Symbiosis \times SA	1,52	0.32	0.567	-	-	-	-
	Symbiosis \times Time	1,52	0.76	0.382	-	-	-	-
	SA \times Time	1,52	1.52	0.217	-	-	-	-
	Symbiosis \times SA \times Time	1,52	1.37	0.241	-	-	-	-
Aphid mass-specific SMR ($\mu\text{L VCO}_2 \text{ h}^{-1} \text{ mg}^{-1}$) ($n = 5$)								
	Symbiosis	1,16	0.28	0.601	5.53	5.13	4.98	4.38
	SA	1,16	0.52	0.479	\pm	\pm	\pm	\pm
	Symbiosis \times SA	1,16	0.07	0.794	0.91	0.56	0.89	0.59

aphid *S. maydis*. However, we found that neither populations nor individuals of this aphid species were affected by the endophyte presence in plants. Despite the fact that aphids are usually controlled by the SA-dependent defence pathway (Thaler, Humphrey & Whiteman,




Figure 3 Population sizes of *Siphya maydis* aphids grown on *Lolium multiflorum* plants exposed to the salicylic acid hormone and symbiotic with the endophyte fungus *Epichloë occultans*. Aphids number were measured at days 10 and 17 post salicylic acid application [treated: SA+ (shaded bars), and untreated: SA- (unshaded bars)] on *L. multiflorum* plants with (E+) and without (E-) the endophyte fungus. Different letters indicate significant differences at $P < 0.05$. The bars represent mean values \pm S.E.M. ($n = 14$).

Full-size DOI: 10.7717/peerj.8257/fig-3

2012), here *S. maydis* aphids were not sensitive to the plant hormone exposure. Consistent with previous reports involving the plant interaction with beneficial microorganisms, the concentration of SA was lower in presence of the *Epichloë* endophyte fungus (Bastías et al., 2018a; Bastías et al., 2018b). However, the concentrations of alkaloids produced by endophytes was not affected by the plant exposure to SA.

The protection that each endophyte alkaloid type confers to host plants depends on, among other factors, the herbivore species (Bastías et al., 2017a; Bastías et al., 2017b). For example, loline fungal alkaloids confer effective protection against *Rhopalosiphum padi* aphids (Wilkinson et al., 2000), but the same alkaloids were ineffective in controlling *Heteronychus arator* beetles (Ball, Miles & Prestidge, 1997). In the present study, the presence of *E. occultans*, a loline producing endophyte fungus, did not confer protection to host *L. multiflorum* plants against *S. maydis* aphids. Similar findings were obtained in previous experiments studying the growth of *S. maydis* populations on the same plant-endophyte species system under field and laboratory conditions (Chaneton & Omacini, 2007; Miranda, Marina & Chaneton, 2011). In meadow fescue grass (*Festuca pratensis*) the presence of *E. uncinatum*, well-known to produce high concentrations of loline alkaloids, did not affect the population growth of *S. maydis* (Sabzalian, Hatami & Mirlohi, 2004). In that same study, however, the endophyte *E. coenophiala* (formerly *Neotyphodium coenophialum*) in tall fescue (*Schedonorus arundinacea*; formerly *F. arundinacea*) did effectively control *S. maydis* populations (Sabzalian, Hatami & Mirlohi, 2004). The

difference between these two grass-endophyte symbioses, in terms of alkaloid profiles, is that while *E. uncinatum* only produces lolines, *E. coenophiala* synthesises ergopeptine and peramine alkaloids (Schardl et al., 2013a; Schardl et al., 2013b). Peramine and ergopeptine alkaloids produced by *Epichloë* endophytes are known for producing bioactivity against insects (Rowan, Hunt & Gaynor, 1986; Fleetwood et al., 2008). Thus, it is possible that the inefficiency of *E. occultans* endophytes in controlling *S. maydis* aphids in *L. multiflorum* is due, in part, to the particular profile of alkaloids produced by this endophyte species (Siegel et al., 1990).

The SA-dependent defence pathway is usually involved in plant responses to aphid attacks (Thaler, Humphrey & Whiteman, 2012; Ballaré, 2014). In the present study, the aphid *S. maydis* was not affected by the induction of the SA pathway (triggered by the plant exposure to the hormone). This insect insensitivity to plant SA-dependent defences has been reported for other aphids species (Bastías et al., 2018a; Bastías et al., 2018b; Onkokesung et al., 2016; Selig et al., 2016). The *S. maydis* insensitivity to SA-dependent defences could be explained by the potential capability of this insect species in detoxify *L. multiflorum* anti-herbivore metabolites. Detoxification of plant toxins is performed by enzymes that can deactivate or neutralize these metabolites (Després, David & Gallet, 2007). The synthesis of these detoxification enzymes is however, generally costly for insects, and these costs can be captured by metabolic measurements (e.g., Castañeda et al., 2010). In our study, however, *S. maydis* aphids grown on SA-treated *L. multiflorum* plants did not show any changes in their standard metabolic rates. To our knowledge, efficient mechanisms of detoxification of plant's toxins have not been described for *S. maydis* aphids, which would be consistent with our findings for SMR. Adjustments in the feeding behaviour is another strategy that *S. maydis* aphids could have used to cope with plant defences (Walling, 2008). For instance, the aphid *Sitobion avenae* can reduce the time spent sucking phloem when feeding on SA-treated *Triticum aestivum* plants (Cao, Wang & Liu, 2014). In addition, it is also possible that an effective defence against *S. maydis* aphids in *L. multiflorum* requires more complex responses than just SA induction. For example, it has been documented that defence responses to aphids in *Arabidopsis thaliana*, *Glycine max*, and *Sorghum bicolor* plant species involves several hormone pathways that are sequentially induced during attacks by aphids (e.g., JA, SA, ethylene) (Moran & Thompson, 2001; Zhu-Salzman et al., 2004; De Vos et al., 2005; Li et al., 2008), and more recently that non-hormonal pathways can also be involved in these plant responses (i.e., methyl-D-erythritol-4-phosphate pathway) (Onkokesung et al., 2019).

Despite the fact that the fungal endophyte did not increase the resistance level of host plants against *S. maydis* aphids, endophyte-symbiotic plants may have been more tolerant to the aphid feeding due to the higher biomass they produced compared to their non-symbiotic counterparts. This growth promotion of *L. multiflorum* plants in presence of fungal endophytes has also been documented in other studies (Vila-Aiub, Gundel & Ghersa, 2005; Ueno et al., 2015; Bastías et al., 2018b). Since we had no aphid-free treatments, we cannot discard the possibility that the growth promotion documented in the present study had been a plant response to the aphid feeding more than a response to the endophyte presence (or a combination of both). However, findings from other studies suggest that

this growth promotion might be a plant response to the endophyte presence. For example, another study using a similar experimental set-up to that used in the present work showed that endophyte-symbiotic *L. multiflorum* plants had a higher biomass than endophyte-free plants, and that this growth enhance was independent of the herbivory by *R. padi* aphids (Ueno et al., 2015).

It has been proposed that beneficial plant symbionts may regulate the SA pathway of the host to facilitate their own growth within plant tissues (Poza & Azcón-Aguilar, 2007; Bastías et al., 2017a; Bastías et al., 2017b). This hypothesis has emerged from studies showing that the SA pathway can regulate the growth of these symbionts (Khaosaad et al., 2007; López-Ráez et al., 2010). In support of this, we observed that the plant SA concentration was reduced in presence of *Epichloë* fungal endophytes (see also Bastías et al., 2018a). In addition to regulating the growth of symbionts within plant tissues, the SA pathway can also modulate the functioning of beneficial symbiotic microorganisms (Hayat et al., 2010; Bastías et al., 2018a; De Vries et al., 2018). For example, the nitrogen fixation gene *NifE* in the beneficial cyanobiont, *Nostoc azollae*, was downregulated when *Azolla filiculoides* host plants were treated exogenously with methyl salicylate (De Vries et al., 2018). Based on these previous studies, we expected that symbiotic plants exposed to SA would show reduced levels of fungal alkaloids compared to untreated plants. Nevertheless, 10-days after the plants were exposed to the hormone, the concentration of loline alkaloids was similar between SA-treated and SA-untreated symbiotic plants. This result does not support our previous work which showed that alkaloid concentrations of endophyte-symbiotic plants were indeed reduced by SA treatment (Bastías et al., 2018a). It may be the case that in the present study, after an initial drop caused by the plant exposure to SA, fungal alkaloids were able to recover to their pre-treatment concentrations. Rapid increases in fungal alkaloids concentrations have been previously reported in plants symbiotic with *Epichloë* fungal endophytes. For example, we previously found that alkaloid concentrations took less than 7-days to respond to aphid herbivory (Bastías et al., 2018a).

CONCLUSIONS

The present study indicates that the aphid *Sipha maydis* is insensitive to the anti-herbivore defences of *L. multiflorum* in symbiosis with *Epichloë occultans*. Our results indicate that neither the presence of *E. occultans* endophytes nor the induction of plant SA pathway regulate *S. maydis* populations. However, endophyte-symbiotic plants may have been more tolerant to the aphid feeding because these plants produced more aboveground biomass. We suggest that this insect insensitivity could be explained by a combination between the ineffectiveness of loline alkaloids (produced by *E. occultans*) in controlling *S. maydis* aphids and the capacity of this herbivore to tolerate the hormone-dependent defences of *L. multiflorum*.

ACKNOWLEDGEMENTS

We gratefully acknowledge Dr. Armen Charchoglyan (University of Guelph) for assistance with GC-MS/MS analysis and Dr. Edmundo L. Ploschuk (Universidad de Buenos Aires) for assistance with metabolic measurements.

ADDITIONAL INFORMATION AND DECLARATIONS

Funding

Daniel A. Bastías was supported by a fellowship from the Agencia Nacional de Promoción Científica and CONICET, and from the Canadian Emerging Leaders in the Americas Program (ELAP). Funding to support this research was provided by the Agencia Nacional de Promoción Científica FONCYT (PICT-2355), Universidad de Buenos Aires UBA (UBACYT 2014-30BA) to M. Alejandra Martínez-Ghersa, and Canadian Natural Science and Engineering Research Council to Jonathan A. Newman. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Grant Disclosures

The following grant information was disclosed by the authors:
CONICET.

Canadian Emerging Leaders in the Americas Program (ELAP).

Agencia Nacional de Promoción Científica FONCYT: PICT-2355.

Universidad de Buenos Aires UBA: UBACYT 2014-30BA.

Canadian Natural Science and Engineering Research Council.

Competing Interests

Daniel A. Bastías, Stuart D. Card and Wade J. Mace are employed by Forage Science, AgResearch Limited.

Author Contributions

- Daniel A. Bastías conceived and designed the experiments, performed the experiments, analyzed the data, prepared figures and/or tables, authored or reviewed drafts of the paper, approved the final draft.
- Maria Alejandra Martínez-Ghersa conceived and designed the experiments, contributed reagents/materials/analysis tools, authored or reviewed drafts of the paper, approved the final draft.
- Jonathan A. Newman conceived and designed the experiments, contributed reagents/materials/analysis tools, prepared figures and/or tables, authored or reviewed drafts of the paper, approved the final draft.
- Stuart D. Card and Wade J. Mace performed the experiments, contributed reagents/materials/analysis tools, authored or reviewed drafts of the paper, approved the final draft.
- Pedro E. Gundel conceived and designed the experiments, prepared figures and/or tables, authored or reviewed drafts of the paper, approved the final draft.

Data Availability

The following information was supplied regarding data availability: The raw data are available on figshare: Bastías, Daniel A; Martínez-Ghersa, Maria Alejandra; Newman, Jonathan A; Card, Stuart D.; Mace, Wade J.; Gundel, Pedro E. (2019): Table S1_Data.xlsx. figshare. Dataset. <https://doi.org/10.6084/m9.figshare.9345524.v1>.

Supplemental Information

Supplemental information for this article can be found online at <http://dx.doi.org/10.7717/peerj.8257#supplemental-information>.

REFERENCES

- Bacon C, White J. 1994.** Stains, media and procedures for analyzing endophytes. In: Bacon C, White J, eds. *Biotechnology of endophytic fungi of grasses*. Boca Raton: CRC Press.
- Ball OJP, Miles CO, Prestidge RA. 1997.** Ergopeptine alkaloids and *Neotyphodium lolii*-mediated resistance in perennial ryegrass against adult *Heteronychus arator* (Coleoptera: Scarabaeidae). *Journal of Economic Entomology* **90**:1382–1391 DOI [10.1093/jee/90.5.1382](https://doi.org/10.1093/jee/90.5.1382).
- Ballaré CL. 2014.** Light regulation of plant defense. *Annual Review of Plant Biology* **65**:335–363 DOI [10.1146/annurev-arplant-050213-040145](https://doi.org/10.1146/annurev-arplant-050213-040145).
- Bastias DA, Martínez-Ghersa MA, Ballaré CL, Gundel PE. 2017a.** *Epichloë* fungal endophytes and plant defenses: not just alkaloids. *Trends in Plant Science* **22**:939–948 DOI [10.1016/j.tplants.2017.08.005](https://doi.org/10.1016/j.tplants.2017.08.005).
- Bastías DA, Martínez-Ghersa MA, Newman JA, Card SD, Mace WJ, Gundel PE. 2018a.** The plant hormone salicylic acid interacts with the mechanism of anti-herbivory conferred by fungal endophytes in grasses. *Plant, Cell & Environment* **41**:395–405 DOI [10.1111/pce.13102](https://doi.org/10.1111/pce.13102).
- Bastías DA, Martínez-Ghersa MA, Newman JA, Card SD, Mace WJ, Gundel PE. 2018b.** Jasmonic acid regulation of the anti-herbivory mechanism conferred by fungal endophytes in grasses. *Journal of Ecology* **106**:2365–2379 DOI [10.1111/1365-2745.12990](https://doi.org/10.1111/1365-2745.12990).
- Bastías DA, Ueno AC, Machado Assefh CR, Alvarez AE, Young CA, Gundel PE. 2017b.** Metabolism or behavior: explaining the performance of aphids on alkaloid-producing fungal endophytes in annual ryegrass (*Lolium multiflorum*). *Oecologia* **185**:245–256 DOI [10.1007/s00442-017-3940-2](https://doi.org/10.1007/s00442-017-3940-2).
- Bedini A, Mercy L, Schneider C, Franken P, Lucic-Mercy E. 2018.** Unraveling the initial plant hormone signaling, metabolic mechanisms and plant defense triggering the endomycorrhizal symbiosis behavior. *Frontiers in Plant Science* **9**:1800–1800 DOI [10.3389/fpls.2018.01800](https://doi.org/10.3389/fpls.2018.01800).
- Bultman TL, McNeill MR, Krueger K, De Nicolo G, Popay AJ, Hume DE, Mace WJ, Fletcher LR, Koh YM, Sullivan TJ. 2018.** Complex interactions among sheep, insects, grass, and fungi in a simple New Zealand grazing system. *Journal of Chemical Ecology* **44**:957–964 DOI [10.1007/s10886-018-0993-6](https://doi.org/10.1007/s10886-018-0993-6).

- Cahill Jr JF, Castelli JP, Casper BB. 2002.** Separate effects of human visitation and touch on plant growth and herbivory in an old-field community. *American Journal of Botany* **89**:1401–1409 DOI [10.3732/ajb.89.9.1401](https://doi.org/10.3732/ajb.89.9.1401).
- Cao HH, Wang SH, Liu TX. 2014.** Jasmonate- and salicylate-induced defenses in wheat affect host preference and probing behavior but not performance of the grain aphid, *Sitobion avenae*. *Insect Science* **21**:47–55 DOI [10.1111/1744-7917.12023](https://doi.org/10.1111/1744-7917.12023).
- Cao Y, Halane MK, Gassmann W, Stacey G. 2017.** The role of plant innate immunity in the legume-*Rhizobium* symbiosis. *Annual Review of Plant Biology* **68**:535–561 DOI [10.1146/annurev-arplant-042916-041030](https://doi.org/10.1146/annurev-arplant-042916-041030).
- Card SD, Rolston MP, Park Z, Cox N, Hume DE. 2011.** Fungal endophyte detection in pasture grass seed utilising the infection layer and comparison to other detection techniques. *Seed Science and Technology* **39**:581–592 DOI [10.15258/sst.2011.39.3.05](https://doi.org/10.15258/sst.2011.39.3.05).
- Castañeda LE, Figueroa CC, Fuentes-Contreras E, Niemeyer HM, Nespolo RF. 2010.** Physiological approach to explain the ecological success of ‘superclones’ in aphids: interplay between detoxification enzymes, metabolism and fitness. *Journal of Insect Physiology* **56**:1058–1064 DOI [10.1016/j.jinsphys.2010.02.019](https://doi.org/10.1016/j.jinsphys.2010.02.019).
- Castañeda LE, Figueroa CC, Nespolo RF. 2010.** Do insect pests perform better on highly defended plants? Costs and benefits of induced detoxification defences in the aphid *Sitobion avenae*. *Journal of Evolutionary Biology* **23**:2474–2483 DOI [10.1111/j.1420-9101.2010.02112.x](https://doi.org/10.1111/j.1420-9101.2010.02112.x).
- Chaneton EJ, Omacini M. 2007.** Bottom-up cascades induced by fungal endophytes in multitrophic systems. In: Price PW, Ohgushi T, Craig TP, eds. *Ecological communities: plant mediation in indirect interaction webs*. Cambridge: Cambridge University Press, 164–187 DOI [10.1017/CBO9780511542701.009](https://doi.org/10.1017/CBO9780511542701.009).
- Clay K. 1988.** Fungal endophytes of grasses: a defensive mutualism between plants and fungi. *Ecology* **69**:10–16 DOI [10.2307/1943155](https://doi.org/10.2307/1943155).
- Corrales CE, Castro AM, Ricci M, Dixon AFG. 2007.** *Sipha maydis*: distribution and host range of a new aphid pest of winter cereals in Argentina. *Journal of Economic Entomology* **100**:1781–1788 DOI [10.1093/jee/100.6.1781](https://doi.org/10.1093/jee/100.6.1781).
- De Vos M, Van Oosten VR, Van Poecke RMP, Van Pelt JA, Pozo MJ, Mueller MJ, Buchala AJ, Métraux J-P, Van Loon LC, Dicke M, Pieterse CMJ. 2005.** Signal signature and transcriptome changes of *Arabidopsis* during pathogen and insect attack. *Molecular Plant-Microbe Interactions* **18**:923–937 DOI [10.1094/MPMI-18-0923](https://doi.org/10.1094/MPMI-18-0923).
- De Vries S, De Vries J, Teschke H, Von Dahlen JK, Rose LE, Gould SB. 2018.** Jasmonic and salicylic acid response in the fern *Azolla filiculoides* and its cyanobiont. *Plant, Cell & Environment* **41**:2530–2548 DOI [10.1111/pce.13131](https://doi.org/10.1111/pce.13131).
- Després L, David J-P, Gallet C. 2007.** The evolutionary ecology of insect resistance to plant chemicals. *Trends in Ecology & Evolution* **22**:298–307 DOI [10.1016/j.tree.2007.02.010](https://doi.org/10.1016/j.tree.2007.02.010).
- Dupont P, Eaton CJ, Wargent JJ, Fechtner S, Solomon P, Schmid J, Day RC, Scott B, Cox MP. 2015.** Fungal endophyte infection of ryegrass reprograms host metabolism and alters development. *New Phytologist* **208**:1227–1240 DOI [10.1111/nph.13614](https://doi.org/10.1111/nph.13614).

- Eichenseer H, Dahlman DL, Bush LP. 1991.** Influence of endophyte infection, plant age and harvest interval on *Rhopalosiphum padi* survival and its relation to quantity of N-formyl and N-acetyl loline in tall fescue. *Entomologia Experimentalis et Applicata* **60**:29–38 DOI [10.1111/j.1570-7458.1991.tb01519.x](https://doi.org/10.1111/j.1570-7458.1991.tb01519.x).
- Fleetwood DJ, Scott B, Voisey CR, Johnson RD. 2008.** Insights into the molecular biology of *Epichloë* endophyte alkaloid biosynthesis. *Proceedings of the New Zealand Grassland Association* **70**:217–220.
- Fournier DA, Skaug HJ, Ancheta J, Ianelli J, Magnusson A, Maunder MN, Nielsen A, Sibert J. 2012.** AD model builder: using automatic differentiation for statistical inference of highly parameterized complex nonlinear models. *Optimization Methods and Software* **27**:233–249 DOI [10.1080/10556788.2011.597854](https://doi.org/10.1080/10556788.2011.597854).
- Fuchs B, Krischke M, Mueller MJ, Krauss J. 2017a.** Herbivore-specific induction of defence metabolites in a grass—endophyte association. *Functional Ecology* **31**:318–324 DOI [10.1111/1365-2435.12755](https://doi.org/10.1111/1365-2435.12755).
- Fuchs B, Krischke M, Mueller MJ, Krauss J. 2017b.** Plant age and seasonal timing determine endophyte growth and alkaloid biosynthesis. *Fungal Ecology* **29**:52–58 DOI [10.1016/j.funeco.2017.06.003](https://doi.org/10.1016/j.funeco.2017.06.003).
- Gundel PE, Garibaldi L, Tognetti P, Aragón R, Ghera C, Omacini M. 2009.** Imperfect vertical transmission of the endophyte *Neotyphodium* in exotic grasses in grasslands of the flooding pampa. *Microbial Ecology* **57**:740–748 DOI [10.1007/s00248-008-9447](https://doi.org/10.1007/s00248-008-9447).
- Gundel PE, Martínez-Ghera MA, Omacini M, Cuyeu R, Pagano E, Ríos R, Ghera CM. 2012.** Mutualism effectiveness and vertical transmission of symbiotic fungal endophytes in response to host genetic background. *Evolutionary Applications* **5**:838–849 DOI [10.1111/j.1752-4571.2012.00261.x](https://doi.org/10.1111/j.1752-4571.2012.00261.x).
- Gundel PE, Rudgers JA, Ghera CM. 2011.** Incorporating the process of vertical transmission into understanding of host–symbiont dynamics. *Oikos* **120**:1121–1128 DOI [10.1111/j.1600-0706.2011.19299.x](https://doi.org/10.1111/j.1600-0706.2011.19299.x).
- Gundel PE, Seal CE, Biganzoli F, Molina-Montenegro MA, Vázquez-de Aldana BR, Zabalgoeazcoa I, Bush LP, Martínez-Ghera MA, Ghera CM. 2018.** Occurrence of alkaloids in grass seeds symbiotic with vertically-transmitted *Epichloë* fungal endophytes and its relationship with antioxidants. *Frontiers in Ecology and Evolution* **6**:Article 211 DOI [10.3389/fevo.2018.00211](https://doi.org/10.3389/fevo.2018.00211).
- Hayat Q, Hayat S, Irfan M, Ahmad A. 2010.** Effect of exogenous salicylic acid under changing environment: a review. *Environmental and Experimental Botany* **68**:14–25 DOI [10.1016/j.envexpbot.2009.08.005](https://doi.org/10.1016/j.envexpbot.2009.08.005).
- Hunt MG, Rasmussen S, Newton PCD, Parsons AJ, Newman JA. 2005.** Near-term impacts of elevated CO₂, nitrogen and fungal endophyte-infection on *Lolium perenne* L. growth, chemical composition and alkaloid production. *Plant, Cell & Environment* **28**:1345–1354 DOI [10.1111/j.1365-3040.2005.01367.x](https://doi.org/10.1111/j.1365-3040.2005.01367.x).

- Johnson LJ, Johnson RD, Schardl CL, Panaccione DG. 2003.** Identification of differentially expressed genes in the mutualistic association of tall fescue with *Neotyphodium coenophialum*. *Physiological and Molecular Plant Pathology* **63**:305–317 DOI [10.1016/j.pmpp.2004.04.001](https://doi.org/10.1016/j.pmpp.2004.04.001).
- Johnson MC, Dahlman DL, Siegel MR, Bush LP, Latch GCM, Potter DA, Varney DR. 1985.** Insect feeding deterrents in endophyte-infected tall fescue. *Applied and Environmental Microbiology* **49**:568–571.
- Jung S, Martinez-Medina A, Lopez-Raez J, Pozo M. 2012.** Mycorrhiza-induced resistance and priming of plant defenses. *Journal of Chemical Ecology* **38**:651–664 DOI [10.1007/s10886-012-0134-6](https://doi.org/10.1007/s10886-012-0134-6).
- Justus M, Witte L, Hartmann T. 1997.** Levels and tissue distribution of loline alkaloids in endophyte-infected *Festuca pratensis*. *Phytochemistry* **44**:51–57 DOI [10.1016/S0031-9422\(96\)00535-3](https://doi.org/10.1016/S0031-9422(96)00535-3).
- Khaosaad T, García-Garrido JM, Steinkellner S, Vierheilig H. 2007.** Take-all disease is systemically reduced in roots of mycorrhizal barley plants. *Soil Biology and Biochemistry* **39**:727–734 DOI [10.1016/j.soilbio.2006.09.014](https://doi.org/10.1016/j.soilbio.2006.09.014).
- Lenth RV. 2016.** Least-squares means: the R package lsmeans. *Journal of Statistical Software* **69**:1–33 DOI [10.18637/jss.v069.i01](https://doi.org/10.18637/jss.v069.i01).
- Li Y, Zou J, Li M, Bilgin DD, Vodkin LO, Hartman GL, Clough SJ. 2008.** Soybean defense responses to the soybean aphid. *New Phytologist* **179**:185–195 DOI [10.1111/j.1469-8137.2008.02443.x](https://doi.org/10.1111/j.1469-8137.2008.02443.x).
- López-Ráez JA, Verhage A, Fernández I, García JM, Azcón-Aguilar C, Flors V, Pozo MJ. 2010.** Hormonal and transcriptional profiles highlight common and differential host responses to arbuscular mycorrhizal fungi and the regulation of the oxylipin pathway. *Journal of Experimental Botany* **61**:2589–2601 DOI [10.1093/jxb/erq089](https://doi.org/10.1093/jxb/erq089).
- Miranda MI, Omacini M, Chaneton EJ. 2011.** Environmental context of endophyte symbioses: interacting effects of water stress and insect herbivory. *International Journal of Plant Sciences* **172**:499–508 DOI [10.1086/658921](https://doi.org/10.1086/658921).
- Moon CD, Scott B, Schardl CL, Christensen MJ. 2000.** The evolutionary origins of *Epichloë* endophytes from annual ryegrasses. *Mycologia* **92**:1103–1118 DOI [10.2307/3761478](https://doi.org/10.2307/3761478).
- Moore JR, Pratley JE, Mace WJ, Weston LA. 2015.** Variation in alkaloid production from genetically diverse *Lolium* accessions infected with *Epichloë* species. *Journal of Agricultural and Food Chemistry* **63**:10355–10365 DOI [10.1021/acs.jafc.5b03089](https://doi.org/10.1021/acs.jafc.5b03089).
- Moran PJ, Thompson GA. 2001.** Molecular responses to aphid feeding in *Arabidopsis* in relation to plant defense pathways. *Plant Physiology* **125**:1074–1085 DOI [10.1104/pp.125.2.1074](https://doi.org/10.1104/pp.125.2.1074).
- Moreira X, Abdala-Roberts L, Castagneyrol B. 2018.** Interactions between plant defence signalling pathways: evidence from bioassays with insect herbivores and plant pathogens. *Journal of Ecology* **106**:2353–2364 DOI [10.1111/1365-2745.12987](https://doi.org/10.1111/1365-2745.12987).
- Navarro-Meléndez AL, Heil M. 2014.** Symptomless endophytic fungi suppress endogenous levels of salicylic acid and interact with the jasmonate-dependent indirect

- defense traits of their host, lima bean (*Phaseolus lunatus*). *Journal of Chemical Ecology* 40:816–825 DOI 10.1007/s10886-014-0477-2.
- Nespolo RF, Roff DA, Fairbairn DJ. 2008.** Energetic trade-off between maintenance costs and flight capacity in the sand cricket (*Gryllus firmus*). *Functional Ecology* 22:624–631 DOI 10.1111/j.1365-2435.2008.01394.x.
- Omacini M, Chaneton EJ, Ghersa CM, Muller CB. 2001.** Symbiotic fungal endophytes control insect host-parasite interaction webs. *Nature* 409:78–81 DOI 10.1038/35051070.
- Onkokesung N, Reichelt M, Van Doorn A, Schuurink RC, Dicke M. 2016.** Differential costs of two distinct resistance mechanisms induced by different herbivore species in *Arabidopsis*. *Plant Physiology* 170:891–906 DOI 10.1104/pp.15.01780.
- Onkokesung N, Reichelt M, Wright LP, Phillips MA, Gershenson J, Dicke M. 2019.** The plastidial metabolite 2-C-methyl-D-erythritol-2, 4-cyclodiphosphate modulates defense responses against aphids. *Plant, Cell & Environment* 42:2309–2323 DOI 10.1111/pce.13538.
- Panaccione DG, Beaulieu WT, Cook D. 2014.** Bioactive alkaloids in vertically transmitted fungal endophytes. *Functional Ecology* 28:299–314 DOI 10.1111/1365-2435.12076.
- Pinheiro J, Bates D, DebRoy S, Sarkar D. 2009.** nlme: linear and nonlinear mixed effects models. R package. Vienna: R Foundation for Statistical Computing.
- Pozo MJ, Azcón-Aguilar C. 2007.** Unraveling mycorrhiza-induced resistance. *Current Opinion in Plant Biology* 10:393–398 DOI 10.1016/j.pbi.2007.05.004.
- R Core Team. 2013.** R: a language and environment for statistical computing. Vienna: R Foundation for Statistical Computing.
- Ramos P, Rivas N, Pollmann S, Casati P, Molina-Montenegro MA. 2018.** Hormonal and physiological changes driven by fungal endophytes increase Antarctic plant performance under UV-B radiation. *Fungal Ecology* 34:76–82 DOI 10.1016/j.funeco.2018.05.006.
- Rasmussen S, Parsons AJ, Bassett S, Christensen MJ, Hume DE, Johnson LJ, Johnson RD, Simpson WR, Stacke C, Voisey CR, Xue H, Newman JA. 2007.** High nitrogen supply and carbohydrate content reduce fungal endophyte and alkaloid concentration in *Lolium perenne*. *New Phytologist* 173:787–797 DOI 10.1111/j.1469-8137.2006.01960.x.
- Rowan DD, Hunt MB, Gaynor DL. 1986.** Peramine, a novel insect feeding deterrent from ryegrass infected with the endophyte *Acremonium loliae*. *Journal of the Chemical Society, Chemical Communications* 142:935–936 DOI 10.1039/C39860000935.
- Ryan GD, Rasmussen S, Xue H, Parsons AJ, Newman JA. 2014.** Metabolite analysis of the effects of elevated CO₂ and nitrogen fertilization on the association between tall fescue (*Schedonorus arundinaceus*) and its fungal symbiont *Neotyphodium coenophialum*. *Plant, Cell & Environment* 37:204–212 DOI 10.1111/pce.12146.
- Sabzalian MR, Hatami B, Mirlohi A. 2004.** Mealybug, *Phenacoccus solani*, and barley aphid, *Sipha maydis*, response to endophyte-infected tall and meadow fescues. *Entomologia Experimentalis et Applicata* 113:205–209 DOI 10.1111/j.0013-8703.2004.00227.

- Saikkonen K, Gundel P, Helander M. 2013. Chemical ecology mediated by fungal endophytes in grasses. *Journal of Chemical Ecology* 39:962–968 DOI 10.1007/s10886-013-0310-3.
- Saikkonen K, Saari S, Helander M. 2010. Defensive mutualism between plants and endophytic fungi? *Fungal Diversity* 41:101–113 DOI 10.1007/s13225-010-0023-7.
- Schardl CL, Florea S, Pan J, Nagabhyru P, Bec S, Calie PJ. 2013a. The Epichloae: alkaloid diversity and roles in symbiosis with grasses. *Current Opinion in Plant Biology* 16:480–488 DOI 10.1016/j.pbi.2013.06.012.
- Schardl CL, Young C, Pan J, Florea S, Takach J, Panaccione D, Farman M, Webb J, Jaromczyk J, Charlton N, Nagabhyru P, Chen L, Shi C, Leuchtman A. 2013b. Currencies of mutualisms: sources of alkaloid genes in vertically transmitted Epichloae. *Toxins* 5:1064–1088 DOI 10.3390/toxins5061064.
- Selig P, Keough S, Nalam VJ, Nachappa P. 2016. Jasmonate-dependent plant defenses mediate soybean thrips and soybean aphid performance on soybean. *Arthropod-Plant Interactions* 10:273–282 DOI 10.1007/s11829-016-9437-9.
- Siegel MR, Latch GCM, Bush LP, Fannin FF, Rowan DD, Tapper BA, Bacon CW, Johnson MC. 1990. Fungal endophyte-infected grasses: alkaloid accumulation and aphid response. *Journal of Chemical Ecology* 16:3301–3315 DOI 10.1007/BF00982100.
- Skvarla MJ, Halbert SE, Footitt RG, Jensen AS, Maw E, Miller GL. 2017. An update to the adventive aphids (Hemiptera: Aphidoidea) of America north of Mexico, with notes on intercepted species. *Proceedings of the Entomological Society of Washington* 119:90–112 DOI 10.4289/0013-8797.119.1.90.
- Stacey G, McAlvin CB, Kim S-Y, Olivares J, Soto MJ. 2006. Effects of endogenous salicylic acid on nodulation in the model legumes *Lotus japonicus* and *Medicago truncatula*. *Plant Physiology* 141:1473–1481 DOI 10.1104/pp.106.080986.
- Sugawara K, Inoue T, Yamashita M, Ohkubo H. 2006. Distribution of the endophytic fungus, *Neotyphodium occultans* in naturalized Italian ryegrass in western Japan and its production of bioactive alkaloids known to repel insect pests. *Grassland Science* 52:147–154 DOI 10.1111/j.1744-697X.2006.00060.x.
- Thaler JS, Humphrey PT, Whiteman NK. 2012. Evolution of jasmonate and salicylate signal crosstalk. *Trends in Plant Science* 17:260–270 DOI 10.1016/j.tplants.2012.02.010.
- Uchitel A, Omacini M, Chaneton EJ. 2011. Inherited fungal symbionts enhance establishment of an invasive annual grass across successional habitats. *Oecologia* 165:465–475 DOI 10.1007/s00442-010-1740-z.
- Ueno AC, Gundel PE, Omacini M, Ghera CM, Bush LP, Martínez-Ghera MA. 2015. Mutualism effectiveness of a fungal endophyte in an annual grass is impaired by ozone. *Functional Ecology* 30:226–232 DOI 10.1111/1365-2435.12519.
- Vila-Aiub MM, Gundel PE, Ghera CM. 2005. Fungal endophyte infection changes growth attributes in *Lolium multiflorum* Lam. *Austral Ecology* 30:49–57 DOI 10.1111/j.1442-9993.2005.01423.x.
- Walling LL. 2008. Avoiding effective defenses: strategies employed by phloem-feeding insects. *Plant Physiology* 146:859–866 DOI 10.1104/pp.107.113142.

- Wilkinson HH, Siegel MR, Blankenship JD, Mallory AC, Bush LP, Schardl CL. 2000.** Contribution of fungal loline alkaloids to protection from aphids in a grass-endophyte mutualism. *Molecular Plant-Microbe Interactions* **13**:1027–1033 DOI [10.1094/MPMI.2000.13.10.1027](https://doi.org/10.1094/MPMI.2000.13.10.1027).
- Zhu-Salzman K, Salzman RA, Ahn J-E, Koiwa H. 2004.** Transcriptional regulation of Sorghum defense determinants against a phloem-feeding aphid. *Plant Physiology* **134**:420–431 DOI [10.1104/pp.103.028324](https://doi.org/10.1104/pp.103.028324).
- Zuur AF, Ieno EN, Walker NJ, Saveliev AA, Smit GM. 2009.** *Mixed effects models and extensions in ecology with R*. New York: Springer Sciences.