







**ORIGINAL RESEARCH:  
EMPIRICAL RESEARCH - QUANTITATIVE**

# Clusters of competence: Relationship between self-reported professional competence and achievement on a national examination among graduating nursing students

Henrietta Forsman RN, PhD, Assistant professor<sup>1</sup>  | Inger Jansson RN, PhD, Senior lecturer<sup>2</sup>  | Janeth Leksell RN, PhD, Associate Professor<sup>1,3</sup>  | Margret Lepp RN, PhD, Professor<sup>2,4,5</sup>  | Christina Sundin Andersson RNT, MSc, Lecturer<sup>6</sup> | Maria Engström RN, PhD, Professor<sup>7,8,9</sup>  | Jan Nilsson RN, PhD, Associate Professor<sup>6,10,11</sup> 

<sup>1</sup>School of Education, Health and Social Studies, Dalarna University, Falun, Sweden

<sup>2</sup>Institute of Health and Care Sciences, Sahlgrenska Academy, University of Gothenburg, Gothenburg, Sweden

<sup>3</sup>Department of Medical Sciences, Clinical Diabetology and Metabolism, Uppsala University, Uppsala, Sweden

<sup>4</sup>Østfold University College, Halden, Norway

<sup>5</sup>School of Nursing and Midwifery, Griffith University, Brisbane, Australia

<sup>6</sup>Department of Health Science, Nursing, Faculty of Health, Science and Technology, Karlstad University, Karlstad, Sweden

<sup>7</sup>Faculty of Health and Occupational Studies, University of Gävle, Gävle, Sweden

<sup>8</sup>Nursing Department, Medicine and Health College, Lishui University, Lishui, China

<sup>9</sup>Department of Public Health and Caring Sciences, Uppsala University, Uppsala, Sweden

<sup>10</sup>Department of Health Promotion Sciences, Sophiahemmet University, Stockholm, Sweden

<sup>11</sup>Japanese Red Cross Institute for Humanitarian Studies, Tokyo, Japan

## Correspondence

Henrietta Forsman, School of Education, Health and Social Studies, Dalarna University, Falun, Sweden.  
Email: hfo@du.se

## Funding information

The study was supported by research allocations from the authors' universities and a scholarship from the The Swedish Society of Nursing named 'The Alice Lindström Scholarship'. The funding sources had no involvement in study design; in the collection, analysis, and interpretation of data; in the writing of the report; or in the decision to submit the article for publication.

## Abstract

**Aims:** To identify clusters based on graduating nursing students' self-reported professional competence and their achievement on a national examination. Furthermore, to describe and compare the identified clusters regarding sample characteristics, students' perceptions of overall quality of the nursing programme, and students' general self-efficacy (GSE).

**Design:** A cross-sectional study combining survey data and results from a national examination.

**Methods:** Data were collected at two universities and one university college in Sweden in January 2017, including 179 students in the final term of the nursing programme. The study was based on the Nurse Professional Competence Scale, the GSE scale, and results from the National Clinical Final Examination. A two-step cluster analysis was used to identify competence profiles, followed by comparative analyses between clusters.

**Results:** Three clusters were identified illustrating students' different competence profiles. Students in Clusters 1 and 2 passed the examination, but differed in their

The peer review history for this article is available at <https://publons.com/publon/10.1111/jan.14222>

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2019 The Authors. *Journal of Advanced Nursing* published by John Wiley & Sons Ltd.

self-assessments of competence, rating themselves under and above the overall median value, respectively. Students in Cluster 3 failed the examination but rated themselves at the overall median level or higher.

**Conclusion:** The study illustrates how nursing students' self-assessed competence might differ from competency assessed by examination, which is challenging for nursing education. Self-evaluation is a key learning outcome and is, in the long run, essential to patient safety.

**Impact:** The study has identified clusters of students where some overestimate and others underestimate their competence. Students who assessed their competence low but passed the exam assessed their GSE lower than other students. The findings illuminate the need for student-centred strategies in nursing education, including elements of self-assessment in relation to examination to make the students more aware of their clinical competence.

#### KEYWORDS

cluster analysis, nursing education, nursing students, professional competence, questionnaires, self-assessment, survey

## 1 | INTRODUCTION

To contribute to a global future workforce of nurses with professional competence in providing safe and high-quality care, it is important to measure and evaluate nursing students' progress and achievements towards educational goals and requirements. Structured methods including different evaluation tools are common in evaluations of nursing students' clinical competence (Lejonqvist, Eriksson, & Meretoja, 2016). However, validity and reliability in assessments of competence may vary and pre-existing expectations from mentors, in addition to their shared understanding of educational goals, are present in this assessment process (Burden, Topping, & O'Halloran, 2018). The ability to identify one's own knowledge gaps and need for competence development is a competence that is clearly formulated in the educational goals for Swedish nursing programmes (Higher Education Ordinance (1993:100)). Self-evaluations can be operationalized in terms of, for example, perceived knowledge or performance/skill evaluation and the overall correspondence between self-evaluated ability and objective performance outcomes is moderate (Zell & Krizan, 2014). Previous studies have shown how assessments of students' achievements might differ between students' self-assessments on the one hand and assessments by preceptors (Ugland Vae, Engström, Mårtensson, & Löfmark, 2018), examiners (Baxter & Norman, 2011), and structured clinical examinations (Sears et al., 2014) on the other hand.

### 1.1 | Background

Quality and safety competencies for nurses have been defined by the Quality and Safety Education for Nurses (QSEN) and the National Advisory Board (Cronenwett et al., 2007) and are based on core

professional competencies described by the Institute of Medicine (2003). QSEN has proposed targets for knowledge, skills, and attitudes to be developed for each of the following competencies: patient-centered care, teamwork and collaboration, evidence-based practice, quality improvement, safety, and informatics. These six competencies can serve as guidelines in the curricular development of formal academic programmes, transition to practice, and continued study programmes (Cronenwett et al., 2007). The Swedish Society of Nursing (2017) has adopted these six core competencies, along with leadership and education, in a description of competence for Swedish nurses. Research shows the association between nurses' competence and patient outcomes (Kendall-Gallagher, Aiken, Sloane, & Cimiotti, 2011) and that academic competence (Aiken et al., 2014) and a higher proportion of professional nurses (Aiken et al., 2017) are associated with better outcomes for patients. A recently published report by The Swedish National Board of Health and Welfare (2018) highlights how a lack of competence among healthcare staff significantly increases the risk of patient injuries, due to patients being exposed to danger. However, the concept of competence lacks a clear, coherent, and comprehensive definition or description (Kajander-Unkuri, Salminen, Saarikoski, Suhonen, & Leino-Kilpi, 2013; Liu & Aunguroch, 2018; Nehrir, Vanaki, Mokthari Nouri, Khademolhosseini, & Ebadi, 2016).

Nursing programmes in Sweden consist of 3 years of full-time studies (180 ECTS credits) with both theoretical and clinical studies included. Nursing programmes are offered at 25 Higher Education Institutions (HEI) and graduating students are awarded with a professional nursing degree and a Bachelor of Science degree. The objectives for Nursing Education are stipulated in the Swedish Higher Education Act (1992:1434) and the Swedish Higher Education Ordinance (1993:100) and are operationalized by each respective HEI.

The Miller's pyramid for clinical competence (Miller, 1990) provides a framework for assessing students' depth of knowledge and performance at four levels; *Knows* refers to the student having sufficient knowledge of the field they will be working in and *Knows how* is when the student can apply this knowledge. These steps can be tested with a written examination. *Shows how* is when the student can demonstrate how a procedure is performed in an artificial environment. *Does* is when students can perform tasks professionally in an authentic context. Levels 3 and 4 can be tested through clinical exams and direct observation in clinical settings.

In Sweden, work on introducing a voluntary national clinical examination started in 2003 (Athlin, Larsson, & Söderhamn, 2012) and now, 14 of the 25 HEIs with nursing programmes conduct The National Clinical Final Examination (NCFE). The aim of the NCFE is to examine third-year nursing students' clinical competence to ensure that they have the clinical knowledge and skills required, as laid out in the national legislation, before they graduate the programme and enter working life. The NCFE has a written and a bedside test and is structured so all the steps in Miller's pyramid (Miller, 1990) can be reached. Evaluation of the NCFE shows that the model measures the level of competence of nursing students and that its design is beneficial for the students' clinical reasoning (Ziegert, Elmquist, Johansson, Larsson, & Andersson, 2014).

While the NCFE represents competence assessed by examination, self-assessed competence is measured using the Nurse Professional Competence Scale (NPC) (Gardulf et al., 2019, 2016; Nilsson et al., 2016, 2014; Theander et al., 2016). This scale has recently been developed into the NPC short form (NPC-SF) that covers six competence areas: Nursing Care, Value-based Nursing Care, Medical and Technical Care, Care Pedagogics, Documentation and Administration of Nursing Care and Development, Leadership and Organization of Nursing Care (Nilsson, Engström, Florin, Gardulf, & Carlsson, 2018).

Self-evaluation of ability can also be operationalized in terms of self-efficacy (Zell & Krizan, 2014). Self-efficacy is the belief in one's capability to execute the behaviour required to produce desired outcomes. Self-efficacy is a concept that includes belief in one's own ability to perform an action; the greater the level of self-efficacy one has, the more likely it is that they will start and continue an activity with a positive result (Bandura, 1997).

Previous research illustrates the complex relationship between self-perceived performance and actual performance and how the ability to make accurate self-assessments might differ between students in nursing education (Burden et al., 2018; Lejonqvist et al., 2016; Ugland Vae et al., 2018). It is essential that this is studied further since correct self-assessments are crucial to both learning and patient safety in clinical practice.

## 2 | THE STUDY

### 2.1 | Aims

The aim of the study was to identify clusters based on graduating nursing students' self-reported professional competence and their

achievement on a national examination. An additional aim was to describe and compare the identified clusters regarding sample characteristics, students' perceptions of overall quality of the BSN programme, and students' general self-efficacy (GSE).

### 2.2 | Design

This was a cross-sectional study, combining survey data and results from a national examination.

### 2.3 | Sample/participants

All students taking the NFCE written and bedside exam and also responding to the questionnaires, in their final term of the nursing programme at two universities and one university college in Sweden were included in this study ( $N = 179$ ).

### 2.4 | Data collection

A research-group representative at each HEI gave written and oral information about the study and issued questionnaires to nursing students during their last 2 weeks of the nursing programme (January, 2017). Students were given an envelope containing information about the study and a coded questionnaire, which was completed individually and then left in a designated box in the classroom. Using a code key, the research-group representative at each HEI linked each questionnaire to a student's NFCE results.

To measure nursing students' self-rated competence, the NPC-SF was used. Data were collected using a 35-item version distributed in six competence areas: Nursing Care (five items, Cronbach's alpha ( $\alpha$ ) in the present study .79); Value-Based Nursing Care (five items,  $\alpha = .79$ ); Medical and Technical Care (six items,  $\alpha = .81$ ); Care Pedagogics (five items,  $\alpha = .87$ ); Documentation and Administration of Nursing Care (eight items,  $\alpha = .79$ ) and Development, Leadership and Organization of Nursing Care (six items,  $\alpha = .80$ ). Response alternatives used a 4-grade scale: 1 = to a very low degree, 2 = to a relatively low degree, 3 = to a relatively high degree, and 4 = to a very high degree. Scores for the respective competence areas were calculated by summing up all items divided by the highest possible score in the competence area and then multiplying by 100, thereby giving 0–100 values.

Self-efficacy was measured using the Swedish version of the 10-item GSE scale (Koskinen-Hagman, Schwarzer, & Jerusalem, 1999; Schwarzer & Jerusalem, 1995). The 10 items are rated on a 4-point Likert scale ranging from 1 = 'not at all true' to 4 = 'exactly true', where higher scores indicate a higher level of self-efficacy.

Sociodemographic data included age, gender, previous education, and work while studying the BSN programme. The students were also asked about the overall quality of the programme and whether they would recommend the programme to others.

To measure the students' competence by examination, results from the NCFE were used (Athlin et al., 2012). The written exam is a modified essay question exam, which means that as the examination

proceeds, situations and conditions change and new questions are added (Khan & Aljarallah, 2011). To pass, students need 33 points of a possible 50 and two questions about drug calculation must be correctly answered. During the bedside examination, students take care of a patient in need of comprehensive medical and nursing care for 3 hr. An experienced Registered Nurse evaluates their performance based on a protocol following the nursing process. A clinical lecturer makes the final assessment that decides whether the student will pass or fail. Students must pass both the written and bedside tests to pass the examination. Students in this present study accomplished their written NCFE in November 2016 (same date and time for all HEIs) and their bedside examination during their final term of the nursing programme, that is, September 2016–January 2017.

## 2.5 | Ethical considerations

The study was carried out in accordance with the World Medical Association declaration of Helsinki (originally adopted in 1964). A local ethics committee reviewed the study (Dnr C2016/567) and it was determined that the project did not fall under the Ethics Assessment Act (2003:460). Participants were informed via a letter of the aim of the study, that their participation was voluntary and that they were entitled to terminate their participation at any time. A response to the questionnaire was interpreted as consent from participants.

## 2.6 | Data analysis

IBM SPSS Statistics version 22 was used for data analysis (SPSS Inc.). Data were screened for missing values and multiple imputation (MI) was used to handle internal missing data. The method generated five imputed datasets together with the original dataset. Students with more than 50% missing data in a factor of the NPC-SF scale were excluded in all further analyses ( $N = 9$ ). The final cluster sample resulted in 170 study participants, whereof 121 had complete responses/no missing data at all in NPC-SF. Descriptive statistics were used to describe sample characteristics, results on the NPC-SF and NCFE.

Cluster analysis was used to identify homogeneous clusters of participants based on their NPC-SF competence profiles and achievement on the NCFE. The two-step cluster analysis (TSCA) procedure was used with log-likelihood distance measures. This method was chosen as it allows the use of both continuous and categorical variables and the method improves the weaknesses of applying a single clustering method. The advantage of the method is that it 'integrates hierarchical and partitioning clustering algorithm with adding attributes to cluster objects' (Shih, Jheng, & Lai, 2010, p.11). The method first pre-clusters cases into small subclasses and then forms final clusters using hierarchical methods. The optimal number of clusters is determined automatically in the TSCA using Schwarz's Bayesian Information Criterion. The Silhouette measure of cohesion and separation is used to determine the quality of the cluster solution (Norusis, 2012). A silhouette value of less than 0.20 indicates

a poor solution, 0.20–0.50 indicates a fair solution and over 0.50 indicates a good solution (Mooi & Sarstedt, 2011). Comparative analyses (one-way ANOVA, Kruskal–Wallis test, Mann–Whitney  $U$  test and chi-square test) were used to test statistical differences between clusters regarding sample characteristics and the variables recommending the BSN programme, overall quality of the BSN programme, and GSE. The significance level was set at  $p < .05$ .

## 2.7 | Validity and reliability

Previous studies have shown that the NPC Scale can be used as a tool for quality assessments and improvements of nursing education programmes (Gardulf et al., 2019). Construct validity of NPC-SF has been tested with principal component analysis and confirmative factor analysis where the factor solution explained 54% of the overall variance. Reliability measured as internal consistency showed  $\alpha$  values  $>.70$  for all competence areas (Nilsson et al., 2018). Also in the present study, the internal consistency of the NPC-SF was high for all competence areas, with  $\alpha$  values ranging between 0.79–0.87. The GSE scale has been validated in several languages and is widely used internationally (Luszczynska, Scholz, & Schwarzer, 2005). Internal consistency has been reported to be  $\alpha = .90$  (Löve, Moore, & Hensing, 2012), with  $\alpha = .89$  in the present study. The silhouette values in the cluster analyses indicated fair cluster solutions and repeated clustering further demonstrated cluster stability. Furthermore, the survey response rate was high and the MI method handling missing data resulted in a maximized sample size. The sample included two universities and one university college which speaks in favour of the generalizability of the results.

## 3 | RESULTS

Sample characteristics and study participants' perceptions of the BSN programme are presented in Table 1. Most (85.2%) of the students in this study were females and their mean age was 27.8 years ( $SD$  6.3). Most students (63.4%) did not have previous experience of higher education before entering the nursing programme and slightly more than half of the sample (57.0%) had studied a theoretical programme at upper secondary school while the rest had studied nursing care (15.2%) or another programme (27.9%). Slightly more than half of the students (54.1%) had prior work experience in healthcare. A large proportion (81.1%) worked in healthcare while they were studying. Most students (82.7%) responded that they would likely or definitely recommend their BSN programme to others and most (73.1%) rated the overall quality of their BSN programme as high. On average, students rated their GSE as 3.2 ( $SD = 0.46$ ). About two-thirds (61.8%) passed the final examination (NCFE). Among the NPC competence areas, Value-based Nursing Care was rated highest while the competence area Development, Leadership and Organization of Nursing Care was rated lowest (Table 2).

A TSCA was performed separately on each of the five MI datasets and showed stability across the different versions. All MI datasets revealed a solution with three clusters with similar patterns. Silhouette

**TABLE 1** Clusters in relation to sample characteristics and students' perceptions of the Bachelor of Science in Nursing (BSN) programme

Variables	Cluster			p values
	1/passed N = 36	2/passed N = 69	3/failed N = 65	
Gender, male/female, N = 169	6/29	8/61	11/54	.623 <sup>b</sup>
Age, years, N = 168, mean (SD)	29.2 (6.9)	26.9 (5.3)	27.9 (6.9)	.200 <sup>a</sup>
General Self-Efficacy Scale, N = 167, mean (SD)	2.8 (0.4)	3.3 (0.4)	3.2 (0.5)	<.001 <sup>a</sup>
Education at upper secondary school level prior to entering the BSN Programme, N = 165, (3-year theoretical programme Natural Science/3-year theoretical programme Social Science/3-year programme Nursing Care/Other programme)	8/14/3/9	9/32/8/19	9/22/14/18	.384 <sup>b</sup>
Higher Education before the nursing programme, N = 167, yes/no	17/18	22/46	22/42	.243 <sup>b</sup>
Work experience in healthcare prior to entering the BSN Programme, N = 170, yes/no	19/17	34/35	39/26	.453 <sup>b</sup>
Paid work experience in healthcare when studying the BSN Programme, N = 169, yes/no	32/4	48/20	57/8	.017 <sup>b</sup>
Recommend the BSN programme to others, N = 168 (no/likely/definitely)	13/19/4	9/29/30	7/32/25	.001 <sup>b</sup>
Overall Quality of the BSN Programme <sup>d</sup> , N = 167, Md (Q1;Q3)	2 (2;3)	3 (3;3)	3 (3;3)	<.001 <sup>c</sup>

Note: Mean, median (Md), standard deviations (SD) and quartiles (Q1, Q3).

<sup>a</sup>One-way ANOVA.

<sup>b</sup>Chi-square test.

<sup>c</sup>Kruskal-Wallis test.

<sup>d</sup>Response alternatives were 1 = very low, 2 = low, 3 = high and 4 = very high.

**TABLE 2** Cluster means and standard deviations in relation to NPC competence areas

Variables	Total sample	Cluster 1/passed	Cluster 2/passed	Cluster 3/failed	P value <sup>a</sup>
Number of participants (%)	170	36 (21.2%)	69 (40.6%)	65 (38.2%)	
Nursing care	84.9 (11.0)	74.2 (10.4)	89.0 (8.6)	86.5 (9.8)	<.001
Value-based nursing care	90.6 (10.1)	79.3 (13.0)	95.4 (5.2)	91.8 (7.2)	<.001
Medical and technical care	82.2 (11.1)	70.7 (10.8)	87.8 (7.5)	82.6 (9.7)	<.001
Care pedagogics	81.8 (12.8)	69.4 (11.1)	87.5 (10.6)	82.7 (11.3)	<.001
Documentation and administration of nursing care	83.5 (9.8)	73.5 (8.8)	88.1 (7.0)	84.2 (8.9)	<.001
Development, leadership, and organization of nursing care	71.7 (12.6)	60.6 (8.9)	77.7 (11.3)	71.5 (11.6)	<.001

Note: Values for NPC scores ranged between 1–100, where 100 correspond to high self-reported competence.


Abbreviation: NPC, Nurse Professional Competence Scale.

<sup>a</sup>One-way ANOVA.


measures of cohesion and separation ranged from 0.455–0.463 (indicating a fair solution). The first MI dataset was then kept for further data analyses. The cases were randomly ordered three times and the stability of the cluster solution was verified. For all solutions, there were three clusters with 36, 65, and 69 students and the silhouette measure of cohesion and separation was 0.459 (a fair solution). The three clusters from MI dataset 1 are presented in Table 2 and Figures 1 and 2. The students in cluster 1 (N = 36) had all passed the national clinical examination test but rated their competence as significantly lower than students in the two other clusters (Post Hoc, Bonferroni adjusted, all p values <.001). Their ratings were below the median for all competence areas (Figure 1; mean values for the competence areas ranged from 60.6–79.3, Table 2). The students in cluster 2 (N = 69) had also

passed the national clinical examination test. They rated themselves as quite good in all competence areas and above the median for the total sample for five of the six competence areas (mean values ranged from 77.7–95.4). The students in cluster 3 (N = 65) had all failed the national clinical examination test. They rated themselves lower than the median for one competence area, higher for one and on the median for the remaining four competence areas (mean values ranged from 71.5–91.8).

Sample characteristics for the clusters and students' perceptions of the overall quality of their BSN programme are presented in Table 1. Cluster 1 scored significantly lower than clusters 2 and 3 regarding GSE (Post Hoc, Bonferroni adjusted, all p values <.001), they rated the overall quality of the BSN programme lower than clusters 2 and 3 (Mann-Whitney U test all p < .001) and fewer in



**FIGURE 1** A comparison of the three clusters. Continuous variables (competence areas) are shown as boxplots with the overall medians and interquartile ranges (white) together with boxplots for each cluster's median and interquartile range. The categorical variable (NCFE) is shown as dot plots (the size indicates the most frequent response for each cluster). Cluster 1 (passed NCFE, dark blue), Cluster 2 (passed NCFE, light blue), and Cluster 3 (failed NCFE, red)



**FIGURE 2** Cluster mean values for NPC competence areas. Cluster 1 (passed NCFE, dark blue), Cluster 2 (passed NCFE, light blue), and Cluster 3 (failed NCFE, red). NPC, Nurse Professional Competence Scale; NCFE, National Clinical Final Examination

cluster 1 would definitely recommend the BSN programme (adjusted residual 3.4). Regarding paid work in healthcare while studying the BSN Programme, the results showed that fewer worked in cluster 2 than in the other two clusters (adjusted residual 2.9).

#### 4 | DISCUSSION

The main result from this study was the identification of three different clusters, which illustrated students' different competence

profiles. Students in cluster 1 passed the NCFE but presented lower self-assessed competence (NPC) than the overall median values in the group. Students in cluster 2 also passed the NCFE and rated themselves as above median in all but one NPC competence area. Students in cluster 3 failed the NCFE but still rated themselves on the median level or higher in all but one of the NPC competence areas. Thus, self-assessed and non-self-assessed competency were concordant among students in cluster 2, whereas students in clusters 1 and 3 differed in these two aspects in different ways. This disparity might have consequences for these students while in nursing

education and also for health care organizations when the students enter working life.

Self-assessment is a critical skill and also complex and challenging for students. In the literature, this has been studied and discussed in relation to the 'impostor syndrome' for example (characterized by chronic degrading feelings) (Villwock, Sobin, Koester, & Harris, 2016), or the opposite, which results in inflated self-assessments (Kruger & Dunning, 1999). This illustrates how people might exhibit different approaches to self-assessments of their abilities. In our sample, the students in cluster 1 might be the ones suffering from the 'impostor syndrome', that is, their self-criticism overrides their actual abilities. Their GSE was lower than for students in the other two clusters and they were also more critical to the quality of their education. They performed 'well enough' but were still dissatisfied. However, the students in cluster 3 seemed to exhibit inflated self-assessments in relation to their NCFE results.

Based on these results, we need to consider what pedagogical methods we use in education and how they can strengthen students' ability to develop their self-assessment competence. We must also consider how we can educate students based on their specific needs. Educational interventions focusing on self-directed and self-regulated learning (Saks & Leijen, 2014) might support students' ability to self-assess and self-reflect, thereby improving their learning conditions. For example, the pedagogical model peer learning, which is characterized by a two-way, reciprocal learning activity supporting abilities like self-assessment and peer assessment as well as communication and articulation of own knowledge, understanding, and skills (Boud, Cohen, & Sampson, 2001; Pålsson, Engström, Leo Swenne, & Mårtensson, 2018). In relation to the clinical education of undergraduate nursing and other health science students, peer learning has demonstrated improved nursing self-efficacy (Pålsson, Mårtensson, Swenne, Adel, & Engström, 2017) and self-evaluation (Secomb, 2008), for example. Another promising learning intervention is drama. Drama pedagogy has been used in nursing education to prepare students on both bachelor and master level for their future nursing roles (Arvekle, 2017). The use of drama in nursing education can provide opportunities to explore interactions with others, which can increase students' self-awareness and their ability to reflect on their future professional identity as nurses. Furthermore, drama allows the students to re-enact situations, ethical dilemmas, and conflicts from the healthcare context to practice, reflect, discuss, and learn about conflict management (Arvekle, Berg, Wigert, Morrison-Helme, & Lepp, 2018).

In previous studies on nurses and competence, factors such as length of work experience, frequent use of competence, factors related to the practice environment, and healthcare context, as well as nurse related and sociodemographic factors have been found to be related to the different areas of competence (Flinkman et al., 2017; Gardulf et al., 2016; Leksell, Gardulf, Nilsson, & Lepp, 2015; Meretoja, Numminen, Isoaho, & Leino-Kilpi, 2015; Nilsson et al., 2016; Salonen, Kaunonen, Meretoja, & Tarkka, 2007). Furthermore, for nursing students, work in healthcare while studying has been found as related

to self-reported competence (Gardulf et al., 2016). Competence is developing over time and depends on work experience, which is also formulated by Benner (1989) who describes the process from novice to expert. The newly graduated nurse is a beginner, which is important to consider in healthcare organizations and for the newly graduated themselves. Results from this study illustrate how students' self-assessments and beliefs in their competence might differ from actual performance. This is also important to consider when the new nurse enters the healthcare organization. Correct assessments of one's own competence/ability are essential for patient safety.

The understanding of lifelong learning in relation to nursing research and curricula is crucial to support the necessary skills and attitudes among the students (Davis, Taylor, & Reyes, 2014). Qalehsari, Khaghanizadeh, and Ebadi (2017) have highlighted the complexity of lifelong learning and concluded that one single strategy cannot lead to lifelong learning alone. The use of strategies for lifelong learning will lead to increased quality of education and of patient care and to the development of nursing competency.

#### 4.1 | Limitations

Analyses were based on overall competency rather than on specific areas of competence, which can be considered as a limitation and this needs further study. However, the cluster approach illustrating different competence profiles for subgroups of students has its advantages. Weaknesses with the TSCA are that the final solution might depend on the order of cases and thereby the method should be tested with the cases in different order (Norusis, 2012). The TSCA was therefore applied three times with the cases in different random order to test the stability of the cluster solution. Generalizability of the results must be done with caution as the sample included only three out of 25 HEI in Sweden.

## 5 | CONCLUSION

This study illustrates how nursing students' self-assessed competence might differ from competency assessed by examination. This is challenging for nursing education, both for theoretical parts of the nursing programme and for clinical education. Pedagogical interventions to support realistic perceptions of own competence are crucial since the perceptions that students and later on registered nurses, have of their own competence might be a matter of patient safety.

#### ACKNOWLEDGEMENTS

The authors would like to thank the participating students for answering the questionnaire.

#### CONFLICT OF INTEREST

No conflict of interest has been declared by the authors.

## AUTHOR CONTRIBUTIONS

All authors have made substantial contributions to all of the following: (1) the conception and design of the study, or acquisition of data, or analysis and interpretation of data, (2) drafting the article or revising it critically for important intellectual content, and (3) final approval of the version to be submitted.

## ORCID

Henrietta Forsman  <https://orcid.org/0000-0002-5892-9897>

Inger Jansson  <https://orcid.org/0000-0002-0211-5077>

Janeth Leksell  <https://orcid.org/0000-0001-8682-2045>

Margret Lepp  <https://orcid.org/0000-0002-3324-0946>

Maria Engström  <https://orcid.org/0000-0002-9912-5350>

Jan Nilsson  <https://orcid.org/0000-0001-6245-1788>

## REFERENCES

- Aiken, L. H., Sloane, D. M., Bruyneel, L., Van den Heede, K., Griffiths, P., Busse, R., ... Sermeus, W. (2014). Nurse staffing and education and hospital mortality in nine European countries: A retrospective observational study. *The Lancet*, *383*, 1824–1830. [https://doi.org/10.1016/S0140-6736\(13\)62631-8](https://doi.org/10.1016/S0140-6736(13)62631-8)
- Aiken, L. H., Sloane, D., Griffiths, P., Rafferty, A. M., Bruyneel, L., McHugh, M., ... Sermeus, W. (2017). Nursing skill mix in European hospitals: Cross-sectional study of the association with mortality, patient ratings and quality of care. *BMJ Quality and Safety*, *26*, 559–568. <https://doi.org/10.1136/bmjqs-2016-005567>
- Arveklef, H. S. (2017). *Drama and Learning in Nursing Education. A study in first and second cycle*. (Diss. Institute of Health and Care Sciences, Sahlgrenska Academy, University of Gothenburg. Gothenburg: BrandFactory. ISBN 978-91-629-0362-6 (PRINT). ISBN 978-91-629-0363-3 (PDF).
- Arveklef, S. H., Berg, L., Wigert, H., Morrison-Helme, M., & Lepp, M. (2018). Learning about conflict and conflict management through drama in nursing education. *Journal of Nursing Education*, *57*(4), 209–216. <https://doi.org/10.3928/01484834-20180322-04>
- Athlin, E., Larsson, M., & Söderhamn, O. (2012). A model for a national clinical final examination in the Swedish bachelor programme in nursing. *Journal of Nursing Management*, *20*, 90–101. <https://doi.org/10.1111/j.1365-2834.2011.01278.x>
- Bandura, A. (1997). *Self-efficacy: The exercise of control*. New York, NY: Freeman.
- Baxter, P., & Norman, G. (2011). Self-assessment or self deception? A lack of association between nursing students' self-assessment and performance. *Journal of Advanced Nursing*, *67*, 2406–2413. <https://doi.org/10.1111/j.1365-2648.2011.05658.x>
- Benner, P. (1989). From novice to expert. *The American Journal of Nursing*, *82*, 402–407.
- Boud, D., R. Cohen, & J. Sampson (Eds.) (2001). *Peer learning in higher education. Learning from and with each other*. New York, NY: Routledge.
- Burden, S., Topping, A. E., & O'Halloran, C. (2018). Mentor judgements and decision-making in the assessment of student nurse competence in practice: A mixed-methods study. *Journal of Advanced Nursing*, *74*, 1078–1089. <https://doi.org/10.1111/jan.13508>
- The World Medical Association Declaration of Helsinki. (1964). Ethical principles for medical research involving human subjects. Retrieved from <https://www.wma.net/policies-post/wma-declaration-of-helsinki>
- nkiethical-principles-for-medical-research-involving-human-subjects/. Accessed 17 October, 2019.
- Cronenwett, L., Sherwood, G., Barnsteiner, J., Disch, J., Johnson, J., Mitchell, P., ... Warren, J. (2007). Quality and safety education for nurses. *Nurse Outlook*, *55*, 122–131. <https://doi.org/10.1016/j.outlook.2007.02.006>
- Davis, L., Taylor, H., & Reyes, H. (2014). Lifelong learning in nursing: A Delphi study. *Nurse Education Today*, *34*, 441–445. <https://doi.org/10.1016/j.nedt.2013.04.014>
- Flinkman, M., Leino-Kilpi, H., Numminen, O., Jeon, Y., Kuokkanen, L., & Meretoja, R. (2017). Nurse Competence Scale: A systematic and psychometric review. *Journal of Advanced Nursing*, *73*, 1035–1050. <https://doi.org/10.1111/jan.13183>
- Gardulf, A., Florin, J., Carlsson, M., Leksell, J., Lepp, M., Lindholm, C., ... Nilsson, J. (2019). The Nurse Professional Competence (NPC) Scale: A tool that can be used in national and international assessments of nursing education programmes. *Nordic Journal of Nursing Research*, *39*, 137–142. <https://doi.org/10.1177/2057158518824530>
- Gardulf, A., Nilsson, J., Florin, J., Leksell, J., Lepp, M., Lindholm, C., ... Johansson, E. (2016). The Nurse Professional Competence (NPC) Scale: Self-reported competence among nursing students on the point of graduation. *Nurse Education Today*, *36*, 165–171. <https://doi.org/10.1016/j.nedt.2015.09.013>
- Institute of Medicine (US) Committee on the health professions education summit, Greiner, A. C., & Knebel, E. (2003). Chapter 3: The core competencies needed for health care professionals. In Institute of Medicine (US) Committee on the health professions education summit, A. C. Greiner, & E. Knebel (Eds.), *Health professions Education: A bridge to quality* (pp. 45–67) Washington, DC: National Academies Press.
- Kajander-Unkuri, S., Salminen, L., Saarikoski, M., Suhonen, R., & Leino-Kilpi, H. (2013). Competence areas of nursing students in Europe. *Nurse Education Today*, *33*, 625–632. <https://doi.org/10.1016/j.nedt.2013.01.017>
- Kendall-Gallagher, D., Aiken, L., Sloane, D. M., & Cimiotti, J. P. (2011). Nurse specialty certification, inpatient mortality and failure to rescue. *Journal of Nursing Scholarship*, *43*, 188–194. <https://doi.org/10.1111/j.1547-5069.2011.01391.x>
- Khan, M. U., & Aljarallah, B. M. (2011). Evaluation of modified essay questions (MEQ) and multiple choice questions (MCQ) as a tool for assessing the cognitive skills of undergraduate medical students. *International Journal of Health Sciences*, *5*, 39–43.
- Koskinen-Hagman, M., Schwarzer, R., & Jerusalem, M. (1999). Swedish version of the general self efficacy scale. Retrieved from <http://userpage.fu-berlin.de/health/swedish.htm>
- Kruger, J., & Dunning, D. (1999). Unskilled and unaware of it: How difficulties in recognizing one's own incompetence lead to inflated self-assessments. *Journal of Personality and Social Psychology*, *77*, 1121–1134. <https://doi.org/10.1037/0022-3514.77.6.1121>
- Lejonqvist, G.-B., Eriksson, K., & Meretoja, R. (2016). Evaluating clinical competence during nursing education: A comprehensive integrative literature review. *International Journal of Nursing Practice*, *22*, 142–151. <https://doi.org/10.1111/ijn.12406>
- Leksell, J., Gardulf, A., Nilsson, J., & Lepp, M. (2015). Self-reported conflict management competence among nursing students on the point of graduation and registered nurses with professional experience. *Journal of Nursing Education and Practice*, *5*(8), 82–89. <https://doi.org/10.5430/jnep.v5n8p82>
- Liu, Y., & Aunguroch, Y. (2018). Current literature review of registered nurses' competency in the global community. *Journal of Nursing Scholarship*, *50*, 191–199. <https://doi.org/10.1111/jnu.12361>
- Löve, J., Moore, C. D., & Hensing, G. (2012). Validation of the Swedish translation of the general self-efficacy scale. *Quality of Life Research*, *21*, 1249–1253. <https://doi.org/10.1007/s11136-011-0030-5>



- Luszczynska, A., Scholz, U., & Schwarzer, R. (2005). The general self-efficacy scale: Multicultural validation studies. *Journal of Psychology, 139*(5), 439–457. <https://doi.org/10.3200/JRPL.139.5.439-457>
- Meretoja, R., Numminen, O., Isoaho, H., & Leino-Kilpi, H. (2015). Nurse competence between three generational nurse cohorts: A cross-sectional study. *International Journal of Nursing Practice, 21*, 350–358. <https://doi.org/10.1111/ijn>
- Miller, G. (1990). The assessment of clinical skills/competence/performance. *Academic Medicine, 65*(9), 63–67. <https://doi.org/10.1097/00001888-199009000-00045>
- Mooi, E., & Sarstedt, M. (2011). Chapter 9: Cluster analysis. In E. Mooi, & M. Sarstedt (Eds.), *A concise guide to market research* (pp. 237–284). Berlin Heidelberg, NY: Springer-Verlag.
- Nehrir, B., Vanaki, Z., Mokhtari Nouri, J., Mohammed Khademolhosseini, S., & Ebadi, A. (2016). Competency in nursing students: A systematic review. *International Journal of Travel Medicine and Global Health, 4*, 3–11. <https://doi.org/10.20286/IJTMGH-04013>
- Nilsson, J., Engström, M., Florin, J., Gardulf, A., & Carlsson, M. (2018). A short version of the nurse professional competence scale for measuring nurses' self-reported competence. *Nurse Education Today, 71*(Dec 2018), 233–239. <https://doi.org/10.1016/j.nedt.2018.09.028>
- Nilsson, J., Johansson, E., Carlsson, M., Florin, J., Leksell, J., Lepp, M., ... Gardulf, A. (2016). Disaster nursing: Self-reported competence of nursing students and registered nurses, with focus on their readiness to manage violence, serious events and disasters. *Nurse Education in Practice, 17*, 102–108. <https://doi.org/10.1016/j.nepr.2015.09.012>
- Nilsson, J., Johansson, E., Egmar, A.-C., Florin, J., Leksell, J., Lepp, M., ... Gardulf, A. (2014). Development and validation of a new tool measuring nurses self-reported professional competence—The nurse professional competence (NPC) Scale. *Nurse Education Today, 34*, 574–580. <https://doi.org/10.1016/j.nedt.2013.07.016>
- Norusis, M. (2012). Chapter 17 cluster analysis. In M. Norusis (Ed.), *IBM SPSS Statistics 19 guide to data analysis* (pp. 375–404). Retrieved from [http://www.norusis.com/pdf/SPC\\_v19.pdf](http://www.norusis.com/pdf/SPC_v19.pdf)
- Pålsson, Y., Engström, M., Leo Swenne, C., & Mårtensson, G. (2018). A peer learning intervention targeting newly graduated nurses: A feasibility study with a descriptive design based on the Medical Research Council framework. *Journal of Advanced Nursing, 74*, 1127–1138. <https://doi.org/10.1111/jan.13513>
- Pålsson, Y., Mårtensson, G., Swenne, C. L., Adel, E., & Engström, M. (2017). A peer learning intervention for nursing students in clinical practice education: A quasi-experimental study. *Nurse Education Today, 51*, 81–87. <https://doi.org/10.1111/jan.13513>
- Qalehsari, M. Q., Khaghanizadeh, M., & Ebadi, A. (2017). Lifelong learning strategies in nursing: A systematic review. *Electronic Physician, 9*, 5541–5550. <https://doi.org/10.19082/5541>
- Saks, K., & Leijen, Ä. (2014). Distinguishing self-directed and self-regulated learning and measuring them in the E-learning context. *Procedia—social and Behavioral Sciences, 112*, 190–198. <https://doi.org/10.1016/j.sbspro.2014.01.1155>
- Salonen, A. H., Kaunonen, M., Meretoja, R., & Tarkka, M. T. (2007). Competence profiles of recently registered nurses working in intensive and emergency settings. *Journal of Nursing Management, 15*, 792–800. <https://doi.org/10.1111/j.1365-2934.2007.00768.x>
- Schwarzer, R., & Jerusalem, M. (1995). Generalized self-efficacy scale. In M. Johnston, J. Weinman, & S. C. Wright (Eds.), *Measures in health psychology: A user's portfolio. Causal and control beliefs* (pp. 35–37). Windsor, UK: Windsor, NFER-NELSON.
- Sears, K., Godfrey, C. M., Luctar-Flude, M., Ginsburg, L., Tregunno, D., & Ross-White, A. (2014). Measuring competence in healthcare learners and healthcare professionals by comparing self assessment with objective structured clinical examinations: A systematic review. *JBI Database of Systematic Reviews and Implementation Reports, 12*, 221–272. <https://doi.org/10.11124/jbisrir-2014-1264>
- Secomb, J. (2008). A systematic review of peer teaching and learning in clinical education. *Journal of Clinical Nursing, 17*, 703–716. <https://doi.org/10.1111/j.1365-2702.2007.01954.x>
- Shih, M.-Y., Jheng, J.-W., & Lai, L.-F. (2010). A two-step method for clustering mixed categorical and numeric data. *Journal of Applied Science and Engineering, 13*, 11–19.
- The Swedish Code of Statutes. (1992). Higher Education Act (1992:1434).
- The Swedish Code of Statutes. (1993). Higher Education Ordinance (1993:100).
- The Swedish Code of Statutes. (2003). Ethics Assessment Act (2003:460).
- The Swedish National Board of Health and Welfare [Socialstyrelsen]. (2018). Kompetensförsörjning och patientsäkerhet. Hur brister i bemanning och kompetens påverkar patientsäkerheten. [In Swedish] Retrieved from <https://www.socialstyrelsen.se/Lists/Artikelkatalog/Attachments/20871/2018-2-15.pdf>.
- The Swedish Society of Nursing [Svensk sjuksköterskeförening]. (2017). Kompetensbeskrivning för legitimerad sjuksköterska. [In Swedish] Retrieved from <https://www.swenurse.se/globalassets/01-svensk-sjukskoterskeforening/publikationer-svensk-sjukskoterskeforening/kompetensbeskrivningar-publikationer/kompetensbeskrivning-legitimerad-sjukskoterska-2017-for-webb.pdf>
- Theander, K., Wilde-Larsson, B., Carlsson, M., Florin, J., Gardulf, A., Johansson, E., ... Nilsson, J. (2016). Adjusting to future demands in healthcare—Curriculum changes and nursing students' self-reported professional competence. *Nurse Education Today, 37*, 178–183. <https://doi.org/10.1016/j.nedt.2015.11.012>
- Ugland Vae, K. J., Engström, M., Mårtensson, G., & Löfmark, A. (2018). Nursing students' and preceptors' experience of assessment during clinical practice: A multilevel repeated-interview study of student-preceptor dyads. *Nurse Education in Practice, 30*, 13–19. <https://doi.org/10.1016/j.nepr.2017.11.014>
- Villwock, J. A., Sobin, L. B., Koester, L. A., & Harris, T. M. (2016). Impostor syndrome and burnout among American medical students: A pilot study. *International Journal of Medical Education, 7*, 364–369. <https://doi.org/10.5116/ijme.5801.eac4>
- Zell, E., & Krizan, Z. (2014). Do people have insight into their abilities? A metasynthesis. *Perspectives on Psychological Science, 9*, 111–125. <https://doi.org/10.1177/1745691613518075>
- Ziegert, K., Elmqvist, M. A., Johansson, U.-B., Larsson, M., & Andersson, P. L. (2014). How the final Swedish clinical exam prepares the nursing students for their future challenges—Qualitative analysis. *Creative Education, 5*, 1887–1894. <https://doi.org/10.4236/ce.2014.521211>

**How to cite this article:** Forsman H, Jansson I, Leksell J, et al. Clusters of competence: Relationship between self-reported professional competence and achievement on a national examination among graduating nursing students. *J Adv Nurs*. 2020;76:199–208. <https://doi.org/10.1111/jan.14222>

The *Journal of Advanced Nursing (JAN)* is an international, peer-reviewed, scientific journal. *JAN* contributes to the advancement of evidence-based nursing, midwifery and health care by disseminating high quality research and scholarship of contemporary relevance and with potential to advance knowledge for practice, education, management or policy. *JAN* publishes research reviews, original research reports and methodological and theoretical papers.

For further information, please visit *JAN* on the Wiley Online Library website: [www.wileyonlinelibrary.com/journal/jan](http://www.wileyonlinelibrary.com/journal/jan)

**Reasons to publish your work in *JAN*:**

- **High-impact forum:** the world's most cited nursing journal, with an Impact Factor of 1.998 – ranked 12/114 in the 2016 ISI Journal Citation Reports © (Nursing (Social Science)).
- **Most read nursing journal in the world:** over 3 million articles downloaded online per year and accessible in over 10,000 libraries worldwide (including over 3,500 in developing countries with free or low cost access).
- **Fast and easy online submission:** online submission at <http://mc.manuscriptcentral.com/jan>.
- **Positive publishing experience:** rapid double-blind peer review with constructive feedback.
- **Rapid online publication in five weeks:** average time from final manuscript arriving in production to online publication.
- **Online Open:** the option to pay to make your article freely and openly accessible to non-subscribers upon publication on Wiley Online Library, as well as the option to deposit the article in your own or your funding agency's preferred archive (e.g. PubMed).