

ORIGINAL RESEARCH ARTICLE

Osteocalcin does not influence acute or chronic inflammation in human vascular cells

Sophie A. Millar  | Ieva Zala | Susan I. Anderson | Saoirse E. O'Sullivan

Division of Graduate Entry Medicine and Medical Sciences, School of Medicine, Royal Derby Hospital, University of Nottingham, Derby, UK

Correspondence

Sophie Millar, Division of Graduate Entry Medicine and Medical Sciences, School of Medicine, Royal Derby Hospital, University of Nottingham, Derby, DE22 3DT UK.
Email: sophieannemillar@gmail.com

Funding information

Biotechnology and Biological Sciences Research Council, Grant/Award Number: BB/M008770/1

Abstract

Some human observational studies have suggested an anti-inflammatory role of osteocalcin (OCN). An inflammatory protocol using interferon- γ and tumor necrosis factor- α (10 ng/ml) was employed to examine the acute (24 hr) and chronic (144 hr) effects of uncarboxylated OCN (ucOCN) in commercial, primary, subcultured human aortic endothelial cells (HAEC), and human smooth muscle cells (HASMCs). The inflammatory protocol increased phosphorylation of intracellular signaling proteins (CREB, JNK, p38, ERK, AKT, STAT3, STAT5) and increased secretion of adhesion markers (vascular cell adhesion molecule-1, intracellular adhesion molecule-1, monocyte chemoattractant protein-1) and proinflammatory cytokines (interleukin-6 [IL-6], IL-8). After acute inflammation, there were no additive or reductive effects of ucOCN in either cell type. Following chronic inflammation, ucOCN did not affect cell responses, nor did it appear to have any pro- or anti-inflammatory effects when administered acutely or chronically on its own in either cell type. Additionally, ucOCN did not affect lipopolysaccharide (LPS)-induced acute inflammation in HAECs or HASMCs. The findings of this study do not support a causal role for OCN within the models of vascular inflammation chosen. Further confirmatory studies are warranted.

KEYWORDS

bone hormone, endothelial, inflammation, osteocalcin, smooth muscle

1 | INTRODUCTION

Atherosclerosis is a progressive inflammatory disease, involving revolving phases of inflammation leading to macrophage infiltration and microcalcifications (Hutcheson & Aikawa, 2014; Ross, 1999). Inflammatory cytokines, chemoattractant proteins, and adhesion molecules are present throughout the atherosclerotic process, of which the endothelium and smooth muscle cell layers are both pivotal. The process of atherosclerotic calcification is also crucially triggered by inflammatory-related pathways (Bessueille & Magne, 2015; Mazzini & Schulze, 2006). A paradox vascular-bone axis exists as while bone formation decreases with age, biomineralization of the

vasculature is increased, which is suggested to be due to a shared etiology of inflammation (Bessueille & Magne, 2015; Sage, Tintut, & Demer, 2010). The endocrine function of bone, particularly osteocalcin (OCN), is increasingly under investigation due to its link with whole body metabolism and far-reaching extra-skeletal effects (Lee et al., 2007; Oldknow, Macrae, & Farquharson, 2015; Oury et al., 2011; Oury et al., 2013).

OCN is produced predominantly by osteoblasts and is the most abundant, noncollagenous protein found in the mineralized matrix of bone (Hauschka, Lian, & Gallop, 1975). Posttranslational γ -carboxylation of three glutamic acid residues within OCN results in carboxylated OCN (cOCN). However, OCN can be

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2019 The Authors. *Journal of Cellular Physiology* published by Wiley Periodicals, Inc.

carboxylated to varying degrees, allowing for undercarboxylated forms to be present in the circulation (one or two carboxylated residues), and uncarboxylated OCN (ucOCN; no carboxylated residues). ucOCN has previously been regarded as the “active form,” and it has less affinity to bind to hydroxyapatite crystals in bone than cOCN.

Cross-sectional observational studies in humans have demonstrated inverse associations between OCN and systemic inflammatory markers, such as C-reactive protein (CRP) and interleukin-6 (IL-6; Bao et al., 2013; Chen et al., 2013; Liao et al., 2015; Lucey et al., 2013; Pittas, Harris, Eliades, Stark, & Dawson-Hughes, 2009; Sarkar & Choudhury, 2013). A study including 108 newly-diagnosed Type 2 diabetic patients reported that total serum OCN levels were inversely associated with IL-6 and CRP (Sarkar & Choudhury, 2013), these results were also evidenced in an older healthy population ($n = 380$; Pittas et al., 2009). Two large studies in Chinese males ($n = 2,043$ and $n = 1,768$) with metabolic syndrome or normal metabolic state also found an inverse association between total serum OCN and CRP (Bao et al., 2013; Liao et al., 2015). In young obese and overweight women, as well as in postmenopausal women, these findings have also been replicated (Lucey et al., 2013; Chen et al., 2013). It is also known that subclinical inflammation is associated with altered bone metabolism, which further potentiates a role for OCN within the vasculature during inflammation (Ding, Parameswaran, Udayan, Burgess, & Jones, 2008; Kim, Kim, & Sohn, 2010; Koh et al., 2005).

In white-adipose tissue from ucOCN-treated obese mice, several inflammatory genes and transcription factors were down-regulated including tumor necrosis factor (TNF); IL-1 β ; IL-6; chemokine (C-C motif) ligand, Ccl2; caspase 1; and NLR family, pyrin domain containing 3, Nlrp3 (Guedes, Esteves, Morais, Zorn, & Furuya, 2017). Within the same study, ucOCN-treated (20 ng/ml) mouse adipocytes similarly displayed a reduction in expression of inflammatory genes following stimulation with TNF- α (Guedes et al., 2017).

OCN has previously been shown to increase nitric oxide production and prevent free fatty acid-induced apoptosis in human aortic endothelial cells (HAECs); increase angiogenesis in a chick embryo in vivo model; increase proliferation in HAECs and HASMCs; and increase glucose metabolism and promote vascular calcification in OCN-over expressing mouse chondrocytes and vascular smooth muscle cells (Cantatore, Crivellato, Nico, & Ribatti, 2005; Idelevich, Rais, & Monsonogo-Ornan, 2011; Jung et al., 2013; Millar, Anderson, & O'sullivan, 2019). Interestingly, Hill et al. (2014) demonstrated an anti-inflammatory role of both cOCN and ucOCN (20 ng/ml) in isolated rat adipocytes by decreasing TNF- α secretion, while cOCN also decreased IL-6 secretion (Hill et al., 2014). Furthermore, in whole tissue extracts, IL-10 (an anti-inflammatory cytokine) was increased (Hill et al., 2014).

The potential role for OCN in the development of atherosclerosis has been recently extensively reviewed (Tacey et al., 2018). The question remains however whether OCN is a pathological bystander or in fact mediator. To our knowledge, no in vitro studies to date

have investigated the inflammatory role of OCN in human vascular cells. The aim of the current experiments was to investigate the distinct role of ucOCN in HAECs and human aortic smooth muscle cells (HASMCs) after acute and chronic administration, with and without stimulated inflammation. We hypothesized that ucOCN may reduce proinflammatory markers when administered on its own and may reduce the stimulated inflammatory responses of HAECs and HASMCs.

2 | MATERIALS AND METHODS

2.1 | Osteocalcin

Human ucOCN (amino acids 1-49, [Glu17,21,24]) was purchased from AnaSpec Inc. CA (AS-65307). The amino acid sequence of purchased OCN was as follows: Tyr-Leu-Tyr-Gln-Trp-Leu-Gly-Ala-Pro-Val-Pro-Tyr-Pro-Asp-Pro-Leu-Glu-Pro-Arg-Arg-Glu-Val-Cys-Glu-Leu-Asn-Pro-Asp-Cys-Asp-Glu-Leu-Ala-Asp-His-Ile-Gly-Phe-Gln-Glu-Ala-Tyr-Arg-Arg-Phe-Tyr-Gly-Pro-Val. The same batch of ucOCN has been previously shown to be biologically active in vascular cells in our previous work (Millar et al., 2019).

2.2 | Cell culture

HAECs and HASMCs were purchased from PromoCell (UK) and maintained at 37°C in a humidified incubator supplemented with 5% CO₂. Cells were cultured in endothelial cell growth media and smooth muscle cell growth media, respectively, containing 1% penicillin-streptomycin (Sigma-Aldrich, UK) and supplemental mix (PromoCell, UK). In all experiments, cells were used between passages 3 and 5. After experimental treatments, cell media was collected and cells were washed once with phosphate-buffered saline (PBS; pH 7.4; Gibco™). Radioimmunoprecipitation buffer (Sigma-Aldrich) with protease and phosphatase inhibitors (A32959; Thermo Fisher Scientific) was added to lyse the cells and the plates were shaken at 4°C for an hour. The cells were then collected and centrifuged at 14,000g for 5 min at 4°C and cell supernatants were frozen at -80°C.

2.3 | Acute and chronic inflammation

Acute inflammation was induced by treating cells with 10 ng/ml interferon- γ (IFN- γ) or vehicle (ethanol) for 8 hr, followed by addition of 10 ng/ml TNF- α or vehicle (1% ethanol) for 16 hr, as described previously (Alhamoruni, Wright, Larvin, & O'sullivan, 2012). Cells were cotreated with or without ucOCN (10 ng/ml; Hannemann et al., 2013; Hu et al., 2013; Withold, Friedrich, & Degenhardt, 1997) for 24 hr in the acute inflammation protocol. The chronic inflammatory protocol included 8 hr of IFN- γ (5 ng/ml) followed by addition of TNF- α (5 ng/ml). After 48 hr, the media was replaced and treatment repeated up until 144 hr. Cells were treated with or without ucOCN (10 ng/mL). Media was collected at each time point and stored at -80°C until analyzed.

To test whether ucOCN could effect the production of endogenously induced inflammation, in a subset of experiments, lipopolysaccharide (LPS; 10 ng/ml) was added to HAECs and HASMCs for 24 hr with and without ucOCN (10 ng/ml). Media was collected after this time and stored at -80°C until analyzed.

2.4 | Total protein content

A bicinchoninic acid protein assay was performed to quantify the total protein content in the cell lysates collected at the end of the experiments (Smith et al., 1985).

2.5 | Cell signaling assay

The MILLIPLEX MAP 9-plex Multi-Pathway Magnetic Bead Signaling Kit (catalog no. 48-680MAG; Merck Millipore) was performed per the manufacturers' instructions to detect changes in phosphorylated ERK/MAP kinase 1/2 (Thr185/Tyr187), AKT (Ser473), STAT3 (Ser727), JNK (Thr183/Tyr185), p70s6 kinase (Thr412), NFkB (Ser536), STAT5A/B (Tyr694/699), CREB (Ser133), and p38 (Thr180/Tyr182) in cell lysates using the Luminex[®] xMAP[®] technology (48-680MAG; Milliplex[™]; Merck Millipore).

2.6 | Enzyme-linked immunosorbent assays

Total OCN, endothelin pan specific, intracellular adhesion molecule-1 (ICAM-1)/CD54, vascular cell adhesion molecule-1 (VCAM-1)/CD106, total MMP-3, CCL2/monocyte chemoattractant protein-1 (MCP-1), IL-8/CXCL8, and IL-17 DuoSet enzyme-linked immunosorbent assays (ELISA) were performed on cell culture media as per the manufacturer's instructions (catalog no. DY1419; R&D systems, DY1160, DY720, DY809, DY513, DY279, DY208, and DY317). IL-6 ELISA Ready-SET-Go! was performed on cell culture media as per the manufacturer's instructions (catalog no. 88-7066-22; Affymetri; eBioscience).

2.7 | Lactate dehydrogenase activity assay

Lactate dehydrogenase activity (LDH) Colorimetric Assay kit (category no. ab102526; Abcam) was performed on cell media from the chronic inflammation experiment as per manufacturer's instructions.

2.8 | Haematoxylin and eosin staining

After 48 hr of the chronic inflammation protocol, a representative selection of HAECs and HASMCs was washed with PBS and fixed with ice cold methanol/acetone (50:50) for 10 min at room temperature. Fixed cells were then washed with PBS and stained with 0.1% Mayer's haematoxylin and counterstained with 1% eosin Y solution to allow visualization of the nuclei and cytoplasm.

2.9 | Statistical analysis

One-way analysis of variances (ANOVAs) were performed to assess differences in protein secretions and protein phosphorylation after 24 hr. Data were mostly normally distributed and nonparametric *t* tests (Mann-Whitney) were performed in a few cases where the data were not normalized. Results were normalized to protein content. Two-way ANOVAs were performed to detect differences in IL-6 secretion and LDH activity using time and treatment as factors for the chronic inflammation experiment. Multiple comparisons were adjusted for by Dunnett's statistical hypothesis test. All statistical analyses were performed using Prism 7 for Windows (Version 7.01; GraphPad Software Inc.). *p* Values were considered significant at $p < .05$.

3 | RESULTS

3.1 | HAEC responses to acute OCN

No changes in protein secretions (VCAM-1, IL-8, ICAM-1, IL-10, IL-6, or MCP-1) were observed in response to ucOCN in HAECs after 24 hr (Figure 1a-f). Furthermore, total protein content levels did not differ between ucOCN and vehicle (Figure S1a). No intracellular signaling protein phosphorylation (CREB, NFkB, p38, JNK, AKT, ERK, STAT3, STAT5, or p70s6k) was affected by ucOCN (Figure 2a-i).

3.2 | HAEC responses to acute inflammation

Acute inflammation significantly increased the secretion of ICAM-1, VCAM-1, IL-6, IL-8, and MCP-1 ($p < .001$; Figure 1a,b,d-f). IL-10 secretion was not affected (Figure 1c). Phosphorylation of CREB, JNK, p38, ERK, AKT, STAT3, and STAT5 was increased with inflammation (Figure 2a,b,d-f,h,i). Phosphorylation of NFkB and p70s6k was not affected (Figure 2c,g). There were no significant differences in cell signaling (CREB, p38, JNK, ERK, AKT, STAT3, STAT5, NFkB, p70s6k), protein secretion, or total protein content between cells treated with or without ucOCN alongside inflammation after 24 hr.

3.3 | Human aortic endothelial cell responses to chronic OCN

In the chronic experiments, IL-6 secretion increased over time but was not affected by OCN (10 ng/ml) after 48, 96, or 144 hr incubation (Figure 3a). Similarly, no effects of ucOCN were observed on LDH activity, IL-10 secretion, or total protein content (Figure 3c-f).

3.4 | Human aortic endothelial cell responses to chronic inflammation

Because we found no anti-inflammatory effects of ucOCN in the acute inflammatory protocol, we next established whether any effects become apparent after prolonged inflammation and ucOCN treatment. IL-6 secretion increased over time with the inflammatory protocol, with




FIGURE 1 Protein secretion responses in HAECs after acute inflammation. Mean % change relative to vehicle and SEM of secreted cell proteins VCAM-1, ICAM-1, IL-6, endothelin, IL-10, IL-8, and MCP-1 when treated with vehicle, ucOCN (10 ng/mL), an inflammatory protocol (IFN- γ and TNF- α , both 10 ng/ml, 8 hr followed by 16 hr, respectively), or inflammatory protocol and ucOCN. Data were normalized to total protein content. Data were analyzed by one-way ANOVA. **** indicates $p < .001$ compared with vehicle. HAEC, human aortic endothelial cell; ICAM-1, intracellular adhesion molecule-1; IL, interleukin; inflm, inflammatory protocol; MCP-1, monocyte chemoattractant protein-1; ucOCN, uncarboxylated osteocalcin; SEM, standard error of mean; VCAM-1, vascular cell adhesion molecule-1

a peak observed at 48 hr, but ucOCN demonstrated no additional effects (Figure 3b). Nicotinamide adenine dinucleotide H concentration was measured as an indicator of LDH activity. LDH activity increased sharply following 48 hr of inflammation in HAECs, following a similar

trajectory as with IL-6 production (Figure 4d). There were no differences detected between cells treated with ucOCN alongside inflammation and those treated with inflammation alone (Figure 3d). IL-10 secretion at 144 hr did not differ between treatment groups




FIGURE 2 Intracellular signaling responses in HAECs. Luminex[®] xMAP[®] technology was used to detect changes in phosphorylated CREB (pS133), JNK (pT183/pY185), NFkB (pS536), p38 (pT180/pY182), ERK (pT185/pY187), Akt (pS473), p70 S6k (pT412), STAT3 (pS727), and STAT5A/B (pY694/699; 48-680MAG; Milliplex[™]; Merck Millipore) in cell lysates when treated with vehicle or ucOCN (10 ng/ml) with and without inflammatory stimulus (IFN- γ and TNF- α) for 24 hr. Data were analyzed by one-way ANOVA with multiple comparisons to vehicle corrected for by Dunnett's test. Data are given as means with error bars representing SEM. *denotes a significant difference compared with vehicle (* $p < .05$, ** $p < .01$, *** $p < .005$, **** $p < .001$). HAEC, human aortic endothelial cell; IFN- γ , interferon- γ ; inflm, inflammatory protocol (8 hr of IFN- γ 10 ng/ml followed by addition of TNF- α 10 ng/mL for 16 hr); SEM, standard error of mean; TNF- α , tumor necrosis factor- α ; ucOCN, uncarboxylated osteocalcin

(Figure 3e). Total protein content (as an indicator of cell death) was decreased with the inflammatory protocol ($p < .05$) but no additional effect of ucOCN was observed (Figure 3f).

After 48 hr of inflammatory protocol, HAECs adopted an activated, spindle-shaped morphology as opposed to the characteristic cobblestone appearance of noninflamed cells (Figure S2).

3.5 | HASMC responses to acute OCN

No changes were observed in the secretion of ICAM-1, IL-8, MCP-1, IL-17, MMP-3, or endothelin after 24 hr treatment with ucOCN

compared with vehicle (Figure 4a–f). Phosphorylation of p38, ERK, AKT, CREB, JNK, NFkB, p70s6k, STAT3, and STAT5 was not affected by ucOCN exposure (Figure 5a–i). Total protein content did not differ with ucOCN treatment (Figure S1B).

3.6 | HASMC responses to acute inflammation

ICAM-1, MCP-1, IL-8, and MMP-3 secretion was significantly increased by inflammation in HASMCs compared to vehicle ($p < .001$), however no effect of ucOCN was observed (Figure 4a–c,e). There was no significant effect of inflammation




FIGURE 3 IL-6 secretion, LDH activity, IL-10 secretion, and total protein content in HAECs after chronic inflammation with or without OCN. The effects of ucOCN on secretion of IL-6 without inflammation (a) and with inflammation (b) after 0, 48, 96, and 144 hr. The effect of ucOCN on LDH activity was measured by NADH concentration without inflammation (c) and with inflammation (d) after 0, 48, 96, and 144 hr. IL-10 secretion was measured after 144 hr (e). Total protein content was measured by BCA assay at the end of the experiment. ucOCN (10 ng/ml) was added every 48 hr. Total $n = 9$ from three experimental repeats. Data are given as means with error bars representing SEM. *denotes a significant difference compared with vehicle, analyzed by one-way ANOVA (** $p < .01$, *** $p < .005$). ANOVA, analysis of variance; HAEC, human aortic endothelial cell; IL, interleukin; IFN- γ , interferon- γ ; inflm, inflammatory protocol (8 hr of IFN- γ followed by addition of TNF- α ; both 5 ng/ml); LDH, lactate dehydrogenase; NADH, nicotinamide adenine dinucleotide H; SEM, standard error of mean; TNF- α , tumor necrosis factor- α ; ucOCN, uncarboxylated osteocalcin

nor OCN on endothelin production or IL-17 secretion (Figure 4d,f). Phosphorylation of CREB, p38, ERK, and STAT3 was increased with inflammation but no additional effect of ucOCN was observed (Figure 5a-i).

3.7 | HASMC cell responses to chronic OCN

In the chronic experiments, IL-6 secretion increased over time but was not affected by OCN after 48, 96, or 144 hr incubation (Figure 6a). LDH activity increased over time, with no additional effect of ucOCN observed (Figure 6c). Similarly, IL-10 secretion and total protein content was not affected by OCN after 144 hr (Figure 6e,f).

3.8 | HASMC responses to chronic inflammation

Following chronic exposure to inflammation, IL-6 secretion increased in HASMCs until 96 hr and remained elevated (Figure 6b). Addition of ucOCN did not affect IL-6 levels. Similarly, LDH activity was increased with inflammation but was not affected by the coadministration of ucOCN (Figure 6d). IL-10 and total protein content were not altered by the inflammation protocol nor ucOCN (Figure 6e,f).

3.9 | OCN secretion is not detected following chronic inflammation

Finally, total OCN was measured in cell media after chronic inflammation (144 hr) to test whether inflammation induced OCN




FIGURE 4 Protein secretions in HASMCs after acute inflammation. Mean % change relative to vehicle and SEM of secreted cell proteins ICAM-1, MCP-1, endothelin, MMP-3, IL-17, and IL-8 when treated with vehicle, ucOCN (10 ng/ml), an inflammatory protocol (IFN- γ and TNF- α , both 10 ng/ml, 8 hr followed by 16 hr, respectively), or inflammatory protocol and ucOCN. Data were normalized to total protein content. Data were analyzed by one-way ANOVA. ****denotes a significant difference compared with vehicle ($p < .005$). ANOVA, analysis of variance; HASMC, human aortic smooth muscle cell; ICAM-1, intracellular adhesion molecule-1; IL, interleukin; IFN- γ , interferon- γ ; MCP-1, monocyte chemoattractant protein-1; MMP-3, matrix metalloproteinase; SEM, standard error of mean; TNF- α , tumor necrosis factor- α ; ucOCN, uncarboxylated osteocalcin; inflm, inflammatory protocol




FIGURE 5 Intracellular signaling responses in HASMCs. Luminex[®] xMAP[®] technology was used to detect changes in phosphorylated CREB (pS133), JNK (pT183/pY185), NFKB (pS536), p38 (pT180/pY182), ERK (pT185/pY187), Akt (pS473), p70 S6K (pT412), STAT3 (pS727), and STAT5A/B (pY694/699; 48-680MAG; Milliplex[™]; Merck Millipore) in cell lysates when treated with vehicle or ucOCN (10 ng/ml) with and without inflammatory stimulus (IFN- γ and TNF- α) for 24 hr. Data were analyzed by one-way ANOVA with multiple comparison to vehicle corrected for by Dunnett's test. Data are given as means with error bars representing SEM. *denotes a significant difference compared with vehicle (* $p < .05$, ** $p < .01$, *** $p < .005$, **** $p < .001$). ANOVA, analysis of variance; HASMC, human aortic smooth muscle cell; IFN- γ , interferon- γ ; inflm, inflammatory protocol (8 hr of IFN- γ 10 ng/ml followed by addition of TNF- α 10 ng/ml for 16 hr); SEM, standard error of mean; TNF- α , tumor necrosis factor- α ; ucOCN, uncarboxylated osteocalcin

secretion. OCN was not detectable in media from HAECs nor HASMCs at the end of the experiments (data not shown).

3.10 | OCN does not prevent LPS-induced inflammation

As ucOCN did not affect inflammation induced by exogenously added cytokines, we then tested whether LPS-induced inflammation elicited a different response. Acute LPS treatment (24 hr) significantly increased secretion of VCAM-1 and ICAM-1 in HAECs and HASMCs,

respectively, compared with control (Figure 1c,d). In both HAECs and HASMCs, there were no differences in VCAM-1 or ICAM-1 secretion respectively between LPS treated and LPS + ucOCN-treated cells (Figure 1c,d).

4 | DISCUSSION

OCN is synthesized predominantly by osteoblast cells, which acts as an extra-skeletal hormone known to effect insulin sensitivity, insulin




FIGURE 6 IL-6 secretion, LDH activity, IL-10 secretion, and total protein content in HASMCs after chronic inflammation with or without OCN. The effects of ucOCN on secretion of IL-6 without inflammation (a) and with inflammation (b) after 0, 48, 96, and 144 hr. The effect of ucOCN on LDH activity was measured by NADH concentration without inflammation (c) and with inflammation (d) after 0, 48, 96, and 144 hr. IL-10 secretion was measured after 144 hr (e). Total protein content was measured by BCA assay at the end of the experiment. ucOCN (10 ng/ml) was added every 48 hr. Total $n = 9$ from three experimental repeats. BCA, bicinchoninic acid; HASMC, human aortic smooth muscle cell; LDH, lactate dehydrogenase; IFN- γ , interferon- γ ; IL, interleukin; inflm, inflammatory protocol (8 hr of IFN- γ followed by addition of TNF- α ; both 5 ng/ml); NADH, nicotinamide adenine dinucleotide H; TNF- α , tumor necrosis factor- α ; ucOCN, uncarboxylated osteocalcin

secretion, energy metabolism, cognition, and fertility (Lee et al., 2007; Oury et al., 2011; Oury et al., 2013). Population-based cross-sectional studies have reported a negative or inverse association between OCN and markers of systemic inflammation, such as IL-6 and CRP, suggesting a potential anti-inflammatory role (Chen et al., 2013; Kim et al., 2010; Liao et al., 2015; Pittas et al., 2009; Sarkar & Choudhury, 2013; Schett et al., 2006). However, the *in vitro* effects of OCN on inflammation within the vasculature have not yet been described. Therefore, we decided to try and reveal mechanisms of action of ucOCN during inflammation, as well as investigate the effects of acute and chronic administration of ucOCN on its own.

Here, we demonstrate for the first time using two human vascular cell types that ucOCN does not regulate or alter the inflammatory responses of HAECs or HASMCs during acute inflammation. Our observational results presented here do not reflect an anti-inflammatory

role of OCN suggested by some cross-sectional, human studies, at least within the experimental models used. ucOCN did not affect inflammatory cytokine production nor inflammatory signaling pathways in HASMCs and HAECs. This may be representative of the nonbinary nature of OCN which cannot be limited to having a positive or negative influence overall in vascular pathology and physiology in humans. It may transpire that OCN plays a protective role in the later stages of atherosclerosis and have more of an impact on the process of calcification.

As no overall anti-inflammatory effects of OCN were reported in the acute inflammatory protocol, we next established whether any effects of OCN become apparent after prolonged inflammation and OCN treatment. Pro- and anti-inflammatory cytokine production as well as LDH activity were not affected by chronic ucOCN treatment, alone or in combination with the inflammatory protocol. Lastly, we further showed that in another model of inflammation using LPS,

ucOCN did not affect VCAM-1 or ICAM-1 secretion in HAECs and HASMCs respectively. An interesting observation from this study was the transition of HAECs after 48 hr of chronic inflammation to an activated morphology that is characteristic of endothelial cells under shear stress and inflammation, representative of the early stages of atherosclerosis (Hunt & Jurd, 1998). This corresponds to the upregulation of activated endothelium markers VCAM-1 and ICAM-1 reported at 24 hr. Additionally we demonstrated that inflammation does not induce secretion of OCN itself.

The role of OCN in cardiovascular disease has had conflicting results in humans. Some longitudinal studies have reported U-shaped associations between OCN and cardiovascular-related or all-cause mortality (Kanazawa, Yamaguchi, & Sugimoto, 2011; Yang et al., 2017; Yeap et al., 2012). However, other studies have found no association between OCN and cardiovascular disease risk (Holvik et al., 2014; Hwang et al., 2015). A systematic review and meta-analysis on the association between OCN and markers of atherosclerosis or calcification in humans found no conclusive relationship (Millar, Patel, Anderson, England, & O'Sullivan, 2017). Although a causal role of ucOCN in vascular inflammation was not identified in this study, further work should explore whether OCN may influence systemic inflammation via directly acting on immune cells. Additionally, the exploration of the possible biological activity of other forms of circulating OCN should not be disregarded. We recognize that phenotypic drift occurs with the isolation and subculturing of cells and we acknowledge this as a limitation of the current study. Further work should explore both the effect of OCN on intact vessels and on cells from different vascular beds including resistance and conduit vessels.

In conclusion, in our models of vascular inflammation, OCN did not display a role in the direct inflammatory responses of primary subcultured HAECs or smooth muscle cells in either an acute or chronic setting.

FUNDING INFORMATION

This work was supported by the Biotechnology and Biological Sciences Research Council (grant number: BB/M008770/1).

CONFLICT OF INTERESTS

The authors declare that there are no conflict of interests.

AUTHOR CONTRIBUTIONS

S. O. S. and S. M. designed the experiments which were carried out by S. M. and I. Z. (I. Z. carried out the IL-6 and L. D. H. measurements for the chronic experiments). S. A. and S. O. S supervised the project. S. M. wrote the manuscript with input from S. O. S. and S. A.

DATA AVAILABILITY STATEMENT

The data that support the findings of this study are available from the corresponding author upon reasonable request.

ORCID

Sophie A. Millar  <http://orcid.org/0000-0003-1318-7671>

REFERENCES

- Alhaboruni, A., Wright, K., Larvin, M., & O'Sullivan, S. (2012). Cannabinoids mediate opposing effects on inflammation-induced intestinal permeability. *British Journal of Pharmacology*, *165*, 2598–2610.
- Bao, Y., Ma, X., Yang, R., Wang, F., Hao, Y., Dou, J., ... Jia, W. (2013). Inverse relationship between serum osteocalcin levels and visceral fat area in Chinese men. *The Journal of Clinical Endocrinology & Metabolism*, *98*, 345–351.
- Bessueille, L., & Magne, D. (2015). Inflammation: A culprit for vascular calcification in atherosclerosis and diabetes. *Cellular and Molecular Life Science*, *72*, 2475–2489.
- Cantatore, F. P., Crivellato, E., Nico, B., & Ribatti, D. (2005). Osteocalcin is angiogenic in vivo. *Cell Biology International*, *29*, 583–585.
- Chen, L., Li, Q., Yang, Z., Ye, Z., Huang, Y., He, M., ... Hu, R. (2013). Osteocalcin, glucose metabolism, lipid profile and chronic low-grade inflammation in middle-aged and elderly Chinese. *Diabetic Medicine*, *30*, 309–317.
- Ding, C., Parameswaran, V., Udayan, R., Burgess, J., & Jones, G. (2008). Circulating levels of inflammatory markers predict change in bone mineral density and resorption in older adults: A longitudinal study. *The Journal of Clinical Endocrinology & Metabolism*, *93*, 1952–1958.
- Guedes, J. A. C., Esteves, J. V., Morais, M. R., Zorn, T. M., & Furuya, D. T. (2017). Osteocalcin improves insulin resistance and inflammation in obese mice: Participation of white adipose tissue and bone. *Bone*, *115*, 68–82.
- Hannemann, A., Friedrich, N., Spielhagen, C., Rettig, R., Ittermann, T., Nauck, M., & Wallaschofski, H. (2013). Reference intervals for serum osteocalcin concentrations in adult men and women from the study of health in Pomerania. *BMC Endocrine Disorders*, *13*, 11.
- Hauschka, P. V., Lian, J. B., & Gallop, P. M. (1975). Direct identification of the calcium-binding amino acid, gamma-carboxyglutamate, in mineralized tissue. *Proceedings of the National Academy of Sciences*, *72*, 3925–3929.
- Hill, H. S., Grams, J., Walton, R. G., Liu, J., Moellering, D. R., & Garvey, W. T. (2014). Carboxylated and uncarboxylated forms of osteocalcin directly modulate the glucose transport system and inflammation in adipocytes. *Hormone and Metabolic Research*, *46*, 341–347.
- Holvik, K., Van Schoor, N. M., Eekhoff, E. M. W., Den Heijer, M., Deeg, D. J. H., Lips, P., & De Jongh, R. (2014). Plasma osteocalcin levels as a predictor of cardiovascular disease in older men and women: A population-based cohort study. *European Journal of Endocrinology*, *171*, 161–170.
- Hu, W. W., Zhang, Z., He, J. W., Fu, W. Z., Wang, C., Zhang, H., ... Zhang, Z. L. (2013). Establishing reference intervals for bone turnover markers in the healthy shanghai population and the relationship with bone mineral density in postmenopausal women. *International Journal of Endocrinology*, *2013*, 513925.
- Hunt, B. J., & Jurd, K. M. (1998). Endothelial cell activation. *BMJ*, *316*, 1328–1329.
- Hutcheson, J. D., & Aikawa, E. (2014). Optical molecular imaging of inflammation and calcification in atherosclerosis. In Aikawa, E. (Ed.), *Cardiovascular Imaging* (pp. 107–120). Cham: Springer.
- Hwang, Y. C., Kang, M., Cho, i J., Jeong, i K., Ahn, k J., Chung, H. Y., & Lee, M. K. (2015). Association between the circulating total osteocalcin level and the development of cardiovascular disease in middle-aged men: A mean 8.7-year longitudinal follow-up study. *Journal of Atherosclerosis and Thrombosis*, *22*, 136–143.
- Idelevich, A., Rais, Y., & Monsonego-Ornan, E. (2011). Bone gla protein increases HIF-1 α -dependent glucose metabolism and induces

- cartilage and vascular calcification. *Arteriosclerosis, Thrombosis, and Vascular Biology*, 31, e55–e71.
- Jung, C. H., Lee, W. J., Hwang, J. Y., Lee, M. J., Seol, S. M., Kim, Y. M., ... Park, J. Y. (2013). The preventive effect of uncarboxylated osteocalcin against free fatty acid-induced endothelial apoptosis through the activation of phosphatidylinositol 3-kinase/Akt signaling pathway. *Metabolism*, 62, 1250–1257.
- Kanazawa, I., Yamaguchi, T., & Sugimoto, T. (2011). Relationship between bone biochemical markers versus glucose/lipid metabolism and atherosclerosis; a longitudinal study in Type 2 diabetes mellitus. *Diabetes Research and Clinical Practice*, 92, 393–399.
- Kim, M., Kim, H., & Sohn, C. (2010). Relationship between vitamin K status, bone mineral density, and hs-CRP in young Korean women. *Nutrition Research and Practice*, 4, 507–514.
- Koh, J. M., Khang, Y. H., Jung, C. H., Bae, S., Kim, D. J., Chung, Y. E., & Kim, G. S. (2005). Higher circulating hsCRP levels are associated with lower bone mineral density in healthy pre- and postmenopausal women: Evidence for a link between systemic inflammation and osteoporosis. *Osteoporosis International*, 16, 1263–1271.
- Lee, N. K., Sowa, H., Hinoi, E., Ferron, M., Ahn, J. D., Confavreux, C., ... Karsenty, G. (2007). Endocrine regulation of energy metabolism by the skeleton. *Cell*, 130, 456–469.
- Liao, M., Huang, L., Mao, Y., Jiang, Y., Yao, Z., Lin, X., ... Mo, Z. (2015). Serum osteocalcin is associated with inflammatory factors in metabolic syndrome: a population-based study in chinese males. *Mediators of Inflammation*, 2015, 683739.
- Lucey, A. J., Paschos, G. K., Thorsdottir, I., Martínez, J. A., Cashman, K. D., & Kiely, M. (2013). Young overweight and obese women with lower circulating osteocalcin concentrations exhibit higher insulin resistance and concentrations of C-reactive protein. *Nutrition Research*, 33, 67–75.
- Mazzini, M. J., & Schulze, P. C. (2006). Proatherogenic pathways leading to vascular calcification. *European Journal of Radiology*, 57, 384–389.
- Millar, S. A., Anderson, S. I., & O'Sullivan, S. E. (2019). Human vascular cell responses to the circulating bone hormone osteocalcin. *Journal of Cellular Physiology*, 234, 21039–21048.
- Millar, S. A., Patel, H., Anderson, S. I., England, T. J., & O'Sullivan, S. E. (2017). Osteocalcin, vascular calcification, and atherosclerosis: A systematic review and meta-analysis. *Frontiers in Endocrinology*, 8, 183.
- Oldknow, K. J., Macrae, V. E., & Farquharson, C. (2015). Endocrine role of bone: Recent and emerging perspectives beyond osteocalcin. *Journal of Endocrinology*, 225, R1–R19.
- Oury, F., Khrimian, L., Denny, C. A., Gardin, A., Chamouni, A., Goeden, N., ... Karsenty, G. (2013). Maternal and offspring pools of osteocalcin influence brain development and functions. *Cell*, 155, 228–241.
- Oury, F., Sumara, G., Sumara, O., Ferron, M., Chang, H., Smith, C. E., ... Karsenty, G. (2011). Endocrine regulation of male fertility by the skeleton. *Cell*, 144, 796–809.
- Pittas, A. G., Harris, S. S., Eliades, M., Stark, P., & Dawson-Hughes, B. (2009). Association between serum osteocalcin and markers of metabolic phenotype. *Journal of Clinical Endocrinology and Metabolism*, 94, 827–832.
- Ross, R. (1999). Atherosclerosis—an inflammatory disease. *New England Journal of Medicine*, 340, 115–126.
- Sage, A. P., Tintut, Y., & Demer, L. L. (2010). Regulatory mechanisms in vascular calcification. *Nature Reviews Cardiology*, 7, 528–536.
- Sarkar, P. D., & Choudhury, A. B. (2013). Relationships between serum osteocalcin levels versus blood glucose, insulin resistance and markers of systemic inflammation in central Indian type 2 diabetic patients. *European Review for Medical and Pharmacological Sciences*, 17, 1631–1635.
- Schett, G., Kiechl, S., Weger, S., Pederiva, A., Mayr, A., Petrangeli, M., ... Willeit, J. (2006). High-sensitivity C-reactive protein and risk of nontraumatic fractures in the Bruneck study. *Archives of Internal Medicine*, 166, 2495–2501.
- Smith, P. K., Krohn, R. I., Hermanson, G. T., Mallia, A. K., Gartner, F. H., Provenzano, M. D., ... Klenk, D. C. (1985). Measurement of protein using bicinchoninic acid. *Analytical Biochemistry*, 150, 76–85.
- Tacey, A., Qaradakh, T., Brennan-Speranza, T., Hayes, A., Zulli, A., & Levinger, I. (2018). Potential role for osteocalcin in the development of atherosclerosis and blood vessel disease. *Nutrients*, 10, 1426.
- Withold, W., Friedrich, W., & Degenhardt, S. (1997). Serum bone alkaline phosphatase is superior to plasma levels of bone matrix proteins for assessment of bone metabolism in patients receiving renal transplants. *Clinica Chimica Acta*, 261, 105–115.
- Yang, S. W., Hennessy, R. R., Khosla, S., Lennon, R., Loeffler, D., Sun, T., ... Lerman, A. (2017). Circulating osteogenic endothelial progenitor cell counts: New biomarker for the severity of coronary artery disease. *International Journal of Cardiology*, 227, 833–839.
- Yeap, B. B., Chubb, S. A. P., Flicker, L., Mccauley, K. A., Ebeling, P. R., Hankey, G. J., ... Norman, P. E. (2012). Associations of total osteocalcin with all-cause and cardiovascular mortality in older men. The Health In Men Study. *Osteoporosis International*, 23, 599–606.

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section.

How to cite this article: Millar SA, Zala I, Anderson SI, O'Sullivan SE. Osteocalcin does not influence acute or chronic inflammation in human vascular cells. *J Cell Physiol*. 2020;235:3414–3424. <https://doi.org/10.1002/jcp.29231>