ERRATUM Open Access

pISSN 1738-6586 / eISSN 2005-5013 / J Clin Neurol 2020;16(1):181-181 / https://doi.org/10.3988/icn.2020.16.1.181



Erratum to: Elevated Serum Uric Acid in Benign Convulsions with Mild Gastroenteritis in Children

II Han Yoo^{a,b}, Woojoong Kim^c, Jaeso Cho^d, Hunmin Kim^a, Byung Chan Lim^c, Hee Hwang^a, Jong-Hee Chae^c, Jieun Choi^e, Ki Joong Kim^c

J Clin Neurol 2019;15(4):496-501 / https://doi.org/10.3988/jcn.2019.15.4.496

Unfortunately, the authors confirmed that some of the data was missing from categories 'Acute gastroenteritis' and 'Febrile seizure' of Table 2 in the original version.

In Table 2, the author wishes to be modified as follows:

Table 2. Demographics of patients with benign CwG, acute gastroenteritis, and febrile seizure

	CwG (n=154)	Acute gastroenteritis (n=2,938)	Febrile seizure (n=154)
Sex, male:female	63:91	1,632:1,306	64:90
Age at onset, years	1.70 (0.73-3.19)	1.71 (0.65–3.25)	1.71 (0.71–3.49)
Serum uric acid, mg/dL	9.79±2.16	6.04±2.38	4.30±1.02
HCO ₃ -, mg/dL	15.72±2.68	16.64±2.69	19.40±1.39

Data are n, mean (range), or mean \pm SD values. CwG: convulsion associated with mild gastroenteritis.

^aDepartment of Pediatrics, Seoul National University Bundang Hospital, Seongnam, Korea

^bDepartment of Pediatrics, St. Vincent's Hospital, College of Medicine, The Catholic University of Korea, Suwon, Korea

^cPediatric Clinical Neuroscience Center, Department of Pediatrics, Seoul National University Children's Hospital, Seoul, Korea

^dDepartment of Biomedical Sciences, College of Medicine, Seoul National University, Seoul, Korea

^eDepartment of Pediatrics, SMG-SNU Boramae Medical Center, Seoul, Korea