

ErbB-3 BINDING PROTEIN 1 Regulates Translation and Counteracts RETINOBLASTOMA RELATED to Maintain the Root Meristem¹[CC-BY]

Ansul Lokdarshi,^{a,2} Csaba Papdi,^{b,2,3} Aladár Pettkó-Szandtner,^c Stefan Dorokhov,^b Ben Scheres,^{d,4} Zoltán Magyar,^e Albrecht G. von Arnim,^{a,f} László Bögre,^{b,5} and Beatrix M. Horváth^{b,d,5,6}

^aDepartment of Biochemistry & Cellular and Molecular Biology, University of Tennessee, Knoxville, Tennessee 37996

^bDepartment of Biological Sciences, Centre for Systems and Synthetic Biology, Royal Holloway, University of London, Egham Hill, Egham TW20 0EX, United Kingdom

^cLaboratory of Proteomics Research, Biological Research Centre, POB 521, H-6701 Szeged, Hungary

^dDepartment of Biology, Utrecht University, 3584 CH Utrecht, The Netherlands

^eInstitute of Plant Biology, Biological Research Centre, POB 521, H-6701 Szeged, Hungary

^fUT-ORNL Graduate School of Genome Science and Technology, Knoxville, Tennessee 37996

ORCID IDs: 0000-0003-2264-2701 (A.L.); 0000-0003-0640-6098 (C.P.); 0000-0003-2968-8542 (S.D.); 0000-0001-5400-9578 (B.S.); 0000-0001-8376-7220 (Z.M.); 0000-0003-3472-3357 (A.G.v.A.); 0000-0002-6083-992X (B.M.H.).

The ErbB-3 BINDING PROTEIN 1 (EBP1) drives growth, but the mechanism of how it acts in plants is little understood. Here, we show that EBP1 expression and protein abundance in *Arabidopsis* (*Arabidopsis thaliana*) are predominantly confined to meristematic cells and are induced by sucrose and partially dependent on TARGET OF RAPAMYCIN (TOR) kinase activity. Consistent with being downstream of TOR, silencing of EBP1 restrains, while overexpression promotes, root growth, mostly under sucrose-limiting conditions. Inducible overexpression of RETINOBLASTOMA RELATED (RBR), a sugar-dependent transcriptional repressor of cell proliferation, depletes meristematic activity and causes precocious differentiation, which is attenuated by EBP1. To understand the molecular mechanism, we searched for EBP1- and RBR-interacting proteins by affinity purification and mass spectrometry. In line with the double-stranded RNA-binding activity of EBP1 in human (*Homo sapiens*) cells, the overwhelming majority of EBP1 interactors are part of ribonucleoprotein complexes regulating many aspects of protein synthesis, including ribosome biogenesis and mRNA translation. We confirmed that EBP1 associates with ribosomes and that EBP1 silencing hinders ribosomal RNA processing. We revealed that RBR also interacts with a set of EBP1-associated nucleolar proteins as well as factors that function in protein translation. This suggests EBP1 and RBR act antagonistically on common processes that determine the capacity for translation to tune meristematic activity in relation to available resources.

Cell growth is generally required for cell proliferation, and although these two processes are separately regulated, they are closely connected and coordinated (Fox et al., 2018). The central conserved regulator of the cell cycle is the cyclin-dependent kinase (CDK), the activity of which (1) sets the unidirectional alternation of cell cycle phases, (2) enables cells to tune the cell cycle to match with developmental and environmental cues, or (3) gives a signal to exit from the cycle to allow cellular differentiation (De Veylder et al., 2007). A pivotal target of CDK that balances cell proliferation with differentiation is the RETINOBLASTOMA RELATED (RBR) protein, whose phosphorylation lifts the repression on the cell-cycle regulatory E2F transcription factors (Magyar et al., 2016). In plants, modulation in the levels of these cell cycle regulators, such as the overexpression of Cyclin D3;1 (Dewitte et al., 2003), silencing of the RBR protein (Gutzat et al., 2011), or overexpression of the E2FB transcription factor (Magyar et al., 2005), results in more but smaller cells and stunted and developmentally arrested plants. This led to the realization

that acceleration of cell proliferation does not necessarily lead to increased plant growth.

The TARGET OF RAPAMYCIN (TOR) kinase pathway is regarded as the central regulator of cell growth in connection with light and sugars by generally boosting anabolic and repressing catabolic processes (Dobrenel et al., 2016a; González and Hall, 2017). Protein synthesis, being an energy and resource demanding cellular process, is one of the major cellular functions controlled by TOR signaling (González and Hall, 2017). S6 kinase (S6K) is a key downstream component of TOR signaling that regulates translation and ribosome biogenesis (Deprout et al., 2007; Schepetilnikov et al., 2013). Thus growth-stimulatory factors such as light, sugar, and auxin activate TOR and affect the translation apparatus in different ways (Liu et al., 2013; Dobrenel et al., 2016b; Chen et al., 2018; Enganti et al., 2018; Schepetilnikov and Ryabova, 2018). Correspondingly, elevated TOR expression boosts plant growth in *Arabidopsis* (*Arabidopsis thaliana*; Deprout et al., 2007) and increases yield

in crop plants, specifically in limiting conditions (Bakshi et al., 2019). On the other hand, TOR silencing or chemical inhibition of TOR activity results in reduced growth (Deprost et al., 2007; Montané and Menand, 2013, 2019).

The growth regulatory TOR signaling drives cell proliferation by controlling the selective translation of key cell-cycle regulatory proteins, such as CLN3 G1 cyclin in yeast (*Saccharomyces cerevisiae*; Barbet et al., 1996) or Cyclin E in human (*Homo sapiens*) cells (Dowling et al., 2010). In plants, however, there appears to be an even more direct connection between growth and cell cycle control, as TOR phosphorylates and thus stimulates E2FA and E2FB activities (Xiong et al., 2013; Wu et al., 2019). More recent research shows that the *YET ANOTHER KINASE1* (*AtYAK1*) gene, encoding a member of the dual-specificity Tyr phosphorylation-regulated kinase family, inhibits growth and cell proliferation by up-regulating the SIAMESE-RELATED (SMR) proteins when TOR activity is reduced (Barrada et al., 2019; Forzani et al., 2019). Moreover, the principal downstream effector of TOR, AtS6K1, interacts with RBR and facilitates its nuclear localization, which is important to impose repression on cell proliferation in sucrose limiting conditions (Henriques et al., 2013). Finally, S6K1, E2FB, and RBR are interconnected through negative feedback loops that might be important to toggle between cell proliferation and quiescence (Henriques et al., 2010).

¹This work was supported by the National Science Foundation (NSF) (grants MCB-1546402 and IOS-1456988), National Institutes of Health (grant R15 GM129672), and the Donald L Akers Jr Faculty Enrichment Fellowship (to A.G.v.A.); the European Commission (EC) (Marie-Curie IEF fellowship, FP7-PEOPLE-2012-IEF-330789 to B.M.H.; FP7-PEOPLE-2012-IEF.330713 to C.P.); the Netherlands Organization for Scientific Research (Spinoza grant to B.S.); Biotechnology and Biological Sciences Research Council (BBSRC) (BBSRC-NSF grant BB/M025047/1 to C.P. and L.B.); and Magyar Tudományok Akadémia (Hungarian Academy of Sciences) (GINOP-2.3.2-15-2016-00032 to A.P.-S. and GINOP-2.3.2-15-2016-00001 to A.P.-S. and Z.M.).

²These authors contributed equally to this work.

³Present address: Institut de Biologie Moléculaire des Plantes, Centre National de la Recherche Scientifique, UPR 2357, Université de Strasbourg, Strasbourg, France.

⁴Present address: Department of Plant Sciences, Wageningen University Research Centre, 6708 PB Wageningen, The Netherlands.

⁵Senior authors.

⁶Author for contact: beatrixmhorvath@gmail.com.

The author responsible for distribution of materials integral to the findings presented in this article in accordance with the policy described in the Instructions for Authors (www.plantphysiol.org) is: Beatrix M. Horváth (beatrixmhorvath@gmail.com).

B.M.H., A.L., C.P., A.P.-S., and Z.M. designed and performed the experiments; A.L., S.D., A.G.v.A., L.B. and B.M.H. analyzed the data; B.S. shared material and gave general support; A.G.v.A., L.B., and B.M.H. conceived and designed the research plan, and wrote the manuscript; all authors saw and commented on the manuscripts.

^[CC-BY]Article free via Creative Commons CC-BY 4.0 license.

www.plantphysiol.org/cgi/doi/10.1104/pp.19.00805

The ErbB-3 BINDING PROTEIN 1 (EBP1) came to the forefront of interest in plants when it was found that it is rapidly induced during early stages of tuber formation, stimulated by sugar in potato (*Solanum tuberosum*; Horváth et al., 2006). Overexpression or silencing of EBP1 in potato could dose-dependently tune leaf and tuber growth and thus potato yield without any obvious developmental abnormalities (Horváth et al., 2006). Studying the cellular basis how EBP1 affects organ growth showed that it boosts cell proliferation in meristematic cells, resulting in more but smaller cells, which correlated with elevated expression of cell cycle regulators. In postmitotic cells, EBP1 enhances cell growth leading to a larger final cell size. On the molecular level EBP1 represses RBR protein abundance, whereas RBR negatively influences the EBP1 level. This antagonism suggested that EBP1 provides an important link between a growth driver and cell cycle regulation (Horváth et al., 2006). The initial finding that EBP1 enhances growth was verified for a number of *EBP1* orthologs identified in different species (Cao et al., 2009; Cheng et al., 2016; Wang et al., 2016); however, another report described an opposite effect (Li et al., 2016). Additionally, *EBP1* expression is induced upon abiotic stress, and when *EBP1* levels are elevated, it can confer stress tolerance (Cao et al., 2009; Cheng et al., 2016). In line with being a positive regulator of growth, in a maize hybrid displaying hybrid vigor, *EBP1* is expressed in an over dominant fashion (Wang et al., 2016). In a recent study, EBP1 was identified as an interactor with the FERONIA receptor-like kinase with a major function in regulating cell expansion. FERONIA directly phosphorylates EBP1, leading to its nuclear localization to regulate genes responsive to FERONIA signaling (Li et al., 2018).

A clue to understand *EBP1*'s molecular function in plants was its remarkable coexpression with genes involved in the regulation of protein synthesis in proliferating cells, the so-called *ribi* regulon (Horváth et al., 2006). In line with this, several additional supporting lines of evidence have been published. First, *EBP1* expression is rapidly induced when growth is stimulated in the shoot meristem of dark-grown etiolated seedlings upon light exposure, correlating with the induction of genes involved in protein synthesis (López-Juez et al., 2008; Mohammed et al., 2018). Second, EBP1 is present in nucleoli in both animals (Squatrito et al., 2004; Liu et al., 2006) and in plants (Pendle et al., 2005), as evidenced by a carboxy-terminal Lys-rich motif (Karlsson et al., 2016) and an N-terminal nucleolar localization signal (Squatrito et al., 2004; Ko et al., 2016). Third, human EBP1 (HsEBP1) has a double-stranded RNA binding activity (Squatrito et al., 2004). It binds to rRNAs and ribosomes (Squatrito et al., 2006) and also supports ribosome biogenesis by interacting with the Pol I transcription factor TIF-IA (Nguyen le et al., 2015; Nguyen et al., 2019) and possibly with nucleolar nucleophosmin (Okada et al., 2007). Fourth, HsEbp1 also stabilizes several mRNAs by binding their 3'UTRs (Bose et al., 2006; Zhou et al., 2010; Pisapia et al., 2015).

Finally, EBP1 has been implicated in translational control in *Leishmania major* (Norris-Mullins et al., 2014) and, repeatedly, in gene specific translation in cap-independent translation initiation of RNA viruses (Pilipenko et al., 2000) on the androgen receptor mRNA (Zhou et al., 2010; Zhou et al., 2011) by affecting eIF2 α phosphorylation (Squatrito et al., 2006).

To gain insight into the growth-promoting function of EBP1, it is paramount to identify the molecular partners. To this end, we performed mass spectrometry to identify EBP1 interactors, which uncovered proteins with a broad range of RNA binding and, to a lesser extent, of DNA binding activities, and also proteins involved in ribosome biogenesis, splicing, translation, and transcription. Here, we confirm that EBP1 can associate with ribosomes and is distributed between cytoplasm, nucleus, and nucleolus, in line with its suggested role in translation and ribosome biogenesis. We demonstrate that the inhibition of *EBP1* expression causes a defect in rRNA processing, which can be rescued by sucrose. Sugar could also rescue the growth defects during root development in the *EBP1* silencing lines, suggesting EBP1 is specifically required to support protein translation in growth limiting conditions. Strikingly, overexpression of EBP1 can suppress both the loss of meristem activity and precocious differentiation that occurs upon induction of elevated RBR expression. In addition, we also found that RBR shares

interacting partners with EBP1, and besides its canonical transcriptional repressor function, it potentially acts on common biological processes with EBP1 in ribosome biogenesis and protein translation. Thus, EBP1 and RBR antagonistically regulate translational capacity to determine the balance between meristematic activity and cellular differentiation.

RESULTS

EBP1 Is Expressed in Root Meristematic Tissues

When ectopically overexpressed or silenced, *EBP1* enhances or reduces the meristematic activity and correspondingly organ growth. However, when and where *EBP1* is naturally expressed is not well documented (Horváth et al., 2006). To study the regulation of *EBP1* expression, we followed the *EBP1* promoter activity driving GUS and cyan-fluorescent protein (CFP) expression. For this, we cloned the entire intergenic region (1193bp) between *EBP1* (At3g51800) and the nearest upstream gene (At3g51810). Around 25 *Arabidopsis* transgenic lines were generated for *pEBP1:GUS* and for *pEBP1:CFP*. For both, five lines were examined in detail, all showing the same expression profile in root tissues. We detected strong GUS activity in the primary meristem (Fig. 1A), in the region




Figure 1. Arabidopsis EBP1 is expressed and localized in actively dividing tissues during root development. Representative differential interference contrast images of *EBP1* expression in the primary root (A) and during lateral root development (B) in the transgenic *pEBP1:GUS* lines after 3h GUS staining. The images of the different stages of lateral root development belong to the same root. C, Confocal microscopy images in the propidium-iodide-stained root sample of the *pEBP1:CFP* lines. D to F, Confocal microscopy images represent the localization of the EBP1-YFP protein in the *pgEBP1:YFP* transgenic line in the primary (D) and secondary (E) meristems. The enlarged sections of the meristematic region in (F) and of a lateral root initial in (E) are showing cytoplasmic localization of the fusion protein (scale bar = 25 μ m), whereas (G) and (H) illustrate the lower level of expression in the slowly dividing stem cells and QC. In each case, 5- to 6-d-old seedling were analyzed. White arrow points toward the QC, and the white arrowhead in (E) labels the occasional nuclear localization of the EBP1-YFP protein in the differentiation zone. The merged image in (A) is a composite of two consecutive overlapping images of the same root, taken with the same median focal plane. The enlarged image in (F) comes from the meristematic region, labeled by two white lines in (D). Scale bars = 50 μ m, unless stated otherwise.

of the rapidly dividing transit amplifying cells and in the emerging lateral root meristematic cells (Fig. 1B). In the elongation and differentiation zones, we observed expression only at a low level, mainly in the vasculature. Similarly, the expression pattern of *pEBP1:CFP* was largely confined to the meristematic cells (Fig. 1C). In this aspect, the expression pattern overlaps with the expression pattern published in Arabidopsis.org or Plant eFP at <https://bar.utoronto.ca/eplant>. To see whether the entire intergenic region is required to regulate the transcription, we studied stepwise-shortened regions of the full promoter. As Supplemental Figure S1 illustrates, the region (-337-0) contains the essential regulatory sequences to induce transcription.

To investigate whether *EBP1* promoter activity and protein accumulation correspond, we fused the genomic coding region of *EBP1* with the coding region of the yellow fluorescent protein (YFP) and expressed this 3'-terminal fusion under its native promoter (*pgEBP1-YFP*). For this, we used the entire intergenic region (1193bp) containing the 5'-upstream region, which also comprises the 5'UTR. Of over 30 regenerated transgenic lines, 5 lines were analyzed in detail and showed the same pattern of *EBP1-YFP* protein accumulation, which overlapped with the *EBP1* promoter activity. *EBP1-YFP* protein was most abundant in root meristematic cells (Fig. 1D), specifically in the region of the transit amplifying cells and emerging lateral root primordia (Fig. 1E). Lower levels of *EBP1-YFP* signal were detected in the slowly dividing stem cells and even less in the quiescent center (QC; Fig. 1H). This pattern at the QC region was also closely mirrored in the *pEBP1:CFP* line (Fig. 1G). The abundance of *EBP1-YFP* protein also corresponds well with transcriptomics data, which demonstrate *EBP1* mRNA is actively translated in the shoot apex (Tian et al., 2019).

In the transgenic lines, *EBP1-YFP* signal was most prominent in the cytoplasm of meristematic cells surrounding the centrally located nuclei (Fig. 1F). Similar cellular localization patterns were shown independently by Palm et al. (2019). The amino acid sequence of *EBP1* shows high conservation to its human counterpart, which contains a motif for nuclear and nucleolar localization near to its C terminus (Squatrito et al., 2004; Karlsson et al., 2016). This motif may be more obscured when the fluorescent protein is fused to the C terminus of *EBP1*. Therefore, we labeled *EBP1* both N- and C-terminally using the Cerulean fluorescent protein and transiently expressed it under the control of the 35S promoter. In the transiently transformed *Nicotiana benthamiana* leaves (Supplemental Fig. S2, A and B), both fusion proteins were found in the nucleoli, the nucleus, and cytosol. Similar localization was observed in onion (*Allium cepa*) epidermal cells where about 10% of Cerulean-*EBP1* localized in the nucleoli with the remainder distributed between the nucleus and cytosol (Supplemental Fig. S2, C and D). Taken together, these data show that both the expression and protein accumulation of *EBP1* are concentrated in the region of rapidly dividing root meristematic cells. Most of the protein is confined to the cytoplasm in meristematic

cells, but it has the ability to enter into the nucleus and nucleolus in different tissue environments.

***EBP1* Supports Root Meristem Activity in Sucrose Limiting Conditions**

Cell cycle activity in the meristem is reliant on the availability of sugar and regulated by the TOR pathway (Ahmad et al., 2019). Therefore, we examined how the external addition of sucrose and chemical inhibition of TOR activity by AZD-8055 influence *EBP1* transcription, *EBP1* promoter activity, and *EBP1-YFP* protein levels. Endogenous *EBP1* mRNA levels were determined in Col-0 seedlings grown on medium without sucrose and then transferred to fresh media with or without sucrose (2% and 0%, respectively) and with or without AZD-8055. *EBP1* transcript levels increased upon sucrose addition but were suppressed by AZD-8055, irrespective of whether sucrose was present in the media (Fig. 2A). To be able to compare the effect of TOR kinase inhibitor in the presence of sucrose on *EBP1* promoter activity and protein accumulation, we used our *pEBP1:CFP* and *pgEBP1-YFP* transgenic lines and followed the amounts of CFP and *EBP1-YFP* on protein blots. Both *EBP1* promoter activity and *EBP1-YFP* protein levels were induced by sucrose and were affected by the TOR inhibitor (Fig. 2, B and C).

To study how *EBP1* level influences root growth, we generated inducible *EBP1* silencing lines and screened them for growth retardation using 17- β -estradiol on a medium without sucrose. Two independent transformants with the *EBP1 RNAi* construct (*iEBP1* line 1 and line 2) showed strongly reduced growth upon 17- β -estradiol treatment as silencing was induced and the *EBP1* mRNA level diminished (Supplemental Fig. S3, A and B). However, from the T3 generation onward the retardation in root growth on sucrose-free media was no longer dependent on 17- β -estradiol, and the *EBP1* level was already reduced without chemical induction, indicating that silencing became constitutive (Supplemental Fig. S4A). Therefore, in subsequent experiments we did not apply the inducer. To ascertain that *EBP1* silencing is not influenced by sucrose, we determined the *EBP1* level both in the presence and absence of sucrose. Although the level of *EBP1* was induced by sucrose as shown above, the silencing remained effective in both *iEBP1* lines (Supplemental Fig. S4A).

Next, we studied how seedling growth was influenced by the addition of external sucrose when *EBP1* was silenced. Although a large portion of the *iEBP1* seedlings (both line 1 and line 2) showed growth arrest and retarded growth on sucrose-free media, this inhibition was largely suppressed when seeds were germinated on 1% sucrose (Fig. 3, A and B). In an insertional mutant, *ebp1-3* (CS854731), we observed an even more pronounced growth retardation in the absence of sucrose and recovery on sucrose-free media (Fig. 3, A and B). Leaf development showed similar effects in the absence and presence of sucrose (Supplemental Fig.




Figure 2. Both the gene expression and the protein abundance of EBP1 depend on sugar availability and TOR kinase activity. A, Endogenous *EBP1* transcript level was analyzed in wild-type seedlings (Col0) grown in sucrose free media (0%) for 6 d in continuous light. Subsequently, seedlings were moved to media with or without sucrose and/or the specific TOR inhibitor, AZD-8055 (0% sucrose, 0% sucrose + 1 μ M AZD-8055, 2% sucrose, 2% sucrose + AZD-8055). Seedlings were harvested after 3-h treatment. *EBP1* expression was set arbitrarily to 1 in the nontreated sample (0% Suc). Significance was determined using the one-way ANOVA and Tukey's test; significant difference a: P -value < 0.05 between 2% versus 0% sucrose treatment, while b: P -value < 0.05 between AZD8055 treated versus nontreated samples. Values represent mean of relative expression of three biological repeats ($n = 3$) with sample size: $n > 100$, error bars: sd (StDv). The transgenic lines *pEBP1:CFP* (B) and *pgEBP1:YFP* (C) were germinated and grown on 0% sucrose, then transferred at 6 d after germination to 2% sucrose containing media or 2% sucrose supplemented with 1 μ M AZD-8055 for 3-h treatment. Protein gel blot analysis was performed using a GFP-specific antibody to detect in *EBP1* promoter activity (B) and in EBP1-YFP protein level (C).

S4, B and C). The level of growth retardation closely matched the decline in *EBP1* expression in the silencing lines (Supplemental Fig. S4A). To test whether the arrested growth upon *EBP1* silencing can be rescued, we transferred seedlings from sucrose-free onto sucrose-containing media and found that root growth indeed recovered (Supplemental Fig. S4, D and E).

Ectopic *EBP1* overexpression can dose-dependently promote growth in potato and Arabidopsis (Horváth et al., 2006). We asked whether this promotion depends on sucrose availability. On sucrose-free medium, we found that roots of the two independent *EBP1* overexpression lines 1 and 2 (Horváth et al., 2006) grew significantly better than Col-0, but the difference in root length between the transgenic lines and the control was less pronounced on sucrose-containing media (Fig. 3C). Taken together, the facts that sucrose rescued the growth retardation caused by *EBP1* silencing and, vice versa, that *EBP1* overexpression overcame the sucrose limitation, suggest that EBP1 has a role in coordinating root growth with sugar availability.

EBP1 Supports Maintenance of the Root Meristem and Counteracts Cell Differentiation Induced by RBR

Sugar availability controls cell proliferation through the RBR pathway (Magyar et al., 2016). Silencing of RBR results in excessive cell proliferation, while elevating the RBR level leads to premature differentiation, exhausting the stem-cell pool, and arresting division of the amplifying cells. As a result, the root meristem is gradually reduced as cells become differentiated (Wildwater et al., 2005). EBP1 and RBR levels are counter-inhibited in cultured cells (Horváth et al., 2006). To follow how the EBP1


level responds to RBR-induced differentiation in the root, we introgressed *pEBP1:CFP* and *pgEBP1:YFP* into the dexamethasone-inducible RBR overexpression line (*RBR^{OE}*). Upon induction of RBR expression, both the *EBP1* promoter activity and EBP1-YFP protein level progressively diminished, together with the meristematic zone, as differentiation progressed (Fig. 4, A and B).

To investigate whether EBP1 has a role in meristem maintenance when RBR is overexpressed, we crossed *EBP1^{OE}* line 1 and 2 with *RBR^{OE}*. Homozygous offspring of three independent introgressed lines were analyzed and showed elevated levels of both *RBR* and *EBP1* transcripts (Fig. 4C). As expected, induction of RBR overexpression reduced root growth and shortened the meristem, as measured by the distance from the tip to the first differentiating root hair. Strikingly, both effects of RBR overexpression were significantly attenuated by EBP1 overexpression (Fig. 4, D to H; Supplemental Fig. S5A). Five days after RBR induction the entire root meristem organization was lost and cells appeared vacuolated, but the elevated level of EBP1 prevented this process and supported meristematic function and root growth for a longer period (Supplemental Fig. S5, B and C). Taken together, the EBP1 level is important to maintain meristematic activity and to counteract the progression of differentiation imposed by RBR overexpression.

EBP1 Predominantly Interacts with Proteins Involved in Ribosome Biogenesis and Protein Translation

To gain insights into the molecular function of EBP1, we searched for EBP1 interacting proteins in a pull-down experiment and identified them by mass spectrometry. To do this we took advantage of the

Figure 3. Sugar can compensate for the lack of *EBP1* expression and an elevated level of *EBP1* can suppress sugar-dependent root growth deficiencies. A, Root growth of two independent *EBP1* RNAi (*iEBP1*) lines and the insertional mutant, *ebp1-3* compared with Col0, germinated and grown (12 d) in the absence (0%) and presence of sucrose (1%). Scale bars = 1 cm. B, Boxplot analyses show the quantification and distribution of the root length (in millimeters) of seedlings grown on 0% and 1% sucrose media at the given time points after sowing. The boxplot gives the mean line and the mean marker (cross); the quartile calculation was done exclusive of the median on root-length measurements from $n = 3$, $n > 35$ in each repeat. Significance was determined by Student *t* test *a*: P -value < 0.01 . C, Quantitative analysis of root growth of the transgenic *EBP1*^{OE} lines (referred as lines 19.2 and 43.2, respectively in Horváth et al., 2006) compared with Col0, germinated and grown on media with or without sucrose (0%, 1%). Boxplot analysis was carried out as in (B); $n = 3$, $n > 35$ in each repeat; significance was determined by Student *t* test *a*: P -value < 0.01 . The level of *EBP1* expression upon silencing and overexpression is shown in Supplemental Figure S4A and Figure 4C, respectively.



transgenic line *pgEBP1-YFP* and used *p35S:GFP* as a control. We identified and quantified *EBP1*-YFP associated proteins against proteins associated with the control GFP alone in six replicates each using label-free mass spectrometry (Hubner et al., 2010). To certify interactors with statistical confidence, we computed the false discovery rate and the amount ratio between proteins identified in *EBP1*-YFP vs GFP pull-downs and established thresholds as visualized in the volcano plot (Fig. 5A; Supplemental Methods). We identified 55 specific interactors with *EBP1*-YFP at high confidence (Supplemental Table S1). Gene ontology overrepresentation analysis based on molecular function placed 52 out of the 55 *EBP1* interactors into different aspects of RNA binding (Fig. 5D; Supplemental Table S1). Classifying them according to biological processes, ribosome functions and regulation of protein translation, ribosome biogenesis, small nuclear RNA synthesis, and modifications and RNA splicing were the only prominent processes (Supplemental Table S1). Taken together, mass spectrometry identification of interactors suggested that *EBP1* has the potential to regulate multiple aspects of ribosome biogenesis and protein translation.

Intrigued by the molecular and functional antagonism between *EBP1* and RBR, we also searched for RBR-interacting proteins (Fig. 5, B and E; Supplemental

Table S2) using the *pgRBR-GFP* line (Magyar et al., 2012). The list of RBR-interactors (55) included known partners, such as E2FB and DPB and additional, previously unidentified interactors. Among which 13 were annotated with the molecular function of RNA-binding, involved in ribosome biogenesis and protein translation (Fig. 5C). Five of these also interacted with *EBP1*, including members of the H/ACA ribonucleoprotein complex, known to catalyze the site-specific pseudo-uridylation and most of the methylation of rRNA (Watkins and Bohnsack, 2012); NOP56, which operates in trans to assist with the maturation of rRNAs (Lykke-Andersen et al., 2018); and finally the pre-mRNA processing protein 40A, which binds the carboxyl-terminal domain of the largest subunit of RNA polymerase II and functions as a scaffold for RNA processing (Morris and Greenleaf, 2000). This set of common interactors points toward an as yet unknown aspect of RBR function that converges with *EBP1* and explaining their opposing functions in the meristem.

***EBP1* Is Associated with the Ribosome and Supports Global Translation in Sucrose Limiting Conditions**

Given that *EBP1* was found to interact with a number of ribosomal proteins by mass spectrometry, we asked




Figure 4. Elevated level of *EBP1* delays differentiation upon induction of RBR expression. A, Confocal microscopy images of the transgenic lines, *RBR^{OE}(pEBP1:CFP)* and *RBR^{OE}(pgEBP1:YFP)*. B, Upon induction of RBR with dexamethasone (dex) treatment (1 μM), at the given time points. Note, the loss of meristem structure and progressive differentiation due to RBR induction and the concomitant drop in *EBP1* transcription and reduction of EBP1-YFP abundance. C, To generate the *RBR^{OE};EBP1^{OE}* lines, the transgenic lines *RBR^{OE}* (Wildwater et al., 2005) and *EBP1^{OE}* (line 1; Horváth et al., 2006) constitutively overexpressing the *Solanum tuberosum EBP1* (*StEBP1*) were introgressed. The expression levels for *RBR* and *EBP1* were analyzed by quantitative reverse transcription-PCR in three independent introgressed lines (*RBR^{OE};EBP1^{OE}* line 10, 68, 78; F3 generation) using *RBR*- and *StEBP1*-specific primers. As a control, the parental lines (*RBR^{OE}* and *EBP1^{OE}*) and Col0 were tested. For the introgressed line 10, the transgenic line *EBP1^{OE}* line 1 was used as a parent, while for the lines 68 and 78, *EBP1^{OE}* line 2. D, Representative root samples of Col0, *EBP1^{OE}*, *RBR^{OE}* and *RBR^{OE};EBP1^{OE}* lines upon 24 h dex (1 μM) induction. See also Supplemental Figure S5A. E, Confocal images of propidium iodide-stained root samples from Col0, *EBP1^{OE}*, *RBR^{OE}* and *RBR^{OE};EBP1^{OE}* lines upon 72 h dex (1 μM) treatment. In (D) and (E), the arrowhead shows the position of the first differentiating epidermal cell, while in (E) the arrow points to the QC. Scale bars = 1 mm (D) and 50 μm (E). F, Root growth (millimeters) after 24, 48, and 72 h dex (1 μM) treatment. Induction with dex started on 6-d-old seedlings. G, Meristem length and number of non-elongating cortical cells (H) of Col0, *EBP1^{OE}*, *RBR^{OE}* and *RBR^{OE};EBP1^{OE}* lines after 0, 24, 48, 72, and 144 h dex (1 μM) induction. The region of the transit amplifying cells in the *RBR^{OE}* root meristem is fully differentiated around 96 h dex treatment; $n > 3$, $n > 15$ seedlings at each repeat for each genotype. In (F), (G), and (H), values represent means with StDv. *a*: *P*-value < 0.01 shows the significance of the measured values of *RBR^{OE};EBP1^{OE}* line compared with *RBR^{OE}* at the given time point.

whether EBP1 is preferentially associated with single ribosomes, their subunits, or with polysomes. In a high-density sucrose gradient (15% to 50%), EBP1 fusion protein cofractionated with polysomes and ribosomal subunits (Fig. 6A), in keeping with its association with 60S subunit proteins in the pull-down experiments (Fig. 5). After separating 40S, 60S, and 80S subunits in a low-sucrose gradient (7% to 20%), EBP1 also appeared in fractions containing only 40S, as well as in the non-ribosomal fraction (Fig. 6A). Taken together, these data

confirm that EBP1 has the potential to interact with the ribosome in polysomal and nonpolysomal contexts.

To test whether EBP1 supports translation, we obtained polysome profiles of 12-d-old *EBP1 RNAi* seedlings grown without sucrose supplementation and after phenotypic rescue using 1% sucrose. When polysomes were normalized by fresh weight, the *EBP1 RNAi* plants grown on 0% sucrose contained fewer ribosomes overall than the wild-type control, whereas at 1% sucrose, this difference disappeared (Fig. 6, B and




Figure 5. Functional classification of the EBP1 and RBR interacting proteins identified by mass-spectrometric analysis. Volcano plots showing the enrichment of EBP1-interacting (A) and RBR-interacting proteins (B) recovered from seedlings expressing the EBP1-YFP (A) and RBR-GFP (B) fusion proteins compared with the proteins identified from the transgenic lines Col0(*p35S::GFP*). The x axis shows the fold enrichment (FC) between the fusion protein and GFP alone (\log_2); whereas the y axis illustrates significance (Sig), the adjusted P -value ($-\log_{10}$) of the pulled-down proteins with anti-GFP antibody from EBP1-YFP (A) and RBR-GFP (B) seedlings compared with GFP-expressing Col0 transgenic plants. The vertical and horizontal stippled lines demarcate a 2-fold enrichment and an adjusted p -value of 0.05, respectively; proteins in the top right portion of the graph are considered to be the candidates for EBP1- (A) and RBR (B)-interactors. C, The table summarizes the list of RBR interactors involved in RNA binding (13) and highlights those that are common in both sets, the RBR and EBP1 interactors (5). D and E, The pie charts represent the major categories by their molecular functions using the gene ontology classification tool PANTHER version 14.1. Notable subcategories are also listed. For details on protein names and functions refer to Supplemental Tables S1 and S2.

C). However, the polysome-to-monosome (80S) ratio was not obviously affected by sucrose. Thus, the deficiency in ribosomes was rescued by sucrose, matching with the effect of sucrose on the growth kinetics shown earlier (Fig. 3; Supplemental Fig. S4).

EBP1 Supports Ribosome Biogenesis

To further delineate the potential molecular functions of EBP1 in plants, especially in light of its association

with proteins involved in ribosome biogenesis in *Arabidopsis* (Supplemental Table S1) and in human (Squatrito et al., 2004), we examined the *iEBP1* lines for defects in rRNA processing. We conducted northern blot analyses of rRNA processing intermediates with probes against the internal transcribed spacers ITS-1 and ITS-2, as well as the mature 18S and 25S rRNA (Fig. 7, A and B). Despite equal loading of all three samples (Fig. 7C), the immature forms of the nuclear rRNAs were elevated in the two *iEBP1* lines compared with the wild-type control (Fig. 7. D and E). In contrast,




Figure 6. EBP1 associates with cytosolic ribosomes and supports ribosome biogenesis. A, Absorbance profile of a sucrose density gradient after fractionation of whole tissue from Col0(*pgEBP1-YFP*) seedlings (12 d). The x axis represents the sedimentation distance in a 15% to 50% (red trace) or 7% to 20% (blue stippled trace) sucrose gradient, while the y axis represents A_{254} (A_{254}). Peaks corresponding to 40S, 60S subunits, and 80S ribosomes are indicated on the individual profiles; polysomes are to the right of the 80S. Under the graph, the immunoblot analysis (EBP1-YFP) illustrates the distribution of EBP1-YFP in the total cell extract (CE) and in the eight fractions of the different gradients, while the ethidium bromide-stained agarose gels (EtBr gel) show the distribution of 18S (40S subunit) and 25S (60S subunit) rRNAs also in the eight fractions from the different gradients as shown in the graph. B and C, Polysome profiling of EBP1 RNAi lines (*iEBP1* line 1 and line 2). Seedlings were grown for 12 d without sucrose (B) and with 1% sucrose (C). Total cell extracts from equal amounts of whole seedlings from Col0 and EBP1 RNAi transgenics were fractionated on 15% to 50% sucrose density gradients.

the abundance of the mature 18S and 25S rRNAs was indistinguishable between the Col-0 and RNAi lines (Fig. 7, F and G). These data suggest that the early processing events take place rapidly, as indicated by small pool sizes of the processing intermediates, whereas the later events (e.g. P[1]-A3 to 20S) appear to be slower, as shown by larger pool sizes (Fig. 7, D and E). Through quantification of the signals (Fig. 7H), we confirmed that the early processing events were particularly sensitive to the reduction in the EBP1 level, suggesting that EBP1 affects early processing steps in rRNA maturation. However, as the later processing intermediates (e.g. P[1]-A3 and 20S) were not depleted in the *EBP1* RNAi lines, we suggest that EBP1 also supports the efficiency of processing steps at later stages. In conclusion, these data implicate EBP1 in ribosome biogenesis and, either directly or indirectly, in

rRNA processing. The molecular defects in ribosome biogenesis may explain the slow growth of the *EBP1* RNAi lines. In the rRNA processing experiments, we did not observe any imbalance in the ratios of specific rRNAs when samples containing equal amounts of total RNA were compared (Fig. 7C). The requirement for EBP1 for rRNA processing was rescued by growing the seedlings on sucrose-containing media (Supplemental Fig. S6, A to E, -S to +S). This result is in keeping with the conclusion that the slow growth of *EBP1* RNAi plants is due, at least in part, to rRNA processing defects.


DISCUSSION

In plants, EBP1 attracted attention as a growth driver both in Arabidopsis and in crop species, where it was shown to increase potato yield (Horváth et al., 2006) and to associate with hybrid vigor in maize (Wang et al., 2016). However, the molecular function and how EBP1 acts to increase plant growth and crop yield remained unknown. Here we show that EBP1 associates with proteins that bind RNA and are involved in a broad range of functions from ribosome biogenesis to transcriptional and translational regulation. Specifically, (1) plant EBP1 can localize to nucleoli, (2) EBP1 is found in cellular fractions containing ribosomal subunits as well as polysomes, and (3) silencing of EBP1 causes imbalances in the pattern of rRNA processing intermediates. In addition, we find that EBP1 antagonizes RBR action to maintain proliferation in the meristem, and sustains root growth, specifically in sucrose-limiting conditions.

Nutrient limiting conditions as well as abiotic stress constrain growth. It is of particular importance that plants maintain the proliferation potential in the meristem under these conditions (Julkowska and Testerink, 2015). For example, drought triggers jasmonate signaling, which counteracts the decline of *EBP1* expression as leaf development progresses and cell proliferation gradually ceases (Kim et al., 2017). The same regulation is also observed for the large majority of genes coregulated with *EBP1* in the *ribi* regulon (Noir et al., 2013). Thus, drought perception through jasmonate signaling has been suggested to establish a “ready-to-go” state that enables rapid recovery of meristematic functions after the stress subsides. Correspondingly, elevated *EBP1* expression confers resistance to abiotic stresses such as cold (Cao et al., 2009) and drought (Cheng et al., 2016). Under stress, survival and growth are separate events, and it is likely that EBP1 is involved in the latter (Skirycz et al., 2011).

Here we show that EBP1 is important to maintain root meristem activity and growth in sucrose-limiting conditions. Thus, as during abiotic stress, EBP1 is also important to sustain growth when assimilates are limited. Consistent with previous data (Deprost et al., 2007), we show here that EBP1 mRNA and protein levels are stimulated by TOR kinase. sucrose stimulates

Figure 7. *EBP1 RNAi* transgenic plants accumulate rRNA processing intermediates. Schematic representation of the Arabidopsis full length 35S pre-rRNA transcript indicating processing sites and the position of the probes used for northern blot analysis (red bars; A) and pre-rRNA processing pathway based on Zakrzewska-Placzek et al. (2010) (B). ETS, external transcribed spacer; ITS, internal transcribed spacer. C, Ethidium bromide-stained 1% (w/v) agarose gel showing equal loading of total RNA (4 μ g) from Col0 and two independent *EBP1 RNAi* transgenic lines 1 and 2 grown without sucrose. Molecular weight marker (M) is shown on left, and the position of the rRNAs is indicated on the right. D to G, northern blot analysis of total RNA from Col0 and *EBP1 RNAi* lines using digoxigenin-labeled specific probes against ITS-1 (D), ITS-2 (E), 25S (F), and 18S (G), visualized by chemiluminescence. Positions of the full length pre-rRNA transcript (35S) and the processed products are indicated on the left. H, Quantification of the signal intensity of the various rRNA transcript species shown in (D) to (G). The x axis indicates the precursors and processed rRNAs, while the y axis shows relative signal intensity. Values represent means and error bars stand for SE from two northern blot experiments carried out with two independent biological materials.



cell proliferation in multiple ways, which include TOR-mediated (Xiong et al., 2013; Dobrenel et al., 2016a) and TOR-independent pathways. For example, sugar controls CDK-driven RBR phosphorylation, which lifts RBR-repression on cell proliferation (Magyar et al., 2012). This event may be TOR-mediated, because TOR, acting through YAK1 kinase, inhibits the expression of a class of CDK inhibitors, the SMRs, to promote root growth (Barrada et al., 2019). We now understand better how EBP1 assists in these events. Specifically, while RBR overexpression halts root growth by depleting stem cells and enhancing differentiation (Wildwater et al., 2005), EBP1 counteracts this RBR-driven repression and allows the maintenance of root meristem activity and growth. Moreover, overexpression of EBP1 bypasses the requirement for sugar to support root growth. Taken together, TOR activity is related to control growth by

regulating both the level of EBP1 and the repressor activity of RBR. A remarkable result is that *EBP1* silencing specifically leads to sensitivity under limited sugar availability, whereas addition of sucrose to the media largely compensates for the loss of EBP1. This result confirms that sucrose, while stimulating *EBP1* expression, also supports growth in other EBP1-independent ways, such as the TOR-dependent pathways regulating E2F and RBR (Magyar et al., 2012; Xiong et al., 2013; Barrada et al., 2019).

Both *EBP1* expression and EBP1 protein are largely confined to meristems, such as the root apical and lateral root meristems. Translational data also demonstrate that *EBP1* mRNA is actively translated in the shoot apex (Tian et al., 2019). Furthermore, *EBP1* expression progressively diminishes as the meristematic cells are exhausted when RBR overexpression is induced. This shows that EBP1 level is tightly linked with

meristematic activity. Fluorescently tagged EBP1 could be detected throughout the cytosol, nucleus, and nucleolus, although the nuclear and nucleolar accumulation were largely masked when EBP1 was tagged at its C terminus. We conclude that EBP1 is likely found in all three compartments as suggested by the N-terminally tagged version, because (1) growth factor signaling promoted nuclear accumulation of EBP1-GFP (Li et al., 2018), (2) others have detected tagged EBP1 proteins in plant nuclei of various cell types (Zhang et al., 2005; Cao et al., 2009; Li et al., 2016; Palm et al., 2019), and (3) native Arabidopsis EBP1 was detected in a purified nucleolar preparation (Pendle et al., 2005).

In line with the localization of EBP1 in the cytoplasm, nucleus, and nucleolus, we identified EBP1-interacting proteins broadly distributed in the cell but remarkably almost exclusively involved in various aspects of protein synthesis. We identified EBP1 in association with ribosomal proteins of the 60S subunit, as well as the eukaryotic translation initiation factor eIF5B. EBP1 was detected in sucrose gradient fractions that contain polysomes, suggesting association of EBP1 with mature cytosolic ribosomes as well as ribosomal subunits, similar to the association pattern of human EBP1 (Squatrito et al., 2004; Squatrito et al., 2006). Related to the nucleolar localization of EBP1, we found EBP1 interaction with ribosome biogenesis factors, such as multiple subunits of the H/ACA ribonucleoprotein complex involved in pre-rRNA processing (Watkins and Bohnsack, 2012), with nucleolar S-adenosyl-L-Met-dependent methyltransferases, with other rRNA-processing proteins (Lykke-Andersen et al., 2018) and the multifunctional protein nucleolin (Jia et al., 2017). Similar to EBP1, nucleolin is also most abundant in meristematic tissues and, in parallel with D-type cyclin, it is rapidly induced when cell proliferation resumes after nutrient starvation (Bögge et al., 1996). Besides the proteins with nucleolar functions, EBP1 also interacted with proteins involved in transcriptional and post-transcriptional regulation. In conclusion, similarly to human EBP1 with RNA binding activity (Squatrito et al., 2004; Squatrito et al., 2006), the common feature of EBP1 interacting proteins is that they function as part of ribonucleoprotein complexes.

Although the precise role of EBP1 in ribosome biogenesis remains unclear, the related metallopeptidase-fold protein Arx1 is a canonical 60S biogenesis factor (Greber, 2016). Seedlings with reduced EBP1 level showed a defect in rRNA processing that affected specifically the early processing intermediates. Just as the growth phenotype was rescued by sucrose, so was the rRNA processing delay in the *EBP1* silencing lines. These results are consistent with the assumption that EBP1 has a role in ribosome biogenesis. Nevertheless, we cannot rule out that Arabidopsis EBP1 also supports rRNA transcription, as was described in human cells (Nguyen le et al., 2015). However, our data point to a defect during rRNA processing, because a transcriptional defect upon *EBP1* silencing would reduce rather

than increase the abundance of rRNA processing intermediates.

In vertebrates EBP1 is expressed in two isoforms; the longer p48 isoform is generally associated with cell proliferation and tumorigenesis, specifically by inhibiting cell death and promoting Pol I-dependent rRNA expression. The shorter p42 isoform lacks the N terminus of p48 and is considered to have a tumor suppressor role. Together with Rb, E2F1, and chromatin modifying proteins, HsEBP1 p42 represses cell proliferation (Liu et al., 2006; Ko et al., 2016; Nguyen et al., 2018; Nguyen et al., 2019). The internal methionine that initiates translation of the p42 isoform is not conserved in plants. Thus, there is no evidence that the shorter suppressive form exists in plants. In line with these findings, the plant EBP1 promotes cell growth and division (Horváth et al., 2006; Cheng et al., 2016).

In addition, in this study we show that *EBP1* overexpression can compensate for the pro-differentiation activity of RBR in the root stem cell niche. Inducible overexpression of *RBR* downregulated *EBP1* expression and rapidly abolished proliferation competence. In contrast, overexpression of *EBP1* countered RBR's pro-differentiation activity and rescued the maintenance of cell proliferation around the stem cell niche. Thus, EBP1 expression is in an antagonistic relationship with the pro-differentiation factor, RBR. Our data illustrate how stem cell fate is maintained by a balance between pro-differentiation signals and the pro-growth agenda of the ribosome biogenesis machinery.

The striking finding that EBP1 and RBR share common interactors involved in ribosome biogenesis and that RBR associated with a number of proteins in the cytoplasmic translation machinery may shed light on how EBP1, in conjunction with RBR, regulates meristem activity. Future research will elaborate on our hypothesis that RBR and EBP1 counteract to regulate meristem function through acting on a common biological process to regulate the capacity for protein synthesis. In conclusion, our findings on the role of EBP1 in root meristem activity and growth could be important in breeding to improve yield and yield stability in crops.

MATERIALS AND METHODS

Plant Material and Growth Conditions

Arabidopsis (*Arabidopsis thaliana*) seeds were sterilized, incubated at 4°C for 2 d, then germinated and grown vertically on half-strength Murashige and Skoog salt plant media (Sigma, cat #2633024) with different concentration of Phytoagar under a long day cycle of 16 h light ($80 \pm 10 \mu\text{mol m}^{-2} \text{s}^{-1}$)/8 h dark at 22°C and 50% humidity. Sucrose was added at 1%, 2% (w/v), or omitted, as specified. Arabidopsis ecotype Columbia (Col-0) was used as the wild-type control as the genetic background to generate transgenic lines and as a source of DNA to clone either promoter or coding regions. Transgenic lines, *EBP1^{OE}* lines 1 and 2, overexpressing the potato (*Solanum tuberosum*) EBP1 (StEBP1, Horváth et al., 2006 referred as line 19 and 43), and *RBR^{OE}* (Wildwater et al., 2005) were described earlier. The insertional line, CS854731, named as *ebp1-3* (Li et al., 2018), was obtained from the Salk Institute (<http://signal.sal.edu>).

Generating Transcriptional and Translational Fusion Proteins

Both transcriptional and translational fusions were constructed by the Multisite Gateway System (Invitrogen) by amplifying the putative regulatory and genomic coding sequences with gene-specific primers. To generate the *p35S: Cerulean-EBP1* reporter construct and the *EBP1* inducible silencing lines, *EBP1 RNAi*, Arabidopsis *EBP1* cDNA was amplified and cloned in the Gateway compatible vectors. The plant binary vector system, *pGREEN* was used in the *Agrobacterium tumefaciens* C589(pMP90)-mediated plant transformation with the floral dip method. Details are described in Supplemental Data S1 and primer sequences summarized in Supplemental Table S3.

Root Measurement and Microscopy

Photographs of vertically or horizontally grown seedlings were taken with a digital camera (Canon). The primary root length was measured using ImageJ version 1.41 software (<http://rsb.info.nih.gov/ij/index.html>) and analyzed statistically with GraphPad Prism 7.0a (GraphPad Software). The root meristematic zone was determined by measuring the distance from the QC to the first elongating cortex cells; the same region was used to count the number of cortex cells with no signs of rapid elongation. For measurements and to follow tissue-specific expression, either light (Zeiss AxioScope with Nomarski optics) or confocal microscopy (Zeiss LSM710 and Leica SP8) was carried out. To take differential interference contrast images, chloral-hydrate treated roots were used; starch granules were visualized with 1% (w/v) lugol solution. Seedlings (T3 generation) were stained for GUS activity for 3–6 h at 37°C in 0.5 mg/mL X-gluc (Biosynth AG), dissolved according to the manufacturer's recommendation. For confocal microscopy, to visualize cell walls, 5–10 µg/mL propidium iodide (Sigma-Aldrich) solution was applied. For analysis, longitudinal, single sections were taken with the same median focal plane. The final images were built and merged from partially overlapping sections with Adobe Illustrator CS6.

Chemical Treatments

The introgression lines, *EBP1^{OE};RBR^{OE}* (F3 generation) were germinated and grown on vertical plates (1% sucrose media). The 5- to 6-d-old seedlings were transferred to 1 µM dexamethasone-containing plates, and growth was scored over a time series (24 h–144 h period). *RBR^{OE}*, *RBR^{OE}* backcrossed to Col-0, *EBP1^{OE}*, and Col-0 were used as a control to follow root growth, meristem size, and level of gene expression. To screen transformants of the silencing lines, *EBP1 RNAi*, seeds (T1 and T2) were germinated and grown on 5 µM 17-β-estradiol-containing medium supplied with 1% sucrose or no sucrose. Next, the candidate lines were grown in normal growth conditions and transferred to 17-β-estradiol medium with or without sucrose. The induction and level of silencing was measured by quantitative reverse transcription PCR.

Immunoblot and AZD Treatment

For immunoblot analysis, the transgenic lines Col-0(*pgEBP1-YFP*) and Col-0(*pEBP1:CFP*) were germinated and grown for 5 d on sucrose-free medium in 12-h light/12-h dark conditions. On the 5th day, at ZT0 timepoint, seedlings were transferred to medium containing either 2% sucrose, 2% sucrose + 1 µM AZD8055 or to fresh sucrose-free medium. Samples were harvested 3 h later by flash-freezing the whole seedlings. Total protein was extracted from samples using extraction buffer containing 75 mM NaCl, 25 mM Tris-HCl (pH = 7.5), 15 mM β-glycerolphosphate, 15 mM EGTA, 15 mM p-nitrophenylphosphate, 10 mM MgCl₂, 1 mM dithiothreitol, 1 mM NaF, 0.5 mM Na₃VO₄, 0.5 mM phenylmethylsulfonyl fluoride, and 0.1% Tween 20 (Magyar et al., 2005). The total protein samples were separated in a 10% acrylamide gel using SDS-PAGE and transferred to polyvinylidene difluoride membrane. Antibodies used for detection: Anti-GFP mouse monoclonal antibody (ROCHE) in 1:1000 dilution and goat anti-mouse IgG-HRP Secondary Antibody (Sigma) in 1:10000 dilution.

Northern Blot Analysis, Polysome Profiling, and Protein Fractionation

Northern blot analyses were performed with 4 µg of total RNA from seedlings grown under the indicated conditions. Blots were probed with digoxigenin-labeled probes corresponding to the internal transcribed spacer

1 and 2, 18S and 25S rRNAs, and signal was detected with chemiluminescent reagent, imaged on a BioRad ChemiDoc instrument, and quantified with ImageJ version 1.41 software (<http://rsb.info.nih.gov/ij/index.html>). Polysome profiles were obtained from 12-d-old Col-0 or *EBP1 RNAi* whole seedlings grown without or with 2% sucrose essentially as described (Kim et al., 2007). Then 100 µL supernatant was layered on either a 2-mL 15–50% or a 2 mL 7% to 20% linear gradient prepared with a Hoefer gradient maker and centrifuged at 50,000 rpm (Beckmann TLS55 rotor) for 1 h 10 min at 4°C. After the UV *A*₂₅₄ and fractionation into eight equal fractions were monitored, *EBP1* protein was detected by immunoblotting with GFP-specific antibody (Lokdarshi et al., 2016). For detailed procedures see Supplemental Data S1.

Protein Complex Isolation, LC-MS/MS Identification, and Statistical Analysis

A detailed description and link to the raw datasets are given in the Supplemental Data S1.

Accession Numbers

Accession numbers are At3g51800, *EBP1*; At3g12280, *RBR*; the other accession numbers are listed in Supplemental Table S3.

Supplemental Data

The following supplemental materials are available.

Supplemental Figure S1. *EBP1* is expressed in actively dividing tissues during root development.

Supplemental Figure S2. *EBP1* localizes to nucleoli, nuclei and cytosol.

Supplemental Figure S3. Reduced level of *EBP1* results in retarded growth in the absence of sucrose.

Supplemental Figure S4. Root and leaf growth deficiency in the absence of sugar and recovery of root growth defects upon sucrose supplementation in the *EBP1* transgenic lines.

Supplemental Figure S5. Elevated level of *EBP1* delays the differentiation induced by *RBR* overexpression.

Supplemental Figure S6. sucrose supplementation alleviates the requirement for *EBP1* in rRNA processing.

Supplemental Table S1. List of *EBP1* interacting proteins.

Supplemental Table S2. List of *RBR* interacting proteins.

Supplemental Table S3. Primers used in this study.

Supplemental Data S1. <https://doi.org/10.6084/m9.figshare.10069247>.

ACKNOWLEDGMENTS

We thank Dr. Patrick Gialvalisco (Max Planck Institute Cologne) for initial discussions on *EBP1*, linking it with protein translation; Dr. Tessa Burch-Smith and Dr. Daniel M Roberts (University of Tennessee) for valuable advice with northern blot analysis and for providing anti-GFP antibody, respectively; Imma Perez-Salamo for valuable discussions (Royal Holloway, University of London); and Christian Bachem (Wageningen University and Research) for critical reading of the manuscript.

Received July 2, 2019; accepted November 22, 2019; published December 9, 2019.

LITERATURE CITED

- Ahmad Z, Magyar Z, Bögre L, Papdi C (2019) Cell cycle control by the target of rapamycin signalling pathway in plants. *J Exp Bot* 70: 2275–2284
- Bakshi A, Moin M, Madhav MS, Kirti PB (2019) Target of rapamycin, a master regulator of multiple signalling pathways and a potential candidate gene for crop improvement. *Plant Biol (Stuttg)* 21: 190–205

- Barbet NC, Schneider U, Helliwell SB, Stansfield I, Tuite MF, Hall MN (1996) TOR controls translation initiation and early G1 progression in yeast. *Mol Biol Cell* 7: 25–42
- Barrada A, Djendli M, Desnos T, Mercier R, Robaglia C, Montané MH, Menand B (2019) A TOR-YAK1 signaling axis controls cell cycle, meristem activity and plant growth in *Arabidopsis*. *Development* 146: dev171298
- Bögre L, Jonak C, Mink M, Meskiene I, Traas J, Ha DT, Swoboda I, Plank C, Wagner E, Heberle-Bors E, Hirt H (1996) Developmental and cell cycle regulation of alfalfa nucMs1, a plant homolog of the yeast Nsr1 and mammalian nucleolin. *Plant Cell* 8: 417–428
- Bose SK, Sengupta TK, Bandyopadhyay S, Spicer EK (2006) Identification of Ebp1 as a component of cytoplasmic bcl-2 mRNP (messenger ribonucleoprotein particle) complexes. *Biochem J* 396: 99–107
- Cao P, Song J, Zhou C, Weng M, Liu J, Wang F, Zhao F, Feng D, Wang B (2009) Characterization of multiple cold induced genes from *Ammopiptanthus mongolicus* and functional analyses of gene AmEBP1. *Plant Mol Biol* 69: 529–539
- Chen GH, Liu MJ, Xiong Y, Sheen J, Wu SH (2018) TOR and RPS6 transmit light signals to enhance protein translation in deetioliating *Arabidopsis* seedlings. *Proc Natl Acad Sci USA* 115: 12823–12828
- Cheng H, Chen X, Zhu J, Huang H (2016) Overexpression of a *Hevea brasiliensis* ErbB-3 Binding protein 1 gene increases drought tolerance and organ size in *Arabidopsis*. *Front Plant Sci* 7: 1703
- De Veylder L, Beeckman T, Inzé D (2007) The ins and outs of the plant cell cycle. *Nat Rev Mol Cell Biol* 8: 655–665
- Deprost D, Yao L, Sormani R, Moreau M, Leterreux G, Nicolaï M, Bedu M, Robaglia C, Meyer C (2007) The Arabidopsis TOR kinase links plant growth, yield, stress resistance and mRNA translation. *EMBO Rep* 8: 864–870
- Dewitte W, Riou-Khamlichi C, Scofield S, Healy JM, Jacqmar A, Kilby NJ, Murray JA (2003) Altered cell cycle distribution, hyperplasia, and inhibited differentiation in *Arabidopsis* caused by the D-type cyclin CYCD3. *Plant Cell* 15: 79–92
- Dobrenel T, Caldana C, Hanson J, Robaglia C, Vincenz M, Veit B, Meyer C (2016a) TOR signaling and nutrient sensing. *Annu Rev Plant Biol* 67: 261–285
- Dobrenel T, Mancera-Martínez E, Forzani C, Azzopardi M, Davanture M, Moreau M, Schepetilnikov M, Chicher J, Langella O, Zivy M, et al (2016b) The Arabidopsis TOR kinase specifically regulates the expression of nuclear genes coding for plastidic ribosomal proteins and the phosphorylation of the cytosolic ribosomal protein S6. *Front Plant Sci* 7: 1611
- Dowling RJ, Topisirovic I, Alain T, Bidnost M, Fonseca BD, Petroulakis E, Wang X, Larsson O, Selvaraj A, Liu Y, et al (2010) mTORC1-mediated cell proliferation, but not cell growth, controlled by the 4E-BPs. *Science* 328: 1172–1176
- Enganti R, Cho SK, Toperzer JD, Urquidi-Camacho RA, Cakir OS, Ray AP, Abraham PE, Hettich RL, von Arnim AG (2018) Phosphorylation of ribosomal protein RPS6 integrates light signals and circadian clock signals. *Front Plant Sci* 8: 2210
- Forzani C, Duarte GT, Van Leene J, Clement G, Huguet S, Paysant-LeRoux C, Mercier R, De Jaeger G, Leprince AS, Meyer C (2019) Mutations of the AtYAK1 Kinase Suppress TOR Deficiency in *Arabidopsis*. *Cell Rep* 27: 3696–3708
- Fox S, Southam P, Pantin F, Kennaway R, Robinson S, Castorina G, Sánchez-Corrales YE, Sablowski R, Chan J, Grieneisen V, et al (2018) Spatiotemporal coordination of cell division and growth during organ morphogenesis. *PLoS Biol* 16: e2005952
- González A, Hall MN (2017) Nutrient sensing and TOR signaling in yeast and mammals. *EMBO J* 36: 397–408
- Greber BJ (2016) Mechanistic insight into eukaryotic 60S ribosomal subunit biogenesis by cryo-electron microscopy. *RNA* 22: 1643–1662
- Gutzat R, Borghi L, Fütterer J, Bischof S, Laizet Y, Hennig L, Feil R, Lunn J, Gruissem W (2011) RETINOBLASTOMA-RELATED PROTEIN controls the transition to autotrophic plant development. *Development* 138: 2977–2986
- Henriques R, Magyar Z, Bögre L (2013) S6K1 and E2FB are in mutually antagonistic regulatory links controlling cell growth and proliferation in *Arabidopsis*. *Plant Signal Behav* 8: e24367
- Henriques R, Magyar Z, Monardes A, Khan S, Zaleski C, Orellana J, Szabados L, de la Torre C, Koncz C, Bögre L (2010) Arabidopsis S6 kinase mutants display chromosome instability and altered RBR1-E2F pathway activity. *EMBO J* 29: 2979–2993
- Horváth BM, Magyar Z, Zhang Y, Hamburger AW, Bakó L, Visser RG, Bachem CW, Bögre L (2006) EBP1 regulates organ size through cell growth and proliferation in plants. *EMBO J* 25: 4909–4920
- Hubner NC, Bird AW, Cox J, Spletstoesser B, Bandilla P, Poser I, Hyman A, Mann M (2010) Quantitative proteomics combined with BAC TransgeneOmics reveals in vivo protein interactions. *J Cell Biol* 189: 739–754
- Jia W, Yao Z, Zhao J, Guan Q, Gao L (2017) New perspectives of physiological and pathological functions of nucleolin (NCL). *Life Sci* 186: 1–10
- Julkowska MM, Testerink C (2015) Tuning plant signaling and growth to survive salt. *Trends Plant Sci* 20: 586–594
- Karlsson T, Altankhuyag A, Dobrovoltska O, Turcu DC, Lewis AE (2016) A polybasic motif in ErbB3-binding protein 1 (EBP1) has key functions in nucleolar localization and polyphosphoinositide interaction. *Biochem J* 473: 2033–2047
- Kim BH, Cai X, Vaughn JN, von Arnim AG (2007) On the functions of the h subunit of eukaryotic initiation factor 3 in late stages of translation initiation. *Genome Biol* 8: R60
- Kim JM, To TK, Matsui A, Tanoi K, Kobayashi NI, Matsuda F, Habu Y, Ogawa D, Sakamoto T, Matsunaga S, Bashir K, Rasheed S, et al (2017) Acetate-mediated novel survival strategy against drought in plants. *Nat Plants* 3: 17097
- Ko HR, Chang YS, Park WS, Ahn JY (2016) Opposing roles of the two isoforms of ErbB3 binding protein 1 in human cancer cells. *Int J Cancer* 139: 1202–1208
- Li C, Liu X, Qiang X, Li X, Li X, Zhu S, Wang L, Wang Y, Liao H, Luan S, Yu F (2018) EBP1 nuclear accumulation negatively feeds back on FERONIA-mediated RALF1 signaling. *PLoS Biol* 16: e2006340
- Li J, Yu G, Sun X, Zhang X, Liu J, Pan H (2016) AcEBP1, an ErbB3-binding protein (EBP1) from halophyte *Atriplex canescens*, negatively regulates cell growth and stress responses in *Arabidopsis*. *Plant Sci* 248: 64–74
- Liu MJ, Wu SH, Wu JF, Lin WD, Wu YC, Tsai TY, Tsai HL, Wu SH (2013) Translational landscape of photomorphogenic *Arabidopsis*. *Plant Cell* 25: 3699–3710
- Liu Z, Ahn JY, Liu X, Ye K (2006) Ebp1 isoforms distinctively regulate cell survival and differentiation. *Proc Natl Acad Sci USA* 103: 10917–10922
- Lokdarshi A, Conner WC, McClintock C, Li T, Roberts DM (2016) Arabidopsis CML38, a calcium sensor that localizes to ribonucleoprotein complexes under hypoxia stress. *Plant Physiol* 170: 1046–1059
- López-Juez E, Dillon E, Magyar Z, Khan S, Hazeldine S, de Jager SM, Murray JA, Beemster GT, Bögre L, Shanahan H (2008) Distinct light-initiated gene expression and cell cycle programs in the shoot apex and cotyledons of *Arabidopsis*. *Plant Cell* 20: 947–968
- Lykke-Andersen S, Ardal BK, Hollensen AK, Damgaard CK, Jensen TH (2018) Box C/D snoRNP Autoregulation by a cis-Acting snoRNA in the NOP56 Pre-mRNA. *Mol Cell* 72: 99–111
- Magyar Z, Bögre L, Ito M (2016) DREAMs make plant cells to cycle or to become quiescent. *Curr Opin Plant Biol* 34: 100–106
- Magyar Z, De Veylder L, Atanassova A, Bakó L, Inzé D, Bögre L (2005) The role of the Arabidopsis E2FB transcription factor in regulating auxin-dependent cell division. *Plant Cell* 17: 2527–2541
- Magyar Z, Horváth B, Khan S, Mohammed B, Henriques R, De Veylder L, Bakó L, Scheres B, Bögre L (2012) Arabidopsis E2FA stimulates proliferation and endocycle separately through RBR-bound and RBR-free complexes. *EMBO J* 31: 1480–1493
- Mohammed B, Biloei SF, Dóczy R, Grove E, Railo S, Palme K, Ditengou FA, Bögre L, López-Juez E (2018) Converging light, energy and hormonal signaling control meristem activity, leaf initiation, and growth. *Plant Physiol* 176: 1365–1381
- Montané MH, Menand B (2013) ATP-competitive mTOR kinase inhibitors delay plant growth by triggering early differentiation of meristematic cells but no developmental patterning change. *J Exp Bot* 64: 4361–4374
- Montané MH, Menand B (2019) TOR inhibitors: From mammalian outcomes to pharmacogenetics in plants and algae. *J Exp Bot* 70: 2297–2312
- Morris DP, Greenleaf AL (2000) The splicing factor, Prp40, binds the phosphorylated carboxyl-terminal domain of RNA polymerase II. *J Biol Chem* 275: 39935–39943
- Nguyen DQ, Hoang DH, Nguyen TTV, Ho HD, Huynh V, Shin JH, Ly QT, Thi Nguyen DD, Ghoda L, Marcucci G, Nguyen LXT (2019) Ebp1 p48 promotes oncogenic activities in human colon cancer cells through

- regulation of TIF-90-mediated ribosomal RNA synthesis. *J Cell Physiol* **234**: 17612–17621
- Nguyen DQ, Hoang DH, Nguyen Vo TT, Huynh V, Ghoda L, Marcucci G, Nguyen LXT** (2018) The role of ErbB3 binding protein 1 in cancer: Friend or foe? *J Cell Physiol* **233**: 9110–9120
- Nguyen XT, Lee Y, Urbani L, Utz PJ, Hamburger AW, Sunwoo JB, Mitchell BS** (2015) Regulation of ribosomal RNA synthesis in T cells: Requirement for GTP and Ebp1. *Blood* **125**: 2519–2529
- Noir S, Bömer M, Takahashi N, Ishida T, Tsui TL, Balbi V, Shanahan H, Sugimoto K, Devoto A** (2013) Jasmonate controls leaf growth by repressing cell proliferation and the onset of endoreduplication while maintaining a potential stand-by mode. *Plant Physiol* **161**: 1930–1951
- Norris-Mullins B, VanderKolk K, Vacchina P, Joyce MV, Morales MA** (2014) LmaPA2G4, a homolog of human Ebp1, is an essential gene and inhibits cell proliferation in *L. major*. *PLoS Negl Trop Dis* **8**: e2646
- Okada M, Jang SW, Ye K** (2007) Ebp1 association with nucleophosmin/B23 is essential for regulating cell proliferation and suppressing apoptosis. *J Biol Chem* **282**: 36744–36754
- Palm D, Streit D, Shanmugam T, Weis BL, Ruprecht M, Simm S, Schleiff E** (2019) Plant-specific ribosome biogenesis factors in *Arabidopsis thaliana* with essential function in rRNA processing. *Nucleic Acids Res* **47**: 1880–1895
- Pendle AF, Clark GP, Boon R, Lewandowska D, Lam YW, Andersen J, Mann M, Lamond AI, Brown JW, Shaw PJ** (2005) Proteomic analysis of the Arabidopsis nucleolus suggests novel nucleolar functions. *Mol Biol Cell* **16**: 260–269
- Pilipenko EV, Pestova TV, Kolupaeva VG, Khitrina EV, Poperechnaya AN, Agol VI, Hellen CU** (2000) A cell cycle-dependent protein serves as a template-specific translation initiation factor. *Genes Dev* **14**: 2028–2045
- Pisapia L, Barba P, Cortese A, Cicatiello V, Morelli F, Del Pozzo G** (2015) EBP1 protein modulates the expression of human MHC class II molecules in non-hematopoietic cancer cells. *Int J Oncol* **47**: 481–489
- Schepetilnikov M, Dimitrova M, Mancera-Martínez E, Geldreich A, Keller M, Ryabova LA** (2013) TOR and S6K1 promote translation reinitiation of uORF-containing mRNAs via phosphorylation of eIF3h. *EMBO J* **32**: 1087–1102
- Schepetilnikov M, Ryabova LA** (2018) Recent discoveries on the role of TOR (Target of Rapamycin) signaling in translation in plants. *Plant Physiol* **176**: 1095–1105
- Skiryicz A, Vandenbroucke K, Clauw P, Maleux K, De Meyer B, Dhondt S, Pucci A, Gonzalez N, Hoeberichts F, Tognetti VB, et al** (2011) Survival and growth of Arabidopsis plants given limited water are not equal. *Nat Biotechnol* **29**: 212–214
- Squatrito M, Mancino M, Donzelli M, Arcesi LB, Draetta GF** (2004) EBP1 is a nucleolar growth-regulating protein that is part of pre-ribosomal ribonucleoprotein complexes. *Oncogene* **23**: 4454–4465
- Squatrito M, Mancino M, Sala L, Draetta GF** (2006) Ebp1 is a dsRNA-binding protein associated with ribosomes that modulates eIF2 α phosphorylation. *Biochem Biophys Res Commun* **344**: 859–868
- Tian C, Wang Y, Yu H, He J, Wang J, Shi B, Du Q, Provart NJ, Meyerowitz EM, Jiao Y** (2019) A gene expression map of shoot domains reveals regulatory mechanisms. *Nat Commun* **10**: 141
- Wang T, Sui Z, Liu X, Li Y, Li H, Xing J, Song F, Zhang Y, Sun Q, Ni Z** (2016) Ectopic expression of a maize hybrid up-regulated gene, ErbB-3 binding Protein 1 (ZmEBP1), increases organ size by promoting cell proliferation in Arabidopsis. *Plant Sci* **243**: 23–34
- Watkins NJ, Bohnsack MT** (2012) The box C/D and H/ACA snoRNPs: Key players in the modification, processing and the dynamic folding of ribosomal RNA. *Wiley Interdiscip Rev RNA* **3**: 397–414
- Wildwater M, Campilho A, Perez-Perez JM, Heidstra R, Blilou I, Korthout H, Chatterjee J, Mariconti L, Gruijssem W, Scheres B** (2005) The RETINOBLASTOMA-RELATED gene regulates stem cell maintenance in Arabidopsis roots. *Cell* **123**: 1337–1349
- Wu Y, Shi L, Li L, Fu L, Liu Y, Xiong Y, Sheen J** (2019) Integration of nutrient, energy, light, and hormone signalling via TOR in plants. *J Exp Bot* **70**: 2227–2238
- Xiong Y, McCormack M, Li L, Hall Q, Xiang C, Sheen J** (2013) Glucose-TOR signalling reprograms the transcriptome and activates meristems. *Nature* **496**: 181–186
- Zakrzewska-Placzek M, Souret FF, Sobczyk GJ, Green PJ, Kufel J** (2010) Arabidopsis thaliana XRN2 is required for primary cleavage in the pre-ribosomal RNA. *Nucleic Acids Res* **38**: 4487–4502
- Zhang WK, Shen YG, He XJ, Du BX, Xie ZM, Luo GZ, Zhang JS, Chen SY** (2005) Characterization of a novel cell cycle-related gene from Arabidopsis. *J Exp Bot* **56**: 807–816
- Zhou H, Mazan-Mamczarz K, Martindale JL, Barker A, Liu Z, Gorospe M, Leedman PJ, Gartenhaus RB, Hamburger AW, Zhang Y** (2010) Post-transcriptional regulation of androgen receptor mRNA by an ErbB3 binding protein 1 in prostate cancer. *Nucleic Acids Res* **38**: 3619–3631
- Zhou H, Zhang Y, Hamburger AW** (2011) EBP1 inhibits translation of androgen receptor mRNA in castration resistant prostate cancer cells. *Anticancer Res* **31**: 3129–3135