

Effects of Radon From Hot Springs on Lymphocyte Subsets in Peripheral Blood

Dose-Response:
An International Journal
January-March 2020:1-7
© The Author(s) 2020
Article reuse guidelines:
sagepub.com/journals-permissions
DOI: 10.1177/1559325820902338
journals.sagepub.com/home/dos



Chunnan Piao¹ , Mei Tian¹, Hongjun Gao², Yanxiao Gao³ ,
Jianlei Ruan¹, Lina Wu¹, Gang Gao¹, Lirong Yi¹, and Jianxiang Liu¹

Abstract

Objective: To analyze changes in immune functions by detecting lymphocyte subsets in the peripheral blood of residents in the vicinity of radon from hot springs.

Methods: Two groups were randomly selected; 61 residents in the vicinity of the hot springs were assigned to the radon group, and 51 residents with a similar lifestyle and habits but no contact with hot springs were assigned to the control group. The percentages of lymphocyte subsets (CD3⁺, CD4⁺CD8⁻, CD4⁻CD8⁺, CD4⁺/CD8⁺, and TCR/CD3) in the 2 groups were evaluated on a FACS Aria flow cytometer. The absolute values of lymphocytes (LYMPH#) and percentages of lymphocytes (LYMPH%) were measured by an automatic blood analyzer.

Results: In the radon group, the numbers of CD3⁺ ($Z = -0.140, P > .05$) and CD4⁺CD8⁻ ($Z = -0.964, P > .05$) T cells were higher, as compared with the controls, but this difference was not significant. In addition, the number of CD4⁻CD8⁺ ($t = -2.141, P < .05$) T cells was significantly lower in the radon group. Furthermore, the average ratios of CD4⁺/CD8⁺ ($t = -2.201, P < .05$) and TCR/CD3 ($t = 2.047, P < .05$) cells were significantly higher in the radon group than in the controls. Compared with the control group, the LYMPH# ($t = -0.485, P > .05$) and LYMPH% ($Z = -0.835, P > .05$) showed no significant change.

Conclusion: Radon-rich hot springs could alter the proportions of lymphocyte subsets and possibly affect immunologic functions.

Keywords

radon hot springs, peripheral blood, lymphocyte subsets, low dose, hormesis

Introduction

Radon is a natural radioactive inert gas that exists widely in the ambient air. The natural radiation dose from radon and its daughters is the largest among normal background radiation, accounting for 54% of the total natural radiation dose. Higher radon levels appear in some mines and indoor environments decorated with radon-containing building materials. Radon has been recognized as the first environmental etiology and second risk factor (after smoking) for lung cancer. Early epidemiological studies have confirmed a significant correlation between radon and lung cancer, even at low levels of exposure. In recent years, health problems caused by long-term radon exposure in daily living have been attracting increased attention. Residents living in the vicinity of radon-rich hot springs for a long duration may be affected by the persistence and accumulation of radon and its daughters.

Animal studies, cell studies, clinical trials, and epidemiological investigations have confirmed that long-term exposure to

high levels of radon and its daughters can initiate biological effects at molecular and cellular levels. Although some studies have suggested that radon and its daughters can pose potential

¹ Key Laboratory of Radiological Protection and Nuclear Emergency, National Institute for Radiological Protection, Chinese Center for Disease Control and Prevention, Beijing, China

² Department of Clinical Laboratory, Emergency General Hospital, Beijing, China

³ School of Public Health (Shenzhen), Sun Yat-sen University, Guangzhou, China

Received 14 October 2019; received revised 18 December 2019; accepted 31 December 2019

Corresponding Author:

Jianxiang Liu, Key Laboratory of Radiological Protection and Nuclear Emergency, National Institute for Radiological Protection, Chinese Center for Disease Control and Prevention, Xinkang Street, Deshengmenwai, XiCheng District, Beijing 100088, China.

Email: liujianxiang@nirp.chinacdc.cn



Creative Commons Non Commercial CC BY-NC: This article is distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 License (<https://creativecommons.org/licenses/by-nc/4.0/>) which permits non-commercial use, reproduction and distribution of the work without further permission provided the original work is attributed as specified on the SAGE and Open Access pages (<https://us.sagepub.com/en-us/nam/open-access-at-sage>).

health risks to the population, it is generally accepted that radon hot springs can relieve pain, alleviate the symptoms of arthritis and skin diseases, and exert specific regulatory effects on physiological and cardiovascular functions.¹⁻⁷ In Japan, many people often go to hot springs to relieve symptoms of osteoarthritis and bronchial asthma. Europeans believe that radon hot springs can treat inflammation and pain.⁸⁻¹⁵ In this study, the lymphocyte subsets in the peripheral blood of residents around radon-rich hot springs were detected and analyzed to explore the effects of the hot springs on the health of the residents in China.

Materials and Methods

Selection of Participants

The inclusion criteria for residents in Pingshan County, Hebei Province, included long-term residence, no history of migration, no recent history of viral or bacterial infections, no history of malignant diseases or the administration of medications, and no exposure to X-rays within the previous 6 months. In the radon hot spring group, 61 local inhabitants were sampled randomly, aged between 29 and 68 years, with an average age of 42.85 years. Moreover, 51 inhabitants from other villages with similar living habits but with no exposure to radon hot springs were sampled randomly as the control group. They were aged between 25 and 71 years, with an average age of 45.90 years. There were no significant differences in age and gender between the radon hot spring group and the control group (max $2 = 0.78$, $P > .05$). This study was approved by the Ethics Committee of the National Institute for Radiological Protection of the Chinese Center for Disease Control and Prevention (Beijing, China), and all participants gave informed consent.

Background Information

The concentration of radon in the hot springs was 102 ± 11.4 Bq/L, the concentration of indoor radon was 41.9 ± 18.6 Bq/m³ over 6 months, and the typical equilibrium factor was 0.61. In some places, the radon concentration in spring water had reached the recommended levels for medical hot springs and radon springs, and the concentration in bathrooms was increased during water use. The measured radon concentration exceeded 200 Bq/m³ (indoor radon limit in China) and 100 Bq/m³ (World Health Organization radon limit value), resulting in an additional annual effective dose of about 0.09 mSv.¹⁶ The distance between the sampling location for the control group and the radon hot springs was about 11.2 km, and there were no other outdoor radon hot springs in the area (the annual indoor radon concentration in Shijiazhuang was 42.4 ± 20.2 Bq/m³).

Bathing or Showering Habits

In the radon hot spring group, 52.38% of households had hot spring baths. The mean number of baths taken was 10.48 a

month per person in the winter and 27.31 in the summer. Furthermore, among the persons in the radon hot spring group, 45.24% took traditional baths and 9.52% took showers. In the control group, no households had hot spring baths. The mean number of baths taken was 3.30 a month per person in the winter and 28.25 in the summer. In the control group, 86.36% of persons took showers, whereas no one took baths.

Instruments and Reagents

Four-color labeled CD3 FITC/CD8 PE/CD45 PerCP/CD4 APC antibody, 2-color labeled CD3 PerCP, and anti-TCR- α -1 FITC antibody were purchased from BD Company (San Jose, California). In addition, BD FAC lysing solution and the FACS Aria system were obtained from BD Company.

Detection of Lymphocyte Subsets, Absolute Values, and Percentages of Lymphocytes

Each specimen was divided into 3 tubes, to which 100 μ L of whole blood was added. In addition, 5 μ L of CD3/CD4/CD8/CD45 antibody and TCR/CD3 antibody were added to each tube. The tubes were then shaken and mixed, and left to react at room temperature for 30 minutes. A volume of 2 mL of prepared erythrocyte lysate was then added, and the tubes were again shaken and mixed, and left to stand at room temperature in the dark for 10 minutes. The tubes were subsequently centrifuged at 1000 rpm for 5 minutes. The supernatant was discarded, and the cells were washed once with $1 \times$ phosphate buffer saline, after which 500 μ L phosphoric acid buffer was added. The cells were then analyzed using a flow cytometer. The various lymphocyte groups were detected and selected, and the percentage of each subset was determined. The absolute values of lymphocytes and percentages of lymphocytes were measured by an automatic blood analyzer.

Epidemiological Investigation on the Prevalence of Rheumatism and Cancer in Radon Hot Spring Area and Control Area

By 2019, the total population of the radon hot spring area was 21 071 and the total population of the control area was 35 762. The prevalence of cancer and rheumatism in the 2 areas was investigated.

Statistical Analyses

SPSS version 21.0 statistical analysis software and the χ^2 test were used to analyze the distribution of basic characteristics between the 2 groups. The χ^2 test was used to analyze the prevalence of rheumatism and cancer between radon hot spring area and control area. The normal distribution of lymphocyte subsets between the 2 groups was evaluated. CD45, CD4⁺/CD8⁺, TCR/CD3 cells, and the number of lymphocytes were

Table 1. Comparison of Basic Data in the 2 Groups.

Group	Number	Gender		Age		Smoking		Alcohol Consumption		BMI (kg/m ²)		
		F	M	≤45	>45	No	Yes	No	Yes	≤18.5	18.5-24	≥24
Control	51	30	21	25	26	40	11	35	16	0	17	34
Radon	61	38	23	35	26	49	12	41	20	0	27	34
χ^2		0.140		0.780		0.061		0.025		1.391		
P		0.708		0.377		0.805		0.873		0.238		

Abbreviations: BMI, body mass index; F, female; M, male.

Table 2. Regression Analysis of Lymphocyte Subsets in Peripheral Blood.

Lymphocyte Subset	Variable	b	Sb	β	t	P
CD45	Constant	96.964	1.041	–	93.186	.000 ^a
	Radon exposure	–0.033	0.014	–0.214	–2.303	–.083
CD3 ⁺	Constant	13.641	0.788	–	17.312	.000 ^a
	Radon exposure	0.001	0.008	0.001	0.001	.999
CD4 [–] CD8 ⁺	Constant	47.366	1.436	–	32.987	.000 ^a
	Radon exposure	–0.038	0.015	–0.232	–2.497	.014 ^a
CD4 ⁺ CD8 [–]	Constant	41.255	1.509	–	27.347	.000 ^a
	Radon exposure	0.019	0.016	0.112	1.177	.242
CD4 ⁺ /CD8 ⁺	Constant	0.966	0.108	–	8.920	.000 ^a
	Radon exposure	0.002	0.001	0.178	1.900	.060
TCR/CD3	Constant	83.782	2.038	–	43.884	.000 ^a
	Radon exposure	–0.055	0.027	–0.192	–2.047	.043 ^a
LYMPH#	Constant	2.275	0.082	–	27.867	.000 ^a
	Radon exposure	0.000	0.001	0.026	0.266	.790
LYMPH%	Constant	32.178	1.143	–	28.147	.000 ^a
	Radon exposure	0.003	0.012	0.026	0.268	.789

Abbreviations: “–” indicates no data; b, regression coefficients; β , standardized partial regression coefficient; Sb, standard error of regression coefficients.

^aA significant difference.

normally distributed so the Student *t* test was applied. The CD4[–]CD8⁺, CD4⁺CD8[–], CD3⁺ cells, and percentage of lymphocytes were not normally distributed; thus, the Mann-Whitney *U* test was used for analysis. Value of *P* < .05 was considered statistically significant.

Results

Basic Data Analysis

The constituent ratios of basic data (gender, age, alcohol consumption, smoking, and body mass index [BMI]) are presented in Table 1. The distribution of basic data showed no significant difference between the 2 groups (*P* > .05).

Multiple Linear Regression Analysis of Lymphocyte Subsets

Multiple linear regression analysis of lymphocyte subsets showed a significant correlation between radon exposure factors and CD4[–]CD8⁺ and TCR/CD3 cells (Table 2).

Table 3. Percentage Change in Lymphocyte Subsets in the Radon Hot Spring Group.

Lymphocyte Subset	Group	N	% ($\bar{x} \pm$ SD)	t	P
CD45	Control	51	97.53 ± 1.272	2.303	–.023 ^a
	Radon	61	94.14 ± 10.446		
CD4 ⁺ /CD8 ⁺	Control	51	0.93 ± 0.387	–2.201	.03 ^a
	Radon	61	1.26 ± 1.041		
TCR/CD3	Control	51	83.78 ± 19.112	2.047	.043 ^a
	Radon	61	89.43 ± 5.259		
LYMPH#	Control	49	2.260 ± 0.578	–0.485	.629
	Radon	56	2.317 ± 0.622		

Abbreviation: SD, standard deviation.

^aA significant difference.

Comparison of Lymphocyte in Peripheral Blood Subsets Between the 2 Groups

The percentage of CD45, CD4⁺/CD8⁺, and TCR/CD3 lymphocyte subsets was significantly higher in the group exposed to radon than that in the control group (*P* < .05; Table 3). Compared with the control group, the percentage of CD4[–]CD8⁺ lymphocytes was significantly lower in the group exposed to radon. In contrast, the percentage of CD4⁺CD8[–]

Table 4. Comparison of the Percentage Change in Lymphocyte Subsets by Mann-Whitney *U* Test.

Lymphocyte Subset	Group	N	P50	Z	P
CD3 ⁺	Control	51	12.90	-0.140	.888
	Radon	61	13.50		
CD4 ⁻ CD8 ⁺	Control	51	48.00	-2.141	.032 ^a
	Radon	61	43.80		
CD4 ⁺ CD8 ⁻	Control	51	38.80	-0.964	.335
	Radon	61	42.30		
LYMPH%	Control	49	50.35	-0.835	.404
	Radon	56	55.32		

^aA significant difference.

Table 5. Comparison of the Prevalence of Rheumatism and Cancer Between Radon Hot Spring Area and Control Area.

Disease Name	Radon	Control	χ^2	P
Rheumatism			14.447	<.001 ^a
Prevalence	34	21		
Nonprevalence	21 037	35 741		
Cancer			6.751	.009 ^a
Prevalence	23	72		
Nonprevalence	21 048	35 690		
Rheumatism + cancer			0.055	.814
Prevalence	57	93		
Nonprevalence	21 014	35 669		

^aA significant difference. Prevalence, number of sick people; nonprevalence, number of people not ill; rheumatism + cancer, sum of the number of patients with rheumatism and cancer.

and CD3⁺ lymphocytes was higher in the radon group, as compared with that in the control group; however, this difference was not significant (Table 4).

Comparison of the Prevalence of Rheumatism and Cancer Between Radon Hot Spring Area and Control Area

Compared with the control group, there was no significant difference in the total prevalence between the 2 area ($P = .814$). The prevalence of rheumatism in radon hot spring area was higher (0.2% vs 0.1%, $P < .001$), the prevalence of cancer was lower (0.1% vs 0.2%, $P = .009$) (Table 5).

Discussion

Studies have shown that low-dose radiation can induce hormetic effects in organisms.¹⁷ Liu¹⁸ observed the results of immune enhancement by low-dose radiation through a high-background radiation study in Yangjiang, China, and a series of animal experiments. They found the change in T lymphocytes was the key factor of immune enhancement by low-dose ionizing radiation, which was conducive to T-cell differentiation toward T-helper (Th) 1 cells and could have induced an adaptive response. At present, the low-dose radiation hormetic

effect is evaluated mainly by low-linear energy transfer (LET) radiation. However, whether low-dose high-LET radiation causes hormetic effects have been less reported.

The impact of radon and its daughters on human health arise mainly from the α particles produced by its decay, which is high-LET irradiation. Studies on α -particles have been conducted mainly in vivo through animal experiments, cell levels in vitro, and clinical treatment of certain human diseases using radon hot spring. Chauhan et al¹⁹ irradiated human pulmonary epithelial cells with different doses of α -particles. Pathway analyses showed that radiation from α -particles could affect cell-cycle arrest, DNA replication, and repair of DNA damage. Those data suggested that high-LET radiation could activate different biological pathways compared with low-LET radiation.

In the present study, the concentration of radon in the hot springs was 102 ± 11.4 Bq/L, the concentration of indoor radon was 41.9 ± 18.6 Bq/m³ over 6 months, and the typical equilibrium factor was 0.61. The radon concentration in the bathroom increases rapidly during water use. The indoor radon concentration increases rapidly from the normal background value (<50 Bq/m³) to more than 200 Bq/m³. After about 3 hours, the indoor radon concentration recovers gradually to a lower level. The variation in the concentration of radon daughters is consistent with that of radon concentration. The additional annual effective dose is about 0.09 mSv.¹⁶ Some residents have been exposed to low-dose ionizing radiation for a long time (more than 5 years). Therefore, radon in hot springs is considered long-term low-dose and high-LET radiation.

Our research team has studied genetic/molecular levels and antioxidant damage in residents of the radon hot spring in Pingshan area.²⁰⁻²⁴ Ruan et al²⁰ compared the micronucleus frequency and micronucleus cell numbers in different age groups of 2 groups of residents. They found that the micronucleus rate and micronucleus cell rate of the radon hot spring group were significantly higher than those of the control group, but that micronucleus rate was within the normal range. Furthermore, they found no significant differences in the chromosome aberration rate compared with that in the control group. In studies by Liu et al²¹⁻²² and Gao et al²³ of the lymphocyte cell cycle and regulatory proteins, upon radon exposure, the S-phase proportion of lymphocytes in peripheral blood was increased; expression of related cyclic regulatory proteins CDK1, CDK6, and cyclin E1 was decreased; expression of microRNA-449a was increased significantly in the radon hot spring group; and expression of cyclin E1 and CDK1 was decreased due to upregulation of microRNA-16 expression. Gao et al²⁴ showed that radon can activate antioxidant function, scavenge oxygen free radicals, and reduce oxidative damage. These phenomena may be related to the hormesis of low-dose radon exposure, which is manifested in various facets. In present study, we observed the effect of long-term low-dose radiation on immune function.

Lymphocytes are radiation sensitive and comprise the main immune cell type. Therefore, they are often used as an important indicator for the early diagnosis of, and dose estimation in

radiation sickness. T lymphocytes are important immunocompetent cells. In addition to mediating cellular immune functions directly, T lymphocytes have key roles in regulating the immune response. The molecules on the surface of T lymphocytes enable these lymphocytes to recognize antigens, interact with other immune cells, and receive signal stimulation, among other functions. These membrane-surface molecules also play a key part in identifying and separating T lymphocytes and T-lymphocyte subsets. The T-lymphocyte subsets play an important role in regulating the stability of immune function.

The distribution of lymphocyte subsets can reflect the immune status of the body. Studies²⁵⁻³⁰ have shown that the distribution of lymphocyte subsets is influenced by age, gender, ethnicity, and region. Therefore, we first compared general data (gender, age, alcohol consumption, smoking, BMI) of the 2 groups in Pingshan County. No significant differences were noted in the constituent ratios between the 2 groups ($P > .05$), and both were comparable.

We found that the percentage of CD3⁺ cells and CD4⁺ cells in the peripheral blood of residents living near the high-radon hot springs in Pingshan was higher than that of residents living in the control areas, but this difference was not significant. In addition, the percentage of CD8⁺ cells was significantly lower, and the percentage of TCR/CD3 complexes, and ratio of CD4⁺/CD8⁺ were significantly higher in the residents living near the high-radon hot springs in Pingshan group than in the control area, and these differences were significant ($P < .05$). Routine blood testing showed that the percentage of lymphocytes and absolute values of lymphocytes in peripheral blood showed no significant change in the residents of the high-radon hot spring area compared with the control group.

Lymphocyte subsets regulate each other in terms of the immune response to maintain normal immune function. Both Th (CD4⁺) cells and Ts (T inhibition cells, CD8⁺) cells play an important regulatory role in the immune response. CD8⁺ cells play an effective cell-mediated immune killing role after being stimulated by foreign antigens, which destroy target cells directly. In addition, CD4⁺ cells, sensitized by foreign antigens, produce lymphokines, which induce the proliferation of T lymphocytes, B lymphocytes, and macrophages. The balance between the proportions of CD4⁺ and CD8⁺ T cells plays an important role in inhibiting tumors, infections, and autoimmune diseases. Immune dysfunction can be caused by an imbalance between these cells, resulting in a series of immunopathological changes that can affect the immune protective mechanisms.^{31,32}

In the present study, the percentage of TCR/CD3 cells in the peripheral blood of residents near the radon hot springs was increased significantly, which may have been related to the hormesis effect caused by low-dose radiation. Low-dose low-LET radiation accelerates the renewal and growth of thymocytes and, thus, increases the number of thymocytes, and accelerates upregulation of TCR/CD3 expression. Therefore, the ability of the thymus to transport T cells to the periphery is enhanced. Simultaneously, cytokine secretion is stimulated, and the intercellular interactions form an enhanced network

which, ultimately, improves the anticancer and anti-infection capabilities of the body. Our study implied that long-term exposure to low-dose high-LET radiation also has the same hormesis effect on the immune system.

Anderson et al³³ suggested that “low-dose” radiation might inhibit Ts with higher sensitivity, thus altering the ratio of Th to Ts, leading to the dominant position of Th, increasing the immunopotentiatory regulation of Th, and enhancing immune function. In their study of the long-term immune effects of high-level natural radiation on residents in Yangjiang, China, Kun et al³⁴ also observed that long-term exposure to low-dose radiation stimulated different CD4⁺ and CD8⁺ T lymphocytes and enhanced immune function. However, few reports have focused on the effects of low-dose and high-LET radiation on immune function. Yamaoka et al³⁵ found that radon treatment and heat treatment could enhance antioxidant capacity, increase the percentage of CD4⁺ cells, and reduce the percentage of CD8⁺ cells among killer T cells and suppressor T cells. Those findings are consistent with our results.

Based on our previous research in this area,^{23,24} 2 factors may be involved in enhancing the immune function among residents of the radon hot springs region in Pingshan County. First, the proportion of S-phase lymphocytes in peripheral blood increased and that of the G0/G1 phase decreased among inhabitants around radon hot springs. These data indicated that the S phase was prolonged, radiation sensitivity of the S phase was poor, and this phase marked a period of DNA repair/synthesis. Lymphocytes might be protected by prolonging the S phase in conditions of long-term low-dose radiation. Second, several studies in China and overseas³⁶⁻⁴¹ have shown that long-term bathing in radon hot springs leads to reductions in oxidative damage and that the ability to scavenge reactive oxygen species increases. The level of 8-hydroxy-2'-deoxyguanosine in the peripheral-blood plasma of residents in the radon hot spring group in Pingshan County was decreased, whereas the level of thioredoxin reductase was increased significantly. These effects may activate antioxidant functions, reduce oxidative damage, and protect radiation-sensitive lymphocytes.

The cancer prevalence in the radon hot springs area was significantly lower than that in the control area (0.1% vs 0.2%, $P = .009$). Combined with laboratory analyses of the immune function in the 2 areas, we postulated that the decrease in cancer prevalence in the radon hot springs area may have been due to long-term low radon exposure, which can stimulate immune function, activate the antioxidant capacity of the body, and reduce the oxidative damage caused by radiation. However, the prevalence of rheumatic diseases in the radon hot springs area was significantly higher than that in the control area. The complex etiology of rheumatic diseases (eg, environmental conditions, living conditions) and many other factors can affect the immune function. The cause of the increased prevalence of rheumatic diseases in radon hot springs areas needs to be studied further. Because the sample size of our study was small, how radon hot springs influence lymphocyte subsets and immune function must be tested further in much larger cohorts.

Conclusions

Radon hot springs in Pingshan County posed no obvious threats to the health of residents and may enhance immune function.

Authors' Note

Chunnan Piao undertook on-site sample collection, processing of laboratory samples data analyses, and wrote the manuscript. Mei Tian and Hongjun Gao guided the research ideas, research programs, experimental methods, and manuscript revision. Jianlei Ruan, Yanxiao Gao, Gang Gao, Lirong Yi, and Lina Wu contributed significantly to on-site investigation, sample collection, and guidance for experimental methods. Jianxiang Liu conceived and designed the work and approved the final version of the manuscript.

Acknowledgments

The authors thank the staff of Hebei CDC, Pingshan CDC, and all participants for their support and cooperation.


Declaration of Conflicting Interests


The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding

The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: This study was sponsored by the National Institute for Radiological Protection, Chinese Center for Disease Control and Prevention.

ORCID iD

Chunnan Piao  <https://orcid.org/0000-0002-4349-9564>

Yanxiao Gao  <https://orcid.org/0000-0002-3552-052X>

References

1. Veiga LH, Amaral EC, Colin D, Koifman S. A retrospective mortality study of workers exposed to radon in a Brazilian underground coal mine. *Radiat Environ Biophys.* 2006;45(2):125-134.
2. Shang YQ. Progress on adverse effects of indoor radon. *Environ Health.* 2002;19:350-352.
3. Shang B, Cui HX, Wen XM, et al. Survey of radiation levels in an abnormally high radon hot spring. *Chin J Radiol Med Protect.* 2011;31:698-702.
4. Alharbi SH, Akber RA. Radon and thoron concentrations in public workplaces in Brisbane, Australia. *J Environ Radioact.* 2015; 144:69-76.
5. Gruber V, Ringer W, Wurm G, Haider W. The Austrian radon activities on the way to the National Radon Action Plan. *Radiat Prot Dosimetry.* 2014;160(1-3):22-26.
6. Valmari T, Arvela H, Reisbacka H, Holmgren O. Radon measurement and mitigation activity in Finland. *Radiat Prot Dosimetry.* 2014;160(1-3):18-21.
7. Ayres da Silva ALM, Eston SM, Iramina WS, Diegues Francisca D. Radon in Brazilian underground mines. *J Radiat Prot.* 2018; 38(2):607-620.
8. Yamaoka K, Mitsunobu F, Hanamoto K, Mori S, Tanizaki Y, Sugita K. Study on biologic effects of radon and thermal therapy on osteoarthritis. *J Pain.* 2004;5(1):20-25.
9. Mitsunobu F, Yamaoka K, Hanamoto K, et al. Elevation of antioxidant enzymes in the clinical effects of radon and thermal therapy for bronchial asthma. *J Radiat Res.* 2003;44(2): 95-99.
10. Shehata M, Schwarzmeier JD, Hilgarth M, et al. Effect of combined spa-exercise therapy on circulating TGF-beta1 levels in patients with ankylosing spondylitis. *Wien Klin Wochenschr.* 2006;118(9-10):266-272.
11. Cucu A, Shreder K, Kraft D, et al. Decrease of markers related to bone erosion in serum of patients with musculoskeletal disorders after serial low-dose radon spa therapy. *Front Immunol.* 2017;8: 882.
12. Annegret F, Thomas F. Long-term benefits of radon spa therapy in rheumatic diseases: results of the randomised, multi-centre IMuRa trial. *Rheumatol Int.* 2013;33(11):2839-2850.
13. Franke A, Reiner L, Pratzel HG, Franke T, Resch KL. Long-term efficacy of radon spa therapy in rheumatoid arthritis—a randomized, sham-controlled study and follow-up. *Rheumatology (Oxford).* 2000;39(8):894-902.
14. Rühle PF, Klein G, Rung T, et al. Impact of radon and combinatory radon/carbon dioxide spa on pain and hypertension: results from the explorative RAD-ON01 study. *Mod Rheumatol.* 2019; 29(1):165-172.
15. Rühle PF, Wunderlich R, Deloch L, et al. Modulation of the peripheral immune system after low-dose radon spa therapy: detailed longitudinal immune monitoring of patients within the RAD-ON01 study. *Autoimmunity.* 2017;50(2):133-140.
16. Zhao MQ, Cui HX, Shang B, et al. Radon level and dose contribution in Pingshan hot spring, Hebei Province [in Chinese]. *Chin J Radiol Health.* 2012;21:30-32.
17. Liu SZ. On radiation hormesis expressed in the immune system. *Crit Rev Toxicol.* 2003;33(3-4):431-441.
18. Liu SZ. Review and prospect of radiation immunology [in Chinese]. *Chin J Radiol Med Prot.* 2005;25:193-200.
19. Chauhan V, Howland M, Mendenhall A, et al. Effects of alpha particle radiation on gene expression in human pulmonary epithelial cells. *Int J Hyg Environ Health.* 2012;215(5):522-535.
20. Ruan JL, Liu CX, Wu LN, et al. Change of micronucleus in peripheral blood lymphocytes of the residents surrounding hot springs with radon. *Int J Radiat Med Nuclear Med.* 2015;39: 363-366.
21. Liu CX, Li XL, Pan Y, et al. Expression of miRNAs in peripheral blood lymphocytes of the residents around hot springs with radon [in Chinese]. *Carcin Teratogen Mut.* 2015;27:446-449.
22. Liu CX, Tian M, Pan Y, Gao G, Liu J. Expression of miRNAs in peripheral blood plasma of the residents surrounding hot springs with radon [in Chinese]. *Chin J Radiol Med Prot.* 2015;35(3): 187-190.
23. Gao YX, Tian M, Gao G, et al. Cell cycle and its regulatory proteins in the peripheral blood lymphocytes of the residents living in a radon hot spring area [in Chinese]. *Chin J Radiol Med Prot.* 2018;38(1):12-16.
24. Gao YX, Tian M, Gao G, Xiaochun W, Jianxiang L. Changes of 8-OHdG and TrxR in the residents who bathe in radon hot springs. *Dose-Response.* 2019;17(1):1559325818820974.

25. Zhan WL, Yang XH, Guo H, et al. Distribution of peripheral blood lymphocyte subsets in 826 healthy children aged 0-6 years [in Chinese]. *Zhongguo Dang Dai Er Ke Za Zhi*. 2019;21(2):180-183.
26. Castilho JL, Shepherd BE, Koethe J, et al. CD4/CD8 ratio, age, and risk of serious noncommunicable diseases in HIV-infected adults on antiretroviral therapy. *AIDS*. 2016;30(6):899-908.
27. Zhao H, Jiao SC, Zhang GQ, Wu LL. Analysis of 21 lymphocytes for the different age and gender healthy population [in Chinese]. *Labeled Immunoassays Clin Med*. 2011;18(2):99-102.
28. Shearer WT, Rosenblatt HM, Gelman RS, et al. Lymphocyte subsets in healthy children from birth through 18 years of age: the pediatric AIDS clinical trials group P1009 study. *J Allergy Clin Immunol*. 2003;112(5):973-980.
29. Ding Y, Zhou L, Xia Y, et al. Reference values for peripheral blood lymphocyte subsets of healthy children in China. *J Allergy Clin Immunol*. 2018;142(3):970-973.
30. Guo ZH, Yao YP, Li XT, et al. Investigation on T lymphocyte subsets in healthy teenagers and adults in Zhejiang areas and establishment of the reference range for normal values of T lymphocyte subsets. [in Chinese] *Chin Prev Med*. 2008;9:32-35.
31. Chen WF. *Medical Immunology*. 3rd ed. Beijing, China: People's Health Publishing House. 2000:94-96.
32. Keane WF, Lyle PA. Recent advances in management of type 2 diabetes and nephropathy: lessons from the RENAAL study. *Am J Kidney Dis*. 2003;41(3 suppl 1):S22-S25.
33. Anderson RE, Williams WL, Tokuda S. Effect of low dose irradiation upon T cell subsets involved in the response of primed A/J mice to SaI cells. *Int J Radiat Biol Relat Stud Phys Chem Med*. 1988;53(1):103-118.
34. Li K, Li W, Jia Y, et al. Long-term immune effects of high-level natural radiation on Yangjiang inhabitants in China. *Int J Radiat Biol*. 2019;95(6):764-770.
35. Yamaoka K, Mitsunobu F, Hanamoto K, et al. Biochemical comparison between radon effects and thermal effects on humans in radon hot spring therapy. *J Radiat Res*. 2004;45(1):83-88.
36. Kataoka T, Nishiyama Y, Yamato K, et al. Comparative study on the inhibitory effects of antioxidant vitamins and radon on carbon tetrachloride-induced hepatopathy. *J Radiat Res*. 2012;53(6):830-839.
37. Kataoka T, Yamato K, Nishiyama Y, et al. Comparative study on the inhibitory effects of A-tocopherol and radon on carbon tetrachloride-induced renal damage. *Ren Fail*. 2012;34(9):1181-1187.
38. Kojima S, Tsukimoto M, Shimura N, Koga H, Murata A, Takara T. Treatment of cancer and inflammation with low-dose ionizing radiation: three case reports. *Dose-Response*. 2017;15(1). doi:10.1177/1559325817697531.
39. Kuciel-Lewandowska JM, Pawlik-Sobecka L, Płaczkowska S, Kokot I, Paprocka-Borowicz M. The assessment of the integrated antioxidant system of the body and the phenomenon of spa reaction in the course of radon therapy: a pilot study. *Adv Clin Exp Med*. 2018;27(10):1341-1346.
40. Yamaoka K, Mifune T, Mitsunobu F, et al. Basic study on radon effects and thermal effects on humans in radon therapy. *Physiol Chem Phys Med NMR*. 2001;33(2):133-138.
41. Chen H, Guo Q, Liu M, et al. The effects of low dose radiation on the levels of oxidative damage and antioxidant in population of high background radiation area of Guangdong [in Chinese]. *Chin J Radiol Med Prot*. 2015;35:83-87.