

Pregnancy, perinatal and postpartum complications as determinants of postpartum depression: the Rhea mother–child cohort in Crete, Greece

K. Koutra^{1,2*}, M. Vassilaki³, V. Georgiou¹, A. Koutis¹, P. Bitsios², M. Kogevinas^{4,5,6,7} and L. Chatzi¹

¹ Department of Social Medicine, Faculty of Medicine, University of Crete, Heraklion, Greece

² Department of Psychiatry and Behavioral Sciences, Faculty of Medicine, University of Crete, Heraklion, Greece

³ Department of Health Sciences Research, Mayo Clinic, Rochester, MN, USA

⁴ National School of Public Health, Athens, Greece

⁵ Centre for Research in Environmental Epidemiology (CREAL), Barcelona, Spain

⁶ Municipal Institute of Medical Research, Barcelona, Spain

⁷ CIBER Epidemiología y Salud Pública (CIBERESP), Madrid, Spain

Aims. Few epidemiological studies evaluated associations between perinatal complications and maternal mood at the early postpartum period and the findings are inconsistent. We aimed at investigating a wide range of complications during pregnancy, at delivery, and at the early postpartum period as determinants of postpartum depression (PPD) at 8 weeks postpartum.

Methods. A total of 1037 women who enrolled in the Rhea mother–child cohort in Crete, Greece participated in the present study. Information on pregnancy, perinatal and postpartum complications was obtained from clinical records or by questionnaires. Postpartum depressive symptoms were assessed at 8 weeks postpartum using the Edinburgh Postnatal Depression Scale (EPDS). Multivariable linear and logistic regression models were fit to estimate the association between pregnancy, perinatal and postpartum complications and maternal depressive symptoms, adjusting also for potential confounders.

Results. The prevalence of women with probable depression (EPDS score ≥ 13) was 13.6% at 8 weeks postpartum. Gestational hypertension and/or preeclampsia (β coefficient 1.86, 95% CI: 0.32, 3.41) and breastfeeding difficulties (β coefficient 0.77, 95% CI: 0.02, 1.53) were significantly associated with higher PPD symptoms. Sleep patterns during pregnancy, such as sleep deprivation (OR=3.57, 95% CI: 1.91, 6.67) and snoring (OR=1.81, 95% CI: 1.11, 2.93), and breastfeeding duration less than 2 months (OR=1.77, 95% CI: 1.19, 2.64) were significantly associated with increase in the odds for PPD. Some other complications, such as unplanned pregnancy and hospitalisation during pregnancy were also associated with EPDS score, but these associations were explained by socio-demographic characteristics of the mother.

Conclusions. We found that several pregnancy, perinatal and postpartum complications may have an adverse effect on maternal mood at the early postpartum period. These findings have considerable implications for developing effective prevention and early psychoeducational intervention strategies for women at risk of developing PPD.

Received 29 June 2016; Accepted 19 November 2016; First published online 22 December 2016

Key words: Perinatal, postpartum complications, postpartum depression, pregnancy.

Introduction

It is estimated that around 10–15% of new mothers experience postpartum depression (PPD) with most cases developing depressive symptoms in the first 3 months after giving birth (O'Hara & Swain, 1996; Evans *et al.* 2001; Josefsson *et al.* 2001; Bennett *et al.*

2004; Gavin *et al.* 2005; Halbreich & Karkun, 2006; Norhayati *et al.* 2015). Depressive symptoms can vary from mild complaints and 'maternity blues' to clinically diagnosed PPD. Depression at the postpartum period is often not diagnosed early enough, resulting in women being depressed for longer periods of time before they are given effective treatment. The prevalence of PPD in Greece (Gonidakis *et al.* 2008; Leonardou *et al.* 2009) is similar to that reported in other countries (Cox *et al.* 1993; Nonacs & Cohen, 2002).

PPD is considered to be a systemic illness affecting the new mother's mental health and functioning, as

* Address for correspondence: K. Koutra, MSc, Ph.D., Department of Social Medicine, Faculty of Medicine, University of Crete, PO Box 2208, Heraklion, 71003 Crete, Greece.
(Email: koutra.k@gmail.com)

well as her relationship with her infant and family (O'Hara, 1997; Grace *et al.* 2003; Howard *et al.* 2014). Furthermore, PPD has significant adverse impact on infant and child development (Cornish *et al.* 2005; Koutra *et al.* 2013). A range of biological and psychosocial risk factors have been identified as important determinants of PPD (Beck, 1996, 2001; O'Hara & Swain, 1996). Previous research suggests that socio-demographic characteristics, such as mother's young age (Rubertsson *et al.* 2003), low educational level (Davis *et al.* 2003) and low socio-economic background (Beck, 2001), as well as psychosocial factors, such as antenatal depression and anxiety (Koutra *et al.* 2014; Tebeka *et al.* 2016), are associated with increased levels of PPD.

While there is an extensive literature on the socio-demographic and the psychosocial determinants of PPD, there are only few epidemiological studies with inconsistent findings that have examined the effects of perinatal complications on PPD (Warner *et al.* 1996; Nielsen Forman *et al.* 2000; Johnstone *et al.* 2001; Blom *et al.* 2010). Some studies found no association between pregnancy or delivery complications and increased risk for developing PPD (Nielsen Forman *et al.* 2000; Johnstone *et al.* 2001), while others revealed a number of obstetric risk factors strongly associated with PPD (Warner *et al.* 1996; Blom *et al.* 2010). More specifically, Blom *et al.* (2010) showed that several pregnancy complications including pre-eclampsia, hospitalisation during pregnancy, emergency caesarean section, concern about fetal distress and newborn's admission to special care were associated with PPD at 2 months postpartum, and the risk of developing PPD increased with the number of complications women experienced. In a similar vein, Warner *et al.* (1996) suggested that unplanned pregnancy, formula feeding instead of breastfeeding, and unemployment in either the mother or the head of household were significant risk factors for PPD at 6–8 weeks postpartum. Two meta-analyses (O'Hara & Swain, 1996; Robertson *et al.* 2004) found that obstetric factors had a significant but small effect on the prevalence of PPD.

From a public health point of view, identifying perinatal risk factors for PPD is of major importance, due to the disorder's high prevalence and its consequences for the new mother and her family. To our knowledge, there are studies which examined a composite score of obstetric complications (O'Hara & Swain, 1996), thus making it difficult to determine which specific complications predict PPD. In addition, some studies reported associations between obstetric complications and PPD that were only minimally adjusted (Warner *et al.* 1996), and thus may have been confounded by unmeasured factors. Moreover, the majority of the

studies examining specific perinatal complications as risk factors for PPD is mainly focused at pregnancy and delivery complications without concurrent consideration of sleep patterns during pregnancy (i.e. sleep deprivation, snoring, daytime sleepiness) and possible complications women experiencing at the early postpartum period (i.e. breastfeeding difficulties and discontinuation). Prior research suggests a possible relationship between sleep patterns during pregnancy and maternal mood during the postpartum period (Lee *et al.* 2000; Wolfson *et al.* 2003; Goyal *et al.* 2007; Chang *et al.* 2010; Tomfohr *et al.* 2015). Furthermore, short duration of breastfeeding or not breastfeeding (Dennis & McQueen, 2007; Watkins *et al.* 2011) and breastfeeding difficulties (Mezzacappa & Katlin, 2002; Ystrom, 2012; Figueiredo *et al.* 2014; Dias & Figueiredo, 2015) have been associated with PPD symptoms.

Within the context of a population-based mother-child cohort study in Crete, Greece (Rhea Study), our study aimed at investigating a wide range of complications during pregnancy, at delivery, or at the early postpartum period as determinants of PPD at 8 weeks postpartum.

Methods

The mother-child cohort in Crete (Rhea Study)

The present study is part of the Rhea Study, a prospective mother-child cohort examining a population sample of pregnant women and their children at the prefecture of Heraklion, Crete, Greece (Chatzi *et al.* 2009a). Female residents (Greek and immigrants) who had become pregnant during the 12-month period starting in February 2007 have been contacted at four maternity clinics in Heraklion and asked to participate in the study. To be eligible for inclusion in the study, women had to have a good understanding of the Greek language and be older than 16 years of age. The first contact was made at the time of the first major ultrasound examination, before week 15 of gestation. Women were informed about the study protocol by trained nurses and midwives and asked to participate in the study. Women were then contacted again at the third trimester of pregnancy, at birth, and at 8 weeks postpartum. Face-to-face structured interviews, together with self-administered questionnaires and medical records, were used to obtain information on several psychosocial, dietary and environmental exposures during pregnancy and postpartum. The study was approved by the Ethical Committee of the University Hospital in Heraklion, Crete, Greece. Written informed consent was obtained from all participants.

Detailed characteristics of the study population have been described elsewhere (Chatzi et al. 2009a). During the study recruitment period 1765 eligible women were approached, 1610 (91%) agreed to participate and 1388 (86%) were followed up until delivery. A total of 1079 women completed the depression scale at 8 weeks postpartum and 1072 women had full data on basic exposure information (gestational age, preterm birth and type of delivery) and PPD. Seven women with a previous diagnosis of severe psychiatric disorders (i.e. schizophrenia, bipolar disorder, etc.) and 28 multiple pregnancies were excluded from the sample, resulting in a cohort of 1037 women available for analyses.

Measures

Pregnancy, perinatal and postpartum complications

The present study examined a wide range of pregnancy, perinatal and postpartum complications as risk factors for PPD. Information on the following complications was obtained through questionnaires administered to pregnant women by trained interviewers during the first or third trimester of pregnancy. Upon enrolment, women reported whether or not the pregnancy was planned, and about how women were trying to get pregnant (i.e. immediately v. with effort, subjective assessment of women). At 30 weeks of gestation, women answered a question about whether they had symptoms like vomit, nausea, etc. during the first two trimesters of pregnancy and a question about whether they had been admitted to hospital for more than 24 h during the first two trimesters of pregnancy. Furthermore, women were asked if they were smoking during pregnancy and if they were drinking alcohol during pregnancy.

Women were screened for gestational diabetes at 24–28 weeks of gestation and were classified as having gestational diabetes in the index pregnancy if two or more of the four plasma glucose values obtained during the 100-g 3-h oral glucose tolerance test (OGTT) were abnormal, as per the criteria proposed by Carpenter & Coustan (1982): fasting blood glucose ≥ 95 mg/dl; 1-h ≥ 180 mg/dl; 2-h values ≥ 155 mg/dl; and 3-h values ≥ 140 mg/dl (Chatzi et al. 2009a). All participants were classified as having metabolic syndrome or not, according to the criteria proposed by the US National Heart, Lung and Blood Institute/American Heart Association (NHLBI/AHA) (Grundy et al. 2005), with some considerations taken into account to adapt to our study population of pregnant women, as described in previous studies from the same cohort (Chatzi et al. 2009a, b). In particular, the metabolic syndrome was diagnosed if three or more of

the following risk factors were present: a pre-pregnancy BMI of >30 kg/m²; a triglyceride level of ≥ 150 mg/dl; a high-density lipoprotein (HDL)-cholesterol level of <50 mg/dl; a fasting glucose level of ≥ 100 mg/dl, and a blood pressure level of $\geq 130/\geq 85$ mm Hg. In this study, we tested metabolic syndrome as both number of symptoms and its components specifically. Information on gestational hypertension and/or pre-eclampsia and thyroid dysfunction during pregnancy was obtained through computer-assisted interviews and medical records at the first and third trimesters. Maternal pre-pregnancy BMI was calculated by height, measured at the first prenatal visit, and pre-pregnancy weight, as reported at the time of the first major ultrasound visit (BMI; weight [kg]/height[m]²). Maternal gestational weight gain was calculated by body weight at delivery (objective) – pre-pregnancy weight (self-reported). Information on sleep patterns, including sleep duration, sleep-disordered breathing (i.e. non-snorers v. snorers) and daytime sleepiness was collected through a computer-assisted interview at the third trimester of pregnancy and was based on the following questions: ‘During the past month, how many hours do you sleep per day? How often do you snore during your sleep? How fresh do you feel when you wake up in the morning, independently of the hours you have slept?’ The answers were classified on 5-point Likert scales (Micheli et al. 2011). Sleep deprivation was defined as five or fewer hours of sleep (Sabanayagam & Shankar, 2010). Snoring symptoms were also recoded into two categories: non-snorers v. snorers. The Epworth Sleepiness Scale was used to determine the level of daytime sleepiness; excessive daytime sleepiness was defined as a total score ≥ 10 (Johns, 1991).

Gestational age, type of delivery and anthropometric measures at birth were collected from clinical records. Gestational age was based on the interval between the last menstrual period and the date of delivery. When the menstrual estimate of gestational age was inconsistent by seven or more days with the ultrasound measurement taken in the first trimester of pregnancy, a quadratic regression formula describing the relation between crown rump length and gestational age was used instead (Chatzi et al. 2009a). Preterm birth was defined as a birth at less than 37 weeks. Type of delivery was divided into vaginal and caesarean section. Birth weight of the child was also obtained from routine midwife and hospital registry records and low birth weight was defined as birth weight below 2500 g.

Information on breastfeeding initiation, duration, and difficulties was collected at 9 months postpartum, but for this analysis only information up to 8 weeks postpartum was used. Mothers were asked if they had ever

breastfed their child (or placed the child on their breast to feed). If they never breastfed their child, the reason was recorded. If women initiated breastfeeding, further information on breastfeeding duration was asked, as well as information regarding the first time they breastfed their infant and the duration of breastfeeding. Finally, at 9 months postpartum, the women reported retrospectively whether or not their *newborn was admitted to Intensive Care Unit (ICU)* immediately after delivery.

Psychological assessment at the postpartum period

Maternal depressive symptoms were assessed at 8 weeks postpartum using the Edinburgh Postnatal Depression Scale (EPDS; Cox *et al.* 1987). The EPDS is a widely used 10-item self-reported questionnaire providing an indication of the severity of mother's mood during the past 7 days. Items are rated on a 4-point Likert scale ranging from 0 (not at all) to 3 (most of the time) and refers to depressed mood, anhedonia, guilt, anxiety and suicidal ideation (possible range 0–30). A cut-off score of 13 or greater on the EPDS has been found to identify probable clinical postnatal depression with a sensitivity of 86% and a specificity of 78% (Cox *et al.* 1987; Matthey *et al.* 2006). This cut-off is also consistent with previous work in our cohort (Koutra *et al.* 2013, 2014). The EPDS has been translated and validated for the Greek population by two research groups (Leonardou *et al.* 2009; Vivilaki *et al.* 2009), and showed a very high overall internal consistency.

Potential confounders

Based on the existing literature, we considered the following non-obstetric factors as possible confounders in the association between pregnancy, perinatal and postpartum complications and PPD: maternal age at delivery; education (low level: ≤ 6 years of school, medium level: 7–12 years of school, and high level: some years in university or university degree, ref: low level); marital status (single/married-engaged); Greek origin (yes/no); parity (primiparous/multiparous); and working during pregnancy (yes/no). Information on the aforementioned characteristics was obtained by questionnaires upon enrolment.

Statistical analysis

Descriptive statistics were used to summarise the baseline characteristics of participants. The primary outcome variable of interest was the PPD score at 8 weeks postpartum. The primary exposures of interest were pregnancy, perinatal and postpartum

complications. EPDS was used as both a continuous and dichotomous variable with a cut-off score of 13 or greater recommended by Cox *et al.* (1987) which appeared to be an effective screening for probable clinical depression.

Bivariate associations between dependent and independent variables were studied using Pearson's χ^2 test or Fisher exact test for categorical variables. The EPDS score failed the normality test; hence, non-parametric tests were applied (Spearman's correlation, Kruskal–Wallis and Mann–Whitney tests). Missing values have been excluded pair-wise. Multivariable linear and logistic regression models were fit to estimate the association between pregnancy, perinatal and postpartum complications with maternal depressive symptoms at 8 weeks postpartum. Potential confounders related with either the outcome or the exposure of interest in the bivariate associations with a p -value < 0.2 (i.e. maternal age, education, marital status and working during pregnancy), as well as *a priori* selected potential confounders (i.e. maternal origin and parity) were included in the multivariable models. To account for the possibility of confounding by antenatal depressive symptoms, we performed additional sensitivity analyses excluding 77 women with high levels of depression during pregnancy (antenatal EPDS ≥ 13) Maternal antenatal depressive symptoms were assessed at 28–32 weeks of gestation using the EPDS (Cox *et al.* 1987). Finally, heterogeneity in associations related to parity (primiparous *v.* multiparous) was evaluated by including interaction terms in the multivariable models (statistically significant effect modification if p -value < 0.10) and by stratifying the sample, accordingly.

Estimated associations are described in terms of β coefficients (linear regression models) with 95% confidence intervals (CIs) or odds ratios (OR) with 95% CIs (logistic regression models). All hypothesis testing was conducted assuming a 0.05 significance level and a two-sided alternative hypothesis. All statistical analyses were performed using SPSS Statistics 20.0 software (IBM, Armonk, NY, USA).

Results

The socio-demographic characteristics of the study population are presented in Table 1. Participating mothers were predominantly of Greek origin (93.0%), married (88.4%) and had a mean age at delivery of 29.51 (s.d. = 4.95) years. More than half of the mothers had medium level of education (52.3%) and were multiparous (58.6%). About half of the infants were boys (51.1%) with a mean gestational age of 38.24 (s.d. = 1.51) weeks. Women of younger age, with low levels

Table 1. Socio-demographic characteristics of participants (Rhea Study, Crete, Greece)

Characteristics	EPDS				EPDS < 13 N (%)	EPDS ≥ 13 N (%)	p-value
	N	%	Mean ± s.d.	p-value			
Maternal origin							
Other	72	7.0	6.89 ± 4.97	0.638	61 (6.9)	11 (7.9)	0.681
Greek	952	93.0	6.70 ± 5.08		823 (93.1)	129 (92.1)	
Maternal education							
Low	187	18.9	7.57 ± 5.21	0.003	157 (18.2)	30 (22.9)	0.196
Medium	519	52.3	6.67 ± 5.03		448 (52.0)	71 (54.2)	
High	286	28.8	6.02 ± 4.71		256 (29.7)	30 (22.9)	
Marital status							
Married	878	88.4	6.51 ± 4.88	0.112	770 (89.2)	108 (83.1)	0.041
Other	115	11.6	7.54 ± 5.76		93 (10.8)	22 (16.9)	
Parity							
Primiparous	409	41.4	6.49 ± 4.67	0.633	356 (41.6)	53 (39.8)	0.705
Multiparous	580	58.6	6.84 ± 5.27		500 (58.4)	80 (60.2)	
Physical activity before pregnancy							
No	775	78.9	6.75 ± 5.05	0.253	670 (78.7)	105 (80.2)	0.710
Yes	207	21.1	6.29 ± 4.87		181 (21.3)	26 (19.8)	
Physical activity during pregnancy							
No	912	92.7	6.67 ± 5.01	0.727	792 (92.8)	120 (91.6)	0.610
Yes	72	7.3	6.47 ± 4.99		61 (7.2)	11 (8.4)	
Working during pregnancy							
No	493	49.7	7.25 ± 5.18	<0.001	415 (48.3)	78 (59.5)	0.016
Yes	498	50.3	6.05 ± 4.75		445 (51.7)	53 (40.5)	
Infant sex							
Male	530	51.1	6.82 ± 4.99	0.207	461 (51.5)	69 (48.9)	0.579
Female	507	48.9	6.58 ± 5.14		435 (48.5)	72 (51.1)	
	N	Mean ± s.d.	rho	p-value	Mean ± s.d.	Mean ± s.d.	p-value
Maternal age at birth (years)	1008	29.51 ± 4.95	-0.074	0.019	29.54 ± 4.92	29.33 ± 5.13	0.693
Gestational age (weeks)	1037	38.24 ± 1.51	-0.010	0.745	38.25 ± 1.47	38.18 ± 1.78	0.837
Birth weight (gr)	1035	3194.29 ± 438.69	0.008	0.807	3193.29 ± 439.90	3200.63 ± 432.39	0.454

EPDS, Edinburgh Postnatal Depression Symptoms; rho, Spearman's correlation coefficient.

Statistical significant differences at $p < 0.05$, based on Mann-Whitney U -test for two independent samples, Kruskal-Wallis one-way analysis of variance by ranks, Spearman's rho correlation coefficient and χ^2 analysis for categorical variables. Missing values have been excluded pair-wise.

Bold font indicates statistically significant differences ($p < 0.05$).

of education, who were not married, and those who did not work during pregnancy had significantly higher mean EPDS scores. The comparison between the 1037 participants and the 402 non-participants (final sample $n=1439$ including only women with live singleton births from the initial sample of eligible women) revealed significant differences between the two groups in terms of age, education, origin, and marital status. Mothers who participated in this study were more likely to be older, to have high levels of education, to be married and of Greek origin as compared with non-participants.

In the present study population, the prevalence of women with probable PPD (EPDS score ≥ 13) was 13.6%, indicating that 141 out of 1037 new mothers

presented high levels of PPD at 8 weeks postpartum. The frequencies of pregnancy, perinatal and postpartum complications between depressed and non-depressed women are presented in Table 2. Women with PPD were more likely to report sleep deprivation, to be snorers and to breastfeed their children for a shorter period than women without PPD.

Table 3 presents the crude and adjusted associations between pregnancy, perinatal and postpartum complications and postpartum EPDS score as continuous variable at 8 weeks postpartum. In the crude models, several complications were associated with postnatal EPDS score, i.e. unplanned pregnancy, gestational hypertension and/or preeclampsia, sleep deprivation, hospitalisation during pregnancy, sleep deprivation

Table 2. Descriptive of pregnancy, perinatal and postpartum complications by score on the Edinburgh Postnatal Depression Scale (Rhea Study, Crete, Greece)

Complications	Total N	Edinburgh Postnatal Depression Scale		
		EPDS < 13 N (%)	EPDS \geq 13 N (%)	<i>p</i> -value
Unplanned pregnancy (yes)	942	333 (40.8)	62 (49.2)	0.075
Trying to get pregnant (with effort)	961	213 (25.5)	30 (23.6)	0.643
Pregnancy induced hypertension and/or pre-eclampsia (yes)	885	34 (4.5)	9 (7.4)	0.164
Gestational diabetes (yes)	892	62 (8.1)	14 (11.4)	0.221
Symptoms during pregnancy (i.e. vomit, nausea) (yes)	969	438 (52.2)	74 (56.9)	0.316
Fasting glucose level (≥ 100 mg/dl)	473	9 (2.2)	0 (0.0)	0.211
HDL cholesterol level (<50 mg/dl)	476	17 (4.1)	2 (3.1)	0.703
Triglyceride level (≥ 150 mg/dl)	475	7 (1.7)	1 (1.6)	0.935
Thyroid dysfunction during pregnancy (yes)	989	135 (15.8)	26 (19.5)	0.272
Hospitalisation during pregnancy (yes)	984	62 (7.3)	16 (12.0)	0.060
Smoking status during pregnancy (smokers)	943	251 (30.6)	46 (37.1)	0.150
Alcohol use during pregnancy (yes)	905	164 (28.9)	28 (30.8)	0.712
Sleep deprivation (≤ 5 h)	835	37 (5.1)	18 (15.8)	<0.001
Sleep-disordered breathing symptoms (snorers)	835	116 (16.1)	28 (24.6)	0.026
Excessive daytime sleepiness score (ESS ≥ 10) (yes)	791	26 (3.8)	5 (4.5)	0.732
Type of delivery (caesarean)	1034	444 (49.7)	75 (53.2)	0.116
Preterm birth (GA < 37 weeks) (yes)	1037	91 (10.2)	20 (14.2)	0.150
Low birth weight (<2500 kg) (yes)	1035	45 (5.0)	8 (5.7)	0.749
Intensive Care Unit (yes)	931	115 (14.1)	22 (18.8)	0.182
Breastfeeding (never)	976	126 (14.8)	22 (17.6)	0.416
Breastfeeding difficulties (yes)	797	253 (36.4)	44 (43.1)	0.189
		Mean \pm S.D.	Mean \pm S.D.	<i>p</i> -value
BMI pre-pregnancy	989	24.16 \pm 4.87	24.18 \pm 4.76	0.791
Weight gain (kg)	823	14.07 \pm 5.90	14.86 \pm 7.00	0.348
Metabolic syndrome (number of symptoms)	978	0.21 \pm 0.47	0.23 \pm 0.50	0.692
Breastfeeding duration up to 2 months (months)	976	0.64 \pm 0.48	0.48 \pm 0.50	0.001

EPDS, Edinburgh Postnatal Depression Symptoms; HDL, High-Density Lipoprotein.

Statistical significant differences at $p < 0.05$, based on Mann-Whitney *U*-test for two independent samples for continuous variables and χ^2 test or Fisher exact test for categorical variables.

Bold font indicates statistically significant differences ($p < 0.05$).

Table 3. Associations of PPD with pregnancy, perinatal and postpartum complications in the mother–child cohort ('Rhea' study) in Crete, Greece

Complications	EPDS		EPDS \geq 13	
	Crude β -coef (95% CI)	Adjusted β -coef (95% CI) ^a	Crude OR (95% CI)	Adjusted OR (95% CI) ^a
Before pregnancy				
Unplanned pregnancy (yes v. no)	0.67 (0.02, 1.33)	0.29 (−0.40, 0.98)	1.40 (0.96, 2.05)	1.25 (0.84, 1.86)
Trying to get pregnant (with effort v. immediately)	−0.31 (−1.04, 0.42)	−0.08 (−0.87, 0.70)	0.90 (0.58, 1.40)	0.94 (0.59, 1.51)
BMI pre-pregnancy	0.01 (−0.06, 0.07)	−0.01 (−0.07, 0.06)	1.00 (0.96, 1.04)	0.99 (0.96, 1.03)
During pregnancy				
Gestational hypertension and/or preeclampsia (yes v. no)	2.02 (0.47, 3.57)	1.86 (0.32, 3.41)	1.71 (0.80, 3.65)	1.69 (0.78, 3.68)
Gestational diabetes (yes v. no)	−0.19 (−1.38, 1.00)	−0.00 (−1.21, 1.21)	1.46 (0.79, 2.71)	1.53 (0.82, 2.87)
Metabolic syndrome (number of symptoms)	0.13 (−0.54, 0.80)	0.15 (−0.54, 0.83)	1.10 (0.76, 1.60)	1.12 (0.76, 1.64)
Fasting glucose level (\geq 100 mg/dl)	−1.43 (−4.77, 1.91)	−0.99 (−4.28, 2.28)	— ^b	— ^b
HDL cholesterol level (<50 mg/dl)	−0.89 (−4.36, 2.59)	0.61 (−3.02, 4.24)	0.92 (0.11, 7.57)	1.47 (0.17, 12.82)
Triglyceride level (\geq 150 mg/dl)	−1.26 (−3.54, 1.02)	−1.14 (−3.37, 1.09)	0.75 (0.17, 3.32)	0.79 (0.17, 3.56)
Symptoms (i.e. vomit, nausea) (yes v. no)	0.24 (−0.40, 0.87)	0.20 (−0.45, 0.84)	1.17 (0.81, 1.70)	1.18 (0.81, 1.73)
Thyroid dysfunction (yes v. no)	0.19 (−0.66, 1.05)	0.40 (−0.46, 1.26)	1.30 (0.81, 1.60)	1.36 (0.84, 2.20)
Hospitalisation (yes v. no)	1.17 (0.00, 2.34)	0.82 (−0.35, 1.99)	1.74 (0.97, 3.12)	1.52 (0.83, 2.78)
Smoking status (smokers v. non-smokers)	0.50 (−0.19, 1.18)	0.50 (−0.20, 1.20)	1.34 (0.90, 1.98)	1.33 (0.89, 2.00)
Alcohol use during pregnancy (yes)	0.03 (−0.82, 0.87)	−0.05 (−0.90, 0.80)	1.09 (0.68, 1.77)	1.06 (0.64, 1.73)
Weight gain (kg)	0.02 (−0.03, 0.08)	0.02 (−0.03, 0.08)	1.02 (0.99, 1.05)	1.03 (0.99, 1.06)
Sleep deprivation (\leq 5 h) (yes v. no)	2.73 (1.36, 4.09)	2.58 (1.21, 3.96)	3.47 (1.90, 6.33)	3.57 (1.91, 6.67)
Sleep-disordered breathing (snorers v. non-snorers)	0.00 (−0.91, 0.90)	0.11 (−0.81, 1.04)	1.70 (1.06, 2.72)	1.81 (1.11, 2.93)
Excessive daytime sleepiness (ESS \geq 10) (yes v. no)	1.09 (0.45, 3.15)	1.28 (0.47, 3.47)	1.19 (0.45, 3.15)	1.28 (0.47, 3.47)
At birth				
Type of delivery (caesarean v. vaginal)	0.37 (−0.25, 0.99)	0.53 (−0.11, 1.17)	1.15 (0.87, 2.46)	1.22 (0.83, 1.78)
Preterm birth (GA < 37 weeks) (yes v. no)	0.44 (−0.55, 1.44)	0.05 (−0.97, 1.08)	1.46 (0.87, 2.46)	1.25 (0.70, 2.22)
Low birth weight (<2500 kg) (yes v. no)	0.52 (−0.88, 1.92)	0.36 (−1.12, 1.84)	1.13 (0.52, 2.46)	1.12 (0.48, 2.58)
Intensive Care Unit (yes v. no)	0.81 (−0.08, 1.70)	0.84 (−0.07, 1.75)	1.39 (0.84, 2.31)	1.48 (0.88, 2.47)
After birth				
Ever breastfeeding (never v. ever)	0.53 (−0.34, 1.39)	0.23 (−0.67, 0.71)	1.23 (0.74, 2.02)	0.98 (0.58, 1.66)
Breastfeeding duration (<2 months v. \geq 2 months)	0.83 (0.19, 1.47)	0.57 (−0.10, 1.25)	1.93 (1.32, 2.82)	1.77 (1.19, 2.64)
Breastfeeding difficulties (yes v. no)	0.79 (0.08, 1.49)	0.77 (0.02, 1.53)	1.32 (0.87, 2.02)	1.27 (0.81, 1.99)

β -coef., beta coefficient; 95% CI, 95% confidence interval; OR, odds ratio; HDL, High-Density Lipoprotein; ESS, Epworth Sleepiness Scale.

Bold font indicates statistically significant differences ($p < 0.05$).

^aAll models are adjusted for maternal age, education, origin, marital status, working during pregnancy and parity.

^bThere were no events of high EPDS scores (EPDS \geq 13) in mothers with Fasting glucose level \geq 100 mg/dl.

during pregnancy, breastfeeding duration less than 2 months, and breastfeeding difficulties. Further adjustment for maternal characteristics, such as maternal age, origin, education, marital status, working during pregnancy, and parity, showed that only gestational hypertension and/or preeclampsia (β coefficient 1.86, 95% CI: 0.32, 3.41), sleep deprivation during pregnancy (β coefficient 2.58, 95% CI: 1.21, 3.96) and breastfeeding difficulties (β coefficient 0.77, 95% CI: 0.02,

1.53) were significantly associated with higher EPDS score. The relationships between postpartum EPDS score and gestational hypertension and/or preeclampsia (β coefficient 2.60, 95% CI 1.07, 4.12), as well as sleep deprivation (β coefficient 2.49, 95% CI 1.10, 3.88), proved remarkably robust when sensitivity analysis was conducted following the exclusion of 77 women with antenatal EPDS \geq 13 to explore the possible impact of antenatal depression. Although this

sensitivity analysis attenuated the observed association between breastfeeding difficulties and postpartum EPDS score, the direction of the association did not change (β coefficient 0.66, 95% CI: -0.09, 1.41).

Associations of pregnancy, perinatal and postpartum complications with high levels of postpartum depressive symptoms (EPDS \geq 13) are also presented in Table 3. Multivariable analysis revealed that sleep patterns during pregnancy, such as sleep deprivation (OR = 3.57, 95% CI: 1.91, 6.67) and snoring (OR = 1.81, 95% CI 1.11, 2.93), were associated with almost fourfold and twofold increase in the odds for PPD, respectively. Furthermore, breastfeeding duration less than 2 months was related to a 77% increase in the odds for PPD (OR = 1.77, 95% CI: 1.19, 2.64). These associations were supported by sensitivity analyses as well, after exclusion of 77 women with antenatal EPDS \geq 13 (sleep deprivation OR = 3.79, 95% CI 1.89, 7.62, snoring OR = 1.98, 95% CI 1.14, 3.42, and breastfeeding duration less than 2 months OR = 1.59, 95% CI 1.01, 2.49).

Finally, we did not find any significant interactions between pregnancy, perinatal and postpartum complications and parity (p for interaction >0.10) (data not shown in tables).

Discussion

In the present study, we investigated the association of various pregnancy, perinatal and postpartum risk factors with maternal mood in the early postpartum period in a population-based mother-child cohort in Crete, Greece. To our knowledge, this is the first study evaluating the effect of specific pregnancy and delivery complications along with sleep patterns in late pregnancy and complications women are experiencing at the early postpartum period (i.e. breastfeeding) on PPD. The results suggest that gestational hypertension and/or preeclampsia and negative early breastfeeding experiences were adversely associated with PPD symptoms at 8 weeks postpartum. Furthermore, sleep deprivation and snoring during pregnancy, as well as breastfeeding duration less than 2 months appeared to increase the odds for the development of PPD. These findings apply equally to primiparous and multiparous women. Our results also indicate that these effects on PPD are not likely to be due to confounding by high levels of antenatal depressive symptoms, even though replication of such analyses in other population samples would be desirable.

According to our findings, higher PPD symptoms were reported by mothers who experienced gestational hypertension and/or preeclampsia. This finding is comparable with a previous study which found an

association between preeclampsia and PPD (Blom *et al.* 2010) and suggested that this association might be explained by hormonal (i.e. serotonin levels) and physical (i.e. morbidity) changes (Blom *et al.* 2010). More specifically, increased blood serotonin levels in pregnant women with preeclampsia (Bolte *et al.* 2001) could be linked with decreased serotonin levels in the brain, thus leading to the development of PPD symptoms. In addition, women who experience complications during pregnancy, including hypertension, are more likely to experience physical morbidity in the postpartum period, which can lead to higher rates of PPD.

Our findings, also, indicated that women with negative early breastfeeding experiences were more likely to have higher depressive symptoms at 8 weeks postpartum. This finding is consistent with the study of Watkins *et al.* (2011), who found that women who disliked breastfeeding in the first week and those who reported severe breastfeeding pain in the first day, the first and the second week, were more likely to experience PPD at 2 months postpartum. These results are also in consistence with the study of Dennis & McQueen (2007) suggesting that mothers who perceived their breastfeeding as terribly progressing at 1 week postpartum were at higher risk to develop PPD at 4 and 8 weeks. Psychological mechanisms, such as women's particular expectations, about breastfeeding might underlie the association between breastfeeding difficulties and PPD. Negative early breastfeeding experiences lead to worries and feelings of disappointment and failure, which may affect a woman's ability to adapt in this situation and thus experience depressive symptoms. When depressive symptomatology at 1 week was controlled for in this latter study (Dennis & McQueen, 2007), the negative impact of perceived breastfeeding difficulties was diminished, thus giving rise to the hypothesis that depressive mood negatively influences cognitions and leading new mothers to have more negative attitudes about breastfeeding and report decreased satisfaction with their breastfeeding experiences. The underlying neuroendocrine mechanism linking breastfeeding difficulties with maternal mental health at the postpartum period is largely unknown. However, neurochemical imbalances in key neurotransmitters, such as serotonin, have been thought to contribute to both feelings of pain and depression (Bair *et al.* 2003).

In our study, shorter duration of breastfeeding up to 2 months postpartum was found to be a risk factor for the development of PPD. Therefore, women who discontinued breastfeeding their infants at 2 months postpartum were at higher risk for developing PPD. This finding is comparable with previous research, which demonstrated an association between longer breastfeeding duration and a lower prevalence of PPD

(Mezzacappa & Katlin, 2002; Ystrom, 2012; Figueiredo *et al.* 2014). This relationship has been suggested to be influenced by biological factors, such as differences in hormone levels between breastfeeding and non-breastfeeding mothers (Groer, 2005), as well as by the mother's intention to breastfeed her baby (Borra *et al.* 2015). Breastfeeding is suggested to attenuate neuroendocrine responses to stress and may act to enhance maternal mood. In addition, breastfeeding may enhance the mother–infant interaction, which could lead to improved maternal mental health.

Finally, our study provides some new evidence that specific sleep patterns during late pregnancy, such as sleep deprivation and snoring, were associated with PPD. Sleep disturbances are prevalent among pregnant women, due to the changes in physiology that occur with pregnancy (Sahota *et al.* 2003). Our findings concur with previous epidemiological studies showing associations between sleep deprivation (Lee *et al.* 2000; Wolfson *et al.* 2003; Goyal *et al.* 2007; Chang *et al.* 2010) and snoring (Bat-Pitault *et al.* 2015) during pregnancy with PPD symptomatology. It has been suggested that sleep disturbances may increase the risk for PPD with inflammation as the underlying mechanism. Specifically, prior research has indicated that sleep deprivation is associated with higher levels of pro-inflammatory serum cytokines (Okun *et al.* 2007), which have been observed in women with PPD symptoms (Irwin *et al.* 2006). However, the proposed pathophysiological explanations (i.e. the neuroendocrine one or the inflammation one) are now only hypotheses.

Finally, some other complications, such as unplanned pregnancy and hospitalisation during pregnancy were positively associated with EPDS score in unadjusted models, but these associations were explained by socio-demographic characteristics of the mother, such as maternal age, origin, education, marital status, working during pregnancy and parity. The following complications were non-significantly associated with PPD symptoms in both unadjusted and adjusted models: trying to get pregnant, metabolic syndrome (defined as number of symptoms) and its components specifically (i.e. BMI pre-pregnancy, triglyceride level, HDL-cholesterol level and fasting glucose level), thyroid dysfunction, gestational diabetes, symptoms during pregnancy (i.e. vomit, nausea, etc.), smoking and alcohol use during pregnancy, weight gain, excessive daytime sleepiness during pregnancy, type of delivery, preterm birth, low birth weight and newborn's admission at ICU. These findings are inconsistent with previous research, which has shown that unintended pregnancy (Mercier *et al.* 2013; Gaillard *et al.* 2014), hospitalisation during pregnancy and emergency caesarean section (Blom *et al.* 2010), preterm birth or low birth weight (Blom *et al.* 2010;

Gulamani *et al.* 2013, Sundaram *et al.* 2014) could be contributing factors for the development of PPD.

Strengths and limitations

The strengths of the present study include the population-based, prospective follow-up design, the large number of women participating, and the availability of detailed information on various pregnancy, perinatal and postpartum complications. The study population included women followed-up since early pregnancy, providing us the opportunity to account prospectively for the effect of exposures during pregnancy. Furthermore, the exclusion of women who gave birth to twins and women with a previous history of psychiatric disorder, as well as adjustment for several socio-demographic variables, reduced the likelihood of confounding. Finally, to evaluate the possibility of confounding by antenatal psychosocial stressors, a sensitivity analysis excluding women with high levels of antenatal depressive symptoms (antenatal EPDS ≥ 13) was carried out and the results remained essentially the same as those from the original analysis, although the possibility of residual confounding cannot be completely eliminated.

There are also some limitations to this study. Despite our large study population, the prevalence of some pregnancy, perinatal and postpartum complications was rather low, and this may have limited the power of our study. Another possible limitation would be the differences found between participants and non-participants. Although we consider that these differences observed at baseline are not likely to have affected appreciably the present results, this limitation should be taken into account when considering study findings. Furthermore, we assessed PPD symptoms with the self-reported EPDS scale rather than definite cases of depression based on clinician-administered structured diagnostic interview. However, this is an epidemiological study assessing the prevalence of PPD and EPDS is an established and widely used screening tool with high specificity and sensitivity. Moreover, although we incorporated extensive information on potential social and environmental factors that are associated with the risk of PPD, we acknowledge that there may be other factors linked with both perinatal complications and PPD that could explain this association. Another possible limitation would be that we did not use a questionnaire to evaluate stressful life events, history of abuse, family functioning and perceived social support, which have been shown to contribute to the development of PPD. Moreover, our results should be interpreted with caution since maternal depression was assessed only once postpartum (at 8 weeks); there is no information about the development

of PPD during the 2 months' period after delivery. More specifically, the study missed cases of depression that had been successfully treated or else the cases of depression developed soon after delivery that after 2 months were at their natural point of attenuation, and this might have affected our findings and possibly underestimated the associations (i.e. based on the positive associations the study has observed). A longitudinal analysis of perinatal risk factors across different stages of the postpartum period is of great value for examining how the effect of various risk factors changes across the postpartum period. Finally, it is difficult to fully decipher the direction of the association between PPD, breastfeeding difficulties and sleep deprivation, since the two latter variables could be risk factors for PPD or else symptoms of the depression itself; thus, we cannot rule out the possibility of a bi-directional interpretation of these two associations in the light of the possible interventions to prevent or treat PPD.

Conclusion

In a large sample of Greek women, we found that specific pregnancy, perinatal and postpartum complications, such as gestational hypertension and/or preeclampsia, sleep deprivation and snoring during pregnancy, negative early breastfeeding experiences and discontinuation of breastfeeding were associated with PPD symptomatology at 8 weeks postpartum. Our findings have considerable implications for developing effective prevention and early psychoeducational intervention strategies for women at risk of developing PPD. Given that women who experience an episode of PPD are more likely to have experienced depression during pregnancy (Koutra *et al.* 2014), and are at increased risk of future episodes (Giallo *et al.* 2014; Woolhouse *et al.* 2015), PPD acts as a marker of risk of depression at other times which in turn might have a significant adverse impact not just on the new mother's sense of well-being, but also on her family as a whole. The causal pathway between these pregnancy, perinatal and postpartum complications and PPD is far from being elucidated. Future longitudinal studies are needed to confirm the findings of our study and better understand the complex underlying processes.

Acknowledgements

We are grateful to all those who participated in the study especially the mothers and their infants, and the whole Rhea team for their contribution and understanding.

Financial Support

The Rhea project was financially supported by European projects (EU FP6-003-Food-3-NewGeneris – Contract no.

16320, EU FP6 STREP Hiwate – Contract no. 36224, EU FP7 ENV.2007.1.2.2.2. – Project no. 211250 Escape, EU FP7-2008-ENV-1.2.1.4 Envirogenomarkers – Contract no. 226756, EU FP7-HEALTH-2009-single stage CHICOS – Contract no. 241604, EU FP7 ENV.2008.1.2.1.6. – Proposal no. 226285 ENRIECO, EU-FP7 – Proposal no. 264357 MeDALL, EU-FP7-HEALTH-2012 – Proposal no. 308333 HELIX) and the Greek Ministry of Health (Program of Prevention of obesity and neurodevelopmental disorders in preschool children, in Heraklion district, Crete, Greece: 2011–2014; 'Rhea Plus': Prevention Program of Environmental Risk Factors for Reproductive Health, and Child Health: 2012–2015).

Conflict of Interest

None.

Ethical Standard

The authors assert that all procedures contributing to this work comply with the ethical standards of the relevant national and institutional committees on human experimentation and with the Helsinki Declaration of 1975, as revised in 2008.

Availability of Data and Materials

Ethical restrictions prevent public sharing of data from the Rhea pregnancy cohort study, as imposed by the Research Ethics Committee of the Rhea Cohort Study. Data can be made available to all interested researchers upon request by contacting the Research Committee at rhea@med.uoc.gr or the Principal Investigators of the study (Professor Manolis Kogevas, kogevas@creal.cat; Dr Leda Chatzi, lchatzi@med.uoc.gr).

References

- Bair MJ, Robinson RL, Katon W, Kroenke K (2003). Depression and pain comorbidity: a literature review. *Archives of Internal Medicine* 163, 2433–2445.
- Bat-Pitault F, Deruelle C, Flori S, Porcher-Guinet V, Stagnara C, Guyon A, Plancoulaine S, Adrien J, Da Fonseca D, Patural H, Franco F (2015). Sleep pattern during pregnancy and maternal depression: study of Aube cohort. *Journal of Sleep Disorders and Management* 1–7.
- Beck CT (1996). A meta-analysis of predictors of postpartum depression. *Nursing Research* 45, 297–303.
- Beck CT (2001). Predictors of postpartum depression: an update. *Nursing Research* 50, 275–285.
- Bennett HA, Einarson A, Taddio A, Koren G, Einarson TR (2004). Prevalence of depression during pregnancy: systematic review. *Obstetrics & Gynecology* 103, 698–709.

- Blom EA, Jansen PW, Verhulst FC, Hofman A, Raat H, Jaddoe VW, Coolman M, Steegers EA, Tiemeier H** (2010). Perinatal complications increase the risk of postpartum depression. The Generation R Study. *BJOG: an International Journal of Obstetrics & Gynaecology* **117**, 1390–1398.
- Bolte AC, van Geijn HP, Dekker GA** (2001). Pathophysiology of preeclampsia and the role of serotonin. *European Journal of Obstetrics and Gynecol and Reproductive Biology* **95**, 12–21.
- Borra C, Iacovou M, Sevilla A** (2015). New evidence on breastfeeding and postpartum depression: the importance of understanding women's intentions. *Maternal & Child Health Journal* **19**, 897–907.
- Carpenter MW, Coustan DR** (1982). Criteria for screening tests for gestational diabetes. *American Journal of Obstetrics & Gynecology* **144**, 768–773.
- Chang JJ, Pien GW, Duntley SP, Macones GA** (2010). Sleep deprivation during pregnancy and maternal and fetal outcomes: is there a relationship? *Sleep Medicine Reviews* **14**, 107–114.
- Chatzi L, Plana E, Daraki V, Karakosta P, Alekakis D, Tsatsanis C, Kafatos A, Koutis A, Kogevas M** (2009a). Metabolic syndrome in early pregnancy and risk of preterm birth. *American Journal of Epidemiology* **170**, 829–836.
- Chatzi L, Plana E, Pappas A, Alekakis D, Karakosta P, Daraki V, Vassilaki M, Tsatsanis C, Kafatos A, Koutis A, Kogevas M** (2009b). The metabolic syndrome in early pregnancy and risk of gestational diabetes mellitus. *Diabetes & Metabolism* **35**, 490–494.
- Cornish AM, McMahon CA, Ungerer JA, Barnett B, Kowalenko N, Tennant C** (2005). Postnatal depression and infant cognitive and motor development in the second postnatal year: the impact of depression chronicity and infant gender. *Infant Behavior and Development* **28**, 407–417.
- Cox JL, Holden JM, Sagovsky R** (1987). Detection of postnatal depression. Development of the 10-item Edinburgh Postnatal Depression Scale. *British Journal of Psychiatry* **150**, 782–786.
- Cox JL, Murray D, Chapman G** (1993). A controlled study of the onset, duration and prevalence of postnatal depression. *British Journal of Psychiatry* **163**, 27–31.
- Davis L, Edwards H, Mohay H, Wollin J** (2003). The impact of very premature birth on the psychological health of mothers. *Early Human Development* **73**, 61–70.
- Dennis CL, McQueen K** (2007). Does maternal postpartum depressive symptomatology influence infant feeding outcomes? *Acta Paediatrica* **96**, 590–594.
- Dias CC, Figueiredo B** (2015). Breastfeeding and depression: a systematic review of the literature. *Journal of Affective Disorders* **171**, 142–154.
- Evans J, Heron J, Francomb H, Oke S, Golding J** (2001). Cohort study of depressed mood during pregnancy and after childbirth. *British Medical Journal* **323**, 257–260.
- Figueiredo B, Canario C, Field T** (2014). Breastfeeding is negatively affected by prenatal depression and reduces postpartum depression. *Psychological Medicine* **44**, 927–936.
- Gaillard A, Le Strat Y, Mandelbrot L, Keita H, Dubertret C** (2014). Predictors of postpartum depression: prospective study of 264 women followed during pregnancy and postpartum. *Psychiatry Research* **215**, 341–346.
- Gavin NI, Gaynes BN, Lohr KN, Meltzer-Brody S, Gartlehner G, Swinson T** (2005). Perinatal depression: a systematic review of prevalence and incidence. *Obstetrics & Gynecology* **106**, 1071–1083.
- Giallo R, Cooklin A, Nicholson JM** (2014). Risk factors associated with trajectories of mothers' depressive symptoms across the early parenting period: an Australian population-based longitudinal study. *Archives of Women's Mental Health* **17**, 115–125.
- Gonidakis F, Rabavilas AD, Varsou E, Kreatsas G, Christodoulou GN** (2008). A 6-month study of postpartum depression and related factors in Athens Greece. *Comprehensive Psychiatry* **49**, 275–282.
- Goyal D, Gay CL, Lee KA** (2007). Patterns of sleep disruption and depressive symptoms in new mothers. *Journal of Perinatal & Neonatal Nursing* **21**, 123–129.
- Grace SL, Eviandar A, Stewart DE** (2003). The effect of postpartum depression on child cognitive development and behavior: a review and critical analysis of the literature. *Archives of Women's Mental Health* **6**, 263–274.
- Groer MW** (2005). Differences between exclusive breastfeeders, formula-feeders, and controls: a study of stress, mood, and endocrine variables. *Biological Research for Nursing* **7**, 106–117.
- Grundy SM, Cleeman JI, Daniels SR, Donato KA, Eckel RH, Franklin BA, Gordon DJ, Krauss RM, Savage PJ, Smith SC Jr., Spertus JA, Costa F, American Heart A, National Heart L, Blood I** (2005). Diagnosis and management of the metabolic syndrome: an American Heart Association/National Heart, Lung, and Blood Institute Scientific Statement. *Circulation* **112**, 2735–2752.
- Gulamani SS, Premji SS, Kanhi Z, Azam SI** (2013). Preterm birth a risk factor for postpartum depression in Pakistani women. *Open Journal of Depression* **2**, 72–81.
- Halbreich U, Karkun S** (2006). Cross-cultural and social diversity of prevalence of postpartum depression and depressive symptoms. *Journal of Affective Disorders* **91**, 97–111.
- Howard LM, Molyneaux E, Dennis CL, Rochat T, Stein A, Milgrom J** (2014). Non-psychotic mental disorders in the perinatal period. *Lancet* **384**, 1775–1788.
- Irwin MR, Wang M, Campomayor CO, Collado-Hidalgo A, Cole S** (2006). Sleep deprivation and activation of morning levels of cellular and genomic markers of inflammation. *Archives of Internal Medicine* **166**, 1756–1762.
- Johns MW** (1991). A new method for measuring daytime sleepiness: the Epworth Sleepiness Scale. *Sleep* **14**, 540–545.
- Johnstone SJ, Boyce PM, Hickey AR, Morris-Yatees AD, Harris MG** (2001). Obstetric risk factors for postnatal depression in urban and rural community samples. *Australian & New Zealand Journal of Psychiatry* **35**, 69–74.
- Josefsson A, Berg G, Nordin C, Sydsjo G** (2001). Prevalence of depressive symptoms in late pregnancy and postpartum. *Acta Obstetrica et Gynecologica Scandinavica* **80**, 251–255.
- Koutra K, Chatzi L, Bagkeris M, Vassilaki M, Bitsios P, Kogevas M** (2013). Antenatal and postnatal maternal mental health as determinants of infant neurodevelopment at 18 months of age in a mother-child cohort (Rhea Study)

- in Crete, Greece. *Social Psychiatry and Psychiatric Epidemiology* **48**, 1335–1345.
- Koutra K, Vassilaki M, Georgiou V, Koutis A, Bitsios P, Chatzi L, Kogevas M** (2014). Antenatal maternal mental health as determinant of postpartum depression in a population based mother-child cohort (Rhea Study) in Crete, Greece. *Social Psychiatry and Psychiatric Epidemiology* **49**, 711–721.
- Lee KA, McEnany G, Zaffke ME** (2000). REM sleep and mood state in childbearing women: sleepy or weepy? *Sleep* **23**, 877–885.
- Leonardou AA, Zervas YM, Papageorgiou CC, Marks MN, Tsartsara EC, Antsaklis A, Christodoulou GN, Soldatos CR** (2009). Validation of the Edinburgh Postnatal Depression Scale and prevalence of postnatal depression at two months postpartum in a sample of Greek mothers. *Journal of Reproductive and Infant Psychology* **27**, 28–39.
- Matthey S, Henshaw C, Elliot S, Barnett B** (2006). Variability in use of cut off scores and formats on the Edinburgh Postnatal Depression Scale – implications for clinical and research practice. *Archives of Women's Mental Health* **9**, 309–315.
- Mercier RJ, Garrett J, Thorp J, Siega-Riz AM** (2013). Pregnancy intention and postpartum depression: secondary data analysis from a prospective cohort. *BJOG: an International Journal of Obstetrics and Gynaecology* **120**, 1116–1122.
- Mezzacappa ES, Katlin ES** (2002). Breast-feeding is associated with reduced perceived stress and negative mood in mothers. *Health Psychology* **21**, 187–193.
- Micheli K, Komninos I, Bagkeris E, Roumeliotaki T, Koutis A, Kogevas M, Chatzi L** (2011). Sleep patterns in late pregnancy and risk of preterm birth and fetal growth restriction. *Epidemiology* **22**, 738–744.
- Nielsen Forman D, Videbech T, Hedegaard M, Dalby Salvig J, Secher N** (2000). Postpartum depression: identification of women at risk. *British Journal of Obstetrics and Gynaecology* **107**, 1210–1217.
- Nonacs R, Cohen LS** (2002). Depression during pregnancy: diagnosis and treatment options. *Journal of Clinical Psychiatry* **63** (Suppl. 7), 24–30.
- Norhayati MN, Hazlina NH, Asrenee AR, Emilin WM** (2015). Magnitude and risk factors for postpartum symptoms: a literature review. *Journal of Affective Disorders* **175**, 34–52.
- O'Hara M, Swain A** (1996). Rates and risk of postpartum depression: a meta-analysis. *International Review of Psychiatry* **8**, 37–54.
- O'Hara MW** (1997). The nature of postpartum depressive disorders. In *Postpartum Depression and Child Development* (ed. L. Murray and P. Cooper), pp. 3–31. Guilford: New York.
- Okun ML, Hall M, Coussons-Read ME** (2007). Sleep disturbances increase interleukin-6 production during pregnancy: implications for pregnancy complications. *Reproductive Sciences* **14**, 560–567.
- Robertson E, Grace S, Wallington T, Steward DE** (2004). Antenatal risk factors for postpartum depression: a synthesis of recent literature. *General Hospital Psychiatry* **26**, 289–295.
- Rubertsson C, Waldenstrom U, Wickberg B** (2003). Depressive mood in early pregnancy: prevalence and women at risk in a national Swedish sample. *Journal of Reproductive and Infant Psychology* **21**, 113–123.
- Sabanayagam C, Shankar A** (2010). Sleep duration and cardiovascular disease: results from the National Health Interview Survey. *Sleep* **33**, 1037–1042.
- Sahota PK, Jain SS, Dhand R** (2003). Sleep disorders in pregnancy. *Current Opinion in Pulmonary Medicine* **9**, 477–483.
- Sundaram S, Harman JS, Cook RL** (2014). Maternal morbidities and postpartum depression: an analysis using the 2007 and 2008 Pregnancy Risk Assessment Monitoring System. *Women's Health Issues* **24**, e381–e388.
- Tebeka S, Le Strat Y, Dubertret C** (2016). Developmental trajectories of pregnant and postpartum depression in an epidemiologic survey. *Journal of Affective Disorders* **203**, 62–68.
- Tomfohr LM, Buliga E, Letourneau NL, Campbell TS, Giesbrecht GF** (2015). Trajectories of sleep quality and associations with mood during the perinatal period. *Sleep* **38**, 1237–1245.
- Vivilaki VG, Dafermos V, Kogevas M, Bitsios P, Lionis C** (2009). The Edinburgh Postnatal Depression Scale: translation and validation for a Greek sample. *BMC Public Health* **9**, 329.
- Warner R, Appleby L, Whitton A, Faragher B** (1996). Demographic and obstetric risk factors for postnatal psychiatric morbidity. *British Journal of Psychiatry* **168**, 607–611.
- Watkins S, Meltzer-Brody S, Zolnoun D, Stuebe A** (2011). Early breastfeeding experiences and postpartum depression. *Obstetrics & Gynecology* **118**, 214–221.
- Wolfson AR, Crowley SJ, Anwer U, Bassett JL** (2003). Changes in sleep patterns and depressive symptoms in first-time mothers: last trimester to 1-year postpartum. *Behavioral Sleep Medicine* **1**, 54–67.
- Woolhouse H, Gartland D, Mensah F, Brown SJ** (2015). Maternal depression from early pregnancy to 4 years postpartum in a prospective pregnancy cohort study: implications for primary health care. *BJOG: an International Journal of Obstetrics & Gynaecology* **122**, 312–321.
- Ystrom E** (2012). Breastfeeding cessation and symptoms of anxiety and depression: a longitudinal cohort study. *BMC Pregnancy Childbirth* **12**, 36.