

Suicidal ideation and subsequent completed suicide in both psychiatric and non-psychiatric populations: a meta-analysis

A. A. M. Hubers^{1*}, S. Moaddine¹, S. H. M. Peersmann¹, T. Stijnen², E. van Duijn^{1,3},
R. C. van der Mast^{1,4}, O. M. Dekkers^{5,6,7} and E. J. Giltay¹

¹ Department of Psychiatry, Leiden University Medical Center, Leiden, The Netherlands

² Department of Medical Statistics & Bioinformatics, Leiden University Medical Center, Leiden, The Netherlands

³ Center for Mental Health Care Delfland, Delft, The Netherlands

⁴ Faculty of Medicine, Collaborative Antwerp Psychiatric Research Institute (CAPRI), University of Antwerp, Antwerp, Belgium

⁵ Department of Epidemiology, Leiden University Medical Center, Leiden, The Netherlands

⁶ Department of Endocrinology, Leiden University Medical Center, Leiden, The Netherlands

⁷ Department of Clinical Epidemiology, Aarhus University Hospital, Aarhus, Denmark

Aims. Several authors claimed that expression of suicidal ideation is one of the most important predictors of completed suicide. However, the strength of the association between suicidal ideation and subsequent completed suicide has not been firmly established in different populations. Furthermore, the absolute suicide risk after expression of suicidal ideation is unknown. In this meta-analysis, we examined whether the expression of suicidal ideation predicted subsequent completed suicide in various populations, including both psychiatric and non-psychiatric populations.

Methods. A meta-analysis of cohort and case-control studies that assessed suicidal ideation as determinant for completed suicide in adults. Two independent reviewers screened 5726 articles for eligibility and extracted data of the 81 included studies. Pooled risk ratios were estimated in a random effects model stratified for different populations. Meta-regression analysis was used to determine suicide risk during the first year of follow-up.

Results. The risk for completed suicide was clearly higher in people who had expressed suicidal ideation compared with people who had not, with substantial variation between the different populations: risk ratio ranging from 2.35 (95% confidence interval (CI) 1.43–3.87) in affective disorder populations to 8.00 (95% CI 5.46–11.7) in non-psychiatric populations. In contrast, the suicide risk after expression of suicidal ideation in the first year of follow-up was higher in psychiatric patients (risk 1.40%, 95% CI 0.74–2.64) than in non-psychiatric participants (risk 0.23%, 95% CI 0.10–0.54). Past suicide attempt-adjusted risk ratios were not pooled due to large underreporting.

Conclusions. Assessment of suicidal ideation is of priority in psychiatric patients. Expression of suicidal ideation in psychiatric patients should prompt secondary prevention strategies to reduce their substantial increased risk of suicide.

Received 6 October 2016; Accepted 21 November 2016; First published online 19 December 2016

Key words: Epidemiology, inpatient psychiatry, outpatient psychiatry, suicide, systematic reviews.

Introduction

Almost 90% of those who died by suicide contacted a health care professional in the 3 months prior to their death (De Leo *et al.* 2013a). Of those who had contact with a health care professional in the four weeks before their death, 22% expressed suicidal intent (Isometsa *et al.* 1995). Several authors claim that expression of suicidal ideation is one of the most important predictors for completed suicide (van Heeringen &

Hengeveld, 2009; Posner *et al.* 2011; Batterham *et al.* 2013), but this association has not been firmly established (Large & Nielssen, 2012) with inconsistent previous reports. Most importantly, the risk of dying by suicide after expression of suicidal ideation is currently unknown.

Some of the previous studies reported a strong association between suicidal ideation and completed suicide (Taiminen *et al.* 2001; McGirr *et al.* 2007; De Leo *et al.* 2013b), whereas others found no association (Bradvik & Berglund, 1993; Bertelsen *et al.* 2007; Jollant *et al.* 2014). Previous meta-analyses (Large *et al.*, 2011a, b; Hawton *et al.*, 2005a, b; 2013; Chapman *et al.*, 2015) that assessed this association also reported inconsistent results, with odds ratios

* Address for correspondence: A. A. M. Hubers, Department of Psychiatry, Leiden University Medical Center, Leiden, The Netherlands.

(Email: a.a.m.hubers@lumc.nl)

ranging from 1.5 in bipolar disorder (Hawton *et al.* 2005b) to 29.8 in schizophrenia (Hawton *et al.* 2005a). These meta-analyses only investigated the effect of suicidal ideation within specific populations, without comparing effects between different populations or exploring other sources of clinical heterogeneity. Furthermore, the effects in the general population and several other psychiatric populations remain unknown (Runeson, 2002; Conwell, 2009).

Suicidal ideation is quite common, with a lifetime prevalence of 10% in the general population (Nock *et al.* 2008). Clinicians can only use suicidal ideation in their clinical risk assessment when they know the absolute risks of suicide for those who do and those who do not express suicidal ideation (Grobbee & Hoes, 2009). However, such summary statistics have not been provided by the aforementioned meta-analyses (Large *et al.*, 2011a, b; Hawton *et al.*, 2005a, b, 2013; Chapman *et al.*, 2015). Furthermore, suicidal ideation is strongly correlated with other predictors of completed suicide like previous suicide attempts (Harris & Barraclough, 1997; ten Have *et al.* 2009) and it is not known whether it independently predicts completed suicide (Large & Nielssen, 2012).

Therefore, we conducted a systematic review and meta-analysis to assess whether the expression of suicidal ideation predicted subsequent completed suicide in various populations, including non-psychiatric and psychiatric populations. Secondly, we aimed to estimate the absolute risks of suicide after expression of suicidal ideation in these populations and to investigate whether the expression of suicidal ideation predicted subsequent completed suicide independent of the presence of past suicide attempts.

Method

Search strategy

Ten electronic databases (PubMed, Embase, Web of science, PsycINFO, PsycARTICLES, Psychology and behavioural sciences collection, Cochrane, CINAHL, Academic search premier and ScienceDirect) were searched until February 5, 2016 without language restrictions. A medical librarian was involved in formulating the search string (Supplement S1).

Eligibility criteria

Only journal articles fulfilling the following inclusion criteria were eligible for inclusion: (1) assessment of presence or absence of suicidal ideation as a distinct determinant (i.e. not combined with suicidal behaviour). Suicidal ideation was considered present when any form of ideation, ranging from death wish to suicide plans or

threats, was expressed; (2) assessment of completed suicide (which could include open verdicts) as a distinct outcome measure; (3) comparison of suicidal ideation *v.* no suicidal ideation with respect to risk of subsequent completed suicide (4) cohort or case-control study design; and (5) mean age of the study population ≥ 18 years.

Next, the following exclusion criteria were applied: (1) presence of suicidal ideation was assessed after a suicide attempt; (2) comparison of suicidal ideation *v.* suicide attempt as determinant in a cohort study; or (3) comparison of those who died by suicide and those who survived an attempt as outcome in a case-control study.

For the assessment of the absolute risks of completed suicide, we included only cohort studies and nested case-control studies (with the size of the source population specified and random selection of controls from the source population) in which the number of suicides in the suicidal ideation group and the exposed person time could be extracted or estimated.

To determine whether suicidal ideation predicted subsequent completed suicide independent of the presence of past suicide attempts, articles that assessed the effect of suicidal ideation on subsequent completed suicide adjusted for previous suicide attempts were selected.

Study selection

All retrieved articles in the original search were screened independently by two of the three reviewers (A.A.M.H., S.M. and S.H.M.P.), first on title, then abstract and subsequently full-text evaluation to consider final eligibility. Disagreements with regard to final eligibility were discussed to reach consensus or, if necessary, another independent reviewer (E.J.G.) got involved. In addition, the reference lists of eligible articles and relevant review articles identified by the search strategy were examined by one of the reviewers to search for eligible studies. When multiple publications used (partially) overlapping study populations only the largest study or, when similar, the most recent study was included.

Data extraction

For each eligible article, two of the three reviewers (A.A.M.H. and S.M. or S.H.M.P.) independently extracted data using a standardised form. Disagreements were discussed or another independent reviewer (E.J.G.) got involved if needed. When information necessary to compute the effect size for the primary aim was missing, a request for the missing numbers was emailed to the corresponding author. In case of no response, the study

was not included. Authors of eleven studies were emailed, two of them provided additional data and could be included in the meta-analysis (Appleby *et al.* 1999b; Dutta *et al.* 2011).

Risk of bias assessment

Two independent reviewers (A.A.M.H. and S.M. or S.H. M.P.) assessed four risk of bias aspects (Supplement S2), judged on the basis of adapted items from the Newcastle–Ottawa scale (Wells *et al.* n.d.) and Altman (2001).

Statistical analyses

A study protocol was written *a priori* (Supplement S1), but was not published or registered. The primary outcome of this meta-analysis was the pooled unadjusted risk ratio for the association between suicidal ideation and completed suicide in a random effects model according to the method of DerSimonian and Laird (Borenstein *et al.* 2009a). Due to the varying absolute risks of suicide among different populations (Nordentoft *et al.* 2011), the pooling of risk ratios was stratified for the following populations: affective disorders (including both in- and outpatients), (former) psychiatric inpatients (mixed diagnoses), schizophrenic patients (including both in- and outpatients), other mixed psychiatric populations (including people with substance use disorders, patients with borderline personality disorder and mixed diagnosis psychiatric outpatients [sometimes combined with inpatients]), general population and a residual category of non-psychiatric study populations that could not be grouped in one of the other categories. Studies in each subgroup were combined using a random effects model with separate estimates of tau-squared (Borenstein *et al.* 2009b). To pool results from different studies, odds ratios, risk ratios and incidence rate ratios were considered to approach the same value, which is reasonable given the low risk of completed suicide. When only χ^2 values or *p*-values were given in combination with a direction of the effect, these were used to estimate risk ratios. If the given *p*-value was <0.05, we assumed a *p*-value of 0.049. There were no eligible studies only reporting a *p*-value >0.05. When articles reported on multiple suicidal ideation determinants (e.g. thoughts and plans separately) we computed a combined effect across these different determinants (Borenstein *et al.* 2009c).

With regard to the risk of suicide after expression of suicidal ideation we were most interested in the suicide risk during the first year of follow-up. Since studies had varying follow-up times, and we did not expect a constant suicide rate over time, we conducted

maximum-likelihood meta-regression analyses with mean study follow-up time as determinant and log_e-transformed rates of completed suicide as outcome. Given the limited number of studies per population subgroup, analyses were stratified for psychiatric and non-psychiatric populations only. When one article assessed suicidal ideation at multiple time points, we only included the determinant that assessed suicidal ideation closest to baseline.

In order to assess whether suicidal ideation predicted completed suicide independently of previous attempts the past suicide attempt-adjusted risk ratios were extracted and used for estimation.

To check the robustness of the results, we restricted the analyses to: (1) studies with a low risk of bias (Supplement S2), (2) cohort studies (for the primary research aim only) and (3) studies that assessed suicidal ideation at baseline or in the preceding month (for the secondary research aim only). Only this last analysis was not pre-specified in the study protocol, but was conducted because of the large variation in suicidal ideation assessment periods, which could have influenced suicide risk. In addition, we stratified analyses of the primary outcome for studies with short (≤ 1 -year) *v.* long (>1-year) timeframe between expression of suicidal ideation and completed suicide and suicidal ideation assessment method. Mean effects within subgroups were compared by testing for total between groups heterogeneity. Maximum likelihood meta-regression was applied to examine the effects of mean age and gender in the study populations.

Heterogeneity was assessed using the I^2 statistic, small study bias by inspecting the funnel plot and Egger's test for funnel plot asymmetry. A *p*-value <0.05 was considered statistically significant. *Post-hoc*, we changed the significance level for Egger's test for funnel plot asymmetry in the different subgroups to 0.10, given the limited number of studies that were included in these analyses and the relatively low power of the test. Statistical analyses were performed using Comprehensive Meta-analysis software version 2.0.064 (www.meta-analysis.com).

Results

Literature search and study characteristics

The literature search identified 5726 unique articles: 80 articles were finally included, which presented results of 81 studies (Fig. 1). The majority of the articles was published in English (*n* = 75), others in German (*n* = 2), Spanish (*n* = 1), French (*n* = 1) or South African (*n* = 1). Together these studies included 4 601 378 participants (median 216; range 14–4 045 993), with a total of 7729




Fig. 1. Flowchart of the study selection process. Si indicates suicidal ideation; cs, completed suicide; sa, suicide attempts. Mixed psychiatric populations: $n = 9$; including three studies on people with substance use disorders, two on patients with borderline personality disorder, two on mixed diagnosis psychiatric in & outpatients, and two on mixed diagnosis psychiatric outpatients. Non-psychiatric residual category: $n = 8$; including two studies on military, one on veterans, one on stalkers, one on HIV infected males, one on survivors of childhood cancer, one on prisoners, and one on emergency department visitors.

completed suicides (median 60; range 3–1429). Suicidal ideation was assessed retrospectively by interviewing a next of kin or clinician of the deceased ($n = 19$), was extracted from medical records ($n = 45$) or was determined by asking the patient him/herself ($n = 16$) (missing: $n = 1$) (Supplement S3).

The majority of included studies were (nested) case-control studies ($n = 51$, 63.0%). In only 26 studies the absolute risk of completed suicide could be extracted. None of the 81 eligible studies assessed the effect of suicidal ideation on completed suicide with sole adjustment for previous suicide attempts. However, there were 46 studies that applied a multivariable model, but only 28 adjusted for previous attempts as a separate determinant in this model.

Association between suicidal ideation and subsequent completed suicide

For all population subgroups investigated in this meta-analysis, the pooled risk of suicide was significantly higher in study participants who had expressed suicidal ideation compared with study participants who had not. Whereas the overall risk ratio (RR) was 4.17 (95% confidence interval (CI) 3.29–5.27), the risk ratios per population subgroup varied substantially. The risk ratio was highest in the non-psychiatric residual subgroup, followed by schizophrenia, mixed psychiatric patients, the general population, psychiatric inpatients and affective disorder (Fig. 2). Overall, the risk ratio was significantly higher in non-




Fig. 2. Forest plot showing the association between suicidal ideation and subsequent completed suicide. CI, confidence interval.
^aAuthors provided additional data to compute effect size.




Fig. 3. Bar chart showing suicide risk ratio and the suicide risk (%) during first year of follow-up. Lines indicate 95% confidence interval. Numbers of studies included in (A) and (B) differ because in only 26 of the 81 studies the absolute risk of completed suicide could be extracted. A. Risk ratio for suicide after expression of suicidal ideation, stratified for psychiatric and non-psychiatric populations. B. Results specified separately for study participants who had expressed suicidal ideation and study participants who had not, stratified for psychiatric and non-psychiatric populations. Percentages calculated using maximum likelihood meta-regression analyses with mean study follow-up time as determinant and log_e-transformed rates of completed suicide as outcome (see Fig. 4).

psychiatric populations (RR 6.61; 95% CI 4.62–9.47) than in psychiatric populations (RR 3.53; 95% CI 2.81–4.44) (p -value = 0.004) (Fig. 3).

Risk of completed suicide

Although the psychiatric subgroups had the lowest risk ratios, meta-regression analyses showed the suicide risk was highest in the psychiatric subgroups who had expressed suicidal ideation (risk during first year of follow-up 1.40%; 95% CI 0.74–2.64). This was clearly higher than the suicide risk in psychiatric patients who had not expressed suicidal ideation (risk during first year of follow-up 0.41%; 95% CI 0.20–0.84). In non-psychiatric study participants the suicide risk during the first year of follow-up was 0.23% (95% CI 0.10–0.54) in participants who had expressed suicidal ideation and 0.04% (95% CI 0.01–0.13) in participants who had not (Fig. 3 and Fig. 4). When grouping all studies, the suicide risk during the first year of follow-up after expression of suicidal ideation was 1.03% (95% CI 0.43–2.44) (data not shown).

Risk of bias assessment and sensitivity analyses

When restricting the analyses to studies that scored 'adequate' on all four risk of bias items ($n=34$, 42.0%), the pooled risk ratio decreased slightly to 3.41 (95% CI 2.67–4.35) and the overall suicide risk

during the first year of follow-up also decreased slightly (risk during first year of follow-up 0.77%; 95% CI 0.27–2.16). In addition, restricting the primary analysis to cohort studies resulted in a slight decrease in the risk ratio (RR 3.10; 95% CI 2.22–4.32). Restricting the secondary analysis to studies that assessed suicidal ideation in the month around baseline resulted in a similar risk (risk during first year of follow-up 0.99%; 95% CI 0.39–2.51).

Is suicidal ideation an independent predictor for completed suicide?

Of all studies that used a multivariable model for the prediction of suicide and adjusted for (among other predictors) the presence of a previous suicide attempt ($n=28$), only 14 reported the adjusted effect size of suicidal ideation. In 11 of these studies, suicidal ideation was a significant independent predictor of completed suicide. The other 14 studies that included previous attempts in their multivariable model either did not include suicidal ideation as a separate predictor ($n=6$), or did include suicidal ideation but did not report the adjusted effect size ($n=8$), which was non-significant in seven of these studies. While in at least 11 studies suicidal ideation was a significant independent predictor of completed suicide, the 14 adjusted effect sizes were not pooled due to large underreporting of mainly non-significant results.




Fig. 4. Meta-regression of mean study follow-up time on suicide rate. Maximum likelihood meta-regression analyses with mean study follow-up time as determinant and \log_e -transformed rates of completed suicide as outcome presented on logarithmic scales. The bubble size is proportional to the study's weight. Vertical line indicates 1 year follow-up. The suicide risk can be calculated using the following formulas: suicide risk (during first year of follow-up) = $1 - e^{-(\text{rate per person-year at year 1})}$. Figure part A, B: Psychiatric subgroup included: Dingman & McGlashan, 1986; Drake & Cotton, 1986; Allebeck *et al.* 1987; Goldstein *et al.* 1991; Bradvik & Berglund, 1993; Stephens *et al.* 1999; Bioulac *et al.* 2000; de Hert *et al.* 2001; Spiessl *et al.* 2002; Coryell & Young, 2005; Dutta *et al.* 2007; Berg, 2010; Dutta *et al.* 2011; Kuo *et al.* 2011; Sani *et al.* 2011; Park *et al.* 2013; Simon *et al.* 2013. Lin *et al.* 2014; Figure part C, D: Non-psychiatric subgroup included: Crandall *et al.* 2006; Khang *et al.* 2010; McEwan *et al.* 2010; Hyman, 2012; Brinkman *et al.* 2014; Kleiman & Liu, 2014; Al-Sayegh *et al.* 2015; Werbeloff *et al.* 2015.

Subgroup analyses

There was considerable heterogeneity (I^2 for overall risk ratio = 84.7%, Q -value = 523), even when stratified for specific populations (I^2 range 39–79%, Q -values 12.7–65.6). The results of subgroup analyses can be found in Supplement S4.

Publication bias

The overall funnel plot (Supplement S5) for the primary research question showed a rather symmetrical funnel plot. However, visual inspection of some of the funnel plots of the subgroups showed there might be missing studies with negative effects. The overall Egger's test indicated no funnel plot asymmetry (two-tailed

p -value = 0.94). Only in the general population subgroup, the Egger's test for funnel plot asymmetry was significant (two-tailed p -value = 0.04).

Discussion

Meta-analysing 81 eligible studies showed that overall, people expressing suicidal ideation are four times more likely to die by suicide than people not expressing suicidal ideation, with the highest relative risks in non-psychiatric populations. However, absolute suicide risks, rather than relative risks, are essential in suicide prediction, and were highest in psychiatric populations, with a suicide risk during the first year of follow-up of 1.4% compared with 0.23% in non-psychiatric populations.

The results of this meta-analysis show an overall modest relationship, with moderate risk ratios for psychiatric subgroups and strong risk ratios for non-psychiatric subgroups (Rosenthal, 1996). The results are in line with previous meta-analyses on suicidal ideation and subsequent completed suicide (Large *et al.*, 2011a, b; Hawton *et al.*, 2005a, 2013; Chapman *et al.*, 2015), which only assessed psychiatric populations. The highest relative risk was found in a meta-analysis restricted to schizophrenic patients (Hawton *et al.* 2005a; Chapman *et al.* 2015) followed by meta-analyses restricted to psychiatric inpatients (Large *et al.*, 2011a, b) and depressed patients (Hawton *et al.* 2013; Chapman *et al.* 2015). Remarkably, in contrast to our study, one recent meta-analysis (Chapman *et al.* 2015) reported that the association between suicidal ideation and completed suicide was not significant in patients with a mood disorder, probably explained by the broader definition of suicidal ideation in our study and the fact that we excluded studies that compared expression of suicidal ideation in suicide completers *v.* attempters. Unfortunately, none of the previous meta-analyses (Hawton *et al.*, 2005a, b; 2013; Large *et al.*, 2011a, b; Chapman *et al.*, 2015) assessed the absolute risks of suicide.

The expression of suicidal ideation was reported as a significant independent predictor of suicide more often than expected by chance. However, underreporting of non-significant effects refrained us from concluding to what extent suicidal ideation is an independent predictor of completed suicide. Many studies included in our meta-analysis did develop a prediction model for suicide, although it is widely accepted that it is very difficult to accurately predict suicide for an individual patient (van Hemert *et al.* 2012), and there is a lot of debate on whether or not clinical risk categorisation is helpful in preventing completed suicide (Large *et al.* 2011c; O'Connor *et al.* 2014). The low baseline rate of suicide results in low positive predictive values and in the majority of suicides occurring in the group that is classified as low-risk according to the prediction model (Large *et al.* 2011b; Madsen & Nordentoft, 2012; Paton *et al.* 2014). It was striking that only 60% of the studies that developed a prediction model for suicide included previous suicide attempts as a predictor, whereas this has been reported as one of the most important independent predictors of completed suicide (Large *et al.* 2011c). Future studies on suicide risk assessment should therefore include known predictors from the literature (e.g. previous suicide attempts) rather than only selecting predictors by univariable screening in the developmental dataset (Steyerberg, 2009).

Strengths and limitations

A major strength of our paper was the attempt to provide risk estimates for different subgroups. Moreover, we

strictly excluded studies that might have combined suicidal ideation with behaviour, and included many studies for the primary research aim. There are also limitations that need to be considered. First, the majority of studies only provided long-term follow-up results, while the short-term risk has most clinical value. Although the provided suicide risks should be interpreted with caution due to the declining rates over time and the limited number of studies, especially with short-term follow-up, that could be included in the meta-regression analyses, results clearly show the risk of suicide is higher after expression of suicidal ideation, especially in psychiatric patients. Future studies should focus on short-term suicide risk in the different psychiatric populations. Second, only part of the large amount of heterogeneity could be explained for. Included studies often did not provide several patient characteristics that would be interesting in exploring/explaining this heterogeneity, like duration of suicidal ideation (ten Have *et al.* 2013) or behavioural traits (Turecki, 2014). We did not find a significant difference between suicidal ideation assessment subgroups, possible due to varying or lacking definitions of suicidal ideation and the large variation in suicidal ideation assessment methods within the subgroups. Unfortunately we could not further explore this variation due to insufficient reporting in the individual studies and study level confounding. Clear definitions of suicidal ideation and its assessment method are necessary (Silverman *et al.* 2007) and validated and standardised assessment methods should be used. Even within studies that used standardised assessment methods, the instruments used varied widely with only a maximum of two studies using the same instrument. This can also be a source of heterogeneity as different validated suicidal ideation assessment methods (Yigletu *et al.* 2004; Vuorilehto *et al.* 2014) can result in different suicide ideation prevalences, even in the same population at the same time point. The considerable heterogeneity that still exists between studies within the different population subgroups, but that supposedly also exists within individual studies, limits the generalisability of our findings to individual patients and will result in higher suicide risks in some patients and lower suicide risks in others. We recommend future studies to explore sources of clinical heterogeneity in more detail, especially with regard to suicidal ideation assessment method. Third, study level confounding (Hingorani *et al.* 2013) might have influenced the results of our subgroup analyses, e.g. resulted in a significant effect of gender on the association between suicidal ideation and completed suicide, as the percentage of females was lower in certain high-risk population subgroups like schizophrenia. Fourth, some required study parameters could not be directly extracted from the original articles. For example the exposed person-years were conservatively estimated when possible,

which could have resulted in an underestimation of the suicide risk. Fifth, reporting bias cannot be ruled out. The power of Egger's test is relatively low (Sterne *et al.* 2011) and effects in the different population subgroups might have been underestimated as we did not attempt to access unpublished data or grey literature and due to underreporting of negative and non-significant effects. Sixth, studies that were published after February 5, 2016 were not included in this meta-analysis. Our meta-analysis especially has limited power on analyses regarding absolute suicide risk in psychiatric and non-psychiatric populations and the independent predictive effect of suicidal ideation. A future update of this meta-analysis should focus on studies that address these research questions.

Implications for clinical practice

For clinical psychiatric care, it is important to inquire about suicidal ideation in psychiatric patients given the substantial risk after expression of suicidal ideation, especially in the first period after expression. Although the relative risk was highest in non-psychiatric populations, the absolute suicide risk after expression of suicidal ideation in non-psychiatric populations was lower than in psychiatric populations, given their lower baseline suicide risk. However, even in psychiatric patients, suicidal ideation assessment should be placed in perspective as the suicide risk during the first year of follow-up among psychiatric patients who had not expressed suicidal ideation was 0.41%, which was higher than the suicide risk in non-psychiatric persons who had expressed suicidal ideation. Expression of suicidal ideation is one of the factors that clinicians take into account when assessing suicide risk in an individual patient (Popovic *et al.* 2014), which should of course not merely be based on a 'tick box' approach that classifies patients as either at high or low risk on the basis of a list of risk factors (van Hemert *et al.* 2012; O'Connor *et al.* 2014). The suicide risks provided in this meta-analysis can help clinicians in their clinical suicide risk assessment and stepped care approach, for example to decide whether burdensome secondary prevention measures such as (compulsory) admission are proportional to the risk of suicide.

In a patient who expresses suicidal ideation, follow-up inquiries are needed about the nature of these thoughts and the assessment of other known risk factors for suicide (Hawton & van Heeringen, 2009). As patients will not always express suicidal ideation spontaneously (van Hemert *et al.* 2012), it is important to ask them about suicidal thoughts. Actually, asking patients about suicidal ideation can reduce future suicidal ideation and follow-up inquiries can benefit long-term mental health (Dazzi *et al.* 2014). Patients often feel

relieved by having an opportunity to talk about their suicidal ideas (Gliatto & Rai, 1999), especially if clinicians take time, show empathy, and acknowledge the suicidal feelings (Talseth *et al.* 1999). Remarkably, in several of the included studies that assessed the presence of suicidal ideation in medical records, this information was missing. This means that it was unavailable to clinicians and appropriate care might have been unnecessarily withheld from these patients (King *et al.* 2001b).

Fortunately, the majority of patients with suicidal ideation do not die by suicide. Nevertheless, assessment of suicidal ideation is of priority in people with psychiatric illnesses and when a patient expresses suicidal ideation, prompt secondary prevention strategies are necessary to reduce their substantial increased risk of suicide. Future suicide prediction studies should focus on absolute suicide risks and the independent predictive effect of suicidal ideation.

Supplementary Material

The supplementary material for this article can be found at <https://doi.org/10.1017/S2045796016001049>

Acknowledgements

We would like to thank medical librarian J.W. Schoones, Walaeus Library, Leiden University Medical Center for his help in conducting the literature search. In addition, we would like to thank the authors that provided additional information on their studies.

Funding

This research received no specific grant from any funding agency, commercial or not-for-profit sectors.

Conflict of Interest

None.

Availability of Data and Materials

Data supporting our findings can be found in Fig. 2 and Supplement S3.

References

- Allebeck P, Varla A, Kristjansson E, Wistedt B (1987). Risk factors for suicide among patients with schizophrenia. *Acta Psychiatrica Scandinavica* 76, 414–419.
- Al-Sayegh H, Lowry J, Polur RN, Hines RB, Liu F, Zhang J (2015). Suicide history and mortality: a follow-up of a national cohort in the United States. *Archives of Suicide Research* 19, 35–47.

- Altman DG (2001). Systematic reviews of evaluations of prognostic variables. *British Medical Journal* **323**, 224–228.
- Appleby L, Dennehy JA, Thomas CS, Faragher EB, Lewis G (1999a). Aftercare and clinical characteristics of people with mental illness who commit suicide: a case-control study. *Lancet* **353**, 1397–1400.
- Appleby L, Cooper J, Amos T, Faragher B (1999b). Psychological autopsy study of suicides by people aged under 35. *British Journal of Psychiatry* **175**, 168–174.
- Baader-Matthei T, Richter P, Mundt C (2004). Suicides of psychiatric hospitalized patients and their risk factors. A case control study. *Revista Chilli Neuro-Psiquiatria* **42**, 293–316.
- Batterham PJ, Calear AL, Mackinnon AJ, Christensen H (2013). The association between suicidal ideation and increased mortality from natural causes. *Journal of Affective Disorders* **150**, 855–860.
- Berg JE (2010). Death by suicide long after electroconvulsive therapy. Is the sense of coherence test of Antonovsky a predictor of mortality from depression? *Mental Illness* **2**, 11–15.
- Bertelsen M, Jeppesen P, Petersen L, Thorup A, Øhlenschlaeger J, le Quach P, Christensen TØ, Krarup G, Jørgensen P, Nordentoft M (2007). Suicidal behaviour and mortality in first-episode psychosis: the OPUS trial. *British Journal of Psychiatry Suppl* **51**, s140–s146.
- Bickley H, Hunt IM, Windfuhr K, Shaw J, Appleby L, Kapur N (2013). Suicide within two weeks of discharge from psychiatric inpatient care: a case-control study. *Psychiatric Services* **64**, 653–659.
- Bioulac S, Bourgeois M, Ekouevi DK, Bonnin JM, Gonzales B, Castello MF (2000). [Predictive factors of suicide? an 8-year-long prospective longitudinal study of 200 psychiatric inpatients]. *Encephale* **26**, 1–7.
- Borenstein M, Hedges LV, Higgins JPT, Rothstein HR (2009a). Chapter 12. Random-effects model. In *Introduction to Meta-Analysis* (ed. Sharples K), pp. 69–75. John Wiley & Sons, Ltd.: Chichester, UK.
- Borenstein M, Hedges LV, Higgins JPT, Rothstein HR (2009b). Chapter 19. Subgroup analyses. In *Introduction to Meta-Analysis* (ed. Sharples K), pp. 149–186. John Wiley & Sons, Ltd.: Chichester, UK.
- Borenstein M, Hedges LV, Higgins JPT, Rothstein HR (2009c). Chapter 24. Multiple outcomes or time-points within a study. In *Introduction to Meta-Analysis* (ed. Sharples K), pp. 225–238. John Wiley & Sons, Ltd.: Chichester, UK.
- Bradvik L, Berglund M (1993). Risk factors for suicide in melancholia. A case-record evaluation of 89 suicides and their controls. *Acta Psychiatrica Scandinavica* **87**, 306–311.
- Brinkman TM, Zhang N, Recklitis CJ, Kimberg C, Zeltzer LK, Muriel AC, Stovall M, Srivastava DK, Sklar CA, Robison LL, Krull KR (2014). Suicide ideation and associated mortality in adult survivors of childhood cancer. *Cancer* **120**, 271–277.
- Brown GK, Beck AT, Steer RA, Grisham JR (2000). Risk factors for suicide in psychiatric outpatients: a 20-year prospective study. *Journal of Consulting and Clinical Psychology* **68**, 371–377.
- Bukstein OG, Brent DA, Perper JA, Moritz G, Baugher M, Schweers J, Roth C, Balach L (1993). Risk factors for completed suicide among adolescents with a lifetime history of substance abuse: a case-control study. *Acta Psychiatrica Scandinavica* **88**, 403–408.
- Castle K, Duberstein PR, Meldrum S, Conner KR, Conwell Y (2004). Risk factors for suicide in blacks and whites: an analysis of data from the 1993 National Mortality Followback Survey. *American Journal of Psychiatry* **161**, 452–458.
- Chapman CL, Mullin K, Ryan CJ, Kuffel A, Nielsens O, Large MM (2015). Meta-analysis of the association between suicidal ideation and later suicide among patients with either a schizophrenia spectrum psychosis or a mood disorder. *Acta Psychiatrica Scandinavica* **131**, 162–173.
- Cheng KK, Leung CM, Lo WH, Lam TH (1990). Risk factors of suicide among schizophrenics. *Acta Psychiatrica Scandinavica* **81**, 220–224.
- Colman I, Kingsbury M, Sareen J, Bolton J, van Walraven C (2016). Migraine headache and risk of self-harm and suicide: a population-based study in Ontario, Canada. *Headache* **56**, 132–140.
- Conlon L, Garland M, Prescott P, Mannion L, Leonard M, Fahy TJ (2007). Psychiatric aftercare and suicide risk: a case-control study using blind rating. *Archives of Suicide Research* **11**, 291–295.
- Conwell Y (2009). Suicide prevention in later life: a glass half full, or half empty? *American Journal of Psychiatry* **166**, 845–848.
- Coryell W, Young EA (2005). Clinical predictors of suicide in primary major depressive disorder. *Journal of Clinical Psychiatry* **66**, 412–417.
- Coryell W, Kriener A, Butcher B, Nurnberger J, McMahon F, Berrettini W, Fiedorowicz J (2016). Risk factors for suicide in bipolar I disorder in two prospectively studied cohorts. *Journal of Affective Disorders* **190**, 1–5.
- Crandall C, Fullerton-Gleason L, Aguero R, LaValley J (2006). Subsequent suicide mortality among emergency department patients seen for suicidal behavior. *Academic Emergency Medicine* **13**, 435–442.
- Dazzi T, Gribble R, Wessely S, Fear NT (2014). Does asking about suicide and related behaviours induce suicidal ideation? What is the evidence? *Psychological Medicine* **44**, 3361–3363.
- De Hert M, McKenzie K, Peuskens J (2001). Risk factors for suicide in young people suffering from schizophrenia: a long-term follow-up study. *Schizophrenia Research* **47**, 127–134.
- De Leo D, Draper BM, Snowden J, Kolves K (2013a). Contacts with health professionals before suicide: missed opportunities for prevention? *Comprehensive Psychiatry* **54**, 1117–1123.
- De Leo D, Draper BM, Snowden J, Kolves K (2013b). Suicides in older adults: a case-control psychological autopsy study in Australia. *Journal of Psychiatric Research* **47**, 980–988.
- Didham R, Dovey S, Reith D (2006). Characteristics of general practitioner consultations prior to suicide: a nested case-control study in New Zealand. *New Zealand Medical Journal* **119**, U2358.
- Dingman CW, McGlashan TH (1986). Discriminating characteristics of suicides. Chestnut Lodge follow-up sample including patients with affective disorder, schizophrenia and schizoaffective disorder. *Acta Psychiatrica Scandinavica* **74**, 91–97.

- Dong JY, Ho TP, Kan CK (2005). A case-control study of 92 cases of in-patient suicides. *Journal of Affective Disorders* 87, 91–99.
- Drake RE, Cotton PG (1986). Depression, hopelessness and suicide in chronic schizophrenia. *British Journal of Psychiatry* 148, 554–559.
- Dutta R, Boydell J, Kennedy N, van Os J, Fearon P, Murray RM (2007). Suicide and other causes of mortality in bipolar disorder: a longitudinal study. *Psychological Medicine* 37, 839–847.
- Dutta R, Murray RM, Allardyce J, Jones PB, Boydell J (2011). Early risk factors for suicide in an epidemiological first episode psychosis cohort. *Schizophrenia Research* 126, 11–19.
- Farberow NL, Shneidman ES, Neuringer C (1966). Case history and hospitalization factors in suicides of neuropsychiatric hospital patients. *Journal of Nervous and Mental Disease* 142, 32–44.
- Farwett J, Scheftner WA, Fogg L, Clark DC, Young MA, Hedeker D, Gibbons R (1990). Time-related predictors of suicide in major affective disorder. *American Journal of Psychiatry* 147, 1189–1194.
- Flood RA, Seager CP (1968). A retrospective examination of psychiatric case records of patients who subsequently committed suicide. *British Journal of Psychiatry* 114, 443–450.
- Fruehwald S, Matschnig T, Koenig F, Bauer P, Frottier P (2004). Suicide in custody: case-control study. *British Journal of Psychiatry* 185, 494–498.
- Funahashi T, Ibuki Y, Domon Y, Nishimura T, Akehashi D, Sugiura H (2000). A clinical study on suicide among schizophrenics. *Psychiatry and Clinical Neurosciences* 54, 173–179.
- Gliatto MF, Rai AK (1999). Evaluation and treatment of patients with suicidal ideation. *American Family Physician* 59, 1500–1506.
- Goldstein RB, Black DW, Nasrallah A, Winokur G (1991). The prediction of suicide. Sensitivity, specificity, and predictive value of a multivariate model applied to suicide among 1906 patients with affective disorders. *Archives General Psychiatry* 48, 418–422.
- Grobbee DE, Hoes AW (2009). Diagnostic research. In *Clinical Epidemiology. Principles, Methods, and Applications for Clinical Research* (ed. Grobbee DE and Hoes AW), pp. 58–102. Jones and Bartlett Publishers: Sudbury, Massachusetts.
- Harris EC, Barraclough B (1997). Suicide as an outcome for mental disorders. A meta-analysis. *British Journal of Psychiatry* 170, 205–228.
- Hawton K, van Heeringen K (2009). Suicide. *Lancet* 373, 1372–1381.
- Hawton K, Sutton L, Haw C, Sinclair J, Deeks JJ (2005a). Schizophrenia and suicide: systematic review of risk factors. *British Journal of Psychiatry* 187, 9–20.
- Hawton K, Sutton L, Haw C, Sinclair J, Harriss L (2005b). Suicide and attempted suicide in bipolar disorder: a systematic review of risk factors. *Journal of Clinical Psychiatry* 66, 693–704.
- Hawton K, Casanas IC, Haw C, Saunders K (2013). Risk factors for suicide in individuals with depression: a systematic review. *Journal of Affective Disorders* 147, 17–28.
- Hingorani AD, Windt DA, Riley RD, Abrams K, Moons KG, Steyerberg EW, Schroter S, Sauerbrei W, Altman DG, Hemingway H, PROGRESS Group (2013). Prognosis research strategy (PROGRESS) 4: stratified medicine research. *British Medical Journal* 346, e5793.
- Hoyer EH, Licht RW, Mortensen PB (2009). Risk factors of suicide in inpatients and recently discharged patients with affective disorders. A case-control study. *European Psychiatry* 24, 317–321.
- Hunt IM, Kapur N, Webb R, Robinson J, Burns J, Turnbull P, Shaw J, Appleby L (2007). Suicide in current psychiatric in-patients: a case-control study The National Confidential Inquiry into Suicide and Homicide. *Psychological Medicine* 37, 831–837.
- Hunt IM, Kapur N, Webb R, Robinson J, Burns J, Shaw J, Appleby L (2009). Suicide in recently discharged psychiatric patients: a case-control study. *Psychological Medicine* 39, 443–449.
- Hunt IM, Bickley H, Windfuhr K, Shaw J, Appleby L, Kapur N (2013). Suicide in recently admitted psychiatric in-patients: a case-control study. *Journal of Affective Disorders* 144, 123–128.
- Hyman JRLL (2012). Suicide incidence and risk factors in an active duty US military population. *American Journal of Public Health* 102 (Suppl. 1), s138–s146.
- Isometsa ET, Heikkinen ME, Marttunen MJ, Henriksson MM, Aro HM, Lonnqvist JK (1995). The last appointment before suicide: is suicide intent communicated? *American Journal of Psychiatry* 152, 919–922.
- Jollant F, Malafosse A, Docto R, Macdonald C (2014). A pocket of very high suicide rates in a non-violent, egalitarian and cooperative population of South-East Asia. *Psychological Medicine* 44, 2323–2329.
- Kasckow J, Liu N, Haas GL, Phillips MR (2010). Case-control study of the relationship of depressive symptoms to suicide in a community-based sample of individuals with schizophrenia in China. *Schizophrenia Research* 122, 226–231.
- Kelly DL, Shim JC, Feldman SM, Yu Y, Conley RR (2004). Lifetime psychiatric symptoms in persons with schizophrenia who died by suicide compared to other means of death. *Journal of Psychiatric Research* 38, 531–536.
- Kessler RC, Warner CH, Ivany C, Petukhova MV, Rose S, Bromet EJ, Brown M III, Cai T, Colpe LJ, Cox KL, Fullerton CS, Gilman SE, Gruber MJ, Heeringa SG, Lewandowski-Romps L, Li J, Millikan-Bell AM, Naifeh JA, Nock MK, Rosellini AJ, Sampson NA, Schoenbaum M, Stein MB, Wessely S, Zaslavsky AM, Ursano RJ, for the Army STARRS Collaborators (2015). Predicting suicides after psychiatric hospitalization in US army soldiers. The army study to assess risk and resilience in servicemembers (army STARRS). *JAMA Psychiatry* 72, 49–57.
- Khang YH, Kim HR, Cho SJ (2010). Relationships of suicide ideation with cause-specific mortality in a longitudinal study of South Koreans. *Suicide Life Threatening Behaviour* 40, 465–475.
- Kim HM, Smith EG, Ganoczy D, Walters H, Stano CM, Ilgen MA, Bohnert AS, Valenstein M (2012). Predictors of suicide in patient charts among patients with depression in

- the Veterans Health Administration health system: importance of prescription drug and alcohol abuse. *Journal of Clinical Psychiatry* 73, e1269–e1275.
- King EA, Baldwin DS, Sinclair JM, Campbell MJ** (2001a). The Wessex Recent In-Patient Suicide Study. 2. Case-control study of 59 in-patient suicides. *British Journal of Psychiatry* 178, 537–542.
- King EA, Baldwin DS, Sinclair JM, Baker NG, Campbell MJ, Thompson C** (2001b). The Wessex Recent In-Patient Suicide Study. 1. Case-control study of 234 recently discharged psychiatric patient suicides. *British Journal of Psychiatry* 178, 531–536.
- Kjelsberg E, Eikeseth PH, Dahl AA** (1991). Suicide in borderline patients—predictive factors. *Acta Psychiatrica Scandinavica* 84, 283–287.
- Kleiman EM, Liu RT** (2014). Prospective prediction of suicide in a nationally representative sample: religious service attendance as a protective factor. *British Journal of Psychiatry* 204, 262–266.
- Kullgren G** (1988). Factors associated with completed suicide in borderline personality disorder. *Journal of Nervous and Mental Disease* 176, 40–44.
- Kuo CJ, Tsai SY, Liao YT, Conwell Y, Lin SK, Chang CL, Chen CC, Chen WJ** (2011). Risk and protective factors for suicide among patients with methamphetamine dependence: a nested case-control study. *Journal of Clinical Psychiatry* 72, 487–493.
- Large M, Smith G, Sharma S, Nielssen O, Singh SP** (2011a). Systematic review and meta-analysis of the clinical factors associated with the suicide of psychiatric in-patients. *Acta Psychiatrica Scandinavica* 124, 18–29.
- Large M, Sharma S, Cannon E, Ryan C, Nielssen O** (2011b). Risk factors for suicide within a year of discharge from psychiatric hospital: a systematic meta-analysis. *Australian & New Zealand Journal of Psychiatry* 45, 619–628.
- Large MM, Nielssen O** (2012). Suicidal ideation and later suicide (letter). *American Journal of Psychiatry* 169, 662.
- Large M, Ryan C, Nielssen O** (2011c). The validity and utility of risk assessment for inpatient suicide. *Australasian Psychiatry* 19, 507–512.
- Li J, Ran MS, Hao Y, Zhao Z, Guo Y, Su J, Lu H** (2008). Inpatient suicide in a Chinese psychiatric hospital. *Suicide Life Threatening Behaviour* 38, 449–455.
- Lin SK, Hung TM, Liao YT, Lee WC, Tsai SY, Chen CC, Kuo CJ** (2014). Protective and risk factors for inpatient suicides: a nested case-control study. *Psychiatry Research* 217, 54–59.
- Lui SY** (2009). Risk factors for deliberate self-harm and completed suicide in young Chinese people with schizophrenia. *Australian & New Zealand Journal of Psychiatry* 43, 252–259.
- Lukaschek K, Baumert J, Krawitz M, Erazo N, Förstl H, Ladwig KH** (2014). Determinants of completed railway suicides by psychiatric in-patients: case-control study. *The British Journal of Psychiatry* 205, 398–406.
- Madsen T, Nordentoft M** (2012). Drs Madsen and Nordentoft reply. *Journal of Clinical Psychiatry* 73, 1034–1035.
- Mahon MJ, Tobin JP, Cusack DA, Kelleher C, Malone KM** (2005). Suicide among regular-duty military personnel: a retrospective case-control study of occupation-specific risk factors for workplace suicide. *American Journal of Psychiatry* 162, 1688–1696.
- McEwan T, Mullen P, MacKenzie R** (2010). Suicide among stalkers. *Journal of Forensic Psychiatry and Psychology* 21, 514–520.
- McGirr A, Renaud J, Seguin M, Alda M, Benkelfat C, Lesage A, Turecki G** (2007). An examination of DSM-IV depressive symptoms and risk for suicide completion in major depressive disorder: a psychological autopsy study. *Journal of Affective Disorders* 97, 203–209.
- McManus H, Petoumenos K, Franic T, Kelly MD, Watson J, O'Connor CC, Jeanes M, Hoy J, Cooper DA, Law MG; Australian HIV, Observational Database** (2014). Determinants of suicide and accidental or violent death in the Australian HIV Observational Database. *PLoS ONE* 9, e89089.
- Mock CN, Grossman DC, Mulder D, Stewart C, Koepsell TS** (1996). Health care utilization as a marker for suicidal behavior on an American Indian Reservation. *Journal of General Internal Medicine* 11, 519–524.
- Murphy GE, Wetzel RD, Robins E, McEvoy L** (1992). Multiple risk factors predict suicide in alcoholism. *Archives General Psychiatry* 49, 459–463.
- Nock MK, Borges G, Bromet EJ, Alonso J, Angermeyer M, Beautrais A, Bruffaerts R, Chiu WT, de Girolamo G, Gluzman S, de Graaf R, Gureje O, Haro JM, Huang Y, Karam E, Kessler RC, Lepine JP, Levinson D, Medina-Mora ME, Ono Y, Posada-Villa J, Williams D** (2008). Cross-national prevalence and risk factors for suicidal ideation, plans and attempts. *British Journal of Psychiatry* 192, 98–105.
- Nordentoft M, Mortensen PB, Pedersen CB** (2011). Absolute risk of suicide after first hospital contact in mental disorder. *Archives of General Psychiatry* 68, 1058–1064.
- Nyman AK, Jonsson H** (1986). Patterns of self-destructive behaviour in schizophrenia. *Acta Psychiatrica Scandinavica* 73, 252–262.
- O'Connor N, Allan J, Scott C** (2014). Debate: clinical risk categorization is valuable in the prevention of suicide and severe violence? Yes. *Australasian Psychiatry* 22, 7–9.
- Palacio C, Garcia J, Diago J, Zapata C, Lopez G, Ortiz J, Lopez M** (2007). Identification of suicide risk factors in Medellin, Colombia: a case-control study of psychological autopsy in a developing country. *Archives of Suicide Research* 11, 297–308.
- Park S, Choi JW, Kyoung Yi K, Hong JP** (2013). Suicide mortality and risk factors in the 12 months after discharge from psychiatric inpatient care in Korea: 1989–2006. *Psychiatry Research* 208, 145–150.
- Paton MB, Large MM, Ryan CJ** (2014). Debate: clinical risk categorisation is valuable in the prevention of suicide and severe violence—no. *Australasian Psychiatry* 22, 10–12.
- Popovic D, Benabarre A, Crespo JM, Goikolea JM, González-Pinto A, Gutiérrez-Rojas L, Montes JM, Vieta E** (2014). Risk factors for suicide in schizophrenia: systematic review and clinical recommendations. *Acta Psychiatrica Scandinavica* 130, 418–426.
- Posner K, Brown GK, Stanley B, Brent DA, Yershova KV, Oquendo MA, Currier GW, Melvin GA, Greenhill L, Shen S, Mann JJ** (2011). The Columbia-Suicide Severity

- Rating Scale: initial validity and internal consistency findings from three multisite studies with adolescents and adults. *American Journal of Psychiatry* **168**, 1266–1277.
- Powell J, Geddes J, Deeks J, Goldacre M, Hawton K** (2000). Suicide in psychiatric hospital in-patients. Risk factors and their predictive power. *British Journal of Psychiatry* **176**, 266–272.
- Roos JL, Boraine H, Bodemer W** (1992). [Suicide in schizophrenic patients]. *South African Medical Journal* **81**, 365–369.
- Rosenthal JA** (1996). Qualitative descriptors of strength of association and effect size. *Journal of Social Service Research* **21**, 37–59.
- Rowe JL, Bruce ML, Conwell Y** (2006). Correlates of suicide among home health care utilizers who died by suicide and community controls. *Suicide Life Threatening Behaviour* **36**, 65–75.
- Runeson BS** (2002). Suicide after parasuicide (editorial). *British Medical Journal* **325**, 1125–1126.
- Salama AA** (1988). Depression and suicide in schizophrenic patients. *Suicide Life Threatening Behaviour* **18**, 379–384.
- Sani G, Tondo L, Koukopoulos A, Reginaldi D, Kotzalidis GD, Koukopoulos AE, Manfredi G, Mazarini L, Pacchiarotti I, Simonetti A, Ambrosi E, Angeletti G, Girardi P, Tatarelli R** (2011). Suicide in a large population of former psychiatric inpatients. *Psychiatry and Clinical Neurosciences* **65**, 286–295.
- Shah AK, Ganesvaran T** (1997). Inpatient suicides in an Australian mental hospital. *Australian & New Zealand Journal of Psychiatry* **31**, 291–298.
- Silverman MM, Berman AL, Sanddal ND, O'Carroll PW, Joiner TE** (2007). Rebuilding the tower of Babel: a revised nomenclature for the study of suicide and suicidal behaviors. Part 1: Background, rationale, and methodology. *Suicide Life Threatening Behaviour* **37**, 248–263.
- Simon GE, Rutter CM, Peterson D, Oliver M, Whiteside U, Operskalski B, Ludman EJ** (2013). Does response on the PHQ-9 Depression Questionnaire predict subsequent suicide attempt or suicide death? *Psychiatric Services* **64**, 1195–1202.
- Spießl H, Hubner-Liebermann B, Cording C** (2002). Suicidal behaviour of psychiatric in-patients. *Acta Psychiatrica Scandinavica* **106**, 134–138.
- Stephens JH, Richard P, McHugh PR** (1999). Suicide in patients hospitalized for schizophrenia: 1913–1940. *Journal of Nervous and Mental Disease* **187**, 10–14.
- Sterne JAC, Egger M, Moher D** (2011). Chapter 10. *Addressing reporting biases*. In *Cochrane Handbook for Systematic Reviews of Interventions* version 5.1.0 (ed. Higgins JPT and Green S). The Cochrane collaboration. Available from www.handbook.cochrane.org
- Steyerberg EW** (2009). Restrictions on candidate predictors. In *Clinical Prediction Models*, pp. 175–189. Springer: New York, USA.
- Taiminen T, Huttunen J, Heilä H, Henriksson M, Isometsä E, Kähkönen J, Tuominen K, Lönnqvist J, Addington D, Helenius H** (2001). The Schizophrenia Suicide Risk Scale (SSRS): development and initial validation. *Schizophrenia Research* **47**, 199–213.
- Talseth AG, Lindseth A, Jacobsson L, Norberg A** (1999). The meaning of suicidal psychiatric in-patients' experiences of being cared for by mental health nurses. *Journal of Advanced Nursing* **29**, 1034–1041.
- ten Have M, de Graaf R, van Dorsselaer S, Verdurmen J, van 't Land H, Vollebergh W, Beekman A** (2009). Incidence and course of suicidal ideation and suicide attempts in the general population. *Canadian Journal of Psychiatry* **54**, 824–833.
- ten Have M, van Dorsselaer S, de Graaf R** (2013). Prevalence and risk factors for first onset of suicidal behaviors in the Netherlands Mental Health Survey and Incidence Study-2. *Journal of Affective Disorders* **147**, 205–211.
- Thong JY, Su AH, Chan YH, Chia BH** (2008). Suicide in psychiatric patients: case-control study in Singapore. *Australian & New Zealand Journal of Psychiatry* **42**, 509–519.
- Thoresen S, Mehlum L** (2004). Risk factors for fatal accidents and suicides in peacekeepers: is there an overlap? *Military Medicine* **169**, 988–993.
- Turecki G** (2014). The molecular basis of suicide. *Nature Reviews Neuroscience* **15**, 802–816.
- van Heeringen K, Hengeveld MW** (2009). Suicidaal gedrag. In *Leerboek Psychiatrie* (ed. Hengeveld MW and van Balkom AJLM), pp. 493–501. De Tijdstroom: Utrecht, The Netherlands.
- van Hemert AM, Kerkhof AJ, de Keijser J, Verwey B, van Boven C, Hummelen JW, de Groot MH, Lucassen P, Meerdinkveldboom J, Steendam M, Stringer B, Verlinde AA, van de Glind G** (2012). *Multidisciplinaire richtlijn diagnostiek en behandeling van suïcidaal gedrag*. Tijdstroom: Utrecht, The Netherlands.
- Vuorilehto M, Valtonen HM, Melartin T, Sokero P, Suominen K, Isometsä ET** (2014). Method of assessment determines prevalence of suicidal ideation among patients with depression. *European Psychiatry* **29**, 338–344.
- Wells GA, Shea B, O'Connell D, Peterson J, Welch V, Losos M, Tugwell P** (n.d.). The Newcastle-Ottawa Scale (NOS) for assessing the quality of nonrandomized studies in meta-analysis. Retrieved 7 July 2015 from http://www.ohri.ca/programs/clinical_epidemiology/nosgen.pdf
- Werbelloff N, Dohrenwend BP, Levav I, Haklai Z, Yoffe R, Large M, Davidson M, Weiser M** (2015). Demographic, behavioural, and psychiatric risk factors for suicide: a 25-year longitudinal cohort study. *Crisis* epub ahead of print. DOI: <https://doi.org/10.1027/0227-5910/a000359>
- Wolfersdorf M, Klinkisch M, Franke C, Keller F, Wurst FM, Dobmeier M** (2003a). In-patient suicide – a comparison of psychiatric in-patient suicides v. a control group of inpatients from the same treatment era without suicide. *Psychiatrische Praxis* **30**, 14–20.
- Wolfersdorf M, Neher F, Arbeitsgemeinschaft Suizidalität und Psychiatrisches Krankenhaus** (2003b). [Schizophrenia and suicide – results of a control group comparison of schizophrenic inpatient suicides with schizophrenic inpatients without suicide]. *Psychiatrische Praxis* **30**, 272–278.
- Yigletu H, Tucker S, Harris M, Hatlevig J** (2004). Assessing suicide ideation: comparing self-report versus clinician report. *Journal of the American Psychiatric Nurses Association* **10**, 9–15.
- Zonda T** (2006). One-hundred cases of suicide in Budapest: a case-controlled psychological autopsy study. *Crisis* **27**, 125–129.