

OPEN

# Predation of young tortoises by ravens: the effect of habitat structure on tortoise detectability and abundance

Amalia Segura<sup>1\*</sup>, José Jimenez<sup>1</sup> & Pelayo Acevedo<sup>1,2</sup>

The predation of young tortoise is considered a major cause of mortality for many tortoise species. The predation by common ravens has been identified as being responsible for significant decreases in tortoise populations. Mediterranean spur-thighed tortoise hatchlings and juveniles in Maamora forest (Morocco) were studied in order to describe the size/age class predation of common ravens on young tortoises and infer the drivers of predation risk and population abundance. The results showed a high level of predation on young tortoises (<75 mm carapace length) attributed to ravens in areas with low vegetation cover, representing 100% of the cases of mortality (n = 147), but it was moderate in covered areas (n = 19), representing 12–27%. The population structure of living juveniles differed significantly between covered and uncovered areas, thus suggesting that raven predation might modify juvenile population structure. Finally, N-mixture models showed a positive relationship between (i) bare cover and tortoise detectability that is only evidenced when the plot is far from a perch and (ii) population abundance and shrub species-richness, being higher in uncovered areas. Our results improve the knowledge on predation and survival on this critical stage in life, which is crucial for the conservation of the Mediterranean spur-thighed tortoise.

Tortoise populations are characterised by high adult survival and low recruitment rates, which probably influence their demographic dynamics<sup>1</sup>. The lack of reliable information on recruitment complicates evaluations of the real importance of young individuals as regards population demography<sup>2</sup>. In this respect, threats that limit recruitment bouts may have an important regulatory effect on population dynamics<sup>3</sup>, especially when these threats continue over long periods of time (e.g.<sup>4</sup>).

Hatchlings and juveniles are more vulnerable than adults owing to their smaller size and higher susceptibility to environmental conditions, such as the temperature or rain, which affect their physiology, but also to vegetation cover, which influences thermoregulation, the availability of food and the risk of predation<sup>3,5,6</sup>. Indeed, the low temperature and high rains of winter<sup>2,7</sup> and the high temperature of summer<sup>8,9</sup> are considered some of the most common causes of mortality in hatchlings. The predation of young tortoises –their shell is soft or not sufficiently solid to protect them from predators– by mammals<sup>7,10,11</sup> and birds<sup>12–14</sup> is also considered a major cause of mortality in many tortoise species, which was particularly studied in North America<sup>7,10,12,15,16</sup>.

The common raven *Corvus corax* is considered to be one of the most relevant predators of tortoises and is responsible for 70–91% of the mortality of desert tortoises *Gopherus agassizii*<sup>4,13,17</sup>. Both single individuals and breeding pairs have been identified as responsible for significant decreases in tortoise populations, also in addition to modifying population structure by affecting juvenile size/age classes<sup>12,18,19</sup>. Indeed, its high numbers and opportunistic feeding habits have severe impacts on its target prey, whose population size may be reduced without affecting the condition of the predator, which will switch to another prey when this resource becomes scarce (e.g.<sup>4</sup>). But the prey species has mechanisms by which to reduce predation. Vegetation cover has been documented as a key determinant as regards reducing young tortoises' detectability, since it facilitates their camouflage<sup>20</sup>. Crypsis has anti-predatory benefits, and refuge areas, such as those areas dominated by vegetation cover, may reduce the risk of predation. It has also been documented that ravens' predation on tortoises differs

<sup>1</sup>Instituto de Investigación en Recursos Cinegéticos, IREC (CSIC-UCLM-JCCM). Ronda de Toledo, 12, 13071, Ciudad Real, Spain. <sup>2</sup>Escuela Técnica Superior de Ingenieros Agrónomos (UCLM), Ronda de Calatrava, 7, 13071, Ciudad Real, Spain. \*email: [amaliasegura@gmail.com](mailto:amaliasegura@gmail.com)

	A	B	C	D
<i>Vegetation</i>				
Bare ground cover	0	0 ± 0.2	17.9 ± 17.4	21.4 ± 13.8
Shrub cover	59.7 ± 21.1	52.0 ± 14.8	45.0 ± 23.0	28.2 ± 26.1
Shrub height	62.1 ± 17.7	71.1 ± 30.6	31.9 ± 27.3	19.8 ± 17.6
Shrub richness	2.0 ± 0.8	1.5 ± 0.7	0.8 ± 0.4	0.7 ± 0.7
<i>Predation risk</i>				
Perch distance	203.6 ± 35.4	510.8 ± 120.8	72.5 ± 36.3	89.2 ± 35.3
Nest distance	999.0 ± 84.5	1076.7 ± 131.1	184.9 ± 89.2	144.7 ± 37.7
<i>Raven occurrence</i>				
Single individuals	5	2	4	5
Breeding pairs	2	1	2	1
Groups	1(28)	1(5)	0	1(17)
Number of live tortoises	39	30	65	39
Number of dead tortoises	11	8	55	92
Raven predated tortoises	3	1	55	92

**Table 1.** Bare ground cover (%), shrub cover (%), shrub height (cm) and shrub species richness; raven perch and nest distance (m) in the four study sites, A, B, C and D (mean ± SD); maximum raven occurrence in the Mediterranean spur-thighed tortoise survey distributed by single individuals, breeding pairs and groups of >2 individuals (number of individuals); live (<100 mm carapace length, CL), dead (<100 mm CL) and raven predated tortoises (<75 mm CL) (in all of them, the number of individuals).

according to their spatial distribution of single individuals or breeding pairs. Tortoise predation is greater in adjacent human developments, which attract large numbers of single ravens, and in adjacent successful nests of breeding pairs throughout human developed and undeveloped areas<sup>4</sup>. Even the success of juvenile tortoise releases is compromised in the head-starting programs included in conservation strategies owing to the fact that certain ravens are attached to the predation of certain tortoise size classes<sup>17,21</sup>. It could, therefore, be hypothesized that the risk of raven predation on young tortoises might be high in areas with high dense tortoise populations and mediated by the overlaid effect of raven presence and vegetation cover.


Maamora forest, an anthropogenic cork oak forest located in northern Morocco, is considered to be close to the optimum niche –the core range– of the Mediterranean spur-thighed tortoise's distribution *Testudo graeca*<sup>22</sup> and one of the areas with the highest density populations documented to date<sup>23</sup>. However, the common raven has, over the last 40 years, increased its numbers and distribution worldwide, and this cork oak forest is no exception<sup>24</sup>. This increase in predators may be a threat to the Mediterranean spur-thighed tortoise, and especially to highly susceptible young individuals. Concretely densities of 0.8 breeding pairs km<sup>-2</sup> had been observed in the lustrum (A. Segura unpublished data). In this context, our specific objectives were to: (i) describe predation and size classes' preference of common ravens on young tortoises, and (ii) infer the main drivers of predation risk and abundance. These goals are relevant as regards improving knowledge on the predation and survival of the Mediterranean spur-thighed tortoise at this critical stage in its life and, therefore, the conservation of this threatened species.

## Results

**Raven predation on hatchlings and juveniles.** Twenty-two occurrences of ravens, not including groups (7, 3, 6 and 6, in A, B, C and D, respectively), were detected on our four study sites (Table 1). Three raven nests with active breeding pairs were found and two breeding pairs had between 1 and 2 chicks (Table 1).

One hundred and sixty-six dead young tortoises (<100 mm Carapace Length; hereafter CL) (11, 8, 55 and 92, in A, B, C and D, respectively) were detected during the surveys. Most of the dead juvenile tortoises found (88%) were located in the uncovered areas (areas C and D), where the main CL size category of dead tortoises corresponded to between 41 and 70 mm (Fig. 1). The cause of mortality in those areas was associated with predation by the common raven (100%, n = 147), where only < 75 mm CL tortoise carcasses were found with signs of raven predation under perch and nest trees. The ravens ate the hatchling and juvenile tortoises by pulling off their head and limbs (6%) or pecking holes through the carapace (60%) or plastron (34%). Indeed, 74 and 15 juvenile tortoises were predated by two active breeding pairs of ravens in spring 2018 (in areas D and C, respectively). Nevertheless, unknown causes of mortality dominated in the covered areas, and only 12–27% were related to raven predation. The threshold size above which young tortoises are safe from raven predation in both covered and uncovered areas was 75 mm CL. Dead young tortoises associated with raven predation differed significantly between covered and uncovered areas (A vs. C:  $X^2 = 4.07$ ,  $p < 0.05$ ,  $n = 66$ ; A vs. D:  $X^2 = 4.25$ ,  $p < 0.05$ ,  $n = 103$ ; B vs. C:  $X^2 = 5.05$ ,  $p < 0.05$ ,  $n = 63$ ; B vs. D:  $X^2 = 5.19$ ,  $p < 0.05$ ,  $n = 100$ ), but did not differ between either covered (A vs. B:  $X^2 = 0.08$ ,  $p = 0.77$ ,  $n = 19$ ) or uncovered areas (C vs. D:  $X^2 = 0$ ,  $p > 0.99$ ,  $n = 147$ ).

**Live hatchlings and juveniles.** One hundred and sixty-three live young tortoises (39, 30, 65, 39 in A, B, C and D, respectively) were found during the surveys. Figure 1 shows the size-class distribution on the four study sites. The juvenile structure of live tortoises did not differ between the covered areas (A vs. B;  $X^2 = 7.68$ ,  $p = 0.26$ ,  $n = 69$ ) but differed significantly between the uncovered areas (C vs. D;  $X^2 = 21.19$ ,  $p < 0.05$ ,  $n = 104$ ) and also



**Figure 1.** Distribution of carapace length (mm; CL) in the four study areas: covered (A,B) and uncovered (C,D). Data represent the frequencies of young tortoises: live shown in black, dead by raven predation in light grey and dead by any other reason in dark grey.

between the covered and the uncovered areas (C vs. A:  $X^2 = 16.06$ ,  $p < 0.05$ ,  $n = 104$ ; C vs. B:  $X^2 = 16.06$ ,  $p < 0.05$ ,  $n = 95$ ; D vs. A:  $X^2 = 36.89$ ,  $p < 0.05$ ,  $n = 78$ ; D vs. B:  $X^2 = 36.77$ ,  $p < 0.05$ ,  $n = 69$ ). The main differences concerned the CL size category of 81–100 mm, which represented 41% versus 16% in the covered and uncovered areas, respectively.

**Tortoise detectability and abundance: risk of predation by ravens.** According to previous results regarding dead animals found on our study sites, only those animals  $< 75$  mm CL were susceptible to predation in Maamora forest. Predation risk analysis was, therefore, restricted to these size classes: 140 detections (A: 27 tortoises in 11 occupied grids; B: 21 tortoises in 10 occupied grids; C: 57 tortoises in 13 occupied grids, and D: 35 tortoises in 7 occupied grids). It varied from a maximum of 3 to 5–11 tortoises per grid in covered and uncovered areas, respectively (Table 1).

In the N-mixture model, we used a negative binomial model (Table 2). The stepwise procedure carried out to select predictors explaining detectability and abundance processes is summarized in Table 2. The bootstrap  $p$ -values for the final model based on the SSE, Freeman-Tukey, and Chi-square statistics were 0.03, 0.00 and 0.01, respectively. The value of  $\hat{c}$  (ratio of observed/expected) was 1.59.

The final model included the interaction between bare ground cover and the distance to the raven perch in the detection process (Fig. 2), and the site and the number of shrub species in the abundance process (Fig. 3, Table 3). The detectability increased with the percentage of bare ground, but this effect was apparent only when the distance to a raven perch was greater. The abundance was higher in uncovered areas and in grids with high shrub species richness.


## Discussion

**Evidence of common raven predation on hatchlings and juveniles.** This study reveals the predation of common ravens on juveniles of Mediterranean spur-thighed tortoises in certain areas of Maamora forest. Both, single individuals and breeding ravens were observed killing, carrying away and consuming juvenile tortoises, their preference being for class sizes 40–70 mm CL. The selection by ravens of certain size classes of tortoises over others has been found in other Testudinidae populations too<sup>18,19</sup>. Despite some tortoises being paint-marked to identify them as recaptures, no evidence of any increased risk of predation was found for the recognisable fraction of the population (but see<sup>25</sup>); only in C uncovered area was one individual found dead, representing 2% of the marked individuals.

Bearing in mind that raven predation might vary between years and among individuals, and the fact that our study comprised only one year, our results still suggest that the predation of juvenile tortoise in the study area was higher when compared with that of populations of Mediterranean spur-thighed tortoises in southern Spain<sup>8</sup>. Indeed in those areas raven predation did not affect tortoise populations at all. Nevertheless, some similarities

Model specification	nPars	AIC	$\Delta AICc < 2$	AICwt	cltvWt
<i>(1) Latent abundance distribution</i>					
Negative binomial	3	541.51	0.00	1.00	1.00
Zero Inflated	3	584.01	42.50	0.00	1.00
Poisson	2	620.03	78.52	0.00	1.00
<i>(2) Covariates of abundance (<math>\gamma</math>) and detection (<math>p</math>)</i>					
$\gamma$ (Shrub richness + Site) $p$ (Bare ground cover*Perch distance)	10	527.82	0.00	0.98	0.98
$\gamma$ (Site) $p$ (Bare ground cover*Perch distance)	9	536.24	8.42	0.01	0.99
$\gamma$ (.) $p$ (Bare ground cover*Perch distance)	6	538.68	10.86	0.00	1.00
$\gamma$ (.) $p$ (.)	3	541.51	13.69	0.00	1.00


**Table 2.** Model selection of Mediterranean spur-thighed tortoise hatchlings and juveniles (<75 mm carapace length; see text for details): (1) Latent abundance distribution and (2) covariates of abundance and detection. Covariates considered: shrub richness; site, bare ground cover (%) and raven perch distance (m). Model selection based on Akaike's Information Criterion (AIC), number of parameters (nPars), the difference in AICc from the best fitted models ( $\Delta AICc < 2$ ), model weights (AICwt), and cumulative model weights (cltvWt).



**Figure 2.** Interaction between two continuous covariates: bare ground cover rate and distance to the nearest raven perch (D, in meters) over the probability of detection.

with other Testudinidae populations affected by raven predation were found<sup>12,16,26</sup>, although the ravens involved in predation in our study appeared to have a slightly lower size threshold above which juveniles are safe from predation (75 mm CL) than reported for other Testudinidae populations (85 and 100 mm CL; 27, 21 respectively). This might be associated with the length of time shell hardening takes in Mediterranean spur-thighed tortoises, which has been documented to limit the probability of predation by ravens (e.g.<sup>27</sup>), and this merits further studies.

Nevertheless, when comparisons of raven predation on tortoises were restricted to covered versus uncovered areas, in the former such occurrences were all anecdotal while high mortality rates were rigorously recorded in the latter. Certainly, we were surprised by the high amount of predation by one pair of breeding ravens, which preyed 74 tortoises of <75 mm CL in a single breeding season. Indeed, raven predation might be modifying juvenile tortoise population structure through altering recruitment (e.g.<sup>4</sup>). Further studies are required to disentangle the role played by predation in tortoise population structure within the Maamora forest.




**Figure 3.** Relationship between sampling sites (A, B, C and D; left graph) and the richness of shrub species (number of species; right graph) with the abundance of Mediterranean spur-thighed tortoise hatchlings and juveniles (<75 mm carapace length).

Process	Variable	Estimate	Standard Error	Z	P (> Z )
Abundance	Intercept	1.025	0.49	2.094	***
	Shrub richness	0.709	0.222	3.196	**
	Site B	-0.419	0.619	-0.677	ns
	Site C	2.697	0.564	4.78	***
	Site D	2.102	0.586	3.588	***
Detection	Intercept	-2.416	0.454	-5.32	***
	Bare ground cover	0.636	0.516	1.23	ns
	Perch distance	1.9	0.588	3.23	**
	Bare ground cover* Perch distance	1.961	0.78	2.52	**

**Table 3.** Summary and statistical parameters of the final N-mixture model parameterized to estimate Mediterranean spur-thighed tortoise hatchlings and juveniles (<75 mm carapace length) as regards both detectability and abundance (significance codes: ns no significant; \*\* < 0.01 and \*\*\* < 0.001).

**Predation risk by ravens, population size and structure of young tortoises.** In this study, tortoise detectability –associated with predation risk– was mediated by the interaction between predator presence and bare or low cover areas; with tortoise detectability increasing with greater areas of bare ground, mainly in locations far from perch trees (e.g. <sup>18,28</sup>). This might suggest that ravens could be modulating the behavioral response of young tortoises, e.g. they will reduce their activity in areas near perch trees in order to be less detectable by the ravens (e.g. <sup>29</sup>). In this respect, it is reasonable to assume that the survivorship of juveniles in areas with higher predation risk might be lower and, therefore, a lower juvenile population size will characterise populations in such areas. Conversely, we found a higher abundance of young tortoises in uncovered areas, where they suffered higher predation, but also in areas where there was high diversity of shrub species in both covered and uncovered areas (e.g. <sup>30</sup>). It is thus plausible to assume that many other factors involving juvenile physiological costs (e.g. <sup>29</sup>) or even female reproduction traits –the number of clutches, clutch size and recruitment success–<sup>3</sup>, might explain part of the variation found in the size of juvenile populations in covered and uncovered areas<sup>3</sup>. However, in higher risk predation areas, it might also be expected that juvenile population structure would be modified and show differences in size/age classes<sup>12,18,19</sup>. Indeed, the higher percentage of longer/older juveniles (76–100 mm) –which are not considered susceptible to predation– found in covered areas compared to in uncovered ones evidenced a threat to juvenile survivorship in the lower size/age classes in those areas where hatchlings are more detectable, e.g. uncovered areas, and whose effects, among others, might be mediated by vegetation cover (e.g. <sup>30</sup>).

Finally, hatchlings are challenging to study since they are rarely encountered in the field<sup>31,32</sup> and, as such, very low capture rates and practically nonexistent recapture rates of Mediterranean spur-thighed tortoise hatchlings have been reported in southern Spain<sup>33,34</sup>. Further studies are required to discover the role played by predation in



**Figure 4.** Location of the study area in Morocco and tortoise populations studied (sites A, B, C and D) is shown. Grey circles size is proportional to the maximum abundance of live young tortoises (<75 mm CL) observed per 1-ha-grid (1, 2–3, 4–5, 6–11 ranges). Small grey squares mark grids where no tortoises were found. Black stars represent the raven perch or nest locations.

the recruitment process, which will have later impacts on the size and structure of tortoise populations. This will also help to disentangling the possible physiological costs associated with predation risk.

## Material and Methods

**Study area.** The study was conducted in areas of low elevation (72–185 m a.s.l.) and sandy soil in Maamora forest (northwest Morocco; 34° 02' 54.19" N, 6° 27' 19.24" W). The climate is Mediterranean, with hot and dry summers, and the annual range of average rainfall is from 300 to 500 mm. Maamora forest is dominated by cork oak trees, *Quercus suber*, with scattered endemic wild pear, *Pyrus mamorensis*, wild olive *Olea europaea*, green olive *Phyllirea latifolia* and mastic *Pistacia lentiscus*, and a sparse understory represented by bush and shrub species, such as Mediterranean broom *Genista linifolia*, *Cytisus arboreus*, *Stauracanthus genistoides*, dwarf palm *Chamaerops humilis*, French lavender *Lavandula stoechas*, sage-leaved rockrose *Cistus salvifolius*, *Halimium halimifolium* and *Thymelaea lythroides*<sup>35</sup>.

The study specifically took place on four sites that were close together (separated by 3 km), but always ensuring that the tortoise populations' territories were separated by a sufficient distance for them to be considered as independent populations<sup>34,36</sup>, Fig. 4. Despite the fact that all the sampling sites were located on private land on which there has been no pet trade (>10 years protected) and that the undergrowth is well represented when compared with other sites in unprotected Maamora, the study sites differed as regards their vegetation cover. Two of them were characterized by high cover and the diversity of their shrub and herbs (hereafter denominated as covered areas, A and B), while the other two were characterized by a lower cover and diversity of shrubs and a high percentage of bare ground (hereafter denominated as uncovered areas, C and D; see Table 1 for further details). This experimental design allowed us to test for differences in tortoise detectability, and, therefore, infer predation risk, in relation to vegetation cover.

**Mediterranean spur-thighed tortoise and common ravens.** The study sites were surveyed during the 2018-breeding period, comprising the end of February until the end of May, covering an area of 15 ha in each of the covered areas and of 18 ha in each of the uncovered ones. Each of the four sites was intensively surveyed in order to detect young tortoises. This was done for four (uncovered areas) or five (covered areas) days by four trained people. The survey consisted of searching the entire territory for the occurrence of individual tortoises, but focusing on the detection of hatchlings and juveniles (<100 mm CL). The tortoises were recorded from 12 h until 16 h on foot and in adequate weather conditions (sunny days with temperatures of between 20 and 24 °C). Each recorded individual was georeferenced using a GPS and the CL was measured using a vernier

calliper (accuracy  $\pm 1$  mm). Size classes were used to characterize the tortoises' population structure in covered and uncovered areas<sup>24</sup>. The Chi-square test was used to assess any differences between covered and uncovered areas in terms of hatchling and juvenile size-classes structure, considering measurements of 30–40, 41–50, 51–60, 61–70, 71–80, 81–90 and 91–100 mm. The tortoises' carapaces were lightly marked with non-toxic paint so as to be able to identify any recaptured individuals. In addition, two additional days per area at the beginning and the end of the raven breeding season were carried out looking for any dead animals, which were collected and measured and, when possible, their cause of death was determined on the basis of a visual examination of the remains of the animal and the area in which it was found. These data were then used to determine the relevance of predation by ravens and to characterize the size-classes of the tortoises that are preyed by this species.

During the tortoise survey, the location of common raven perches and nests were registered with a GPS. For this purpose, we also surveyed a buffer area of 1.5 km around the sampling areas, which is slightly greater than the average distance between the ravens' territories<sup>4</sup>. The number of single individuals and breeding pairs of ravens was also recorded, along with their breeding status and their recruitment success (chicks or fledglings).

**Modelling tortoise detectability and abundance.** The recorded data were referred to 1-ha grids, which were our territorial unit for analytical purposes. The number of young tortoises susceptible to predation (according to our data  $< 70$  mm CL, see below; but see<sup>21,37</sup>) was, therefore, quantified (our response variable) for each grid and survey. In addition, each grid was characterized during the survey in relation to vegetation cover, and specifically to the shrub cover (%), shrub height (cm) and species richness, and the bare ground cover (%). In order to incorporate the effect of the common raven into species detectability, the mean distance to the nearest raven perch and nest (m; two variables) in each grid was also calculated.

We used N-mixture models to model the detectability and abundance of the hatchlings and juveniles that were, according to the count data, susceptible to predation, while we accounted for imperfect detection<sup>38</sup> using the unmarked package<sup>39</sup> in R<sup>40</sup>. Our assumption is that the detectability of young tortoises during the surveys can be considered a proxy of young individuals' detectability by the ravens. Hatchlings and juveniles are well camouflaged and difficult to follow in their environment. Their detectability is very low<sup>8,10,16,41</sup>, since it is linked to the central hours of the day when animals are active. We, therefore, assume that detectability by observers is a proxy of detectability by ravens in order to be able to explore the effect of different factors on predation risk. This does not mean that the observer is able to detect the same number of young tortoises, but that the higher the detectability for the observer, the higher detectability for the ravens.

We used Akaike's Information Criterion corrected for small sample sizes (AICc)<sup>41</sup> to select the most appropriate error distribution by comparing the performance of Poisson, zero-inflated Poisson and Negative Binomial models. The detection and abundance processes were then modelled sequentially. A forward stepwise procedure was used to identify the most relevant predictors explaining the detection process. The same procedure was subsequently followed in order to identify the predictors explaining the abundance process. AICc was also used to compare models in the stepwise procedures by following the  $\Delta\text{AICc} < 2$  rules<sup>42</sup>. N-mixture models require an integer value that specifies the upper bound used in the integration (K). In our study, this upper bound was set at  $K = 120$ , which is sufficiently large for it not to have an effect on the model results. We used the parametric bootstrap approach (1000 samples) in unmarked to attain p-values from the sums of squares, along with Chi-square and Freeman-Tukey fit statistics as a measure of the goodness of fit of the final model.

**Ethic statements.** Sampling of Mediterranean spur-thighed tortoises in Maamora forest was conducted under the authorization and following the protocols approved by Le Haut-Commissariat aux Eaux et Forêts et à la Lutte Contre la Désertification of Morocco (High Commission for Waters and Forests and the Fight against Desertification). The field surveys were done in accordance with the guidelines and regulations. There was non-invasive sampling.

### Data availability

The datasets generated during and/or analyzed during the current study are available from the corresponding author on reasonable request.

Received: 22 October 2019; Accepted: 22 January 2020;

Published online: 05 February 2020

### References

- Epperson, D. M. & Colleen, D. H. Nestling and hatchling ecology of Gopher tortoises (*Gopherus polyphemus*) in southern Mississippi. *J. Herpetol.* **37**, 315–324 (2003).
- Campbell, S. P., Steidl, R. J. & Zylstra, E. R. Recruitment of desert tortoises (*Gopherus agassizii* and *G. morafkai*): a synthesis of reproduction and first-year survival. *Herpetol. Conserv. Biol.* **10**, 583–591 (2015).
- Díaz-Paniagua, C., Keller, C. & Andreu, A. C. Long-term demographic fluctuations of the spur-thighed tortoise, *Testudo graeca* in SW Spain. *Ecography* **24**, 707–721 (2001).
- Kristan, W. B. & Boarman, W. I. Spatial pattern of risk of common raven predation on desert tortoises. *Ecology* **84**, 2432–2443 (2003).
- Ben Kaddour, K., Slimani, T., El Mouden, E. H., Lagarde, F. & Bonnet, X. Population structure, population density and individual catchability of *Testudo graeca* in the central Jbilet (Morocco). *Vie et Milieu* **56**, 49–54 (2006).
- Raymond, K. M. N. Determining habitat preferences of the juvenile *Gopher polyphemus* using spatially model vegetation on central Florida sandhill. Graduate theses and dissertations (2007).
- Butler, J. A. & Sowell, S. Survivorship and predation of hatchling and yearling gopher tortoises. *Gopherus polyphemus*. *Chelon. Conserv. Biol.* **1**, 173–180 (1996).
- Keller, C., Díaz-Paniagua, C. & Andreu, A. C. Survival rates and mortality causes of *Testudo graeca* hatchlings in southwestern Spain. *J. Herpetol.* **32**, 238–243 (1998).

9. Nafus, M. G., Esque, T. C., Averill-Murray, R. C., Nussear, K. E. & Swaisgood, R. R. Habitat drives dispersal and translocated juvenile desert tortoises. *J. Applied Ecol.* **54**, 430–438 (2017).
10. Tom, J. Microhabitats and use of burrows of Bolson tortoise hatchlings in Biology of North American tortoises (ed. Bury, R. B. & Germano, D. J.) 137–146 (National Biological Survey, 1994).
11. Smith, L. L., Steen, D. A., Conner, L. M. & Rutledge, J. C. Effects of predator exclusion on nest and hatchling survival in the gopher tortoise. *J. Wildl. Manage.* **77**, 352–358 (2013).
12. Wilson, D. S. Estimates for survival gopher tortoises. *Gopherus polyphemus*. *J. Herpetol.* **25**, 376–379 (1991).
13. Camp, R. J., Knight, R. L., Knight, H. A. L., Sherman, M. W. & Kawashima, J. Y. Food habits of nesting common ravens in the eastern Mojave Desert. *Southwest. Nat.* **38**, 163–165 (1993).
14. Barje, F., Slimani, T., El Mouden, E. H., Lagarde, F. & Bonnet, X. Shrewd shrikes and spiny shrubs: a calamity for hatchling Moorish tortoise. *Amphibia-Reptilia* **26**, 113–115 (2005).
15. Germano, D. J. Comparative life histories of North American tortoises in Biology of North American tortoises (ed. Bury, R. B. & Germano, D. J.) 175–185 (National Biological Survey, 1994).
16. Pike, A. Movement patterns, habitat use and growth hatchling tortoises. *Gopherus polyphemus*. *Copeia* **1**, 68–76 (2006).
17. Nagy, K. A., Hillard, L. S., Tuma, M. W. & Morafka, D. J. Head-started desert tortoises (*Gopherus agassizii*): movements, survivorship and mortality causes following their release. *Herpetol. Conserv. Biol.* **10**, 203–215 (2015).
18. Berry, K. H., Yee, J. L., Coble, A. A., Perry, W. M. & Shields, T. A. Multiple factors affect a population of Agassiz's desert tortoise (*Gopherus agassizii*) in the northwestern Mojave desert. *Herpetological Monographs* **27**, 87–109 (2013).
19. Boarman, W.I. Problems with management of a native predator on a threatened species: raven predation on desert tortoises in Proc. 15th Vertebrate Pest Conf. (ed. Borrecco, J.E., & Marsh R.E.). University of California, Davis (1992).
20. Nafus, M. G. *et al.* Hiding in plain sight: a study on camouflage and habitat selection in slow-moving desert herbivore. *Behav. Ecol.* **26**, 1389–1394 (2015).
21. Hazard, L. C. Post-release dispersal and predation of head-started juvenile desert tortoises (*Gopherus agassizii*): Effect of release site distance on homing behaviour. *Herpetol. Conserv. Biol.* **10**, 504–515 (2015).
22. Anadón, J. D. *et al.* Distribution of *Testudo graeca* in the western Mediterranean according to climatic factors. *Amphibia-Reptilia* **33**, 285–296 (2012).
23. Segura, A. & Acevedo, P. The importance of protected and unprotected areas for the Mediterranean spur-thighed tortoise demography in northwest Morocco. *Amphibia-Reptilia* **40**, 369–371 (2019).
24. Birdlife International. *Corvus corax* (amended version published in 2016) The IUCN Red List of threatened Species 2017: e. T22706068A113271893 (2017).
25. Calderon-Chalco, K. A. & Putman, B. J. The effect of paint marking on predation risk in western fence lizards: a test using clay models. *Herpetol. Conserv. Biol.* **14**, 80–90 (2019).
26. Berry, K. H., *et al.* Changes in desert tortoise populations at the desert tortoise research natural area between 1979 and 1985. Desert tortoise council proceedings of the 1986 symposia: 100–123 (1986).
27. Nagy, K. A., Tuma, M. W. & Hillard, L. S. Shell hardness measurement in juvenile desert tortoises *Gopherus agassizii*. *Herpetol. Review* **42**, 191–195 (2011).
28. Hansen, N. A., Sato, C. F., Michael, D. R., Lindenmayer, D. B. & Driscoll, D. A. Predation risk for reptiles is highest at remnant edges in agricultural landscapes. *J. Applied Ecol.* **56**, 31–43 (2019).
29. Martin, J. When hiding from predators is costly: optimization of refuge use in lizards. *Etologia* **9**, 9–13 (2001).
30. Schneider, M. F. Habitat loss, fragmentation and predator impact: spatial implications for prey conservation. *J. Applied Ecol.* **38**, 720–735 (2001).
31. Dunham, A. E., Morin, P. J. & Wilbur, H. R. Methods for the study of reptile populations in Biology of the reptilia vol 16. Defense and life history (ed. Gans C., & Huey R.B.) 331–385 (Alan Liss, 1988).
32. Morafka, D. J. Neonates: missing links in the life histories of North American tortoises in Biology of North American tortoises (ed. Bury, R. B. & Germano, D. J.) 161–173 (National Biological Survey, 1994).
33. Braza, F., Delibes, M. & Castroviejo, J. Estudio biométrico y biológico de la Tortuga mora (*Testudo graeca*) en la Reserva Biológica de Doñana, Huelva. *Doñana Acta Vertebr.* **3**, 15–41 (1981).
34. Andreu, A. C. Ecología y dinámica poblacional de la Tortuga mora, *Testudo graeca*, en Doñana. Tesis doctoral. Univ. Sevilla (1987).
35. Aafi, A. Etude de la diversité floristique de l'écosystème de chene-liège du foret de la Maamora. These, Institut. Agron. et Vétér. Hassan II, Rabat (2007).
36. Anadón, J. D. *et al.* Factors determining the distribution of the spur thighed tortoise *Testudo graeca* in south east Spain: a hierarchical approach. *Ecography* **29**, 339–346 (2006).
37. Boarman, W. I. Managing a subsidized predator population: reducing common raven predation on desert tortoises. *Environ. Manage.* **32**, 205–217 (2003).
38. Royle, J. A. *N*-mixture models for estimating population size from spatially replicated counts. *Biometrics* **60**, 108–115 (2004).
39. Fiske, I. & Chandler, R. Unmarked: An R package for fitting hierarchical models of wildlife occurrence and abundance. *J. Stat. Softw.* **43**, 1–23 (2011).
40. R Core Team. R: A language and environment for statistical computing. R. Foundation for Statistical Computing, Vienna, Austria, [www.R-project.org/](http://www.R-project.org/) (2019).
41. Hellgrem, E. C., Kazmaier, R. T., Ruthven, D. C. & Synatzske, D. R. Variation in tortoise life history: demography of *Gopherus berlandieri*. *Ecology* **81**, 1297–1310 (2000).
42. Burnham, K. P. & Anderson, D. R. Model Selection and Multimodel Inference: A Practical Information-Theoretic Approach (Springer Verlag, 2002).

## Acknowledgements

We are very grateful to Oscar Rodríguez, Hassan Belhajjamia, Absallam Belhajjamia and Bouhali Kaddouri for their field assistance. We truly appreciate their commitment to the surveys and their ability to detect hatchlings. We would like to thank HCEFLCD services for their guidance. We would also like to thank Greg Trollip and Jacob Mwanzia for their support and interest in wild species conservation. PA is supported by MINECO-UCLM through a “Ramón y Cajal” contract (RYC-2012–11970).

## Author contributions

A.S. and P.A. conceived the initial ideas, A.S. did the field surveys, A.S., J.J. and P.A. analysed the data and led the writing. All the authors discussed the findings.

## Competing interests

The authors declare no competing interests.



## Additional information

**Correspondence** and requests for materials should be addressed to A.S.

**Reprints and permissions information** is available at [www.nature.com/reprints](http://www.nature.com/reprints).

**Publisher's note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2020