



RESEARCH ARTICLE

**REVISED** Biometric and genetic differences in kelabau (*Osteochilus spp.*) as revealed using cytochrome c oxidase subunit 1 [version 3; peer review: 3 approved]

Nur Asiah <sup>1,2</sup>, Junianto Junianto<sup>1</sup>, Ayi Yustiati<sup>1</sup>, Sukendi Sukendi<sup>2</sup>, Melta Rini Fahmi<sup>3</sup>, Zainal A. Muchlisin <sup>4</sup>, Muhamad Kadapi<sup>5</sup>, Windarti Windarti<sup>6</sup>

<sup>1</sup>Department of Fisheries and Marine Science, Padjadjaran University, Sumedang, West Java, 45363 Sumedang, Indonesia

<sup>2</sup>Aquaculture Department, Riau University, Pekanbaru, Riau, 28293, Indonesia

<sup>3</sup>Aquaculture Department, Research Center for Ornamental Fish Culture, Depok, West Java, 16436, Indonesia

<sup>4</sup>Aquaculture Department, Syiah Kuala University, Banda Aceh, Aceh Darussalam, 23111, Indonesia

<sup>5</sup>Agronomy Department, Padjadjaran University, Sumedang, West Java, 45363, Indonesia

<sup>6</sup>Aquatic Resource Management Department, Riau University, Pekanbaru, Riau, 28293, Indonesia

**v3** First published: 12 Feb 2019, 8:177 (<https://doi.org/10.12688/f1000research.17319.1>)  
 Second version: 23 Jan 2020, 8:177 (<https://doi.org/10.12688/f1000research.17319.2>)  
 Latest published: 10 Mar 2020, 8:177 (<https://doi.org/10.12688/f1000research.17319.3>)

**Abstract**

**Background:** Kelabau (*Osteochilus spp.*) is a freshwater fish commonly found in the rivers of Riau, Indonesia. Researchers believe that these are *Osteochilus kelabau*; however, accurate taxonomic determination of these fish in Riau waters has not been made. The purpose of this study was to facilitate the identification of the kelabau based on its morphology and genetics using biometric and cytochrome c oxidase subunit 1 (*CO1*) analyses, respectively.

**Methods:** Fish samples were collected from the Siak, Kampar and Rokan rivers in Riau Province, Indonesia. The DNA of 90 fish was extracted from the caudal fins using a DNA extraction kit, after which it was amplified using primers Fish-F1 and Fish-R1. Sequencing was conducted by Applied Biosystems Macrogen Korea, and the DNA sequences were then edited and aligned using MEGA v. 7. All samples were BLAST-searched for identification using the National Center for Biotechnology Information and BOLD System. Phylogenetic trees were constructed, and the similarity index was calculated using accession numbers AP011385.1 and KC631202.1 in GenBank.

**Results:** Analysis of the consensus barcode sequence for 86 species revealed a high percentage of barcode matches (96%–97% in GenBank and 96.6%–96.76% in the BOLD System). The nucleotide distance between groups of kelabau from the different rivers based on the Kimura 2-parameter model gave the following results: 0.05% between groups from the Siak and Kampar rivers, 0.09% between those from the Siak and Rokan rivers and 0.05% between those from the Kampar and Rokan rivers. The nucleotide distance between the groups in the Siak (0.09%), Kampar (0.00%) and Rokan (0.10%) Rivers indicated that the kelabau in those

**Open Peer Review**

Reviewer Status

|   | Invited Reviewers |            |            |
|---|-------------------|------------|------------|
|   | 1                 | 2          | 3          |
| <b>version 3</b><br>(revision)<br>10 Mar 2020 |                   | <br>report |            |
| <b>version 2</b><br>(revision)<br>23 Jan 2020 |                   | <br>report | <br>report |
| <b>version 1</b><br>12 Feb 2019               | <br>report        | <br>report | <br>report |

- Indra Junaidi Zakaria** , Andalas University, Padang, Indonesia
- Ja'afar Nuhu Ja'afar** , Modibbo Adama University of Technology (MAUTECH), Yola, Nigeria  
Modibbo Adama University of Technology (MAUTECH), Yola, Nigeria
- Pallipuram Jayasankar**, ICAR-Central Institute of Freshwater Aquaculture, Bhubaneswar, India

rivers were related to each other.

**Conclusions:** Based on the results of the research data using *CO1* and biometric analyses, the kelabau were confirmed to be *O. melanopleurus*.

### Keywords

DNA barcoding, Kelabau Fish, Common Rivers of Riau, Population Structure

Any reports and responses or comments on the article can be found at the end of the article.

**Corresponding authors:** Melta Rini Fahmi ([m\\_rinif@yahoo.com](mailto:m_rinif@yahoo.com)), Windarti Windarti ([windarti.unri@gmail.com](mailto:windarti.unri@gmail.com))

**Author roles:** **Asiah N:** Data Curation, Formal Analysis, Investigation, Project Administration, Software, Writing – Original Draft Preparation, Writing – Review & Editing; **Junianto J:** Conceptualization, Supervision, Validation, Writing – Review & Editing; **Yustiati A:** Conceptualization, Methodology, Supervision, Validation, Writing – Review & Editing; **Sukendi S:** Conceptualization, Methodology, Supervision, Validation, Writing – Review & Editing; **Fahmi MR:** Conceptualization, Formal Analysis, Investigation, Methodology, Validation, Writing – Review & Editing; **Muchlisin ZA:** Validation, Writing – Review & Editing; **Kadapi M:** Methodology, Supervision, Validation; **Windarti W:** Writing – Review & Editing

**Competing interests:** No competing interests were disclosed.

**Grant information:** The study was supported by Doctor's Dissertation Research SP DIPA-042.061.401516/2018, Ministry of Research, Technology and Higher Education (MRTHE), the Republic of Indonesia; therefore, the authors thank MRTHE for the financial collaboration support with the Research Institution and Community Service of Riau University.

*The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.*

**Copyright:** © 2020 Asiah N *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

**How to cite this article:** Asiah N, Junianto J, Yustiati A *et al.* **Biometric and genetic differences in kelabau (*Osteochilus spp.*) as revealed using cytochrome c oxidase subunit 1 [version 3; peer review: 3 approved]** F1000Research 2020, 8:177 (<https://doi.org/10.12688/f1000research.17319.3>)

**First published:** 12 Feb 2019, 8:177 (<https://doi.org/10.12688/f1000research.17319.1>)

**REVISED Amendments from Version 2**

In this new version of the manuscript we have revised as follows:

1. We have added three citations in the discussion - "35,36" become "35 to 39".
2. In paragraph three in the discussion we have added reasons for limited variation and a discussion of data [Table 3](#).
3. Dr Windarti was added as a co-author for her work in improving the manuscript

**Any further responses from the reviewers can be found at the end of the article**

## Introduction

Kelabau are ancient fish belonging to genus *Osteochilus* of family Cyprinidae. Kelabau fish are distributed throughout Thailand, Vietnam, Peninsular Malaysia, Borneo and Sumatra<sup>1,2</sup>. In Sumatra Island in Indonesia, these fish is commonly found in the Siak, Kampar and Rokan rivers in Riau Province<sup>3,4</sup>.

According to local fishers in Riau, kelabau are divided into two types on the basis of morphology; although, there is no detailed information about these fish types. The local fishers said that the first type of kelabau fish is the fish found in this study. The morphological characteristic of this fish is brownish body, with brighter bottom, dark hazy blotches present above the pectoral fins, and the size ranges from 15 to 2,778.93 gr. The second type of fish is larger sized and yellowish color. However, during the study time, from 2017 to 2018, this type was not found. Thus, a study was needed to identify the species using morphological and molecular methods to determine these types in the Siak, Rokan and Kampar rivers. Identification of any species using morphological traits can be difficult and can lead to errors<sup>5</sup>; therefore, owing to morphological similarities among *Osteochilus* spp., molecular markers, such as DNA barcodes, are important to identify the fish species uses a specific sequence region (i.e. *cytochrome c oxidase subunit 1 (COI)*) to identify a species and is a technique that can identify taxonomic units as well as biodiversity for determining species of several organisms<sup>5-9</sup>. Unlike molecular phylogeny used to determine relationships among species, the purpose of DNA barcoding is to identify unknown or undetermined species into phylogeny<sup>10</sup>. The common mitochondrial (mt) DNA region used as a barcode in protists and animals comprises 600 bp. In addition, *COI* is one of the genetic markers used to identify insects, birds, primates and fish to species<sup>11-13</sup>. MtDNA *COI* is selected as a target in DNA barcoding because it is a highly conserved site. This method has advantages over the morphological identification approach in that it is fast, reliable and it can be used for all types of samples because it uses a single gene along with mutations in the nucleotides to acknowledge the taxonomic features of each species<sup>13</sup>.

The study on DNA barcoding for freshwater fish has been widely practiced in various countries, including Nigeria<sup>14</sup>, India<sup>15</sup>, Philippines<sup>16</sup>, Canada<sup>9</sup> and Indonesia<sup>5,17-19</sup>. The method has been successfully validated for the taxonomic status within

*Rasbora* in Lake Laut Tawar<sup>20</sup>; Anguillidae in Aceh waters<sup>21</sup>; Ornamental fish from Peat lands<sup>3</sup>; Channidae in Peninsular Malaysia, Sarawak, Sumatera, Borneo, Myanmar, Vietnam, India, Germany, Singapore and the United Kingdom<sup>22</sup> and Cichlidae in northeastern Nigeria<sup>23</sup>; therefore, it can be used to equally successfully validate the taxonomic unit of the kelabau using its morphology supported by molecular data. This information is crucial for designing a remedial course of action about the conservation strategy for this species in the Siak, Kampar and Rokan rivers in Riau Province, Indonesia.

## Methods

### Ethics

The study population was collected and sampled according to the guidelines on the use of living organisms for research from the Laboratory of the Faculty of Fisheries and Marine, Riau University, Indonesia<sup>24</sup>

### Sampling sites and collection

A total of 90 kelabau (30 fish from each river) were collected from the Siak, Kampar and Rokan rivers ([Figure 1](#)). Fish were caught using a gill net 3 m deep and 20 m long with a 12.7-cm mesh. The gill nets were installed in the river water close to the riverbank and remained for 24 h from 08:00 to 08:00 the following day. The fish collected were counted using hand-counter and cleaned using freshwater. A number of 50-mm caudal fin tissue samples were taken using a sterile scissors and preserved in ethanol, after which a photo of each fish sample was taken for documentation using a digital camera.

All samples were preserved in 3-kg sample bags which were labeled according to site location, date and serial number. Before preservation, the fish samples were injected with 10% formalin. The fish samples were then transported to the laboratory for further evaluation. The morphologies of the collected fish were identified up to species level using the identification book produced by the Indonesian Institute of Sciences ichthyology museum<sup>1,25</sup>. The fish morphologies observed were length, color, shape of scales, mouth shape, barbels, number of fins and special marks on the body.


### Biometrics

Biometric analyses were used to measure morphological characteristics in this study<sup>1</sup>. This tool is considered conventional for identifying organisms. Molecular identification using *COI* gene sequences has been supported for providing additional organism classification.

### DNA isolation and amplification

DNA was extracted using the spin-column method from the gSYNC DNA Extrusion Kit (Geneaid Catalogue No. GS 300, Taiwan). The extracted DNA was then transferred to a 1X Tris-borate ethylenediaminetetraacetic acid (TBE) solution with a 1.5% agarose gel and PEGreen gel dye (PEQLAB Biotechnologies GmbH, Erlangen, Germany)<sup>19</sup>. The quantity of DNA was visualized with the help of a GeneQuant Spectrophotometer by adding 78 µL nuclease-free water in a cuvette along with 2 µL DNA. The DNA was then amplified using the universal primer

### Research Location Map



**Figure 1.** Sample sites for *Osteochilus melanopleurus* from the Siak (N: 00°39'22.28" and E: 101°17'28.67"), Kampar (N: 00°22'13.64" and E: 101°54'11.97") and Rokan rivers (N: 01°22'33.65" and E: 100°58'26.76"), Riau Province, Indonesia.

Fish-F1(5'-TCA-ACC-AAC-CAC-AAA-GAC-ATT-GGC-AC-3') and Fish-R1 (5'-TAG-ACT-TCT-GGG-TGG-CCA-AAG-AAT-CA-3') with a target of 707 bp and 655 bp<sup>26</sup>, respectively. The amplification thermocycling conditions as follow: the PCR condition using pra PCR (94°C for 5 min), 35 cycles of denaturation (94°C for 30 s), annealing (56.6°C for 30 s) and extension (72°C for 30 s), followed by post-PCR extension (72°C for 5 min) and hold (4°C for 5 min). PCR results were analyzed using 1.5% agarose gel at 100 V to assess the bands, and only the clear products were sent to Applied Biosystems Macrogen Korea for sequencing.

#### Controlling molecular samples and sequence quality

The PCR amplicon was 707 bp, which implied that no sequence of DNA was derived from mtDNA nuclear mitochondrial DNA segments (NUMTs), because a NUMT barely reaches 600 bp<sup>9</sup>.

The selected *COI* sequences were entered into GenBank and the BOLD System databases to compare the alignment of nucleotide sequences and 99%–100% values with that with no insertions/deletions. All sequences were aligned using ClustalW with MEGA v.7<sup>27</sup>.

#### Data analysis

##### Blasting of *COI* by NCBI (GenBank) and BOLD System (online)

The entire nucleotide sequence obtained from the sequence chromatogram was assembled using DNA Baser Assembler, aligned and then analyzed using MEGA 7. It was further aligned (multiple alignments) using the reference NCBI GenBank accession numbers AP011385.1 and KC631202.1. Similarly, the percentages of *COI* sequences were blasted using NCBI Blast and BOLD Systems databases.

### Nucleotide variations

Nucleotide variations among samples were analyzed using dnaSP v.5<sup>28</sup>. The parameters of these calculations were haplotype number, variable site, parsimony site, haplotype diversity and nucleotide diversity.

### Phylogenetic tree

Phylogenetic trees were estimated using all samples from the three populations and calculated according to the Tamura-Nei model<sup>29</sup> using MEGA 7<sup>27</sup>.

### Nucleotide distance

The distance among the nucleotide bases of the mtDNA *COI*s was analyzed using the Kimura 2-parameter model<sup>30</sup>. The nucleotide distances between and within the populations were examined according to the model based on the similarity of frequencies and ratios of transition to transversion (Ti:Tv) using MEGA 7<sup>27</sup>.

## Results

### Morphological identification


The morphological traits of all kelabau used in this study matched those of *O. melanopleurus*. We used the important morphological traits to identify these fish according to Kottelat *et al.*<sup>1</sup>. The morphological characteristics measurement of *O. melanopleurus* showed that the fish have 16–19 branched dorsal rays, the number of scales was ranged from 10.5 to 12.5 in between dorsal origin and lateral line, the number of circum peduncular rows of scale was ranged from 20 to 24 and lips covered with folds and plicae and there was no hard tubule at the tip of the mouth (Figure 2a). This species has one pair of barbels at above and one pair at bottom, dark hazy blotches near above of the pectoral fins. The body is brownish, with the bottom brighter than the top and the type of steroid scales (Figure 2b). Raw biometric data are available on OSF<sup>31</sup>, as ~707 bp.

### Genetic analysis

A sequence amplified by Fish-F1 primer was successfully identified in 86 of 90 samples of mtDNA fish. The base length of the *COI* nucleotide obtained from the formulation process and electrophoresis (Figure 3)

Based on genetic analysis using the Tamura-Nei model, there was an unequal distribution of all nucleotides with the following frequencies: adenine (A), 26.73%; thymine (T), 30.44%; cytosine (C), 25.93% and guanine (G), 16.90% (Table 1). The rates ratio between transition and transversion was 10.257 purines and 1.915 pyrimidines, and the overall transition and transversion bias were  $R = 2.499$  based on Tamura-Nei model<sup>32</sup>. The pattern of nucleotide distribution of A, T, C and G was  $T > A > C > G$ , but the carps which cultivated in India have the following pattern of nucleotide distribution:  $T > C > A > G$ <sup>33</sup>.

The nucleotide distances between nucleotide bases within the groups indicated that the values of the nucleotide base sequences within the fish population were 0.0009, 0.0000 and



**Figure 2.** (a) Kelabau (*O. melanopleurus*) and (b) ctenoid scale of *O. melanopleurus*.

0.0010 in the Siak, Kampar and Rokan rivers, respectively. The evolutionary distance between the nucleotides of the groups had a comparative difference in the nucleotide sequences of 0.0005 for fish from the Siak and Kampar rivers, of 0.0009 in those from the Siak and Rokan rivers and of 0.0005 for those from the Rokan and Kampar rivers. Based on the nucleotide distance, the fish were identified as being from the same species (0.06%) (Table 2).

*COI* had 612 conserved sites (98%), 9 variable sites (1.45%), 4 informative parsimony (0.64%) and 5 singleton sites (0.81%). The highest nucleotide variation was in the Rokan river population ( $0.00100 \pm 0.00032$ ); whereas, the Kampar river population had no nucleotide variation based on DnaSP5 calculations (Table 3). Using the NCBI database with accession numbers AP011385.1 and KC631202.1, the DNA sequence of Kelabau was identified as belonging to *O. melanopleurus* with 96%–97% accuracy, query coverage of 99%–100% and an E-value of 0.0. While based on the BOLD System, the identity of all samples was 96.60%–96.93% accurate.

In the phylogenetic tree consisted of two major groups (Figure 4). The first group was *Cirrhinus moltonela* (Gen-Bank KC631192.1) and it was divided from *O. melanopleurus*. The second group was differentiated into two sub groups, *O. melanopleurus* from GenBank (AP011385.1 and C631202.1) and 86 fish samples from Kampar, Siak and Rokan rivers. The 86 samples have BLASTN similarity values with *O. melanopleurus* of 96%–97%.

## Discussion

Overall, the morphological traits and DNA barcoding showed that the majority of, if not all, kelabau fish in the three rivers at





**Figure 3.** DNA amplified using Fish-F1 and Fish-R1 primers; M= Marker 100 bp (Vivantis, Malaysia); 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209 and 210= Rokan River; 211= Kampar River.

**Table 1.** Maximum composite likelihood estimates of the pattern of nucleotida substitution.

|   | A            | T    | C           | G            |
|---|--------------|------|-------------|--------------|
| A | -            | 4.03 | 3.34        | <b>22.95</b> |
| T | 3.54         | -    | <b>6.57</b> | 2.24         |
| C | 3.54         | 7.72 | -           | 2.24         |
| G | <b>36.29</b> | 4.03 | 3.34        | -            |

i Bold: different transitional substitutions; italic: transversal substitutions.

**Table 2.** Nucleotide distances among the populations.

|                      | 1      | 2      | 3      | 4      |
|----------------------|--------|--------|--------|--------|
| Siak                 | -      |        |        |        |
| Kampar               | 0.0005 | -      |        |        |
| Rokan                | 0.0009 | 0.0005 | -      |        |
| GenBank (AP011385.1) | 0.0367 | 0.0366 | 0.0368 | -      |
| GenBank (KC631202.1) | 0.0351 | 0.0350 | 0.0351 | 0.0048 |

i Source: Kimura estimation of 1980: "Evolutionary Divergence over Sequence Pair Between Groups"<sup>30</sup>.

**Table 3.** Nucleotide variation in mtDNA CO1 of *Osteochilus melanopleurus* by DnaSP5.

| Sampling location | Number of sites | Number of sequences | Haplotype number | Variable site | Parsimony site | Haplotype diversity | Nucleotide diversity |
|-------------------|-----------------|---------------------|------------------|---------------|----------------|---------------------|----------------------|
| Siak              | 621             | 28                  | 6                | 6             | 2              | 0.439 ± 0.114       | 0.00090 ± 0.00029    |
| Kampar            | 621             | 30                  | 1                | 0             | 0              | 0                   | 0                    |
| Rokan             | 621             | 28                  | 6                | 6             | 2              | 0.437 ± 0.113       | 0.00100 ± 0.00032    |

i Source: Nei, 1987 for haplotype and nucleotide diversity<sup>34</sup>.


Riau Province were *O. melanopleurus*. there were 2 types of kelabau fish present in Riau. The first type is relatively small (15 gr to 2,778.93 gr) fish with dark colored dorsal and whitish ventral. This type of fish is used in this study. The second type of kelabau fish is larger, with yellowish color. This fish was not found during the study.

Environmental changes can cause fish death or migration to suitable habitats. Overfishing using both legal and illegal methods has also triggered the decline of certain species<sup>35-39</sup>. In addition, our results suggested that there was little nucleotide diversity among *O. melanopleurus* in the Siak, Rokan and Kampar rivers in Riau Province, particularly the fish in the Kampar river

The lack nucleotide diversity of *O. melanopleurus* from the three rivers was likely to be caused by limited opportunities for kelabau migration, so that the genetic exchanges with other populations are very small<sup>36</sup>; moreover, the lack nucleotide diversity is believed to be caused by inbreeding, and

overfishing<sup>5,40</sup>. Variation in nucleotide diversity of the fish from the sampling areas is different. In the Kampar River, the fish was sampled from the relatively narrow area and it may cause the lowness of the nucleotide diversity. The diversity was 0 and it means that the nucleotide of the fish samples was almost identical. The low nucleotide diversity may occur as the habitat of the fish is very suitable and the fish may not face environmental related problem that trigger any genetic changing<sup>41</sup>. However the nucleotide diversity of the fish from the Siak and Rokan River was higher than Kampar River. Siak and Rokan Rivers have relatively low water quality that was caused mainly by anthropogenic activities. Changing in water quality may trigger the fish to adapt with that environmental condition and it may cause changes in genetic variation. Freeland *et al.*<sup>41</sup> stated that fish population with high genetic variation may be able to face problems related to environmental changing

In addition, Kelabau from the Siak and Rokan rivers were designated as one sub-sub group in group two (Figure 4)



**Figure 4.** Phylogenetic tree of kelabau (*Osteochilus melanopleurus*) based on the neighbour-joining model.

because both rivers are geographically connected, allowing for hybridisation, whereas, there is inbreeding of these fish in the Kampar River, which causes a nucleotide diversity value of 0.0. Geographically, these three rivers (Siak, Kampar and Rokan rivers) have different environmental condition, that cause variations in morphology in fish (*O. melanopleurus*) based on truss morphometric<sup>42</sup>. Factors causing morphological variations

are environmental differences, habitat pollution levels, long-term isolation and crossbreeding in populations<sup>43,44</sup>. The same deductions were drawn from previous studies on *Desmopuntius pentazona* and *D. rombochelatus*, although the taxonomies of the two fish are different. Nevertheless, based on a genetic difference of only 0.4%, the two species were grouped into one cluster<sup>5</sup>. These are distributed throughout Asia, the Mekong

and Chao Praya river basins, Peninsular Malaysia, Sumatra and Borneo<sup>1,2</sup>.

The identity of a species was derived using the morphological identification method to distinguish between species or individuals<sup>45,46</sup>. Basically, the genetic identification of a species can be done using mtDNA *COI*, a more effective approach than using rRNA<sup>5,33</sup>. The nucleotide locus and mutations were used as references to conduct DNA barcoding in all fish samples<sup>13</sup>. Previous studies have identified several species using DNA bar-coding, such as ornamental fish of wetlands<sup>5</sup>, wetland fish larvae<sup>46</sup>, rainbow fish<sup>47</sup>, Cyprinidae fish<sup>48</sup>, salmon and trout<sup>7</sup> and freshwater fish<sup>9,19</sup>. Furthermore, the phylogenetics of *COI* sequences can effectively show congeneric and confamilial species<sup>9</sup>.

The phylogenetic trees could describe the line of biological evolution from species or organisms with a different ancestry<sup>49</sup>. Nonetheless, the results of all these species did not show a 100% undistinguishable identity. The branch length between species leading to a gap in the pairwise distance distribution is referred to as the barcoding gap in *COI*<sup>23</sup>. Intra-species relationships were quite high in all samples, which confirmed that kelabau (*melanopleurus*) were native in the three rivers and could to adapt to changes in environmental conditions<sup>40</sup>.

Moreover, the existence of inter-nucleotide patterns and distances between A, T, C and G in the chromosomes showed the characteristics and genetic signs that distinguished each of the individuals, even though they belong to the same species<sup>50</sup>. This is reinforced by referencing the phylogenetic tree made using the neighbor-joining model.

The identification of fish species is normally conducted using morphological characteristics such as dorsal fins, pelvic fins, pectoral fins, anal fins, linea lateral fins, upper linea lateral fins, lower linea lateral fins, around body fins, and caudal peduncle fins; however, in this study, we used 12 morphological traits as described by the classification system of Kottelat *et al.*<sup>1</sup>. These results supported the classification using biometric data that all fish in the three rivers were *O. melanopleurus*. The morphological characteristics were consistent with the species having a relatively large body with a standard length of 119–560 mm, lips covered with folds and plicae, no tubercles on the snout, a pair of maxillary barbels, and a pair of lower jaw barbels. The body is brownish, with the bottom brighter than the top. Dark hazy blotches near above of the pectoral fins, which is a special trait of *O. melanopleurus*.

However, this method can be difficult, and molecular identification is necessary. In particular, using mtDNA *COI* was

an effective approach<sup>9,17</sup>. The results from nucleotide distance data based on the Kimura 2-parameter model indicated that the nucleotide distance among the fish was short in intraspecific species using mtDNA *COI*<sup>40</sup>, which was supported by data showing that the percentage identity in *O. melanopleurus* species ranged between 96% and 97%. The Kelabau fish from the three sample sites were identified as *O. melanopleurus* by percentage identity, supported by an E-value of 0.0 and a 99%–100% query cover. The p-value indicated that the BLASTN results contained no errors. Besides, the low nucleotide distance values (<3%–5%) among the samples of *O. melanopleurus* from the Siak, Kampar and Rokan rivers, indicated that all samples were monophyletic.

## Conclusion

Based on our findings, we concluded that 86 of the 90 samples of kelabau from the Siak, Kampar and Rokan rivers in Riau were *O. melanopleurus*, as revealed by their morphological traits and the molecular analyses.

## Data availability

### Underlying data

*COI* gene sequences and raw biometric data of *Osteochilus melanopleurus* from Riau rivers can be found on OSF.

*COI* gene sequence DOI: <https://doi.org/10.17605/OSF.IO/XGEZD><sup>51</sup>

Raw biometric data DOI: <https://doi.org/10.17605/OSF.IO/CFGM8><sup>31</sup>.

Data are available under the terms of the [Creative Commons Zero “No rights reserved” data waiver](#) (CC0 1.0 Public domain dedication).

The raw *COI* sequence data were also deposited in GenBank and can be found under sequential accession numbers [MH430827-MH430854](#) and [MH459085-MH459142](#).

## Acknowledgements

The authors thank Neli Safrina as a staff Laboratory of Fisheries and Marine Faculty, Riau University and Gema Wahyu Dewantoro, as a staff researcher at the Laboratory of Ichthyology in the Field of Zoology, Research Center for Biology of the Indonesian Institute of Sciences (LIPI) on the identification of the Kelabau fish using morphological traits. Also, thanks to Ornamental Fish Aquaculture Research and Development Center, Ministry of Marine and Fisheries Republic of Indonesia, Depok for providing laboratory facilities.



## References

- Kottelat M, Whitten AJ, Kartikasari SN, *et al.*: **Ikan air tawar Indonesia bagian Barat dan Sulawesi**. *Periplus E Jakarta Indonesia*. 1993.  
[Reference Source](#)
- Kottelat M: **The Fishes of the Inland Waters of Southeast Asia: a Catalogue and Core Bibliography of the Fishes Known To Occur in Freshwaters, Mangroves and Estuaries**. *Raffles Bull Zool*. 2013.  
[Reference Source](#)
- Asiah N, Juniarto J, Yustiati A, *et al.*: **Morfometrik dan Meristik Ikan Kelabau (*Osteochilus melanopleurus*) dari Sungai Kampar, Provinsi Riau**. *J Perikan dan Kelaut*. 2018; **23**(1): 47–56.  
[Reference Source](#)
- Pulungan CP: **Ikan-ikan air tawar dari sungai Ukui, anak sungai Siak, Riau**. *Terubuk*. 2011; **39**(1): 24–32.  
[Reference Source](#)
- Fahmi MR, Prasetyo AB, Kusumah RV, *et al.*: **Barcoding DNA ikan Hias Lahan Gambut**. *Aquac Res J*. 2016; **11**(13): 137–145.  
[Publisher Full Text](#)
- Hajibabaei M, Singer GA, Hebert PD, *et al.*: **DNA barcoding: how it complements taxonomy, molecular phylogenetics and population genetics**. *Trends Genet*. 2007; **23**(4): 167–172.  
[PubMed Abstract](#) | [Publisher Full Text](#)
- Rasmussen RS, Morrissey MT, Hebert PD: **DNA barcoding of commercially important salmon and trout species (*Oncorhynchus* and *Salmo*) from North America**. *J Agric Food Chem*. 2009; **57**(18): 8379–8385.  
[PubMed Abstract](#) | [Publisher Full Text](#)
- Taylor HR, Harris WE: **An emergent science on the brink of irrelevance: a review of the past 8 years of DNA barcoding**. *Mol Ecol Resour*. 2012; **12**(3): 377–388.  
[PubMed Abstract](#) | [Publisher Full Text](#)
- Hubert N, Hanner R, Holm E, *et al.*: **Identifying Canadian freshwater fishes through DNA barcodes**. *PLoS One*. 2008; **3**(6): e2490.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Kress WJ, Wurdack KJ, Zimmer EA, *et al.*: **Use of DNA barcodes to identify flowering plants**. *Proc Natl Acad Sci U S A*. 2005; **102**(23): 8369–8374.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Dawnay N, Ogden R, McEwing R, *et al.*: **Validation of the barcoding gene COI for use in forensic genetic species identification**. *Forensic Sci Int*. 2007; **173**(1): 1–6.  
[PubMed Abstract](#) | [Publisher Full Text](#)
- Jeong TJ, Jun J, Han S, *et al.*: **DNA barcode reference data for the Korean herpetofauna and their applications**. *Mol Ecol Resour*. 2013; **13**(6): 1019–32.  
[PubMed Abstract](#) | [Publisher Full Text](#)
- Hebert PD, Cywinska A, Ball SL, *et al.*: **Biological identifications through DNA barcodes**. *Proc Biol Sci*. 2003; **270**(1512): 313–321.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Falade MO, Opene AJ, Benson O: **DNA barcoding of *Clarias gariepinus*, *Coptodon zillii* and *Sarotherodon melanothron* from Southwestern Nigeria [version 1; referees: 2 approved]**. *F1000Res*. 2016; **5**: 1268.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Bright AR, Sathyan N, Antony SP, *et al.*: **Phylogeny and genetic divergence of three phenotypic variants of the ornamental goldfish, *Carassius auratus* (Linnaeus, 1758) based on CO1 gene**. *Int J Res Zool*. 2013; **3**(1): 4–9.  
[Reference Source](#)
- Guino RS II, Quilang JP, Geraldino JPL, *et al.*: **DNA Barcoding of Philippine Mullet (Family Mugilidae: Osteochthyidae)**. 2017; 1–23.  
[Reference Source](#)
- Fahmi MR, Kusumah RV, Ardi I, *et al.*: **DNA Barcoding Ikan Hias Introduksi**. *J Ris Akuakultur*. 2017; **12**(13): 29–40.  
[Publisher Full Text](#)
- Muchlisin ZA, Thomy Z, Fadli N, *et al.*: **DNA Barcoding of Freshwater Fishes from Lake Laut Tawar, Aceh Province, Indonesia**. *Acta Ichthyol Piscat*. 2013; **43**(1): 21–29.  
[Publisher Full Text](#)
- Rosnaeni A, Elfidasari D, Fahmi MR: **Dna Barcodes of the Pleco (Loricariidae, Pterygoplichthys) in the Ciliwung River**. *Int J Adv Res*. 2014; **5**(2): 33–45.  
[Publisher Full Text](#)
- Muchlisin ZA, Fadli N, Siti-Azizah MN: **Genetic variation and taxonomy of Rasbora group (Cyprinidae) from Lake Laut Tawar, Indonesia**. *J Ichthyol*. 2012; **52**(4): 284–290.  
[Publisher Full Text](#)
- Muchlisin ZA, Batubara AS, Fadli N, *et al.*: **Assessing the species composition of tropical eels (Anguillidae) in Aceh Waters, Indonesia, with DNA barcoding gene *cox1***. [version 1; referees: 1 approved, 2 approved with reservations]. *F1000Res*. 2017; **6**: 258.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Conte-Grand C, Britz R, Dahanukar N, *et al.*: **Barcoding snakeheads (Teleostei, Channidae) revisited: Discovering greater species diversity and resolving perpetuated taxonomic confusions**. *PLoS One*. 2017; **12**(9): e0184017.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Sogbesan OA, Sanda MK, Ja'afar JN, *et al.*: **DNA Barcoding of Tilapia Species (Pisces: Cichlidae) from North-Eastern Nigeria**. *J Biotechnol Biomater*. 2017; **7**(4): 1–4.  
[Publisher Full Text](#)
- Windarti: **Keterampilan Dasar Laboratorium Biologi Perikanan**. Unri Pres, 2018.  
[Reference Source](#)
- Robert TR: **The Freshwater Fishes of Western Borneo (Kalimantan Barat, Indonesia)**. *California Academic of Sciences*. San Francisco. 1989; 210.  
[Reference Source](#)
- Ward RD, Zemlak TS, Innes BH, *et al.*: **DNA barcoding Australia's fish species**. *Philos Trans R Soc Lond B Biol Sci*. 2005; **360**(1462): 1847–1857.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Kumar S, Stecher G, Tamura K: **MEGA7: Molecular Evolutionary Genetics Analysis Version 7.0 for Bigger Datasets**. *Mol Biol Evol*. 2016; **33**(7): 1870–1874.  
[PubMed Abstract](#) | [Publisher Full Text](#)
- Librado P, Rozas J: **DnaSP v5: a software for comprehensive analysis of DNA polymorphism data**. *Bioinformatics*. 2009; **25**(11): 1451–1452.  
[PubMed Abstract](#) | [Publisher Full Text](#)
- Saitou N, Nei M: **The neighbor-joining method: a new method for reconstructing phylogenetic trees**. *Mol Biol Evol*. 1987; **4**(4): 406–425.  
[PubMed Abstract](#) | [Publisher Full Text](#)
- Kimura M: **A simple method for estimating evolutionary rates of base substitutions through comparative studies of nucleotide sequences**. *J Mol Evol*. 1980; **16**(2): 111–120.  
[PubMed Abstract](#) | [Publisher Full Text](#)
- Asiah N: **Biometric of Kelabau Fish in Riau**. 2019.  
<http://www.doi.org/10.17605/OSF.IO/CFGM8>
- Tamura K, Nei M, Kumar S: **Prospects for inferring very large phylogenies by using the neighbor-joining method**. *Proc Natl Acad Sci U S A*. 2004; **101**(30): 11030–5.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Mohanty M, Jayasankar P, Sahoo L, *et al.*: **A comparative study of COI and 16 S rRNA genes for DNA barcoding of cultivable carps in India**. *Mitochondrial DNA*. 2013; **26**(1): 1–9.  
[PubMed Abstract](#) | [Publisher Full Text](#)
- Nei M: **Analysis of gene diversity in subdivided populations**. *Proc Natl Acad Sci U S A*. 1973; **70**(12): 3321–3.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Nasution S, Nuraini: **Grant of Feed Containing Vitamin E in Home Fish Kelabau (*Osteochilus kelabau*) to Improve Quality Eggs and Larvae**. *Int J Sci Eng Res*. 2014; **2**(4): 4–9.  
[Reference Source](#)
- Kusmini II, Mulyasari RGD: **Karakterisasi Genetik Ikan Kalabau (*Osteochilus kelabau*) dari Berbagai Lokasi di Kalimantan Barat Menggunakan Metode RAPD (Random Amplified Polymorphism DNA)**. *Ber Biol*. 2011; **10**(4): 449–454.  
[Reference Source](#)
- Fauzi M, Dahelmi D, Zakaria IJ, *et al.*: **Biological aspects of Ielan fish, *Diplocheilichthys pleurotaenia* (Cyprinidae) from the upstream and downstream of the Kampar River, Riau Province, Indonesia**. *AAAL Biolux*. 2016; **9**: 305–315.  
[Reference Source](#)
- Munir W, Tan UM, Zakaria IJ: **Developmental stages of endemic bilih fish larvae (*Mystacoleucus padangensis*) from Singkarak Lake, West Sumatra, Indonesia**. 2016; **9**: 965–975.  
[Reference Source](#)
- West L, Indonesia S, Zeswita AL, *et al.*: **Study Population Of Freshwater Shellfish *Corbicula sumatrana* In Singkarak Lake West Sumatra Indonesia**. *Res J Pharm Biol Chem Sci*. 2016; **7**(6): 1435–1441.  
[Reference Source](#)
- Frankham R, Ballou JD, Briscoe DA, *et al.*: **Introduction to conservation genetics**. 2002.  
[Reference Source](#)
- Freeland JR, Kirk H, Petersen SD: **Molecular Ecology 2nd ed**. West Sussex (GB): Wiley-Blackwell. 2011.  
[Publisher Full Text](#)
- Asiah N, Sukendi S, Juniarto J, *et al.*: **Trussmorphometric and meristic characters of kelabau fish (*Osteochilus melanopleurus* Bleeker, 1852) from three populations in Kampar, Siak, and Rokan Rivers, Riau Province**. *J Iktiologi Indones*. 2019; **19**(2): 283–295.  
[Publisher Full Text](#)
- Rawat S, Benakappa S, Kumar J, *et al.*: **Identification of fish stocks based on Truss Morphometric: A review**. *J Fish Life Sci*. 2017; **2**(1): 9–14.  
[Reference Source](#)
- Takács P, Vitál Z, Ferencz Á, *et al.*: **Repeatability, Reproducibility, Separative Power and Subjectivity of Different Fish Morphometric Analysis Methods**. *PLoS One*. 2016; **11**(6): e0157890.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Zhang H, Yu H, Gao T, *et al.*: **Analysis of genetic diversity and population**

- structure of *Pleuronectes yokohamae* indicated by AFLP markers. *Biochem Syst Ecol.* 2012; 44: 102–108.  
[Publisher Full Text](#)
46. Hubert N, Kadarusman AW, Busson F, *et al.*: **DNA barcoding Indonesian freshwater fishes: challenges and prospects.** *DNA Barcodes.* 2015; 3(1): 144–169.  
[Publisher Full Text](#)
47. Kadarusman, Hubert N, Hadiaty RK, *et al.*: **Cryptic diversity in Indo-Australian rainbowfishes revealed by DNA barcoding: implications for conservation in a biodiversity hotspot candidate.** *PLoS One.* 2012; 7(7): e40627.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
48. Collins RA, Armstrong KF, Meier R, *et al.*: **Barcoding and border biosecurity: identifying cyprinid fishes in the aquarium trade.** *PLoS One.* 2012; 7(1): e28381.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
49. Pramono TB, Artiaty D, Widodo MS, *et al.*: **Identifikasi Ikan Genus *Mystus* Dengan Pendekatan Genetik.** *J Sumberd Akuatik Indopasifik.* 2017; 1(2): 123–132.  
[Publisher Full Text](#)
50. Afreixo V, Bastos CA, Pinho AJ, *et al.*: **Genome analysis with inter-nucleotide distances.** *Bioinformatics.* 2009; 25(23): 3064–70.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
51. Asiah N: **raw data CO1 gene in Kelabau fish.** 2019.  
<http://www.doi.org/10.17605/OSF.IO/XGEZD>

# Open Peer Review

Current Peer Review Status:   

---

## Version 3

Reviewer Report 12 March 2020

<https://doi.org/10.5256/f1000research.24696.r61112>

© 2020 Ja'afar J. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



**Ja'afar Nuhu Ja'afar** 

<sup>1</sup> Department of Biotechnology, Modibbo Adama University of Technology (MAUTECH), Yola, Nigeria

<sup>2</sup> Chevron Biotechnology Center, Modibbo Adama University of Technology (MAUTECH), Yola, Nigeria

Please find my highlighted copy of the manuscript [here](#). The areas highlighted in the manuscript are supposed to portray the integrity and diligence of both the journal and reviewers, thus, maintaining scientific standards. Aside from this, the manuscript is ready to be indexed.

**Competing Interests:** No competing interests were disclosed.

**I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.**

---

## Version 2

Reviewer Report 10 February 2020

<https://doi.org/10.5256/f1000research.24129.r59013>

© 2020 Zakaria I. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



**Indra Junaidi Zakaria** 

Departement of Biology, Faculty of Math and Science, Andalas University, Padang, Indonesia

**Competing Interests:** No competing interests were disclosed.

**Reviewer Expertise:** Fish Biology and Fisheries biology

**I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.**

Author Response 08 Feb 2020

**Nur Asiah**, Padjadjaran University, Indonesia, Sumedang, Indonesia

Dear Mr. Indra Junaidi Zakaria,

Thank you for your suggestions on our manuscript. Through this letter I would like to explain the revision of my article titled "*Biometric and genetic differences in kalabau (Osteochillus spp) as revealed using Cytochrome C Oxidase subunit 1*". I have revised these following points:

- Grammatical and spelling mistakes throughout the paper have been revised.
- The introduction has been improved by adding information on the difference of the characteristics of two species of *kelabau* fish (*Osteochillus spp*). As there are 2 types of fish that are named as "kelabau fish" by local people, information on the difference of those fishes was obtained by interviewing local fishermen. The presence of this information may be useful to give more explanation about the difference and the similarity of *kelabau* fish present in Riau in general.
- A phrase "*The demand for it as ...*" until "*... the population of these as well as many other fish*" has been deleted as it is not correlate with the theme of this manuscript.
- Finally, the author has also added a quote in the following paragraph: "*The study population was collected and sampled ...*" until "*... of the Faculty of Fisheries and Marine Science, Riau University, Indonesia*".
- The discussion section has been improved by comparing the result of this study with more articles available.

Thank you for your suggestions and we hope that our article will be approved.

Best Regards,

Nur Asiah and the authors

**Competing Interests:** No competing interests were disclosed.

Reviewer Report 07 February 2020

<https://doi.org/10.5256/f1000research.24129.r59015>

© 2020 Jayasankar P. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



**Pallipuram Jayasankar**

Fish Genetics and Biotechnology Division, ICAR-Central Institute of Freshwater Aquaculture, Bhubaneswar, Odisha, India

I have gone through the revision/corrections effected by the authors in the manuscript "Biometric and genetic differences in kelabau (*Osteochilus* spp.) as revealed using cytochrome c oxidase subunit 1". The manuscript has been improved as per my suggestions. I recommend to consider its indexing.

**Competing Interests:** No competing interests were disclosed.

**I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.**

Author Response 08 Feb 2020

**Nur Asiah**, Padjadjaran University, Indonesia, Sumedang, Indonesia

Dear Mr. Pallipuram Jayasankar,

Thank you very much for your approved on our article, and thank you for your suggestions for the good of our article.

Best regards,

Nur Asiah and the authors

**Competing Interests:** No competing interests were disclosed.

Reviewer Report 03 February 2020

<https://doi.org/10.5256/f1000research.24129.r59014>

© 2020 Ja'afar J. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

? **Ja'afar Nuhu Ja'afar** 

<sup>1</sup> Department of Biotechnology, Modibbo Adama University of Technology (MAUTECH), Yola, Nigeria

<sup>2</sup> Chevron Biotechnology Center, Modibbo Adama University of Technology (MAUTECH), Yola, Nigeria

In version 1 of the manuscript, I have highlighted some sections that may require the authors' attention. However, none of those have been effected. Perhaps, the authors did not get the script I have corrected. It is available if needed.

**Competing Interests:** No competing interests were disclosed.

**Reviewer Expertise:** Molecular diagnostics

**I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.**



Author Response 08 Feb 2020

**Nur Asiah**, Padjadjaran University, Indonesia, Sumedang, Indonesia

Dear Mr. Ja'afar Nuhu Ja'afar,,

Thank you for your suggestions on our manuscript and we apologize for the late to response your suggestion because we need to discuss with other authors. Here we would like to inform that we already revised paragraph three by adding reasons for limited variation such as inbreeding within the species due to limited migration and we also added discussion about data on table 3.

Variation in nucleotide diversity of the fish from the sampling areas are different. In the Kampar River, the fish was sampled from the relatively narrow area and it may cause the lowness of the nucleotide diversity. The diversity was 0 and it means that the nucleotide of the fish samples was almost identical. Freeland *et al.*, (2011) stated that the low nucleotide diversity may occur as the habitat of the fish is very suitable and the fish may not face environmental related problem that trigger any genetic changing. However, the nucleotide diversity of the fish from the Siak and Rokan River, was higher than Kampar River. Siak and Rokan Rivers have relatively low water quality that was caused mainly by anthropogenic activities. Changing in water quality may trigger the fish to adapt with that environmental condition and it may cause changes in genetic variation. Freeland *et al.*, (2011) stated that fish population with high genetic variation may be able to face problems related to environmental changing.

Finally, we want to thank you very much for your suggestion on our manuscript and we hope that our manuscript will be approved.

Best regards,

Nur Asiah

**Competing Interests:** No competing interests were disclosed.

---

### Version 1

Reviewer Report 04 November 2019

<https://doi.org/10.5256/f1000research.18938.r55212>

© 2019 **Jayasankar P.** This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



**Pallipuram Jayasankar**

Fish Genetics and Biotechnology Division, ICAR-Central Institute of Freshwater Aquaculture, Bhubaneswar, Odisha, India

DNA barcoding is an important and useful tool to ratify species status in circumstances where morphology-based taxonomy fails or nearly-fails to perform unambiguous identification of species. The authors in the present study have taken up an important group of freshwater fish. There are many grammatical errors and some technical flaws in the manuscript. I felt one major drawback could be their total silence on the nominal species of *Osteochilus kelabu*. Of course using barcoding they have proved that the species is *O. melanopleurus*. However, they should have shown a phylogenetic comparison of these two nominal species to rule out confusion in identification. Morphological method followed does not appear to be robust. They are advised to use TRUSS MORPHOMETRICS which can be performed using user-friendly software nowadays. In the phylogenetic tree the majority of bootstrap values do not appear to be robust; hence it could reflect a spurious relationship.

**Is the work clearly and accurately presented and does it cite the current literature?**

Partly

**Is the study design appropriate and is the work technically sound?**

Partly

**Are sufficient details of methods and analysis provided to allow replication by others?**

Partly

**If applicable, is the statistical analysis and its interpretation appropriate?**

Yes

**Are all the source data underlying the results available to ensure full reproducibility?**

Partly

**Are the conclusions drawn adequately supported by the results?**

No

**Competing Interests:** No competing interests were disclosed.

**Reviewer Expertise:** I have been working on molecular taxonomy of fish, shellfish and marine mammals since 1996, thus 23 years of experience in this field; also gained expertise in freshwater aquaculture technology during the past 11 years.

**I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.**

Reviewer Report 04 October 2019

<https://doi.org/10.5256/f1000research.18938.r53842>

© 2019 Ja'afar J. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



**Ja'afar Nuhu Ja'afar** 

<sup>1</sup> Department of Biotechnology, Modibbo Adama University of Technology (MAUTECH), Yola, Nigeria

<sup>2</sup> Chevron Biotechnology Center, Modibbo Adama University of Technology (MAUTECH), Yola, Nigeria

Generally, the article needs to be proofread by an expert. There are a lot of areas I have **highlighted** to guide the authors.

In paragraph three of the discussion section, the authors should be more scientific in providing reasons for limited variation such as inbreeding within the species due to limited migration. More so, if COI is known to be conserved within a species, it is not expected to change within a small time frame of genetic event. In general, the discussion should be rewritten to explain the results and interpretations of the study.

In addition, I suggest the authors contact an expert that can explain/interpret the results obtained using DNASP. The explanation of the results from the software should be more than presenting the values.

The gel result will be more appreciable if the marker has been properly labelled.

**Is the work clearly and accurately presented and does it cite the current literature?**

Yes

**Is the study design appropriate and is the work technically sound?**

Partly

**Are sufficient details of methods and analysis provided to allow replication by others?**

Yes

**If applicable, is the statistical analysis and its interpretation appropriate?**

Partly

**Are all the source data underlying the results available to ensure full reproducibility?**

Yes

**Are the conclusions drawn adequately supported by the results?**

Yes

**Competing Interests:** No competing interests were disclosed.

**Reviewer Expertise:** Molecular diagnostics

**I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.**

Reviewer Report 04 March 2019

<https://doi.org/10.5256/f1000research.18938.r44418>

© 2019 Zakaria I. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



**Indra Junaidi Zakaria**

Departement of Biology, Faculty of Math and Science, Andalas University, Padang, Indonesia

#### Generally:

- There are still a lot of grammatical and spelling mistakes throughout the paper. Therefore, the authors should remove all of the grammatical and typographical errors as well as spelling mistakes. Use past tense and rephrase the sentences.

#### Introduction:

- Inconsistency in writing in the first paragraph, the first line: it is written as “genus” but in the second row it is written as “species”. *Kelabau* are ancient fish belonging to the genus *Osteochilus* of family Cyprinidae. The species are distributed throughout Thailand, Borneo and Sumatra (etc.).
- According to local fishers in Riau, *kelabau* are divided into two types on the basis of morphology. Although there is no detailed information about these fish types, please, the authors should explain what the differences in the characteristics are of the two species which the fishermen know.
- Sentences from: “The demand for it as...” until “...the population of these as well as many other fish.” I suggest deleting it, because it has no correlation with the theme of this manuscript which explains the verification of the species of fish *kelabau* using cytochrome c oxidase - it is not about increasing demand and the problem of decreasing populations of *kelabau* fish.

#### Methods:

- Cite references in the following paragraph: “The study population was collected and sampled...” until “...of the Faculty of Fisheries and Marine, Riau University, Indonesia.”

#### Results:

- Ok.

#### Discussion:

- The Discussion is not up to the mark, especially in paragraphs one and two; please rewrite back with an explanation of how the characteristics of the two species of fish differ according to fishermen and the reason that these differences were caused by environmental changes and overfishing.

**Is the work clearly and accurately presented and does it cite the current literature?**

Partly

**Is the study design appropriate and is the work technically sound?**

Yes

**Are sufficient details of methods and analysis provided to allow replication by others?**

Yes

**If applicable, is the statistical analysis and its interpretation appropriate?**

Yes

**Are all the source data underlying the results available to ensure full reproducibility?**

Yes

**Are the conclusions drawn adequately supported by the results?**

Yes

**Competing Interests:** No competing interests were disclosed.

**Reviewer Expertise:** Fish Biology and Fisheries biology

**I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.**

---

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact [research@f1000.com](mailto:research@f1000.com)

**F1000Research**