



Published in final edited form as:

Psychiatry Res. 2020 February ; 284: 112682. doi:10.1016/j.psychres.2019.112682.

Social anhedonia, social networks, and psychotic-like experiences: A test of social deafferentation

David Dodell-Feder^{a,b,*}, Bridget Shovestul^a, Jessica Woodyatt^a, Victoria Popov^b, Laura Germiné^{c,d}

^aDepartment of Psychology, University of Rochester, Rochester, NY, USA

^bDepartment of Neuroscience, University of Rochester Medical Center, Rochester, NY, USA

^cInstitute for Technology in Psychiatry, McLean Hospital, Belmont, MA, USA

^dDepartment of Psychiatry, Harvard Medical School, Boston, MA, USA

Abstract

Social connection is robustly associated with physical and mental health. So important is social connection that it features prominently in several etiological theories of serious psychopathology. Most notably, the social deafferentation hypothesis of schizophrenia posits that social anhedonia (SA) and isolation cause neural changes that produce psychosis. Here, we test several tenants of this theory by examining the relation between SA, psychotic-like experiences (PLE), and social networks. We find that SA and PLE are related to social networks, and that the relation between SA and PLE can be explained, in part, by the impact of SA on social networks.

Keywords

social deafferentation; social networks; schizophrenia

1. Introduction

Social connectedness carries profound consequences for our mental and physical health (Holt-Lunstad et al 2015; Kawachi, 2001; Yang et al., 2016). So consequential is our extent of social connection that a lack thereof has been proposed as an etiological factor in the development of psychopathology. Most notably, Hoffman (2007) describes a *social deafferentation hypothesis* of schizophrenia akin to neurological syndromes involving the loss of sensory input, and subsequent cortical reorganization that produces aberrant sensory experiences (e.g., phantom limb). According to this hypothesis, predisposing factors for

*Corresponding author: d.dodell-feder@rochester.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Declaration of interests

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

social withdrawal, such as social anhedonia (SA), lead to social isolation and a loss of social input, which curtails information to the social brain. Neural changes occur as a result, causing the brain to endogenously produce aberrant experiences and cognitions with social meaning (i.e., psychotic experiences). Beyond neural changes, social isolation may foster psychotic experiences by removing a buffer from stress (Nuechterlein and Dawson, 1984) and a source for reality-testing (Garety et al., 2001). Whatever the mechanism, a key factor in this model is a characteristic or experience (e.g., SA) that leads to diminished social networks.

Support for the social deafferentation hypothesis comes from several lines of work. People with psychotic disorders and psychotic-like experiences (PLE) have smaller social networks (Gayer-Anderson and Morgan, 2013) and fewer social interactions (Granholtm et al., 2019). Moreover, SA and accompanying withdrawal and isolation often precede and predict the onset of psychotic disorders (Gooding et al., 2005; Kwapil, 1998; Matheson et al., 2013; Tarbox and Pogue-Geile, 2008; Velthorst and Meijer, 2012; Wiles et al., 2006), proving themselves to be robust risk factors. Finally, among those diagnosed with a psychotic disorder, psychotic symptoms are more likely to occur when one is alone (Myin-Germeys et al., 2001).

If loss of social connectedness, by way of SA, contributes to PLE, it would bolster characterizations of social withdrawal and isolation as a public health crisis and calls for improving social connection as a public health priority (Holt-Lunstad et al., 2017). Thus, here, we evaluate the relations between SA, PLE, and social networks in a large international sample from the general population assuming a continuous, dose- response relation between these factors (van Os et al., 2009). Further, we conduct a formal test of a major component of the social deafferentation hypothesis by evaluating whether SA impacts PLE through its effect on social networks.

2. Methods

2.1. Participants

Participants were an international sample of 2,512 individuals who visited [TestMyBrain.org](https://www.testmybrain.org). Inclusion criteria were 18-65 years of age and fluent/native English- speaking. Participants were on average in their early 30s ($M=33.4\pm 13.1$), predominantly female (59.2%), White (68.7%), and from the United States (51.5%). Consent and study procedures were reviewed by the Harvard Committee for the Use of Humans Subjects.

2.2. Measures

Participants completed three self-report measures: the Social Network Index (SNI; Cohen, 1997), the short form of the Revised Social Anhedonia Scale (RSAS; Winterstein et al., 2011), and the Prodromal Questionnaire—Brief (PQB; Loewy et al., 2011). We selected these measures because they are widely used in their respective fields, and exhibit adequate psychometric properties (e.g., Loewy et al., 2011; Winterstein et al., 2011). Critically, elevated scores on the RSAS and PQB are associated with increased risk for psychosis-risk syndromes and psychotic-spectrum disorders (e.g., Kwapil, 1998; Savill et al., 2018). The

SNI assesses engagement in 12 different relationships from which three social network metrics are derived: number of people in one's social network (*number of people*), number of high-contact social roles in which the participant has regular contact (*network diversity*), and the number of social domains in which the participant is highly active (i.e., has at least 4 high-contact people in a given domain; *embedded networks*). The RSAS is a 15-item *true/false* questionnaire that assess social amotivation and lack of social pleasure ($\alpha=.83$). The PQB is a 21-item questionnaire that assess the frequency and associated distress of 21 positive symptoms of psychotic disorders. Participants who endorse an item on the frequency scale ($\alpha=.87$) are asked a follow-up question about distress using a 5-point Likert scale (*strongly disagree*—scored 1—to *strongly agree*—scored 5; $\alpha=.93$).

2.3. Analysis

Data were analyzed in R (R Core Team, 2018). We performed a 5% Winsorization of the three SNI metrics and PQB-Distress scores after identifying outlying scores ($>3 SD+M$) on these measures. Pearson r correlations are accompanied by 95% bias-corrected-and-accelerated (BCa) CIs generated from 10,000 bootstrap samples with the package *boot* (Canty and Ripley, 2017; Davison and Hinkley, 1997). Correlations are interpreted using the guidelines described in Gignac and Szodorai (2016). We used the package *psych* (Revelle, 2018) to conduct a multiple mediation analysis evaluating the effect of social anhedonia on psychotic-like experiences (separately for frequency and distress) through the three SNI metrics; that is, the indirect effect of RSAS on PQB through SNI-Number of People, SNI-Embedded Networks, and SNI-Network Diversity. We provide a bootstrapped estimate of the total indirect effect and specific indirect effects (RSAS→SNI→PQB) along with 95% BCa CIs derived from 10,000 bootstrap samples (Preacher and Hayes, 2008). As recommended by others (Miošević et al., 2018), we report fully standardized estimates which can be interpreted as the SDs change in PQB for every 1 SD increase in RSAS indirectly through SNI.

3. Results

First, we examined the relation between the measures of psychosis-proneness and social networks (Table 1). The association between SA and the social network metrics were large and similar across the SNI metrics. These correlations were larger than those observed for PLE and the social network metrics, which were relatively small, similar between the three social network metrics, and slightly higher for frequency versus distress. Thus, both SA and PLE are related to one's social network such that greater SA and PLE are associated with smaller and less diverse social networks, and fewer highly active social domains.

Next, we used mediation analysis to evaluate the hypothesis that, consistent with the social deafferentation hypothesis, SA would impact PLE through its deleterious impact on social networks. In line with this idea, for both frequency and distress, the bootstrapped CI of the total indirect effect did not contain zero indicating that social networks mediated the association between SA and PLE, with the mediation effect being slightly larger for frequency (Figure 1). Examination of the specific indirect effects revealed that for frequency,

only network diversity was a mediator; for distress, none of the individual indirect effects were different from zero.

4. Discussion

Here, we find that SA and PLE are related to smaller, less diverse, and less embedded social networks. These findings converge with those from other studies evaluating patient, at-risk, and healthy samples (Horan et al., 2006; Robustelli et al., 2017; Velthorst and Meijer, 2012). Further, in support of the social deafferentation hypothesis, we find that the relation between SA and PLE can be explained, in part, by the impact of SA on social networks. Interestingly, for at least the frequency of PLE, this effect was largely driven by network diversity; that is, the number of different social roles in which the participant has regular social contact. This finding converges with a number of brain imaging studies suggesting that network diversity—as opposed to the number of people in one’s network—may be particularly important for the structural and functional development of the social brain (Dziura and Thompson, 2014; Molesworth et al., 2015). Thus, the diversity of social roles one plays in their social network may be a better proxy for social support, social connectedness, or other factors that might protect against neural changes and concomitant PLE.

Beyond the inherent limitations of performing mediation analyses with cross-sectional data (Maxwell and Cole, 2007), we note that these data cannot be taken to demonstrate causal relations among the variables, including the hypothesized direction of causation (i.e., SA→social networks→PLE). Indeed, other research suggests reverse causation such that psychotic symptoms precipitate a social “network crisis” (Lipton et al., 1981) possibly by way of odd/eccentric behavior, paranoia, or social skills deficits that ultimately serve to isolate an individual. Conversely, other studies suggest that disturbed social networks and social withdrawal predate the onset of psychotic symptoms (Gayer-Anderson and Morgan, 2013; Gooding et al., 2005; Kwapil, 1998; Matheson et al., 2013; Tarbox and Pogue-Geile, 2008; Wiles et al., 2006). Of course, causation may be a dynamic, reciprocal process operating in both directions, such that social withdrawal contributes to the onset of symptoms, and the onset of symptoms (odd/eccentric behavior, paranoia, poor social skills) further negatively impacts social networks. The present findings cannot adjudicate between these possibilities; longitudinal data are needed. We also note that just under half of the participants (49.5%) were within the peak age range of risk for psychotic disorders (i.e., 29 years of age or younger; van der Werf et al., 2014) meaning that the majority of our participants would not be expected to develop a psychotic disorder. This also means that the interplay of SA, social networks, and PLE is still detectable in a non-high-risk, general population sample. Finally, the mediation effect was extremely small suggesting that processes related to PLE, but not explicitly tested here (e.g., social defeat; Selten et al., 2013) may be at work or that aspects of our study sample (e.g., a non-high-risk, general population sample) may have attenuated the relations among these variables.

Notwithstanding these considerations, the data from the current study add to a growing body of literature demonstrating the importance of social networks for psychological health. Together, these findings suggest that social isolation, and factors that contribute to social isolation, whether through deliberate social withdrawal as in SA or through exclusion and

social defeat (e.g., Selten et al., 2013) should be a public health priority (Holt-Lunstad et al., 2017).

Acknowledgments

We thank the [TestMyBrain.org](https://www.testmybrain.org) volunteers for their participation. This work was supported indirectly by a grant from National Institutes of Health to D.D.-F. (1L30MH117569-01).

References

- Canty A, Ripley B, 2017 boot: Bootstrap R (S-Plus) Functions. R package version 1.3–20.
- Cohen S, 1997 Social Ties and Susceptibility to the Common Cold. *JAMA J. Am. Med. Assoc* 277, 1940. [10.1001/jama.1997.03540480040036](https://doi.org/10.1001/jama.1997.03540480040036)
- Davison AC, Hinkley DV, 1997 Bootstrap methods and their application. Cambridge University Press, Cambridge; New York, NY, USA.
- Dziura SL, Thompson JC, 2014 Social-Network Complexity in Humans Is Associated With the Neural Response to Social Information. *Psychol. Sci* 25, 2095–2101. <https://doi.org/10.1177/0956797614549209> [PubMed: 25253279]
- Garety PA, Kuipers E, Fowler D, Freeman D, Bebbington PE, 2001 A cognitive model of the positive symptoms of psychosis. *Psychol. Med* 31 <https://doi.org/10.1017/S0033291701003312>
- Gayer-Anderson C, Morgan C, 2013 Social networks, support and early psychosis: a systematic review. *Epidemiol. Psychiatr. Sci* 22, 131–146. [10.1017/S2045796012000406](https://doi.org/10.1017/S2045796012000406) [PubMed: 22831843]
- Gignac GE, Szodorai ET, 2016 Effect size guidelines for individual differences researchers. *Personal. Individ. Differ* 102, 74–78. <https://doi.org/10.1016/j.paid.2016.06.069>
- Gooding DC, Tallent KA, Matts CW, 2005 Clinical Status of At-Risk Individuals 5 Years Later: Further Validation of the Psychometric High-Risk Strategy. *J. Abnorm. Psychol* 114, 170–175. <https://doi.org/10.1037/0021-843X.114.1.170> [PubMed: 15709824]
- Granhölm E, Holden JL, Mikhael T, Link PC, Swendsen J, Depp C, Moore RC, Harvey PD, 2019 What Do People With Schizophrenia Do All Day? Ecological Momentary Assessment of Real-World Functioning in Schizophrenia. *Schizophr. Bull*, 10.1093/schbul/sbz070
- Hoffman RE, 2007 A Social Deafferentation Hypothesis for Induction of Active Schizophrenia. *Schizophr. Bull* 33, 1066–1070. <https://doi.org/10.1093/schbul/sbm079> [PubMed: 17631618]
- Holt-Lunstad J, Robles TF, Sbarra DA, 2017 Advancing social connection as a public health priority in the United States. *Am. Psychol* 72, 517–530. <https://doi.org/10.1037/amp0000103> [PubMed: 28880099]
- Holt-Lunstad J, Smith TB, Baker M, Harris T, Stephenson D, 2015 Loneliness and Social Isolation as Risk Factors for Mortality: A Meta-Analytic Review. *Perspect. Psychol. Sci* 10, 227–237. [10.1177/1745691614568352](https://doi.org/10.1177/1745691614568352) [PubMed: 25910392]
- Horan WP, Subotnik KL, Snyder KS, Nuechterlein KH, 2006 Do Recent-Onset Schizophrenia Patients Experience a “Social Network Crisis”? *Psychiatry Interpers. Biol. Process* 69, 115–129. <https://doi.org/10.1521/psyc.2006.69.2.115>
- Kawachi I, 2001 Social Ties and Mental Health. *J. Urban Health Bull. N. Y. Acad. Med* 78, 458–467. <https://doi.org/10.1093/jurban/78.3.458>
- Kwapil TR, 1998 Social anhedonia as a predictor of the development of schizophrenia-spectrum disorders. *J. Abnorm. Psychol* 107, 558–565. [10.1037/0021-843X.107.4.558](https://doi.org/10.1037/0021-843X.107.4.558) [PubMed: 9830243]
- Lipton FR, Cohen CI, Fischer E, Katz SE, 1981 Schizophrenia: A Network Crisis. *Schizophr. Bull* 7, 144–151. <https://doi.org/10.1093/schbul/7.1.144> [PubMed: 7233102]
- Loewy RL, Pearson R, Vinogradov S, Bearden CE, Cannon TD, 2011 Psychosis risk screening with the Prodromal Questionnaire — Brief Version (PQ- B). *Schizophr. Res* 129, 42–46. <https://doi.org/10.1016/j.schres.2011.03.029> [PubMed: 21511440]
- Matheson SL, Vijayan H, Dickson H, Shepherd AM, Carr VJ, Laurens KR, 2013 Systematic meta-analysis of childhood social withdrawal in schizophrenia, and comparison with data from at-risk

- children aged 9–14 years. *J. Psychiatr. Res* 47, 1061–1068. <https://doi.org/10.1016/j.jpsychires.2013.03.013> [PubMed: 23628387]
- Maxwell SE, Cole DA, 2007 Bias in cross-sectional analyses of longitudinal mediation. *Psychol. Methods* 12, 23–44. [10.1037/1082-989X.12.1.23](https://doi.org/10.1037/1082-989X.12.1.23) [PubMed: 17402810]
- Mio evi M, O'Rourke HP, MacKinnon DP, Brown HC, 2018 Statistical properties of four effect-size measures for mediation models. *Behav. Res. Methods* 50, 285–301. [10.3758/s13428-017-0870-1](https://doi.org/10.3758/s13428-017-0870-1) [PubMed: 28342072]
- Molesworth T, Sheu LK, Cohen S, Gianaros PJ, Verstynen TD, 2015 Social network diversity and white matter microstructural integrity in humans. *Soc. Cogn. Affect. Neurosci* 10, 1169–1176. [10.1093/scan/nsv001](https://doi.org/10.1093/scan/nsv001) [PubMed: 25605966]
- Myin-Germeys I, Nicolson NA, Delespaul PA, 2001 The context of delusional experiences in the daily life of patients with schizophrenia. *Psychol. Med* 31, 489–498. <https://doi.org/10.1017/S0033291701003646> [PubMed: 11305857]
- Nuechterlein KH, Dawson ME, 1984 A Heuristic Vulnerability/Stress Model of Schizophrenic Episodes. *Schizophr. Bull* 10, 300–312. <https://doi.org/10.1093/schbul/10.2.300> [PubMed: 6729414]
- Preacher KJ, Hayes AF, 2008 Asymptotic and resampling strategies for assessing and comparing indirect effects in multiple mediator models. *Behav. Res. Methods* 40, 879–891. <https://doi.org/10.3758/BRM.40.3.879> [PubMed: 18697684]
- R Core Team, 2018 R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria.
- Revelle W, 2018 psych: Procedures for Personality and Psychological Research, Northwestern University, Evanston, Illinois, USA, <https://CRAN.R-project.org/package=psych>. Version = 1.8.12
- Robustelli BL, Newberry RE, Whisman MA, Mittal VA, 2017 Social relationships in young adults at ultra high risk for psychosis. *Psychiatry Res.* 247, 345–351. <https://doi.org/10.1016/j.psychres.2016.12.008> [PubMed: 27987484]
- Savill M, D'Ambrosio J, Cannon TD, Loewy RL, 2018 Psychosis risk screening in different populations using the Prodromal Questionnaire: A systematic review: SAVILL et al. *Early Interv. Psychiatry* 12, 3–14. [10.1111/eip.12446](https://doi.org/10.1111/eip.12446)
- Selten J-P, van der Ven E, Rutten BPF, Cantor-Graae E, 2013 The Social Defeat Hypothesis of Schizophrenia: An Update. *Schizophr. Bull* 39, 1180–1186. <https://doi.org/10.1093/schbul/sbt134> [PubMed: 24062592]
- Tarbox SI, Pogue-Geile MF, 2008 Development of social functioning in preschizophrenia children and adolescents: A systematic review. *Psychol. Bull* 134, 561–583. <https://doi.org/10.1037/0033-2909.134.4.561> [PubMed: 18605820]
- van der Werf M, Hanssen M, Kohler S, Verkaaik M, Verhey FR, RISE Investigators, van Winkel R, van Os J, Allardyce J, 2014 Systematic review and collaborative recalculation of 133 693 incident cases of schizophrenia. *Psychol. Med* 44, 9–16. [10.1017/S0033291712002796](https://doi.org/10.1017/S0033291712002796) [PubMed: 23244442]
- van Os J, Linscott RJ, Myin-Germeys I, Delespaul P, Krabbendam L, 2009 A systematic review and meta-analysis of the psychosis continuum: evidence for a psychosis proneness-persistence-impairment model of psychotic disorder. *Psychol. Med* 39, 179–195. [10.1017/S0033291708003814](https://doi.org/10.1017/S0033291708003814) [PubMed: 18606047]
- Velthorst E, Meijer C, 2012 The association between social anhedonia, withdrawal and psychotic experiences in general and high-risk populations. *Schizophr. Res* 138, 290–294. <https://doi.org/10.1016/j.schres.2012.03.022> [PubMed: 22484023]
- Wiles NJ, Zammit S, Bebbington P, Singleton N, Meltzer H, Lewis G, 2006 Self-reported psychotic symptoms in the general population: Results from the longitudinal study of the British National Psychiatric Morbidity Survey. *Br. J. Psychiatry* 188, 519–526. [10.1192/bjp.bp.105.012179](https://doi.org/10.1192/bjp.bp.105.012179) [PubMed: 16738341]
- Winterstein BP, Silvia PJ, Kwapił TR, Kaufman JC, Reiter-Palmon R, Wigert B, 2011 Brief assessment of schizotypy: Developing short forms of the Wisconsin Schizotypy Scales. *Personal. Individ. Differ* 51, 920–924. <https://doi.org/10.1016/j.paid.2011.07.027>

Yang YC, Boen C, Gerken K, Li T, Schorpp K, Harris KM, 2016 Social relationships and physiological determinants of longevity across the human life span. *Proc. Natl. Acad. Sci* 113, 578–583. 10.1073/pnas.1511085112 [PubMed: 26729882]

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Highlights

- Social anhedonia and psychotic-like experiences are related to social networks
- Social anhedonia may impact psychotic-like experiences through social networks
- Social anhedonia and isolation should be a public health priority




Figure 1. Results of the multiple mediation analysis. Values represent fully standardized point estimates of the indirect effect and their bias-corrected-and-accelerated 95% CIs derived from 10,000 bootstrap samples. The top panel (A) depicts the results for psychotic-like experiences (Prodromal Questionnaire—Brief) frequency; the bottom panel (B) depicts the results for psychotic-like experiences distress.

Table 1.

Descriptive statistics and measure correlations.

	<i>M</i>	<i>SD</i>	Min- Max	Pearson <i>r</i> [95% CI]*				
				2.	3.	4.	5.	6.
1. SNI-Network Diversity	4.6	2.0	0-12	.77 [.75, .78]	.69 [.67, .71]	-.18 [-.22, -.14]	-.14 [-.18, -.11]	-.35 [-.39, -.32]
2. SNI-Number of People	15.0	8.4	2-37		.89 [.88, .89]	-.18 [-.22, -.15]	-.15 [-.19, -.11]	-.38 [-.41, -.35]
3. SNI-Embedded Networks	1.6	1.2	0-5			-.17 [-.21, -.13]	-.12 [-.16, -.08]	-.37 [-.40, -.34]
4. PQB-Frequency	7.6	5.1	0-21				.78 [.76, .80]	.31 [.27, .34]
5. PQB-Distress	29.3	22.4	0-75					.27 [.23, .30]
6. RSAS	5.3	3.8	0-15					

* All correlations $p < .0001$.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript