

Review



Cite this article: Whitfield TT. 2019

Cilia in the developing zebrafish ear.

Phil. Trans. R. Soc. B **375**: 20190163.

<http://dx.doi.org/10.1098/rstb.2019.0163>

Accepted: 8 August 2019

One contribution of 17 to a Theo Murphy meeting issue 'Unity and diversity of cilia in locomotion and transport'.

Subject Areas:

developmental biology, cellular biology, genetics, neuroscience

Keywords:

zebrafish, otic vesicle, kinocilia, motile cilia, otolith, sensory hair cell

Author for correspondence:

Tanya T. Whitfield

e-mail: t.whitfield@sheffield.ac.uk

Cilia in the developing zebrafish ear

Tanya T. Whitfield

Department of Biomedical Science, University of Sheffield, Sheffield S10 2TN, UK

TTW, 0000-0003-1575-1504

The inner ear, which mediates the senses of hearing and balance, derives from a simple ectodermal vesicle in the vertebrate embryo. In the zebrafish, the otic placode and vesicle express a whole suite of genes required for ciliogenesis and ciliary motility. Every cell of the otic epithelium is ciliated at early stages; at least three different ciliary subtypes can be distinguished on the basis of length, motility, genetic requirements and function. In the early otic vesicle, most cilia are short and immotile. Long, immotile kinocilia on the first sensory hair cells tether the otoliths, biomineralized aggregates of calcium carbonate and protein. Small numbers of motile cilia at the poles of the otic vesicle contribute to the accuracy of otolith tethering, but neither the presence of cilia nor ciliary motility is absolutely required for this process. Instead, otolith tethering is dependent on the presence of hair cells and the function of the glycoprotein Otogelin. Otic cilia or ciliary proteins also mediate sensitivity to ototoxins and coordinate responses to extracellular signals. Other studies are beginning to unravel the role of ciliary proteins in cellular compartments other than the kinocilium, where they are important for the integrity and survival of the sensory hair cell.

This article is part of the Theo Murphy meeting issue 'Unity and diversity of cilia in locomotion and transport'.

1. Introduction

The cilia of vertebrate sensory organs are extraordinarily diverse in morphology and function, forming an integral part of cellular receptors for light, odorants, sound, gravity and motion. The zebrafish, *Danio rerio*, is a widely used vertebrate model system for development and disease, with particular advantages for combining imaging in the live animal with genetic or pharmacological manipulation of protein function. Cilia in the zebrafish retina, nose,¹ inner ear and lateral line have all been the subject of intense study; in this review, I focus on cilia in the inner ear, with occasional reference to the lateral line. From the earliest stages of its development, the inner ear is a major site of expression for genes coding for ciliary proteins. Analysis of the different ciliary types in the early otic vesicle has revealed some fascinating biology linking both immotile kinocilia and motile cilia-driven fluid flow with the formation of the otoliths. In the lateral line, a sensory system closely related to the inner ear, kinocilia of the sensory hair cells in superficial neuromasts are uniquely accessible for study. Both the ear and lateral line in zebrafish form important models for our understanding of hearing, balance and deafness in humans.

2. Three distinct types of cilia in the zebrafish otic vesicle

There are at least three distinct types of cilia in the zebrafish ear at early stages. From the onset of hollowing of the otic placode to form a vesicle at 18 h post fertilization (hpf), most cells of the otic epithelium bear a single immotile cilium on their apical (luminal) side [2]. Each sensory hair cell also bears a single cilium, the kinocilium, located on one side of the stereociliary bundle, an apical collection of microvillar-like projections that respond to mechanical deflection. In the early otic vesicle (20–25-somite stage), hair cells (tether cells) form in two pairs, one at each pole of the otic vesicle [3,4], and thus initially there are only four kinocilia among the 200 or so cilia in total in each ear (figure 1*a,b*). Despite




Figure 1. Cilia and otolith formation in the zebrafish ear. (a) Differential interference contrast image of the wild-type otic vesicle at the 25-somite stage, showing the two birefringent otoliths. Scale bar, 40 μm . (b) Lumen of the otic vesicle at the 27-somite stage, stained with an anti-acetylated Tubulin antibody. The arrowheads mark longer tether kinocilia and motile cilia at the anterior and posterior poles of the otic lumen. Scale bar, 20 μm . (c) Time-to-colour merged image of the nascent otolith in a phenotypically wild-type embryo at the 25-somite stage (sibling of the embryo in d). The colour (arrowhead) indicates a motile cilium near the otolith. Scale bar, 10 μm (also applies to d). (d) Time-to-colour merged image of the nascent otolith in a homozygous *Irrc50* (*dnaaf1*) mutant. The otolith has adhered to the tips of two kinocilia (clearly visible in this image), despite the lack of ciliary motility. The arrowhead marks an otolith precursor particle. (e) Phenotypically wild-type embryo (sibling of the embryo in f) lacking maternal *ift88* contribution but with normal zygotic *ift88* function at 30 hpf. Two otoliths are present, attached to kinocilia (arrowhead, inset). Scale bar, 30 μm (also applies to f). (f) Typical otolith phenotype for a ciliary mutant, in this case an embryo lacking both maternal and zygotic *ift88* function. Three otoliths are present, but still localize to the anterior and posterior poles of the ear. The anterior otolith sits directly on the hair cell stereociliary bundles (arrowhead, inset); all cilia are absent. (g) Transgenic zebrafish otic vesicle at 56 hpf expressing GFP in hair cells. Note that the kinocilia of hair cells in the utricular macula do not all contact the anterior otolith (bottom left). (h) Image of a crista in the same transgenic line at 5 dpf. Note the long straight kinocilia. Scale bar, 10 μm . a, b, e, f and g are lateral views with anterior to the left. Abbreviations: dpf, days post fertilization; hpf, hours post fertilization; M, maternal; MZ, maternal-zygotic; som, somite stage; WT, phenotypically wild-type. a, c, d and e are reproduced from [5]; b and f are reproduced from [2]; g is reproduced from [6]; h is reproduced from [7].

the prefix *kino-*, indicating movement, zebrafish hair cell kinocilia are not thought to generate active motility. However, motile cilia (distinct from the kinocilia and the immotile cilia) are also found in the ear, with most concentrated at the poles in close proximity to the hair cells [2,8,9]. Although some reports have suggested otherwise, motile cilia only appear to make up a small fraction of the total number of cilia in the ear, with the vast majority (92–98%) being immotile [2].

3. Ultrastructure of cilia in the zebrafish ear

Transmission electron microscopy of kinocilia from zebrafish inner ear hair cells reveals a 9+2 axoneme (nine doublet microtubules surrounding a central pair of single microtubules) [10], as described for the supernumerary ear kinocilia in homozygous *mib^{ta52b}* mutants [9]. Although based on very few examples in zebrafish, these studies corroborate findings for utricular and lateral line kinocilia in another teleost fish, the burbot (*Lota lota*) [11–13]. Fish sensory hair cell kinocilia thus make up a class of cilium (9+2, but immotile) often ignored in overviews of ciliary ultrastructure. Hair cell kinocilia are thought to have outer dynein arms and radial spokes, but to lack inner dynein arms and nexin links, as summarized by McHenry & van Netten [14], although this has not been shown definitively for the zebrafish. It will be exciting to see whether the relatively new technique of cryo-electron tomography, already so successful in revealing spectacular ultrastructural detail in the motile 9+2 flagellum of the unicellular alga *Chlamydomonas* [15,16], and of the actin filaments in mammalian vestibular hair cell stereocilia [17,18], can be applied to zebrafish hair cell kinocilia.

Kinocilia of amphibian vestibular hair cells also have the 9+2 pattern (for example, those of the marsh frog, *Pelophylax ridibundus* [19]). However, kinocilia of mammalian (rat and guinea pig) vestibular hair cells have been described to have a 9+0 axoneme, with electron-dense material (but no clear microtubules) at the centre [20,21]. In amniote auditory hair cells, there are further variations: the transient kinocilia of immature hair cells in cochleae of the chick and mouse, for example, have a 9+0 arrangement near the base, but a pattern of eight doublet microtubules surrounding one central doublet (presumed to be one of the nine outer doublets displaced inward) further up the ciliary shaft [22,23]. The axonemal configuration of the short immotile cilia and the motile cilia present at early stages in the zebrafish otic vesicle has not yet been described. Given the similarity of the immotile short cilia to primary cilia, and the rotational movement of the motile cilia (see below), it might be expected that both types have a 9+0 axoneme.

4. Ciliary motility and cilia-driven fluid flow in the zebrafish ear

Motile cilia in the zebrafish ear move in a rotational fashion, with an estimated beat frequency of 28–44 Hz, depending on stage (and possibly other factors such as strain and temperature). Beating cilia can be imaged using high-speed video microscopy at 300 frames per second and above, and are found to be concentrated at the anterior and posterior poles of the early otic vesicle, close to, but distinct from, the immotile tether kinocilia [2,8,9]. In the early otic vesicle, fluid flow

driven by ciliary movement can be inferred from observations of the free otolith precursor particles within the vesicle lumen, which move in vortices near the anterior and posterior poles [3]. The flow has been mapped using particle image velocimetry, by collecting otolith precursor particles in an optical trap and releasing them into the flow near motile cilia [8]. These measurements, together with a physical model simulating cilia-driven flow in a spherical cavity, indicated that the flow field near the tether kinocilia (where most motile cilia are concentrated) is advective, whereas the flow field further from the otolith is dominated by diffusion [8].

In contrast to the motile cilia in the ear, the kinocilia of zebrafish ear and lateral line hair cells are immotile and rigid in the living animal [2,14]. This is beautifully illustrated by live imaging of the *Tg(pou4f3:gfp)* line, which expresses a membrane-tethered GFP specifically in hair cells, revealing the long straight kinocilium, bright stereociliary bundle and hair cell soma of crista hair cells [7] (figure 1*h*). Tether cell kinocilia at the poles of the early otic vesicle show some passive displacement as a result of the beating of nearby motile cilia, but do not display active movement [2].

5. Expression of genes required for ciliary motility in the zebrafish ear

The transcription factor Foxj1 is required for the expression of a suite of genes coding for proteins expressed in motile cilia [9,24,25]. There are two *foxj1* genes in the zebrafish genome: the ear is a major site of expression for *foxj1b*, but not *foxj1a* [24–26]. *foxj1b* is initially expressed throughout the otic placode at 10 hpf, some 8 h before a lumen forms in the vesicle and motile cilia are apparent. In the otic vesicle, *foxj1b* expression becomes restricted to the poles at 22 hpf, apparently marking the first hair cells (the tether cells) developing there [9]. (Expression in hair cells is surprising, as the kinocilia, as discussed above, are immotile.) Otic expression of *foxj1b* is dependent on Wnt signalling: it is strongly downregulated after heat shock-induced mis-expression of the Wnt inhibitor Dkk [27].

In a screen for genes regulated by Foxj1 transcription factors, over 600 differentially expressed genes were uncovered in zebrafish embryos overexpressing *foxj1a* [28]. In this study, the authors assayed for otolith abnormalities (see below) as one of five tissue-specific phenotypic criteria to indicate a successful disruption of motile cilia. Of 50 randomly selected genes tested for function by morpholino-mediated knock-down, two genes appeared to be required for the function of motile cilia in all tissues except the otic vesicle, whereas knockdown of another two (*arhgef18b* and *ect2l*, coding for guanine nucleotide exchange factors) gave defects that were specific to the ear and Kupffer's vesicle [28]. These results highlight potential commonalities between the motile cilia in the ear and those in Kupffer's vesicle, both of which have rotational movement.

As expected for tissues expressing *foxj1b*, the zebrafish otic placode and vesicle are prominent sites of expression for many genes known to be required for ciliogenesis and ciliary motility (see [2] and references therein). These include genes coding for Dyneins and Dynein Axonemal Assembly Factors (DNAAFs) [24,29–34]. A comprehensive expression analysis of dynein axonemal heavy chain (*dnah*) genes in

the zebrafish revealed several expressed in the ear, including one (*dnah91*) that was ear-specific [29]. The tissue specificity of the *dnah* expression patterns appears to correlate with the type of ciliary movement patterns present (oscillatory or rotational): notably, the subset of *dnah* genes expressed in the ear largely seems to overlap with that expressed in the floorplate and Kupffer's vesicle, again highlighting tissues with rotational ciliary movement [29]. All four zebrafish PIH (Protein Interacting with Hsp90) genes are also expressed in the otic placode or vesicle, some very strongly [29]. PIH gene products are DNAAFs implicated in the pre-assembly of dynein arms prior to their transport into cilia. Transcripts for components of the radial spokes of the axoneme (leucine-rich repeat-containing protein 23, *Lrrc23*) and radial spoke head proteins (*Rsph9* and *Rsph4a*) are also expressed in the zebrafish otic placode during somite stages [35,36].

6. The role of zebrafish otic cilia in otolith formation

Many organisms, both invertebrate and vertebrate, sense gravity and other linear acceleration through an inertial mass coupled to a mechanosensitive sensory cell type, which is often ciliated. In ctenophores (comb jellies), for example, lithocytes—entire cells whose cytoplasm harbours a membrane-bound concretion—are actively transported from the base to the tip of sensory cilia in a truly astonishing example of ciliary surface transport (see [37] and references therein). In teleost fish, the inertial mass is the otolith, a dense crystalline aggregate of calcium carbonate and protein that sits over hair cells in the sensory maculae, embedded within an extracellular matrix, the otolithic membrane (for a recent review, see [38]). Otoliths grow by daily incremental deposition of otolithic material: sectioning through an otolith thus reveals rings that can be used to determine the age and life history of a fish [39], just as in the trunk of a tree.

Both kinocilia and motile cilia are implicated in the earliest stages of otolith formation in the zebrafish, as summarized in figure 2. Otolith precursor particles are initially found distributed throughout the otic vesicle lumen, but adhere specifically to the tips of kinocilia on the first hair cells to differentiate in the ear (named tether cells for this reason) [2,3,9]. The composition of the particles has not been fully described, but they are likely to contain Cadherin11 [40], Otoconin90 [41,42] and Starmaker [43,44]. At the kinociliary tip, the bound otolith precursor particles serve as a nucleation site for deposition of further layers of protein and mineral (calcium carbonate), and thus the tether kinocilia become partially embedded within the structure of the otolith. Efficient production or nucleation of the precursor particles, in both zebrafish and medaka, appears to require the activity of the enzyme polyketide synthase [41,45,46]. Growing zebrafish otoliths incorporate a number of proteins, including Otolith Matrix Protein-1, Sparc, Starmaker and others [47–50]. Exciting progress is being made in elucidating the mechanism of otolith biomineralization, which is dependent on the activity of the calcium-binding protein Starmaker [43,50,51] and the conserved proton-selective ion channel Otopetrin 1 [44,52–54].

A defect in otolith formation is a signature phenotype of almost all zebrafish ciliary mutants and morphants described to date, including those with a loss of ciliary motility [2,9,36,55–61]. This observation led to the idea that ciliary




Figure 2. Schematic diagram summarizing the main early steps in zebrafish otolith formation discussed in this review. A single sensory hair cell is shown for simplicity, but tether cells normally form in pairs (figure 1*d*). Not to scale. SC, stereocilia. For details and supporting references, see the text. Key: 1, short, immotile cilia, present on most otic epithelial cells; 2, tether kinocilium, present on each of the first sensory hair cells to develop in the ear; 3, motile cilia—these contribute to the accuracy of otolith precursor particle tethering; 4, extracellular proteins, possibly bound to the kinociliary tip, required for otolith precursor particle tethering: Otogelin (and others?); 5, otolith precursor particles, thought to contain Cadherin11, Otoconin90 and Starmaker; 6, otolith proteins incorporated after initial nucleation, including Otolith Matrix Protein-1, Sparc, Starmaker and others; 7, biomineralization: deposition of crystalline calcium carbonate, requiring the activity of Starmaker and Otopetrin 1; 8, further layers added daily throughout life.

motility must somehow be critical for otolith formation. As discussed above, ciliary flow can readily be demonstrated in the zebrafish ear, and various interpretations have been proposed for its function: that it agitates and distributes the otolith precursor particles, preventing their premature aggregation [3,8,9]; that it creates an advection zone to draw free precursor particles towards the nucleating otolith [8]; or that it contributes to otolith shape [8]. Nevertheless, the otolith phenotype of ciliary mutants is often mild, variable and incompletely penetrant: a typical defect is the presence of three otoliths rather than two, which may resolve over time to give the normal pattern.

A breakthrough in understanding came from the unexpected observation that homozygous maternal-zygotic (MZ) mutants for an intraflagellar transport (IFT) gene, *ift88*, which lack cilia altogether [62], have a similarly mild otolith phenotype [2] (figure 1*e,f*). On close inspection, the otoliths in MZ *ift88*^{otz288b} mutants formed close to the surface of the sensory epithelium, but still nucleated on time at the poles of the ear, in close proximity to the tether cells, which are present but lack kinocilia in this mutant. This suggested that both the immotile tether kinocilia and ciliary motility are, in fact, dispensable for otolith formation. Notably, premature aggregation of otolith precursor

particles was not observed in either MZ*ift88*^{tz288b} mutants (lacking cilia altogether) or in the *lrrc50* (*dnaaf1*) mutant, which lacks ciliary motility (figure 1*c,d*) [2,31,32], and so the proposition that ciliary flow acts to prevent premature aggregation of otolith precursor particles seems unlikely.

A much more striking otolith phenotype—quite distinct from that of the ciliary or biomineralization mutants—results from the lack of tether cells in the zebrafish ear. Injection of an *atoh1b* morpholino into wild-type embryos, which prevents the formation of tether cells, but does not compromise ciliogenesis or ciliary motility, disrupts the early stages of otolith formation completely: otolith precursor particles remain distributed throughout the ear. Here, they become larger than normal and appear birefringent, demonstrating that the process of otolith biomineralization does not require attachment to kinocilia. Eventually, the crystalline particles coalesce to form a single untethered otolith [2]. This observation strongly suggested that it is the presence of tether cells, rather than cilia or cilia motility *per se*, that is essential for otolith development, implicating the function of a hair cell-specific factor or factors in the initial tethering step. Such a factor, perhaps produced by the tether cells, might become transported to the kinociliary tip.

In a quest to identify the otolith precursor-binding factor, we and others turned to mutants that phenocopied the *atoh1b* morphant. Isolated in forward mutagenesis screens on the basis of their phenotype—a single otolith, rather than two, in each ear—there were several promising candidates, initially given the charming names of *einstein* (German: one stone) [63], *rock solo* [64] and *monolith* [65]. Both the *einstein* [66] and *rock solo* [64] mutants were found to disrupt the gene *otogelin*, which codes for a large glycoprotein. *Otogelin* remains the best candidate to date for an otolith precursor-binding factor. Although expression of *otogelin* mRNA is not restricted to hair cells [66], staining with an anti-*Otogelin* antibody suggests that the protein itself becomes concentrated at the distal half of the tether kinocilia [41]. How *Otogelin* gets there is not currently understood. It is possible that one or more adhesion proteins, trafficked to the kinociliary tip, help to localize it there. In this context, it is interesting to note that the integrin *Itga8* and protocadherin *Pcdh15* localize in a complex at the kinociliary tip of zebrafish lateral line hair cells [67]. In chick auditory hair cells, transport of *Pcdh15* to the kinociliary tip is dependent on the IFT machinery and *FGFR1* activity [22]. Zebrafish lateral line and amniote auditory kinocilia, of course, do not bind otoliths, but it is possible that a tether cell-specific factor contributes to the localization of both *Otogelin* protein and otolith precursor particles at the tips of tether kinocilia via a similar mechanism.

Note that ear and otolith morphology vary enormously between different teleosts, and it is not known whether otoliths are nucleated in the same way in other species. Otolith formation was reported to be normal in the medaka mutant *kintoun*, which lacks ciliary motility as a result of mutation in a gene coding for a dynein arm pre-assembly factor [45]. As in zebrafish, this argues against an essential requirement for ciliary motility in otolith formation in this species. In the zebrafish, after the initial nucleation step, it is unlikely that cilia play a further role in otolith growth. Additional hair cell kinocilia do not all contact the otolith directly as the tether kinocilia do (figure 1*g*), and total otolith growth is unaffected by focal laser ablation of motile cilia after otolith nucleation [8]. As the sensory patch and otolith develop further, a layer of extracellular matrix, the otolithic membrane, is deposited on

the apical surface of the sensory patch, and it is this structure that now tethers the otolith to its cognate sensory epithelium [66]. Several protein components of the otolith and otolithic membrane in zebrafish are conserved with those of otoconia and the otoconial membrane, respectively, in vestibular end organs of the mammalian inner ear (reviewed in [68]). Here, multiple small otoconia are present rather than the large individual otoliths found in bony fish. A role for cilia in the initial formation of mammalian otoconia has not been described, to my knowledge.

7. Length control of cilia in the zebrafish ear and lateral line

The length of cilia in the zebrafish ear is both dynamic and sensitive to perturbation. All cilia in the ear are initially of similar length at the very earliest otic vesicle stages (19 somites), but rapidly become differentiated from one another [2]. By the 21-somite stage, cilia at the poles of the ear (tether cell kinocilia and motile cilia) are longer than other cilia throughout the vesicle lumen. Within a few hours, the motile cilia have regressed in length, and by 24 hpf are about half the length of the kinocilia [2]. Kinocilia of different sensory hair cell types can also be distinguished on the basis of length. At 24–25 hpf, the tether kinocilia in the presumptive sensory maculae are about 8 µm long, with the distal 3 µm embedded in the otolith [2]. Kinocilia of the crista hair cells are the longest, projecting up to 30 µm into the lumen of the ear at 4 days post fertilization (dpf) [7,69] (figure 1*h*). Kinocilia on lateral line hair cells are 14–20 µm long in the zebrafish larva, occupying about half the length of the gelatinous cupula that covers them [14,69,70]. In adult teleost fish, kinociliary length can also vary in different regions of the sensory maculae, as shown for the blue gourami, *Trichogaster trichopterus* [71].

The length of a cilium depends on a balance between kinesin-driven anterograde (IFT-B) and dynein-driven retrograde (IFT-A) transport along the axoneme. Mutations in genes coding for proteins involved in ciliogenesis or IFT can thus disrupt cilia length, resulting in either shorter or longer cilia than normal. Disruption of *foxj1b* results in shorter motile cilia and kinocilia in the ear, whereas transgenic mis-expression results in the development of long motile cilia throughout the otic vesicle [9]. Ubiquitous transgenic expression of a GFP-tagged murine *Arl13b* protein in the zebrafish, while labelling most cilia very effectively [72], also increases their length, and is reported to disrupt both ciliary motility and otolith formation in the ear [73]. Mutation of *arl13b* results in shorter lateral line kinocilia; cilia in the ear were not examined [74]. Cilia in all tissues, including the ear, are missing in MZ mutants for the IFT-B gene *ift88* [62] (see above) and the ciliogenesis gene *ta3* [75], while disruption of another IFT-B gene, *ift46* [76], or the IFT-A gene *ift122* [77] results in fewer or abnormal cilia in many tissues, including the ear and lateral line.

Different ciliary types in the zebrafish ear also have different genetic requirements for some gene products. Mutation of the *abcc4* gene, which codes for a transporter involved in prostaglandin signalling, appears to disrupt short cilia, but not tether kinocilia, in the early otic vesicle [60]. The long kinocilia of the cristae seem particularly resistant to genetic perturbation of kinesin subunit genes, which may indicate the existence of additional crista-specific IFT-B components. Crista cilia were unaffected by mutation of *kif3b*, which codes for a

heterotrimeric kinesin II subunit, whereas cilia in maculae were lost altogether [78]. Short crista kinocilia were also able to develop in a strong loss-of-function *kif3a* mutant, which otherwise eliminates cilia in maculae of the ear and in several other tissues, including the lateral line [79]. Cilia length in the zebrafish ear is also governed by the activity of Crumbs genes [80,81]. mRNA for *crb3a* is strongly and almost exclusively expressed in the ear; the Crb3a protein localizes to the base of hair cell kinocilia at the otic vesicle stage, and to the basal body of kinocilia in adult hair cells. Morpholino-mediated knockdown of *crb3a* results in shorter kinocilia in maculae [80], but interestingly, in loss-of-function mutants for *crb3a* or *crb2b*, cilia in the cristae get longer [81].

8. The role of kinocilia and ciliary proteins in hair cell planar polarity, stereociliary bundle assembly and mechanosensitivity

The eccentric position of the kinocilium on the hair cell apical surface defines the directional sensitivity of the cell: deflection of the stereociliary bundle in the direction of the kinocilium depolarizes the cell, whereas deflection in the opposite direction hyperpolarizes the cell (see [70] and references therein). This can be demonstrated in the zebrafish larva by imaging of lateral line hair cells labelled with the calcium indicator GCaMP6s after deflection of kinocilia in the neuromast cupula with a glass fibre or fluid jet [70,82]. The kinociliary axoneme is arranged in a polar fashion with respect to the hair bundle, with the first doublet microtubule (number 1) located on the side facing the stereociliary bundle [11].

In mammalian auditory hair cells, the kinocilium plays a role in orientation of the hair cell stereociliary bundle, together with planar cell polarity genes (see [83] and references therein). However, there is currently no strong evidence to suggest a role for kinocilia in establishing hair bundle polarity in zebrafish hair cells, which are thought to be more closely related to mammalian vestibular hair cells. Normal stereocilia polarity was observed in both ear and lateral line hair cells in the zebrafish *ift88^{tz288b}* mutant, which has missing or stunted kinocilia [5,70]. This contrasts with the obvious disruption of planar cell polarity in mammalian auditory hair cells in the murine *Ift88* mutant, where mis-oriented and even circular stereociliary bundles were observed in the organ of Corti [83]. Although hair bundle polarity is not affected in zebrafish zygotic *ift88^{tz288b}* mutants, hair bundle morphology is abnormal and fewer hair cells form stereocilia [5,84]. Analysis of individual mutant phenotypes and assays for protein colocalization revealed that the Ift88 protein helps to stabilize a complex of the Usher syndrome 1 (USH1) proteins Cdh23, Harmonin and MyoVIIa in the endoplasmic reticulum (ER), ensuring their correct trafficking for assembly of the stereociliary bundle. Disruption of any one component of the complex can result in ER stress and eventual hair cell death [84]. Thus, Ift88 and possibly other ciliary proteins have additional roles in protein or vesicular traffic within hair cells.

Kinocilia of immature zebrafish lateral line hair cells also have an early role in mediating hair cell mechanosensitivity, a property restricted to the stereociliary bundle in mature hair cells [70]. Small but robust responses were measured in hair cells with morphologically immature stereociliary bundles, but with an opposite polarity to responses found in mature

hair cells. By using the *ift88^{tz288b}* mutant, in which cilia are either absent or stunted, the authors were able to show that these early responses were mediated by kinocilia. The loss of responses in mutants for the cadherin genes *cdh23* and *pcdh15*, together with pharmacological disruption, demonstrated that the early polarity-reversed responses were dependent on the integrity of kinociliary links between the kinocilium and adjacent stereocilia [70].

9. The role of kinocilia and ciliary proteins in sensitivity to aminoglycoside ototoxicity

Aminoglycoside antibiotics are potent ototoxins, killing sensory hair cells in both zebrafish and mammals (see [85] and references therein). As a result, off-target damage to the auditory and vestibular systems can be a considerable clinical problem in patients treated with aminoglycosides for serious acute or chronic bacterial infections. Work in the zebrafish has revealed that cilia and ciliary proteins appear to play a role in the uptake and/or toxicity of these compounds. Mutations in anterograde IFT-B, retrograde IFT-A and ciliary transition zone genes in the zebrafish all confer protection against neomycin ototoxicity [74,86,87]. In most IFT mutants, where ciliary morphology and hair cell mechanotransduction are compromised, neomycin uptake into the cell was disrupted [74,87]. However, mutations in transition zone genes (*cc2d2a*, *mks1*, *cep290*)—where ciliogenesis, mechanotransduction and neomycin uptake were unaffected—still conferred partial protection, suggesting that transition zone proteins may modulate trafficking or metabolism of neomycin once it has entered the hair cell [74]. These studies are important for our understanding of aminoglycoside uptake and toxicity in hair cells, with implications for the search for and design of new otoprotective compounds or alternative antibiotic agents.

10. Role of cilia in signalling in the zebrafish ear

Cilia are well known to act as cellular antennae for the reception of various extracellular signals (reviewed in [88]). As the zebrafish ear is a prominent site of expression for many ciliary genes, it is perhaps surprising that few studies to date have addressed the contribution of cilia to cell signalling in this tissue. In zebrafish, the complete loss of cilia in MZ*ift88^{tz288b}* mutants, or depletion of cilia in *igu* (*dzip1*) mutants, reduces levels but expands the reach of Hedgehog pathway activity, rather than eliminating it [62]. This finding corroborates preliminary observations of the later ear phenotype of MZ*ift88* mutants, which have morphological abnormalities that resemble those of *igu* (*dzip1*) mutants [5], but are not a close phenocopy of the mirror-symmetric double-anterior ears generated by a strong loss of Hh signalling (see [89] and references therein). Signalling via cilia or cilia-driven flow, therefore, does not appear to play a major role in otic axial patterning in the otic vesicle. This contrasts with the similarly sized Kupffer's vesicle, where ciliary-driven flow is essential for left-right patterning of the embryo (see [90] and references therein). However, given the mix of motile and non-motile ciliary types in both the early otic vesicle and Kupffer's vesicle [90], their shared dependence on Wnt signalling [27], and the overlap in their ciliary transcriptome profile [28,29], this remains an interesting area to explore further.

The active phosphorylated form of type II calcium/calmodulin kinase CaMK-II is expressed in the zebrafish ear in hair cell kinocilia, most strongly at their base, where it may localize to the ciliary transition zone and/or stereociliary bundle [91]. Interestingly, disruption of CaMK-II activity—either pharmacologically or via morpholino-mediated knock-down of *camk2g1*—resulted in a disruption of Delta–Notch signalling, and the development of supernumerary tether cells. Morphants and treated embryos had otolith defects that resembled those of *mib1^{ta52b}* (loss of Notch signalling) mutants at 24 hpf, where multiple tether kinocilia make contact with otolithic material [2,91]. At later stages (30–72 hpf), *camk2g1* morphants or treated embryos showed a phenotype characteristic of ciliary mutants, with ectopic fused or third otoliths in the ear [91]. Thus, signalling via calcium, CaMK-II, Delta and cilia appears to be closely linked in the zebrafish ear.

11. Conclusion

Our understanding of ciliary structure and function has come a long way in the last 50 years, although many of the stunning electron micrographs of kinocilia from the 1960s [11,12,20] are still among the clearest in the literature, and repay careful attention. There remains enormous interest in cilia, not least for their important link to rare genetic disease: mutations in the human orthologues of many of the genes discussed here result in ciliopathies that affect the quality of life and are challenging to diagnose and treat. Sensorineural deafness is a feature of several of these syndromes, underlining the need to understand cilia function in the inner ear. Here, the zebrafish

holds promise as a model system, partly because the otoliths are a highly visible proxy readout of ciliary integrity. As technologies to achieve tissue-specific disruption of zebrafish gene function improve (e.g. [92–94]), new ways of working to analyse ciliary function in the ear will become possible, where—unlike in the brain or kidney—loss of function should not compromise adult viability. Cilia in the zebrafish ear are also interesting in their own right; key among the problems remaining to be solved is an understanding of the unique protein or lipid composition at the tip of the tether kinocilium that distinguishes it from other cilia, allowing otolith precursor particles to bind there so precisely and robustly. Comparison of the rotational motile cilia in the otic vesicle to those in Kupffer's vesicle or the floorplate may also lead to general insights into mechanisms of cilia-driven flow and its consequences for developmental patterning.

Data accessibility. This article has no additional data.

Competing interests. I declare I have no competing interests.

Funding. Work in the Whitfield lab is funded by the BBSRC (grant no. BB/S007008/1).

Acknowledgements. Thanks are due to Raymond Goldstein, Gáspár Jékely and Kirsty Wan for organizing a hugely enjoyable and thought-provoking meeting, which provided the springboard for this review. I am grateful to Georgina Stooke-Vaughan for comments on the manuscript.

Endnote

¹For a review of motile cilia in the olfactory system, see reference [1].

References

- Ringers C, Olstad EW, Jurisch-Yaksi N. 2020 The role of motile cilia in the development and physiology of the nervous system. *Phil. Trans. R. Soc. B* **375**, 20190156. (doi:10.1098/rstb.2019.0156)
- Stooke-Vaughan GA, Huang P, Hammond KL, Schier AF, Whitfield TT. 2012 The role of hair cells, cilia and ciliary motility in otolith formation in the zebrafish otic vesicle. *Development* **139**, 1777–1787. (doi:10.1242/dev.079947)
- Riley BB, Zhu C, Janetopoulos C, Aufderheide KJ. 1997 A critical period of ear development controlled by distinct populations of ciliated cells in the zebrafish. *Dev. Biol.* **191**, 191–201. (doi:10.1006/dbio.1997.8736)
- Haddon C, Lewis J. 1996 Early ear development in the embryo of the zebrafish, *Danio rerio*. *J. Comp. Neurol.* **365**, 113–123. (doi:10.1002/(SICI)1096-9861(19960129)365:1<113::AID-CNE9>3.0.CO;2-6)
- Stooke-Vaughan GA. 2013 The role of cilia in inner ear development and otolith formation in the zebrafish otic vesicle. PhD thesis, University of Sheffield. See <http://etheses.whiterose.ac.uk/id/eprint/4992>.
- Maier EC, Saxena A, Alsina B, Bronner ME, Whitfield TT. 2014 Sensational placodes: neurogenesis in the otic and olfactory systems. *Dev. Biol.* **389**, 50–67. (doi:10.1016/j.ydbio.2014.01.023)
- Stawicki TM, Owens KN, Linbo T, Reinhardt KE, Rubel EW, Raible DW. 2014 The zebrafish *merovingian* mutant reveals a role for pH regulation in hair cell toxicity and function. *Dis. Models Mech.* **7**, 847–856. (doi:10.1242/dmm.016576)
- Wu D, Freund JB, Fraser SE, Vermot J. 2011 Mechanistic basis of otolith formation during teleost inner ear development. *Dev. Cell* **20**, 271–278. (doi:10.1016/j.devcel.2010.12.006)
- Yu X, Lau D, Ng CP, Roy S. 2011 Cilia-driven fluid flow as an epigenetic cue for otolith biomineralization on sensory hair cells of the inner ear. *Development* **138**, 487–494. (doi:10.1242/dev.057752)
- Leventea E, Hazime K, Zhao C, Malicki J. 2016 Analysis of cilia structure and function in zebrafish. *Methods Cell Biol.* **133**, 179–227. (doi:10.1016/b9.mcb.2016.04.016)
- Flock Å, Duvall AJ. 1965 The ultrastructure of the kinocilium of the sensory cells in the inner ear and lateral line organs. *J. Cell Biol.* **25**, 1–8. (doi:10.1083/jcb.25.1.1)
- Flock Å, Wersäll J. 1962 A study of the orientation of the sensory hairs of the receptor cells in the lateral line organ of fish, with special reference to the function of the receptors. *J. Cell Biol.* **15**, 19–27. (doi:10.1083/jcb.15.1.19)
- Flock Å. 1964 Structure of the macula utriculi with special reference to directional interplay of sensory responses as revealed by morphological polarization. *J. Cell Biol.* **22**, 413–431. (doi:10.1083/jcb.22.2.413)
- McHenry MJ, van Netten SM. 2007 The flexural stiffness of superficial neuromasts in the zebrafish (*Danio rerio*) lateral line. *J. Exp. Biol.* **210**, 4244–4253. (doi:10.1242/jeb.009290)
- Nicastro D. 2009 Cryo-electron microscope tomography to study axonemal organization. *Methods Cell Biol.* **91**, 1–39. (doi:10.1016/S0091-679X(08)91001-3)
- Bui KH, Pigino G, Ishikawa T. 2011 Three-dimensional structural analysis of eukaryotic flagella/cilia by electron cryo-tomography. *J. Synchrotron Radiat.* **18**, 2–5. (doi:10.1107/S0909049510036812)
- Metlagel Z *et al.* 2019 Electron cryo-tomography of vestibular hair-cell stereocilia. *J. Struct. Biol.* **206**, 149–155. (doi:10.1016/j.jsb.2019.02.006)
- Sazzed S, Song J, Kovacs JA, Wriggers W, Auer M, He J. 2018 Tracing actin filament bundles in three-dimensional electron tomography density maps of hair cell stereocilia. *Molecules* **23**, 882. (doi:10.3390/molecules23040882)
- Nagel G, Neugebauer D-C, Schmidt B, Thurm U. 1991 Structures transmitting stimulatory force to

- the sensory hairs of vestibular ampullae of fishes and frog. *Cell Tiss. Res.* **265**, 567–578. (doi:10.1007/BF00340881)
20. Hamilton DW. 1969 The cilium on mammalian vestibular hair cells. *Anat. Rec.* **164**, 253–258. (doi:10.1002/ar.1091640301)
 21. Kikuchi T, Takasaka T, Tonosaki A, Watanabe H. 1989 Fine structure of guinea pig vestibular kinocilium. *Acta Otolaryngol.* **108**, 26–30. (doi:10.3109/00016488909107388)
 22. Honda A, Kita T, Seshadri SV, Misaki K, Ahmed Z, Ladbury JE, Richardson GP, Yonemura S, Ladher RK. 2018 FGFR1-mediated protocadherin-15 loading mediates cargo specificity during intraflagellar transport in inner ear hair-cell kinocilia. *Proc. Natl Acad. Sci. USA* **115**, 8388–8393. (doi:10.1073/pnas.1719861115)
 23. Sobkowicz HM, Slapnick SM, August BK. 1995 The kinocilium of auditory hair cells and evidence for its morphogenetic role during the regeneration of stereocilia and cuticular plates. *J. Neurocytol.* **24**, 633–653. (doi:10.1007/BF01179815)
 24. Yu X, Ng CP, Habacher H, Roy S. 2008 Foxj1 transcription factors are master regulators of the motile ciliogenic program. *Nat. Genet.* **40**, 1445–1453. (doi:10.1038/ng.263)
 25. Hellman NE *et al.* 2010 The zebrafish *foxj1a* transcription factor regulates cilia function in response to injury and epithelial stretch. *Proc. Natl Acad. Sci. USA* **107**, 18 499–18 504. (doi:10.1073/pnas.1005998107)
 26. Tian T, Zhao L, Zhao X, Zhang M, Meng A. 2009 A zebrafish gene trap line expresses GFP recapturing expression pattern of *foxj1b*. *J. Genet. Genomics* **36**, 581–589. (doi:10.1016/S1673-8527(08)60150-2)
 27. Caron A, Xu X, Lin X. 2012 Wnt/ β -catenin signaling directly regulates Foxj1 expression and ciliogenesis in zebrafish Kupffer's vesicle. *Development* **139**, 514–524. (doi:10.1242/dev.071746)
 28. Choksi SP, Babu D, Lau D, Yu X, Roy S. 2014 Systematic discovery of novel ciliary genes through functional genomics in the zebrafish. *Development* **141**, 3410–3419. (doi:10.1242/dev.108209)
 29. Yamaguchi H, Oda T, Kikkawa M, Takeda H. 2018 Systematic studies of all PIH proteins in zebrafish reveal their distinct roles in axonemal dynein assembly. *eLife* **7**, e36979. (doi:10.7554/eLife.36979)
 30. Jaffe KM *et al.* 2016 *c21orf59/kurly* controls both cilia motility and polarization. *Cell Rep.* **14**, 1841–1849. (doi:10.1016/j.celrep.2016.01.069)
 31. van Rooijen E, Giles RH, Voest EE, van Rooijen C, Schulte-Merker S, van Eeden FJ. 2008 LRRC50, a conserved ciliary protein implicated in polycystic kidney disease. *J. Am. Soc. Nephrol.* **19**, 1128–1138. (doi:10.1681/ASN.2007080917)
 32. Sullivan-Brown J, Schottenfeld J, Okabe N, Hostetter CL, Serluca FC, Thiberge SY, Burdine RD. 2008 Zebrafish mutations affecting cilia motility share similar cystic phenotypes and suggest a mechanism of cyst formation that differs from *pkd2* morphants. *Dev. Biol.* **314**, 261–275. (doi:10.1016/j.ydbio.2007.11.025)
 33. Hjej R *et al.* 2014 *CCDC151* mutations cause primary ciliary dyskinesia by disruption of the outer dynein arm docking complex formation. *Am. J. Hum. Genet.* **95**, 257–274. (doi:10.1016/j.ajhg.2014.08.005)
 34. Jerber J, Baas D, Soulavie F, Chhin B, Cortier E, Vesque C, Thomas J, Durand B. 2014 The coiled-coil domain containing protein CCDC151 is required for the function of IFT-dependent motile cilia in animals. *Hum. Mol. Genet.* **23**, 563–577. (doi:10.1093/hmg/ddt445)
 35. Sedykh I, TeSlaa JJ, Tatarsky RL, Keller AN, Toops KA, Lakkaraju A, Nyholm MK, Wolman MA, Grinblat Y. 2016 Novel roles for the radial spoke head protein 9 in neural and neurosensory cilia. *Scient. Rep.* **6**, 34437. (doi:10.1038/srep34437)
 36. Han X, Xie H, Wang Y, Zhao C. 2018 Radial spoke proteins regulate otolith formation during early zebrafish development. *FASEB J.* **32**, 3984–3992. (doi:10.1096/fj.201701359R)
 37. Noda N, Tamm SL. 2014 Lithocytes are transported along the ciliary surface to build the statolith of ctenophores. *Curr. Biol.* **24**, R951–R952. (doi:10.1016/j.cub.2014.08.045)
 38. Schulz-Mirbach T, Ladich F, Plath M, Heß M. 2019 Enigmatic ear stones: what we know about the functional role and evolution of fish otoliths. *Biol. Rev. Camb. Philos. Soc.* **94**, 457–482. (doi:10.1111/brv.12463)
 39. Campana SE, Neilson JD. 1985 Microstructure of fish otoliths. *Can. J. Fish. Aquat. Sci.* **42**, 1014–1032. (doi:10.1139/f85-127)
 40. Clendenon SG, Shah B, Miller CA, Schmeisser G, Walter A, Gattone 2nd VH, Barald KF, Liu Q, Marrs JA. 2009 Cadherin-11 controls otolith assembly: evidence for extracellular cadherin activity. *Dev. Dyn.* **238**, 1909–1922. (doi:10.1002/dvdy.22015)
 41. Thiessen KD, Grzegorski SJ, Chin Y, Higuchi L, Wilkinson CJ, Shavit JA, Kramer KL. 2019 Zebrafish otolith biomineralization requires polyketide synthase. *Mech. Dev.* **157**, 1–9. (doi:10.1016/j.mod.2019.04.001)
 42. Petko JA, Kabbani N, Frey C, Woll M, Hickey K, Craig M, Canfield VA, Levenson R. 2009 Proteomic and functional analysis of NCS-1 binding proteins reveals novel signaling pathways required for inner ear development in zebrafish. *BMC Neurosci.* **10**, 27. (doi:10.1186/1471-2202-10-27)
 43. Söllner C, Burghammer M, Busch-Nentwich E, Berger J, Schwartz H, Riekel C, Nicolson T. 2003 Control of crystal size and lattice formation by Starmaker in otolith biomineralization. *Science* **302**, 282–286. (doi:10.1126/science.1088443)
 44. Söllner C, Schwarz H, Geisler R, Nicolson T. 2004 Mutated *otopetrin 1* affects the genesis of otoliths and the localization of Starmaker in zebrafish. *Dev. Genes Evol.* **214**, 582–590. (doi:10.1007/s00427-004-0440-2)
 45. Hojo M *et al.* 2015 Unexpected link between polyketide synthase and calcium carbonate biomineralization. *Zool. Lett.* **1**, 3. (doi:10.1186/s40851-014-0001-0)
 46. Lee MS, Philippe J, Katsanis N, Zhou W. 2019 Polyketide synthase plays a conserved role in otolith formation. *Zebrafish* **16**, 363–369. (doi:10.1089/zeb.2019.1734)
 47. Kang YJ, Stevenson AK, Yau PM, Kollmar R. 2008 Sparc protein is required for normal growth of zebrafish otoliths. *J. Ass. Res. Otolaryngol.* **9**, 436–451. (doi:10.1007/s10162-008-0137-8)
 48. Rotllant J, Liu D, Yan YL, Postlethwait JH, Westerfield M, Du SJ. 2008 Sparc (Osteonectin) functions in morphogenesis of the pharyngeal skeleton and inner ear. *Matrix Biol.* **27**, 561–572. (doi:10.1016/j.matbio.2008.03.001)
 49. Murayama E, Herbolme P, Kawakami A, Takeda H, Nagasawa H. 2005 Otolith matrix proteins OMP-1 and Otolin-1 are necessary for normal otolith growth and their correct anchoring onto the sensory maculae. *Mech. Dev.* **122**, 791–803. (doi:10.1016/j.mod.2005.03.002)
 50. Kalka M, Markiewicz N, Ptak M, Sone ED, Ozyhar A, Dobryszczycki P, Wojtas M. 2019 *In vivo* and *in vitro* analysis of starmaker activity in zebrafish otolith biomineralization. *FASEB J.* **33**, 6877–6886. (doi:10.1096/fj.201802268R)
 51. Wojtas M, Hotubowicz R, Poznar M, Maciejewska M, Ozyhar A, Dobryszczycki P. 2015 Calcium ion binding properties and the effect of phosphorylation on the intrinsically disordered Starmaker protein. *Biochemistry* **54**, 6525–6534. (doi:10.1021/acs.biochem.5b00933)
 52. Saotome K, Teng B, Tsui CCA, Lee WH, Tu YH, Kaplan JP, Sansom MSP, Liman ER, Ward AB. 2019 Structures of the otopetrin proton channels Otop1 and Otop3. *Nat. Struct. Mol. Biol.* **26**, 518–525. (doi:10.1038/s41594-019-0235-9)
 53. Tu YH *et al.* 2018 An evolutionarily conserved gene family encodes proton-selective ion channels. *Science* **359**, 1047–1050. (doi:10.1126/science.aao3264)
 54. Hughes I *et al.* 2004 Otopetrin 1 is required for otolith formation in the zebrafish *Danio rerio*. *Dev. Biol.* **276**, 391–402. (doi:10.1016/j.ydbio.2004.09.001)
 55. Gao C, Wang G, Amack JD, Mitchell DR. 2010 Oda16/Wdr69 is essential for axonemal dynein assembly and ciliary motility during zebrafish embryogenesis. *Dev. Dyn.* **239**, 2190–2197. (doi:10.1002/dvdy.22355)
 56. Panizzi JR, Jessen JR, Drummond IA, Solnica-Krezel L. 2007 New functions for a vertebrate Rho guanine nucleotide exchange factor in ciliated epithelia. *Development* **134**, 921–931. (doi:10.1242/dev.02776)
 57. Neugebauer JM, Amack JD, Peterson AG, Bisgrove BW, Yost HJ. 2009 FGF signalling during embryo development regulates cilia length in diverse epithelia. *Nature* **458**, 651–654. (doi:10.1038/nature07753)
 58. Wilkinson CJ, Carl M, Harris WA. 2009 Cep70 and Cep131 contribute to ciliogenesis in zebrafish embryos. *BMC Cell Biol.* **10**, 17. (doi:10.1186/1471-2121-10-17)
 59. Pathak N, Austin CA, Drummond IA. 2011 Tubulin tyrosine ligase-like genes *ttl3* and *ttl6* maintain zebrafish cilia structure and motility. *J. Biol. Chem.* **286**, 11 685–11 695. (doi:10.1074/jbc.M110.209817)

60. Jin D *et al.* 2014 Prostaglandin signalling regulates ciliogenesis by modulating intraflagellar transport. *Nat. Cell Biol.* **16**, 841–851. (doi:10.1038/ncb3029)
61. Jurisch-Yaksi N *et al.* 2013 Rer1p maintains ciliary length and signaling by regulating γ -secretase activity and Foxj1a levels. *J. Cell Biol.* **200**, 709–720. (doi:10.1083/jcb.201208175)
62. Huang P, Schier AF. 2009 Dampened Hedgehog signaling but normal Wnt signaling in zebrafish without cilia. *Development* **136**, 3089–3098. (doi:10.1242/dev.041343)
63. Whitfield TT *et al.* 1996 Mutations affecting development of the zebrafish inner ear and lateral line. *Development* **123**, 241–254.
64. Roberts R, Elsner J, Bagnall MW. 2017 Delayed otolith development does not impair vestibular circuit formation in zebrafish. *J. Ass. Res. Otolaryngol.* **18**, 415–425. (doi:10.1007/s10162-017-0617-9)
65. Riley BB, Grunwald DJ. 1996 A mutation in zebrafish affecting a localized cellular function required for normal ear development. *Dev. Biol.* **179**, 427–435. (doi:10.1006/dbio.1996.0272)
66. Stooke-Vaughan GA, Obholzer ND, Baxendale S, Megason SG, Whitfield TT. 2015 Otolith tethering in the zebrafish otic vesicle requires Otogelin and α -Tectorin. *Development* **142**, 1137–1145. (doi:10.1242/dev.116632)
67. Goodman L, Zallocchi M. 2017 Integrin α 8 and Pcdh15 act as a complex to regulate cilia biogenesis in sensory cells. *J. Cell Sci.* **130**, 3698–3712. (doi:10.1242/jcs.206201)
68. Lundberg YW, Xu Y, Thiessen KD, Kramer KL. 2014 Mechanisms of otoconia and otolith development. *Dev. Dyn.* **244**, 239–253. (doi:10.1002/dvdy.24195)
69. Imtiaz A *et al.* 2018 CDC14A phosphatase is essential for hearing and male fertility in mouse and human. *Hum. Mol. Genet.* **27**, 780–798. (doi:10.1093/hmg/ddx440)
70. Kindt KS, Finch G, Nicolson T. 2012 Kinocilia mediate mechanosensitivity in developing zebrafish hair cells. *Dev. Cell* **23**, 329–341. (doi:10.1016/j.devcel.2012.05.022)
71. Popper AN, Hoxter B. 1981 The fine structure of the sacculus and lagena of a teleost fish. *Hear. Res.* **5**, 245–263. (doi:10.1016/0378-5955(81)90049-6)
72. Borovina A, Superina S, Voskas D, Ciruna B. 2010 Vangl2 directs the posterior tilting and asymmetric localization of motile primary cilia. *Nat. Cell Biol.* **12**, 407–412. (doi:10.1038/ncb2042)
73. Lu H, Toh MT, Narasimhan V, Thamilselvam SK, Choksi SP, Roy S. 2015 A function for the Joubert syndrome protein Arl13b in ciliary membrane extension and ciliary length regulation. *Dev. Biol.* **397**, 225–236. (doi:10.1016/j.ydbio.2014.11.009)
74. Stawicki TM *et al.* 2016 Cilia-associated genes play differing roles in aminoglycoside-induced hair cell death in zebrafish. *G3* **6**, 2225–2235. (doi:10.1534/g3.116.030080)
75. Ben J, Elworthy S, Ng AS, van Eeden F, Ingham PW. 2011 Targeted mutation of the *talpid3* gene in zebrafish reveals its conserved requirement for ciliogenesis and Hedgehog signalling across the vertebrates. *Development* **138**, 4969–4978. (doi:10.1242/dev.070862)
76. Lee MS *et al.* 2015 IFT46 plays an essential role in cilia development. *Dev. Biol.* **400**, 248–257. (doi:10.1016/j.ydbio.2015.02.009)
77. Boubakri M, Chaya T, Hirata H, Kajimura N, Kuwahara R, Ueno A, Malicki J, Furukawa T, Omori Y. 2016 Loss of ift122, a retrograde intraflagellar transport (IFT) complex component, leads to slow, progressive photoreceptor degeneration due to inefficient opsin transport. *J. Biol. Chem.* **291**, 24 465–24 474. (doi:10.1074/jbc.M116.738658)
78. Zhao C, Omori Y, Brodowska K, Kovach P, Malicki J. 2012 Kinesin-2 family in vertebrate ciliogenesis. *Proc. Natl Acad. Sci. USA* **109**, 2388–2393. (doi:10.1073/pnas.1116035109)
79. Pooranachandran N, Malicki JJ. 2016 Unexpected roles for ciliary kinesins and intraflagellar transport proteins. *Genetics* **203**, 771–785. (doi:10.1534/genetics.115.180943)
80. Omori Y, Malicki J. 2006 *oko meduzy* and related *crumbs* genes are determinants of apical cell features in the vertebrate embryo. *Curr. Biol.* **16**, 945–957. (doi:10.1016/j.cub.2006.03.058)
81. Hazime K, Malicki JJ. 2017 Apico-basal polarity determinants encoded by *crumbs* genes affect ciliary shaft protein composition, IFT movement dynamics, and cilia length. *Genetics* **207**, 1041–1051. (doi:10.1534/genetics.117.300260)
82. Lukasz D, Kindt KS. 2018 In vivo calcium imaging of lateral-line hair cells in larval zebrafish. *J. Vis. Exp.* **141**, e58794. (doi:10.3791/58794)
83. Jones C, Roper VC, Foucher I, Qian D, Banizs B, Petit C, Yoder BK, Chen P. 2008 Ciliary proteins link basal body polarization to planar cell polarity regulation. *Nat. Genet.* **40**, 69–77. (doi:10.1038/ng.2007.54)
84. Blanco-Sánchez B, Clément A, Fierro Jr J, Washbourne P, Westerfield M. 2014 Complexes of Usher proteins preassemble at the endoplasmic reticulum and are required for trafficking and ER homeostasis. *Dis. Models Mech.* **7**, 547–559. (doi:10.1242/dmm.014068)
85. Harris JA, Cheng AG, Cunningham LL, MacDonald G, Raible DW, Rubel EW. 2003 Neomycin-induced hair cell death and rapid regeneration in the lateral line of zebrafish (*Danio rerio*). *J. Ass. Res. Otolaryngol.* **4**, 219–234. (doi:10.1007/s10162-002-3022-x)
86. Owens KN, Santos F, Roberts B, Linbo T, Coffin AB, Knisely AJ, Simon JA, Rubel EW, Raible DW. 2008 Identification of genetic and chemical modulators of zebrafish mechanosensory hair cell death. *PLoS Genet.* **4**, e1000020. (doi:10.1371/journal.pgen.1000020)
87. Stawicki TM, Linbo T, Hernandez L, Parkinson L, Bellefeuille D, Rubel EW, Raible DW. 2019 The role of retrograde intraflagellar transport genes in aminoglycoside-induced hair cell death. *Biol. Open* **8**, bio038745. (doi:10.1242/bio.038745)
88. Nachury MV, Mick DU. 2019 Establishing and regulating the composition of cilia for signal transduction. *Nat. Rev. Mol. Cell Biol.* **20**, 389–405. (doi:10.1038/s41580-019-0116-4)
89. Hartwell RD, England SJ, Monk NAM, van Hateren NJ, Baxendale S, Marzo M, Lewis KE, Whitfield TT. 2019 Anteroposterior patterning of the zebrafish ear through Fgf- and Hh-dependent regulation of *hmx3a* expression. *PLoS Genet.* **15**, e1008051. (doi:10.1371/journal.pgen.1008051)
90. Sampaio P *et al.* 2014 Left-right organizer flow dynamics: how much cilia activity reliably yields laterality? *Dev. Cell.* **29**, 716–728. (doi:10.1016/j.devcel.2014.04.030)
91. Rothschild SC, Lahvic J, Francescato L, McLeod JJ, Burgess SM, Tombes RM. 2013 CaMK-II activation is essential for zebrafish inner ear development and acts through Delta–Notch signaling. *Dev. Biol.* **381**, 179–188. (doi:10.1016/j.ydbio.2013.05.028)
92. Yamaguchi N, Colak-Champollion T, Knaut H. 2019 zGrad is a nanobody-based degron system that inactivates proteins in zebrafish. *eLife* **8**, e43125. (doi:10.7554/eLife.43125)
93. Kirchgeorg L, Felker A, van Oostrom M, Chivacci E, Mosimann C. 2018 Cre/*lox*-controlled spatiotemporal perturbation of FGF signaling in zebrafish. *Dev. Dyn.* **247**, 1146–1159. (doi:10.1002/dvdy.24668)
94. Savage AM *et al.* 2019 *tmem33* is essential for VEGF-mediated endothelial calcium oscillations and angiogenesis. *Nat. Commun.* **10**, 732. (doi:10.1038/s41467-019-08590-7)