



# Algorithm for the management of patients at low, high and very high risk of osteoporotic fractures

J. A. Kanis<sup>1,2</sup> · N. C. Harvey<sup>3</sup> · E. McCloskey<sup>1,4</sup> · O. Bruyère<sup>5</sup> · N. Veronese<sup>6</sup> · M. Lorentzon<sup>2,7,8</sup> · C. Cooper<sup>3,9</sup> · R. Rizzoli<sup>10</sup> · G. Adib<sup>11</sup> · N. Al-Daghri<sup>12</sup> · C. Campusano<sup>13</sup> · M. Chandran<sup>14</sup> · B. Dawson-Hughes<sup>15</sup> · K. Javaid<sup>9</sup> · F. Jiwa<sup>16</sup> · H. Johansson<sup>1,2</sup> · J. K. Lee<sup>17</sup> · E. Liu<sup>2</sup> · D. Messina<sup>18</sup> · O. Mkinsi<sup>19</sup> · D. Pinto<sup>20,21</sup> · D. Prieto-Alhambra<sup>9,22</sup> · K. Saag<sup>23</sup> · W. Xia<sup>24</sup> · L. Zakraoui<sup>25</sup> · J. -Y. Reginster<sup>12,26</sup>

Received: 26 June 2019 / Accepted: 18 September 2019 / Published online: 13 November 2019  
© The Author(s) 2019, corrected publication 2020

## Abstract

**Summary** Guidance is provided in an international setting on the assessment and specific treatment of postmenopausal women at low, high and very high risk of fragility fractures.

**Introduction** The International Osteoporosis Foundation and European Society for Clinical and Economic Aspects of Osteoporosis and Osteoarthritis published guidance for the diagnosis and management of osteoporosis in 2019. This manuscript seeks to apply this in an international setting, taking additional account of further categorisation of increased risk of fracture, which may inform choice of therapeutic approach.

**Methods** Clinical perspective and updated literature search.

**Results** The following areas are reviewed: categorisation of fracture risk and general pharmacological management of osteoporosis.

**Conclusions** A platform is provided on which specific guidelines can be developed for national use to characterise fracture risk and direct interventions.

**Keywords** Anabolic agents · Fracture risk assessment · FRAX · Inhibitors of bone resorption · Treatment of osteoporosis

## Introduction

In 2018, the International Osteoporosis Foundation (IOF) and the European Society for Clinical and Economic Evaluation of Osteoporosis and Osteoarthritis (ESCEO) updated guidelines for the diagnosis and management of postmenopausal osteoporosis, subsequently published in full in 2019 [1] and as executive summaries [2, 3]. Translation of this guideline into easy to use, practical algorithms is needed to facilitate the recognition and

treatment of women at increased risk of fracture. This translation could also enable the incorporation of several recent developments that significantly impact on strategies for the management of patients. The first is the widespread recognition that the risk of a subsequent osteoporotic fracture is particularly acute immediately after an index fracture and wanes progressively with time [4–9]. This very high fracture risk and the consequent further utility loss immediately after a subsequent fracture (often termed “imminent risk” because of the temporal association) suggests that preventive treatment given as soon as possible after fracture would avoid a higher number of new fractures and reduce the attendant morbidity, compared with treatment given later. This provides the rationale for very early intervention immediately after a sentinel fracture and necessitates treatment with agents that have the most rapid effect on fracture reduction. A further recent development is the demonstration of a more rapid and greater fracture risk reduction of anabolic compared with antiresorptive treatments [10–14], with the potential to revolutionise treatment strategies, particularly in individuals at very high fracture risk [15, 16].

---

Outcomes of an experts’ consensus meeting jointly organised by the European Society for Clinical and Economic Aspects of Osteoporosis, Osteoarthritis and Musculoskeletal Diseases (ESCEO) and the International Osteoporosis Foundation (IOF)

---

✉ J. A. Kanis  
w.j.Pontefract@shef.ac.uk

Extended author information available on the last page of the article

The scope of the present report is to review and update the assessment of osteoporosis, in particular the categorisation of risk to better target therapeutic interventions for the prevention of fragility fracture in postmenopausal women. The guideline is intended for all healthcare professionals involved in the management of osteoporosis. Where available, systematic reviews, meta-analyses and randomised controlled trials have been used to provide the evidence-base with the available literature updated using PubMed to identify systematic reviews and meta-analyses from January 2017 to December 2018, subsequent to the generation of the recent European Guidelines. The recommendations in this guidance have been endorsed by the Scientific Advisory Board of ESCEO and the Committee of Scientific Advisors and the Committee of National Societies of the IOF.


## Risk assessment

The IOF and ESCEO recommend that risk of fracture should be expressed as an absolute risk, i.e. probability of fracture over a ten-year interval [1]. The absolute risk of fracture depends upon age and life expectancy as well as the current fracture risk. The period of 10 years was chosen to cover the likely length of treatment and the time over which benefits may continue or risks arise if treatment is stopped [17]. Algorithms that integrate the weight of clinical risk factors for fracture risk, with or without information on BMD, were developed in 2007 by the then WHO Collaborating Centre for Metabolic Bone Diseases at Sheffield. The resulting FRAX tool ([www.shef.ac.uk/FRAX](http://www.shef.ac.uk/FRAX)) computes the 10-year probability of hip fracture or a major osteoporotic fracture, the latter comprising a clinical spine, hip, forearm or humerus fracture. The tool has been externally validated in independent cohorts [18] and calibrated to the epidemiology of fracture and death in 67 countries covering more than 80% of the world population at risk [19].

## Intervention and assessment thresholds

FRAX has been incorporated into more than 100 guidelines worldwide but the approach to intervention thresholds has varied widely [19, 20]. For the purposes of this report, the guidance of the IOF and ESCEO [1] is used as an example and shown in Fig. 1.



In the European guidance, it is recommended that postmenopausal women with a prior fragility fracture should be treated without further assessment, although BMD measurement and incorporation into the FRAX calculation is sometimes appropriate, particularly in younger postmenopausal women. In women without a previous fragility fracture, the management strategy should be based on assessment of the ten-year probability of a major osteoporotic fracture (clinical spine, hip,



**Fig. 1** Assessment guidelines based on the ten-year probability of a major osteoporotic fracture (%). The dotted line denotes the intervention threshold. Where assessment is made in the absence of BMD, a BMD test is recommended for individuals where the probability assessment lies in the orange region i.e. between the lower assessment threshold (LAT) and the upper assessment threshold (UAT). The intervention threshold and BMD assessment thresholds used are those derived from [1] and reproduced in the Appendix, Table 5, with kind permission from Springer Science and Business Media

forearm or humerus). Women with probabilities below the lower assessment threshold can be considered at low risk. Women with probabilities above the upper assessment threshold can be considered for treatment. Women with probabilities between the upper and lower assessment threshold should be referred for BMD measurements and their fracture probability reassessed [1]. The age-dependent intervention threshold is set at a risk equivalent to that associated with a prior fracture in a woman of the same age with average BMI and, therefore, rises with age [21]. The same thresholds are used in men since the cost-effectiveness of interventions is broadly similar in men and women [22, 23].


**Fig. 2** Infographic outlining of the characterisation of fracture risk by FRAX major osteoporotic fracture probability in postmenopausal women. Initial risk assessment uses FRAX with clinical risk factors alone. FRAX probability in the red zone indicates very high risk and that an initial course of anabolic treatment followed by antiresorptive therapy may be appropriate. FRAX probability in the green zone suggests low risk, with advice to be given on lifestyle, calcium and vitamin D nutrition and menopausal hormone treatment considered. FRAX probability in the intermediate (orange) zone should be followed by BMD assessment and recalculation of FRAX probability including femoral neck BMD. After recalculation, risk may be in the red zone (very high risk), orange zone (high risk, which suggests initial antiresorptive therapy) or green zone (low risk, either in the original green zone or in the original orange zone but below the intervention threshold). *Note that patients with a prior fragility fracture are at least designated at high risk and possibly at very high risk dependent on the FRAX probability*



Some agencies have been reluctant to reimburse treatments on the basis of fracture probability, particularly at younger ages where the 10-year probability of a major osteoporotic


fracture is less than 10%. In the UK, for example, the intervention threshold for women aged 50–54 years is 8.18%. At the same age, however, the remaining lifetime probability of a

**Fig. 3** Treatment pathways according to the categorisation of fracture risk. For treatment modalities (inhibitors of bone resorption, anabolic agents, etc.), see Appendix, Table 6



MHT, menopausal hormone therapy;  
 SERM, selective estrogen receptor modulator;  
 LOEP, local osteo-enhancement procedure  
 \* See Appendix, table A2

**Fig 4** Proportion (%) of postmenopausal women by age in a simulated normal population of 50633 women from the UK [51] characterised at low, high and very high risk. The high-risk category includes women with a prior fracture not characterised at very high risk. Numbers in the high and very high risk categories refer to the percentage so characterised at each age interval



major fracture is high (57%). Moreover, there are cost-effective scenarios for treatment available at these levels of risk [1].

In addition to the 10-year probability of a major osteoporotic fracture, the European guidance also provides intervention thresholds that are based on the 10-year probability of hip fracture. Either or both thresholds can be used; indeed, the screening for prevention of fractures in older women (SCOOP) trial showed that a screening strategy decreased the incidence of hip fracture (but not other fractures), based on treatment targeted by hip fracture probability [24].

To enhance fracture risk assessment, relatively simple arithmetic adjustments have been developed, which can be applied to conventional FRAX estimates of probabilities of hip fracture and a major fracture to adjust the probability assessment with knowledge of

- High, moderate and low exposure to glucocorticoids [25]
- Concurrent data on lumbar spine BMD [26, 27]
- Trabecular bone score of the lumbar spine [28–30]
- Hip axis length [31]
- Falls history [32]
- Immigration status [33]
- Type 2 diabetes [34, 35]
- Chronic kidney disease [36]
- Recency of fracture [9, 37]

### Categorisation of risk

The assessment strategy above permits the classification of risk. In addition to the categories of low and high risk espoused in the current IOF-ESCEO guideline, very high risk can be identified as outlined in Fig. 2. Very high risk is defined as a fracture probability that lies above the upper assessment threshold after a FRAX assessment, with or without the inclusion of BMD, i.e. where BMD testing is unavailable, the same probability threshold can be used. The numerical data for

**Table 1** Examples of risk assessment in women from the UK (BMI set to 25 kg/m<sup>2</sup>). Risk factors include prior fracture (of uncertain recency), prior clinical vertebral fracture within the past two years, family history of

Age (years)	Prior fracture	Recent spine fracture	Family history	GC	GC high dose	BMD (T-score)	10-year probability (%)	Category of risk
70	Yes					-	20	Low <sup>1</sup>
70			Yes			-	17	Low
70	Yes		Yes			-	30	Very high
70		Yes				-	30	Very high
60				Yes		-1.5	10	Low
60				Yes		-2.0	13	High
60				Yes	Yes	-2.0	15	Very high

<sup>1</sup> Qualifies for treatment by virtue of a prior fracture

**Table 2** 10-year probability of major osteoporotic fracture (MOF) for Icelandic women at different ages, categorised by (A) a clinical vertebral fracture within the previous 2 years and (B) a prior fracture of undetermined recency. The right-hand column provides the ratio by which to adjust FRAX probabilities by virtue of a recent clinical vertebral fracture. From [59], with kind permission from Springer Science and Business Media

Age	10-year probability of MOF		Ratio
	(A) Recent vertebral fracture	(B) Prior fracture in adult life	
50	29.0	11.7	2.47
60	36.1	19.4	1.86
70	41.9	27.6	1.52
80	42.5	34.2	1.24
90	34.7	33.3	1.04

these thresholds are given in the Appendix (Table 5) based on the weighted average of 5 European countries (Germany, France, Italy, Spain and the UK) but will vary by country.

### Impact on treatment

The rationale for the more refined characterisation of risk is to direct appropriate interventions. Thus, initial treatment recommendations for women at high risk might most usually start with an inhibitor of bone resorption. For example, the UK National Institute for Clinical Excellence (NICE) and the IOF/ESCEO guidelines recommend oral bisphosphonates [1, 38, 39] but a very large range of pharmacological interventions are recommended worldwide (Appendix, Tables 6 and 7). In contrast, women at very high risk might be more suitably treated with an anabolic treatment followed thereafter by an inhibitor of bone resorption [40] (Fig. 3).

Non-pharmacological management should be considered for all patients but may be adapted according to the category

hip fracture, exposure to glucocorticoids, exposure to higher than average doses of glucocorticoids and bone mineral density (BMD) T-score at the femoral neck

**Table 3** Unadjusted probabilities of a major fracture in women with a prior fragility fracture by age using the UK FRAX model together with the categories of risk and adjusted FRAX probabilities for women with a recent clinical vertebral fracture. BMI set at 24 kg/m<sup>2</sup>

Age	Prior fracture in adult life		Recent clinical vertebral fracture	
	Probability (%)	Category of risk	Probability (%)	Category of risk
50	7.3	High	18.0	Very high
60	12.2	High	22.7	Very high
70	20.3	High	30.9	Very high
80	27.6	Very high	34.2	Very high
90	33.9	Very high	35.3	Very high

of fracture risk [41, 42]. For all patients, education on osteoporosis (e.g. knowledge of osteoporosis, medications, diet and exercise) and advice for daily weight-bearing physical activity are appropriate [43–45]. Where indicated, the addition of fall prevention measures, including supervised exercise and/or rehabilitation, has been shown to be useful in reducing falls [46, 47] but the effects on fracture risk remain uncertain [47]. Programmes should continue over a duration of at least 50 hours, be progressive in nature and include strength and balance training components [48–50].

The implications of the categorisation of risk on the use of anabolic regimens in this way are shown in Fig. 4 as applied to the age-specific NOGG guidance [21]. As would be expected, the proportion of women characterised at low risk decreased with age and, conversely, those at high risk increased with age. The proportion of women characterised at very high risk increased with age though the quantum of effect was modest. Overall, in women aged 50 years or more, 64.8% were categorised at low risk, 19.7% at high risk and 15.6% at very high risk. Numerical data by age are given in the Appendix (Table 6).

**Table 4** The effect on hip fracture (number/1000 patient years) of an anabolic agent (AA) given for the first 18 months followed by an antiresorptive (AR) for a total of 10 years. The clinical scenario is a postmenopausal woman from the UK with a recent major osteoporotic fracture. The efficacy (RRR) of the anabolic agent is modelled at 70% and

	Untreated	AA/AR RRR 70/40% <sup>a</sup>	Fractures saved <sup>b</sup>	AR/AN RRR 40/70%	Fractures saved <sup>b</sup>
Age	N/1000	N/1000	N/1000	N/1000	N/1000
50	8.1	2.4	5.7	4.7	3.4
55	12.8	3.8	9.0	7.5	5.3
60	20.4	6.1	14.3	11.9	8.5
65	31.3	9.4	21.9	18.3	13.0
70	48.3	14.5	33.8	28.3	20.0
75	73.6	22.1	51.5	43.1	30.5
80	104.7	31.4	73.3	61.2	43.5
85	160.4	48.1	112.3	93.8	66.6
90	180.9	54.3	126.6	105.8	75.1

<sup>a</sup> It is assumed that the effect of the anabolic agent is maintained with the subsequent antiresorptive agent

<sup>b</sup> First fractures

## Examples of very high risk

A prior fragility fracture provides informative examples of the categorisation of risk and recommendations for treatment which are illustrated for a woman age 70 years in Table 1. A prior fragility fracture of undetermined recency is associated with a 10-year probability of a major osteoporotic fracture of 16%. Whereas this probability is designated as low, treatment with an inhibitor of bone resorption is indicated by the IOF-ESCEO guidance and many other guidelines by virtue of the prior fracture [19]. A family history of hip fracture in the absence of any other risk factors provides a similar fracture probability is characterised at low risk and lifestyle advice recommended. The combination of a prior fragility fracture of uncertain recency and a family history of hip fracture is associated with a much higher risk than either risk factor alone or falls into the category of very high risk where an anabolic regimen might be recommended.

A further example of the interaction of risk factors on the categorisation of risk is provided in the context of exposure to glucocorticoids (see Table 1). A woman aged 60 years exposed to average doses of glucocorticoids and a femoral neck T-score of –

that of the antiresorptive at 40%. The time course of a subsequent hip fracture is non-linear as given in [58]. The two right-hand columns show the effects of an antiresorptive followed by an anabolic agent for the last 18 months of a 10-year treatment

1.5 would be characterised as being at low risk in the absence of other clinical risk factors. A lower T-score of  $-2.0$  would place her in the category of high risk. However, for high doses ( $> 7.5$  mg daily), probabilities should be upward-revised by about 15% [25, 52] which would place the patient at very high risk.

There is now a substantial body of evidence that the risk of a subsequent osteoporotic fracture is particularly acute immediately after the index fracture and wanes progressively with time [4–8, 53–57]. Thus, the incidence of a second fracture in those who will sustain a further fracture is particularly high in the first 2 years after the index event [58]. In the case of hip fracture, 61% of subsequent fractures over a 10-year time horizon will occur within the first 2 years. For forearm, spine and humerus fractures the recurrence within two years is 54, 42 and 53%, respectively. A recent population-based study demonstrated that the phenomenon of immediate risk was also age dependent [9]. For a woman at age 70 years, a prior clinical vertebral fracture within the past 2 years is associated with a 1.52-fold higher fracture probability than for a woman of the same age with a prior fragility fracture of uncertain recency [59] (Table 2). Thus, a recent clinical vertebral fracture uplifts the fracture probability from 16 to 24% and would place the woman in the category of very high risk (see Table 1).

Risk stratification with NOGG guidelines\* BMI set at  $25 \text{ kg/m}^2$

The ratios given in Table 2 can be used to adjust fracture probabilities derived from FRAX for recency of clinical vertebral fracture. Table 3 gives the current probabilities of a major fracture in women with a prior fragility fracture of uncertain recency using the UK FRAX model together with the adjusted probabilities for women with a clinical vertebral fracture within the past two years.

## Impact of sequential treatment on fracture

In patients at very high risk of fracture, starting treatment with an anabolic agent seems most appropriate to promptly reduce the fracture risk [60–62]. Given that treatments with anabolic agents are limited to 12–24 months and that efficacy will wane once treatment is stopped, the real potential of the anabolic treatments is that their greater effect on BMD and fracture can be maintained with the inhibitors of bone turnover once anabolic treatment is stopped [11, 63, 64]. Take for example a hypothetical anabolic agent that reduced the risk of hip fracture by 70% (relative risk reduction, RRR = 70%). In this case, the anabolic agent, followed by an antiresorptive to maintain the effect for a total of 10 years, might be expected to save 33.8 hip fractures/1000 patient years in women age 70 years with a recent fragility fracture. In contrast, an antiresorptive (RRR = 40%) followed by an anabolic regimen for the last 18 months of a 10-year treatment would save only 20.0 hip fractures/1000 patient years. The difference illustrates the

importance of the sequence. The assumptions used and data for other ages are given in Table 4.

## Conclusion

The risk categorisation of individuals deemed to merit treatment into high and very high risk and aids the targeting of anabolic therapy followed by antiresorptive medications.

**Acknowledgements** We are grateful to the Scientific Advisory Board of ESCEO and the Committee of Scientific Advisors and the Committee of National Societies of the IOF for their review and endorsement of the recommendations in this guidance.

## Compliance with ethical standards

**Conflicts of interest** NM Al-Daghri G Adib, C Campusano, M Chandran, F Jiwa, H Johansson, JK Lee, E Liu, D Pinto, N Veronese, W Xia, L Zakraoui have no conflicts of interest to declare.

O Bruyère received research grants from Biophytis, IBSA, MEDA, Servier and SMB and consulting or lecture fees from Amgen, Biophytis, IBSA, MEDA, Servier, SMB, TRB Chemedica and UCB.

C Cooper reports personal fees from Alliance for Better Bone Health, Amgen, Eli Lilly, GSK, Medtronic, Merck, Novartis, Pfizer, Roche, Servier, Takeda and UCB.

B. Dawson-Hughes has received grant support from Pfizer and DSM and consulting fees from TTY Biopharma Co, Ltd, Intrinsic Therapeutics and Agnovos.

NC Harvey has received consultancy/lecture fees/honoraria/grant funding from Alliance for Better Bone Health, Amgen, MSD, Eli Lilly, Radius Health, Servier, Shire, UCB, Consilient Healthcare and Internis Pharma.

MK Javaid has received honoraria, unrestricted research grants, travel and/or subsistence expenses from Amgen, Lilly UK, Internis, Consilient Health, Zebra Medical Vision, Kyowa Kirin Hakin and UCB.

JA Kanis reports grants from Amgen, Eli Lilly and Radius Health and consulting fees from Theramex. JAK is the architect of FRAX® but has no financial interest.

M Lorentzon has received lecture fees from Amgen, Lilly, Meda, Renapharma and UCB Pharma and consulting fees from Amgen, Radius Health, UCB Pharma, Renapharma and Consilient Health, all outside the presented work.

EV McCloskey has received consultancy/lecture fees/grant funding/honoraria from ActiveSignal, AgNovos, Amgen, AstraZeneca, Consilient Healthcare, Fresenius Kabi, Gilead, GSK, Hologic, Internis, Lilly, Medtronic, Merck, Novartis, Pfizer, Radius Health, Redx Oncology, Roche, SanofiAventis, Servier, Synexus, Tethys, UCB, Viiv, Warner Chilcott, I3 Innovus and Unilever.

OD Messina has received honoraria from Amgen, Lilly, Novartis and Pfizer

O Minski received honorary fees for lectures, from Abbvie, Roche, Pfizer, Pierre Fabre, Janssen, Novartis and MSD.

D Prieto-Alhambra's research group has received research grants from Amgen, Servier and UCB; speaker fees from Amgen and UCB; educational grants from Johnson & Johnson and consultancy fees from Amgen and UCB.

J-Y Reginster has received advisory board or consulting fees from IBSA-Genévrier, Pierre Fabre, Radius Health, TEVA and Mylan; lecture fees from Agnovos, IBSA-Genévrier, Mylan, CNIEL, Dairy Research Council (DRC) and Theramex and institutional grant support from IBSA-Genévrier, Mylan, CNIEL and Radius Health.

R Rizzoli has received consulting fees or advisory board fees from Radius Health, Labatec, Danone, Nestlé, CNIEL and Sandoz.

K Saag reports grant support from Amgen and Radius and consulting fees from Amgen, Radius and Roche.

## Appendix

**Table 5** Intervention thresholds as set by FRAX-based 10-year probability (%) of a major osteoporotic fracture equivalent to women with a previous fracture (no other clinical risk factors, a body mass index of 24 kg/m<sup>2</sup> and without BMD). The lower assessment thresholds set by FRAX is based on the 10-year probability (%) of a major

osteoporotic fracture equivalent to women without clinical risk factors (a body mass index of 24 kg/m<sup>2</sup> and without BMD). The upper assessment threshold is set at 1.2 times the intervention threshold. The UK FRAX model is used. From [1], with kind permission from Springer Science and Business Media.

Age range (years)	Ten-year fracture probability (%)			Lifetime risk at intervention threshold*
	Intervention threshold*	Lower assessment threshold	Upper assessment threshold**	
50–54	7.8	4.0	9.4	57
55–59	11	5.3	13.2	54
60–64	14	7.3	16.8	50
65–69	19	9.8	22.8	47
70–74	22	12	26.4	43
75–79	26	16	31.2	39
80–84	31	20	37.2	36
85–89	33	18	39.6	34

\*Threshold for high risk

\*\*Threshold for very high risk

**Table 6** Pharmaceutical interventions used in the management of postmenopausal osteoporosis. The list is not comprehensive but includes agents approved in Europe, the USA and member countries represented by the authors

Inhibitors of bone resorption	Stimulators of bone formation
Vitamin D derivatives	
Alfacalcidol	Abaloparatide
Calcidiol	Teriparatide (including biosimilars)
Calcitriol	Romosozumab
Bisphosphonates	
Alendronate (including effervescent formulation)	
Clodronate	Uncertain action
Neridronate	Strontium ranelate
Risedronate (including gastric resistant formulation)	
Ibandronate	
Zoledronate	
MHT and SERMs	
Oestrogen only MHT	
Opposed MHT (with progestogen)	
Tibolone	
Bazedoxifene	
Raloxifene	
Other	
Vitamin K	
Calcitonin	
Denosumab	

MHT, menopause hormonal treatment; SERMs, selective oestrogen receptor modulator



**Table 7** Number of postmenopausal women by age in a simulation cohort of 50,633 women from the UK [51] and the proportion (%) characterised at low, high and very high risk. The high risk category includes women with a prior fracture not characterised at very high risk

	Age (years)								Total
	50–54	55–59	60–64	65–69	70–74	75–79	80–84	85+	
Number	8799	8384	8552	6671	5792	4784	3807	3844	50,633
Low risk (%)	73.0	70.1	67.1	65.1	61.4	58.0	54.8	52.0	64.8
High risk (%)	13.4	15.9	17.0	19.1	22.2	24.7	28.2	31.1	19.7
Very high risk (%)	13.7	14.0	16.0	16.0	16.3	17.3	17.1	16.9	15.6

**Open Access** This article is licensed under a Creative Commons Attribution-NonCommercial 4.0 International License, which permits any non-commercial use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by-nc/4.0/>.

## References


- Kanis JA, Cooper C, Rizzoli R, Reginster J-Y, Scientific Advisory Board of the European Society for Clinical and Economic Aspects of Osteoporosis (ESCEO) and the Committees of Scientific Advisors and National Societies of the International Osteoporosis Foundation (IOF) (2019) European guidance for the diagnosis and management of osteoporosis in postmenopausal women. *Osteoporos Int* 30:3–44
- Kanis JA, Cooper C, Rizzoli R, Reginster J-Y (2019) Executive summary of European guidance for the diagnosis and management of osteoporosis in postmenopausal women. *Aging Clin Exp Res* 31: 15–17
- Kanis JA, Cooper C, Rizzoli R, Reginster J-Y (2019) Executive summary of the European guidance for the diagnosis and management of osteoporosis in postmenopausal women. *Calcif Tissue Int* 104:235–238
- Johnell O, Oden A, Caulin F, Kanis JA (2001) Acute and long-term increase in fracture risk after hospitalization for vertebral fracture. *Osteoporos Int* 12:207–214
- Johnell O, Kanis JA, Oden A, Sernbo I, Redlund-Johnell I, Pettersen C, De Laet C, Jonsson B (2004) Fracture risk following an osteoporotic fracture. *Osteoporos Int* 15:175–179
- Nymark T, Lauritsen JM, Ovesen O, Röck ND, Jeune B (2006) Short time-frame from first to second hip fracture in the Funen County Hip Fracture Study. *Osteoporos Int* 17:1353–1357
- Giangregorio LM, Leslie WD (2010) Manitoba bone density program. Time since prior fracture is a risk modifier for 10-year osteoporotic fractures. *J Bone Miner Res* 25:1400–1405
- van Geel TACM, van Helden S, Geusens PP, Winkens B, Dinant G-J (2009) Clinical subsequent fractures cluster in time after first fractures. *Ann Rheum Dis* 68:99–102
- Johansson H, Siggeirsdóttir K, Harvey NC, Odén A, Gudnason V, McCloskey E, Sigurdsson G, Kanis JA (2017) Imminent risk of fracture after fracture. *Osteoporos Int* 28:775–780
- Kendler DL, Marin F, Zerbinì CAF, Russo LA, Greenspan SL, Zikan V, Bagur A, Malouf-Sierra J, Lakatos P, Fahrleitner-Pammer A, Lespessailles E, Minisola S, Body JJ, Geusens P, Möricke R, López-Romero P (2018) Effects of teriparatide and risedronate on new fractures in post-menopausal women with severe osteoporosis (VERO): a multicentre, double-blind, double-dummy, randomised controlled trial. *Lancet* 391(10117):230–240
- Saag KG, Petersen J, Brandi ML, Karaplis AC, Lorentzon M, Thomas T, Maddox J, Fan M, Meisner PD, Grauer A (2017) Romosozumab or alendronate for fracture prevention in women with osteoporosis. *N Engl J Med* 377:1417–1427
- Barrionuevo P, Kapoor E, Asi N, Alahdab F, Mohammed K, Benkhadra K, Almasri J, Farah W, Sarigianni M, Muthusamy K, Al Nofal A, Haydour Q, Wang Z, Murad MH (2019) Efficacy of pharmacological therapies for the prevention of fractures in postmenopausal women: a network meta-analysis. *J Clin Endocrinol Metab* 104:1623–1630
- Lou S, Lv H, Yin P, Li Z, Tang P, Wang Y (2019) Combination therapy with parathyroid hormone analogs and antiresorptive agents for osteoporosis: a systematic review and meta-analysis of randomized controlled trials. *Osteoporos Int* 30:59–70
- Diez-Pérez A, Marin F, Eriksen EF, Kendler DL, Krege JH, Delgado-Rodríguez M (2019) Effects of teriparatide on hip and upper limb fractures in patients with osteoporosis: a systematic review and meta-analysis. *Bone* 120:1–8. <https://doi.org/10.1016/j.bone.2018.09.020>
- Russow G, Jahn D, Appelt J, Märdian S, Tsitsilonis S, Keller J (2018) Anabolic therapies in osteoporosis and bone regeneration. *Int J Mol Sci* 26:20(1)
- Khosla S, Hofbauer LC (2017) Osteoporosis treatment: recent developments and ongoing challenges. *Lancet Diabetes Endocrinol* 5: 898–907
- Kanis JA on behalf of the World Health Organization Scientific Group (2008) Assessment of osteoporosis at the primary health-care level. Technical Report. WHO Collaborating Centre, University of Sheffield, UK. Available at: [https://www.sheffield.ac.uk/FRAX/pdfs/WHO\\_Technical\\_Report.pdf](https://www.sheffield.ac.uk/FRAX/pdfs/WHO_Technical_Report.pdf) Accessed 2 April 2019
- Kanis JA, Oden A, Johnell O, Johansson H, De Laet C, Brown J, Burckhardt P, Cooper C, Christiansen C, Cummings S, Eisman JA, Fujiwara S, Gluer C, Goltzman D, Hans D, Krieg MA, La Croix A, McCloskey E, Mellstrom D, Melton LJ 3rd, Pols H, Reeve J, Sanders K, Schott AM, Silman A, Torgerson D, van Staa T, Watts NB, Yoshimura N (2007) The use of clinical risk factors enhances the performance of BMD in the prediction of hip and osteoporotic fractures in men and women. *Osteoporos Int* 18:1033–1046
- Kanis JA, Harvey NC, Cooper C, Johansson H, Odén A, McCloskey EV, the Advisory Board of the National Osteoporosis

- Guideline Group (2016) A systematic review of intervention thresholds based on FRAX. A report prepared for the National Osteoporosis Guideline Group and the International Osteoporosis Foundation. *Arch Osteoporos* 11:25
20. Clark P, Denova-Gutiérrez E, Zerbini C, Sanchez A, Messina O, Jaller JJ, Campusano C, Orces CH, Riera G, Johansson H, Kanis JA (2018) FRAX-based intervention and assessment thresholds in seven Latin American countries. *Osteoporos Int* 29:707–715
  21. Kanis JA, EV MC, Johansson H, Strom O, Borgstrom F, Oden A, the National Osteoporosis Guideline Group (2008) Case finding for the management of osteoporosis with FRAX® - assessment and intervention thresholds for the UK. *Osteoporos Int* 19:1395–1408
  22. Tosteson AN, Melton LJ 3rd, Dawson-Hughes B et al (2008) Cost-effective osteoporosis treatment thresholds: the United States perspective. *Osteoporos Int* 19:437–447
  23. Kanis JA, Stevenson M, McCloskey EV, Davis S, Lloyd-Jones M (2007) Glucocorticoid-induced osteoporosis: a systematic review and cost-utility analysis. *Health Technol Assess*; 11: iii–iv, ix–xi, 1–231
  24. Shepstone L, Lenaghan E, Cooper C, Clarke S, Fong-Soe-Khioe R, Fordham R, Gittos NJ, Harvey I, Harvey N, Heawood A, Holland R, Howe A, Kanis J, Marshall T, O'Neill T, Peters T, Redmond N, Torgerson D, Turner D, McCloskey E (2018) Screening in the community to reduce fractures in older women (SCOOP): a randomised controlled trial. *Lancet* 391(10122):741–747
  25. Kanis JA, Johansson H, Oden A, McCloskey EV (2011) Guidance for the adjustment of FRAX according to the dose of glucocorticoids. *Osteoporos Int* 22:809–816
  26. Leslie WD, Lix LM, Johansson H, Odén A, McCloskey E, Kanis JA (2011) Spine-hip discordance and fracture risk assessment: a physician-friendly FRAX enhancement. *Osteoporos Int* 22:839–847
  27. Johansson H, Kanis JA, Odén A, Leslie WD, Fujiwara S, Glüer CC, Kroger H, LaCroix AZ, Lau E, Melton LJ 3rd, Eisman JA, O'Neill TW, Goltzman D, Reid DM, McCloskey E (2014) Impact of femoral neck and lumbar spine BMD discordances on FRAX probabilities in women: a meta-analysis of international cohorts. *Calcif Tissue Int* 95:428–435
  28. McCloskey EV, Odén A, Harvey NC, Leslie WD, Hans D, Johansson H, Barkmann R, Boutroy S, Brown J, Chapurlat R, Elders PJM, Fujita Y, Glüer CC, Goltzman D, Iki M, Karlsson M, Kindmark A, Kurumatani N, Kwok T, Leung J, Lippuner K, Ljunggren Ö, Lorentzon M, Mellström D, Merlijn T, Oei L, Ohlsson C, Pasco J, Rivadeneira F, Rosengren B, Sornay-Rendu E, Szulc P, Tamaki J, Kanis JA (2016) A meta-analysis of trabecular bone score in fracture risk prediction and its dependence on FRAX. *J Bone Miner Res* 31:940–948
  29. Leslie WD, Johansson H, Kanis JA, Lamy O, Odén A, McCloskey EV, Hans D (2014) Lumbar spine texture enhances 10-year fracture probability assessment. *Osteoporos Int* 25:2271–2277
  30. Leslie WD, Shevroja E, Johansson H, McCloskey EV, Harvey NC, Kanis JA, Hans D (2018) Risk-equivalent T-score adjustment using lumbar spine trabecular bone score (TBS): the Manitoba BMD Registry. *Osteoporos Int* 29:751–758
  31. Leslie WD, Lix LM, Morin SN, Johansson H, Odén A, McCloskey EV, Kanis JA (2015) Adjusting hip fracture probability in men and women using hip axis length: the Manitoba Bone Density Database. *J Clin Densitom* 19:326–331
  32. Masud T, Binkley N, Boonen S, Hannan MT, on behalf of the FRAX Position Conference members (2011) Can falls and frailty be used in FRAX? *J Clin Densitom* 14:194–204
  33. Johansson H, Odén A, Lorentzon M, McCloskey E, Kanis JA, Harvey NC, Karlsson MK, Mellstrom D (2015) Is the Swedish FRAX model appropriate for immigrants to Sweden? *Osteoporos Int* 26:2617–2622
  34. Leslie WD, Rubin MR, Schwartz AZ, Kanis JA (2012) Perspective: diabetes and bone. *J Bone Miner Res* 27:2231–2237
  35. Leslie WD, Johansson H, McCloskey EV, Harvey NC, Kanis JA, Hans D (2018) Comparison of methods for improving fracture risk assessment in diabetes: the Manitoba BMD Registry. *J Bone Miner Res* 33:1923–1930
  36. Whitlock RH, Leslie WD, Shaw J, Rigatto C, Thorlacius L, Komenda P, Collister D, Kanis JA, Tangri N (2019) The Fracture Risk Assessment Tool (FRAX®) predicts fracture risk in patients with chronic kidney disease. *Kidney Int* 95:447–454
  37. Kanis JA, Johansson H, Harvey NC, Lorentzon M, Liu E, Borgström F, McCloskey EV (2019) Correspondence in response to OSIN-D-18-00831 quantifying imminent risk. *Osteoporos Int* 30:525–526
  38. National Institute for Health and Care Excellence (NICE) (2017) Bisphosphonates for treating osteoporosis. Technology appraisal guidance 464. National Institute for Health and Care Excellence, London
  39. Compston J, Cooper A, Cooper C, Gittos N, Gregson C, Harvey N, Hope S, Kanis JA, McCloskey EV, KES P, Reid DM, Selby P, Thompson F, Thurston A, Vine N, The National Osteoporosis Guideline Group (NOGG) (2017) UK clinical guideline for the prevention and treatment of osteoporosis. *Arch Osteoporos* 12:43
  40. Kanis JA, Rizzoli R, Cooper C, Reginster J-Y (2014) Challenges for the development of bone forming agents in Europe. *Calcif Tissue Int* 94:469–473
  41. Beck BR, Daly RM, Singh MA, Taaffe DR (2017) Exercise and Sports Science Australia (ESSA) position statement on exercise prescription for the prevention and management of osteoporosis. *J Sci Med Sport* 20:438–445
  42. Giangregorio L, El-Kotob R (2017) Exercise, muscle, and the applied load-bone strength balance. *Osteoporos Int* 28:21–33
  43. Bian S, Hu J, Zhang K, Wang Y, Yu M, Ma J (2018) Dairy product consumption and risk of hip fracture: a systematic review and meta-analysis. *BMC Public Health* 18:165. <https://doi.org/10.1186/s12889-018-5041-5>
  44. Jensen AL, Lomborg K, Wind G, Langdahl BL (2014) Effectiveness and characteristics of multifaceted osteoporosis group education - a systematic review. *Osteoporos Int* 25:1209–1224
  45. Morfeld JC, Vennedey V, Müller D, Pieper D, Stock S (2017) Patient education in osteoporosis prevention: a systematic review focusing on methodological quality of randomised controlled trials. *Osteoporos Int* 28:1779–1803
  46. Zhou X, Deng H, Shen X, Lei Q (2018) Effect of balance training on falls in patients with osteoporosis: a systematic review and meta-analysis. *J Rehabil Med* 50:577–581
  47. de Souto BP, Rolland Y, Vellas B (2018 Dec 28) Maltais M (2019) Association of long-term exercise training with risk of falls, fractures, hospitalizations, and mortality in older adults: a systematic review and meta-analysis. *JAMA Intern Med*. <https://doi.org/10.1001/jamainternmed.2018.5406>
  48. Kemmler W, von Stengel S, Kohl M (2017) Exercise frequency and fracture risk in older adults -how often is enough? *Curr Osteoporos Rep* 15:564–570
  49. Sherrington C, Michaleff ZA, Fairhall N, Paul SS, Tiedemann A, Whitney J, Cumming RG, Herbert RD, Close JCT, Lord SR (2017) Exercise to prevent falls in older adults: an updated systematic review and meta-analysis. *Br J Sports Med* 51:1750–1758
  50. Tricco AC, Thomas SM, Veroniki AA, Hamid JS, Cogo E, Striffler L, Khan PA, Robson R, Sibley KM, MacDonald H, Riva JJ, Thavorn K, Wilson C, Holroyd-Leduc J, Kerr GD, Feldman F, Majumdar SR, Jaglal SB, Hui W, Straus SE (2017) Comparisons of interventions for preventing falls in older adults: a systematic review and meta-analysis. *JAMA* 318:1687–1699
  51. McCloskey E, Kanis JA, Johansson H, Harvey N, Odén A, Cooper A, Cooper C, Francis R, Reid D, Selby P, Davies C, Bowring C,

- Compston J (2015) FRAX-based assessment and intervention thresholds—an exploration of thresholds in women aged 50 years and older in the UK. *Osteoporos Int* 26:2091–2099
52. Lekamwasam S, Adachi JD, Agnusdei D, Bilezikian J, Boonen S, Borgström F, Cooper C, Diez Perez A, Eastell R, Hofbauer L, Kanis JA, Langdahl BL, Lesnyak O, Lorenc R, McCloskey E, Messina OD, Napoli N, Obermayer-Pietsch B, Ralston SH, Sambrook PN, Silverman S, Sosa M, Stepan J, Suppan G, Wahl DA, Compston JE, for the Joint IOF-ECTS GIO Guidelines Working Group (2012) A framework for the development of guidelines for the management of glucocorticoid-induced osteoporosis. *Osteoporos Int* 23:2257–2276
  53. Dretakis KE, Dretakis EK, Papakitsou EF, Psarakis S, Steriopoulos K (1998) Possible predisposing factors for the second hip fracture. *Calcif Tissue Int* 62:366–369
  54. Lindsay R, Silverman SL, Cooper C, Hanley DA, Barton I, Broy SB, Licata A, Benhamou L, Geusens P, Flowers K, Stracke H, Seeman E (2001) Risk of new vertebral fracture in the year following a fracture. *JAMA* 285:320–323
  55. Ryg J, Rejnmark L, Overgaard S, Brixen K, Vestergaard P (2009) Hip fracture patients at risk of second hip fracture: a nationwide population-based cohort study of 169,145 cases during 1977–2001. *J Bone Miner Res* 24:1299–1307
  56. Banefelt J, Åkesson KE, Spångéus A, Ljunggren O, Karlsson L, Ström O, Orsäter G, Libanati C, Toth E (2019) Risk of imminent fracture following a previous fracture in a Swedish database study. *Osteoporos Int* 30:601–609
  57. Balasubramanian A, Zhang J, Chen L, Wenkert D, Daigle SG, Grauer A, Curtis JR (2019) Risk of subsequent fracture after prior fracture among older women. *Osteoporos Int* 30:79–92
  58. Kanis JA, Johansson H, Odén A, Harvey NC, Gudnason V, Sanders K, Sigurdsson G, Siggeirsdottir K, Borgström F, McCloskey EV (2018) Characteristics of recurrent fractures. *Osteoporos Int* 29:1747–1757
  59. Johansson H, Siggeirsdottir K, Harvey NC, McCloskey EV, Gudnason V, Sigurdsson G, Lorentzon M, Liu E, Kanis JA (2019) Adjusting FRAX for imminent risk. *Osteoporos Int*, abstract in press.
  60. Cosman F, Nieves JW, Dempster DW (2017) Treatment sequence matters: anabolic and antiresorptive therapy for osteoporosis. *J Bone Miner Res* 32(198–202):270
  61. Kanis JA, Cooper C, Rizzoli R, Abrahamsen B, Al-Daghri NM, Brandi ML, Cannata-Andia J, Cortet B, Dimai HP, Ferrari S, Hadji P, Harvey NC, Kraenzlin M, Kurth A, McCloskey E, Minisola S, Thomas T, Reginster JY (2017) Identification and management of patients at increased risk of osteoporotic fracture: outcomes of an ESCEO expert consensus meeting. *Osteoporos Int* 28:2023–2034
  62. Kanis JA, Cooper C, Rizzoli R, Reginster J-Y (2018) Review of the guideline of the American College of Physicians on the treatment of osteoporosis. *Osteoporos Int* 29:1505–1510
  63. Leder BZ, Tsai JN, Neer RM, Uihlein AV, Wallace PM, BurnettBowie SA (2016) Response to therapy with teriparatide, denosumab, or both in postmenopausal women in the DATA (denosumab and teriparatide administration) study randomized controlled trial. *J Clin Densitom* 19:346–351
  64. Bone HG, Cosman F, Miller PD, Williams GC, Hattersley G, Hu MY, Fitzpatrick LA, Mitlak B, Papapoulos S, Rizzoli R, Dore RK, Bilezikian JP, Saag KG (2018) ACTIVEExtend: 24 months of alendronate after 18 months of abaloparatide or placebo for postmenopausal osteoporosis. *J Clin Endocrinol Metab* 103:2949–2957

**Publisher's note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

## Affiliations

J. A. Kanis<sup>1,2</sup>  · N. C. Harvey<sup>3</sup> · E. McCloskey<sup>1,4</sup> · O. Bruyère<sup>5</sup> · N. Veronese<sup>6</sup> · M. Lorentzon<sup>2,7,8</sup> · C. Cooper<sup>3,9</sup> · R. Rizzoli<sup>10</sup> · G. Adib<sup>11</sup> · N. Al-Daghri<sup>12</sup> · C. Campusano<sup>13</sup> · M. Chandran<sup>14</sup> · B. Dawson-Hughes<sup>15</sup> · K. Javaid<sup>9</sup> · F. Jiwa<sup>16</sup> · H. Johansson<sup>1,2</sup> · J. K. Lee<sup>17</sup> · E. Liu<sup>2</sup> · D. Messina<sup>18</sup> · O. Mkinsi<sup>19</sup> · D. Pinto<sup>20,21</sup> · D. Prieto-Alhambra<sup>9,22</sup> · K. Saag<sup>23</sup> · W. Xia<sup>24</sup> · L. Zakraoui<sup>25</sup> · J. -Y. Reginster<sup>12,26</sup>

<sup>1</sup> Centre for Metabolic Bone Diseases, University of Sheffield Medical School, Beech Hill Road, S10 2RX Sheffield, UK

<sup>2</sup> Mary McKillop Health Institute, Australian Catholic University, Melbourne, Australia

<sup>3</sup> MRC Lifecourse Epidemiology Unit, University of Southampton, Southampton, UK

<sup>4</sup> MRC and Arthritis Research UK Centre for Integrated Research in Musculoskeletal Ageing Mellanby, Sheffield, UK

<sup>5</sup> World Health Organization Collaborating Center for the Public Health Aspects of Musculoskeletal Health and Aging, Department of Public Health, Epidemiology and Health Economics, University of Liège, Liège, Belgium

<sup>6</sup> National Research Council, Neuroscience Institute, Aging Branch, Via Giustiniani, 2, 35128 Padova, Italy

<sup>7</sup> Geriatric Medicine, Department of Internal Medicine and Clinical Nutrition, Institute of Medicine and Clinical Nutrition, Sahlgrenska Academy, University of Gothenburg, Gothenburg, Sweden

<sup>8</sup> Region Västra Götaland, Geriatric Medicine Clinic, Sahlgrenska University Hospital, Gothenburg, Sweden

<sup>9</sup> NIHR Musculoskeletal Biomedical Research Unit, University of Oxford, Oxford, UK

<sup>10</sup> Service of Bone Diseases, Geneva University Hospitals and Faculty of Medicine, Geneva, Switzerland

<sup>11</sup> Syrian National Osteoporosis Society, Damascus, Syria

<sup>12</sup> Chair for Biomarkers of Chronic Diseases, Biochemistry Dept., College of Science, King Saud University, Riyadh, Kingdom of Saudi Arabia

<sup>13</sup> Clinica Universidad de los Andes and Faculty of Medicine, Universidad de los Andes, Santiago, Chile

<sup>14</sup> Osteoporosis and Bone Metabolism Unit, Department of Endocrinology, Singapore General Hospital, ACADEMIA, 20, College Road, Singapore 169856, Singapore

- <sup>15</sup> Jean Mayer USDA Human Nutrition Research Center on Aging at Tufts University, Boston, MA, USA
- <sup>16</sup> Chair of the Committee of Patients Societies at the International Osteoporosis Foundation, Osteoporosis Canada, Toronto, Canada
- <sup>17</sup> Beacon International Specialist Centre, Petaling Jaya, Malaysia
- <sup>18</sup> IRO Medical Research Center, Buenos Aires and Rheumatology section, Cosme Argerich, Buenos Aires, Argentina
- <sup>19</sup> Department of Rheumatology, Ibn Rochd University Hospital, Casablanca, Morocco
- <sup>20</sup> Department of Physical Therapy, Marquette University, Milwaukee, WI, USA
- <sup>21</sup> Department of Medical Social Sciences, Feinberg School of Medicine, Northwestern University, Chicago, IL, USA
- <sup>22</sup> GREMPAL Research Group, CIBERFes and Idiap Jordi Gol, Instituto de Salud Carlos III and Universitat Autònoma de Barcelona, Barcelona, Spain
- <sup>23</sup> University of Alabama at Birmingham, Birmingham, AL, USA
- <sup>24</sup> Department of Endocrinology, Key Laboratory of Endocrinology, Ministry of Health, Peking Union Medical College Hospital, Chinese Academy of Medical Sciences & Peking Union Medical College, Shuaifuyuan No. 1, Wangfujing, Dongcheng District, Beijing 100730, China
- <sup>25</sup> Service de Rhumatologie, University Tunis Manar and Hôpital Mongi-Slim, la Marsa, Tunisia
- <sup>26</sup> Department of Public Health, Epidemiology and Health Economics, University of Liège, Liège, Belgium