



Published in final edited form as:

Environ Sci Pollut Res Int. 2020 February ; 27(6): 6476–6487. doi:10.1007/s11356-019-07066-x.

Insecticide and Metal Exposures are Associated with a Surrogate Biomarker for Non-alcoholic Fatty Liver Disease in the National Health and Nutrition Examination Survey 2003–2004

Banrida Wahlang^{1,2}, Savitri Appana^{3,†}, Keith C. Falkner⁴, Craig J. McClain^{1,2,4,5}, Guy Brock^{3,‡}, Matthew C. Cave^{1,2,4,5,*}

¹UofL Superfund Research Center, University of Louisville, Louisville, KY, USA

²Department of Pharmacology and Toxicology, School of Medicine, University of Louisville, Louisville, KY, USA

³Department of Bioinformatics and Biostatistics, School of Public Health, University of Louisville, Louisville, KY, USA

⁴Department of Medicine, Division of Gastroenterology and Hepatology, Department of Medicine, School of Medicine, University of Louisville, Louisville, KY, USA

⁵The Robley Rex Veterans Affairs Medical Center

Abstract

Non-alcoholic fatty liver disease (NAFLD), the most common form of liver disease, affects over 30% of the US population. Our group and others have previously demonstrated that low-level environmental pollutant exposures were associated with increased odds ratios for unexplained alanine aminotransferase (ALT) elevation, a surrogate biomarker for NAFLD, in the adult National Health and Nutrition Examination Survey (NHANES). However, recently, more sensitive and lower ALT cut-offs have been proposed. The objective of this observational study is to utilize these ALT cut-offs to determine new associations between environmental chemicals and the surrogate NAFLD biomarker. Adult NHANES 2003–2004 participants without viral hepatitis, hemochromatosis, or alcoholic liver disease were analyzed in this cross-sectional study. ALT elevation was defined as >30 IU/L in men and >19 IU/L in women. Odds ratios adjusted for potential confounders for ALT elevation were determined across exposure quartiles for 17 pollutant subclasses comprised of 111 individual pollutants. The overall prevalence of ALT elevation was 37.6%. Heavy metal and organochlorine insecticide subclasses were associated with dose-dependent increased adjusted odds ratios for ALT elevation of 1.6 (95% CI 1.2–2.3) and 3.5 (95% CI 2.3–5.5) respectively, for the highest vs. lowest exposure quartiles ($p_{\text{trend}} < 0.01$). Within these subclasses, increasing whole blood levels of lead and mercury, and lipid-adjusted serum levels of dieldrin, and the chlordane metabolites, heptachlor epoxide and trans-nonachlor, were associated with increased odds ratios for ALT elevation. In conclusion, organochlorine insecticide,

*To whom correspondence should be addressed: Matthew C. Cave, M.D., Kosair Charities Clinical & Translational Research Building, 505 South Hancock Street, Louisville, KY, 40202, Phone: (502) 852-6189, Fax: (502) 852-8927, matt.cave@louisville.edu.

†Current affiliation: Department of Biostatistics, MD Anderson Cancer Center, Houston, TX, USA

‡Current affiliation: Department of Biomedical Informatics, The Ohio State University College of Medicine, Columbus, OH, USA

Competing Interests Declaration: The authors have no competing interests to disclose.

lead, and mercury exposures were associated with ALT elevation and suspected NAFLD in adult NHANES 2003–2004.

Keywords

insecticides; metals; ALT; toxicant-associated steatohepatitis (TASH); non-alcoholic fatty liver disease (NAFLD); NHANES

Introduction

Non-alcoholic fatty liver disease (NAFLD), and its more advanced form, non-alcoholic steatohepatitis (NASH), are currently the most common forms of liver disease in the United States with a prevalence of approximately 30% in the general population (Le et al. 2017). NAFLD is a spectrum of disorders in the liver ranging from lipid accumulation (steatosis) to steatohepatitis, which is often accompanied by inflammation and hepatocyte death (Aguilera-Mendez 2019). NASH may progress to cirrhosis with severe clinical sequela including hepatocellular carcinoma, death, and liver transplantation. Non-invasive biomarkers of NAFLD and NASH, among others, include elevated activity levels of liver enzymes namely alanine aminotransferase (ALT) and aspartate transaminase (AST); as well as investigational biomarkers such as cytokeratin 18 (Cave et al. 2011, Zhou et al. 2019).

In general, in randomly selected individuals without a clinical diagnosis of NAFLD, ALT activity level greater than two standard deviations of the mean value of 100 is considered the upper limit of normal laboratory references. Previously, numerous epidemiologic studies on the National Health and Nutrition Examination Survey (NHANES) data adopted the following criteria for this range: >40 IU/L for men and >31 IU/L for women, to estimate the prevalence of liver disease in the general US population (Cave et al. 2010a, Clark 2006, Clark et al. 2003, Lazo et al. 2008, Liangpunsakul & Chalasani 2004, 2005). Unexplained ALT elevation which is liver enzyme elevation that is not due to either viral hepatitis, excessive ethanol consumption, or iron overload has become an accepted surrogate biomarker for NAFLD in NHANES studies (Clark et al. 2003). Prati *et al.* performed a four-year, comprehensive study to define ALT cutoffs for NAFLD diagnosis by eliminating participants who had ultrasound readings suggestive of fatty liver, and then re-evaluated ALT ranges in that population (Prati et al. 2002). These ALT reference ranges, namely >30 IU/L for men and >19 IU/L for women, were much lower than previously used ranges and have been increasingly recognized by many research groups (Dunn et al. 2008, Kwo et al. 2017, Martin-Rodriguez et al. 2017, Takyar et al. 2017). Therefore, it becomes important to consider these revised ALT cut-offs in environmental liver disease population studies.

Historically, most cases of NAFLD have been attributed to obesity, insulin resistance and the metabolic syndrome. With time, an increasing number of studies have reported that exposure to several industrial chemicals and environmental toxicants is associated with NAFLD and NASH, and termed as toxicant-associated steatohepatitis (TASH) (Al-Eryani et al. 2015, Bassler et al. 2019, Cave et al. 2010b, Guardiola et al. 2016, Wahlang et al. 2013). Additionally, utilizing previous ALT reference ranges: 48 IU/L for men and 31 IU/L for women (< 21 years of age); Cave *et al.* demonstrated that chronic low-level exposures to

environmental toxicants were dose-dependently associated with increased odds ratios for ALT elevation and suspected NAFLD in the general US adult population (NHANES 2003–2004) (Cave et al. 2010a). Specifically, polychlorinated biphenyl (PCB) exposures were implicated, as well as lead and mercury. In addition, Serdar *et al.* confirmed that liver enzymes levels were significantly higher in the highest exposure groups of PCBs and organochlorine pesticides in this population (NHANES 2003–2004). Moreover, additional epidemiological and toxicological studies have confirmed the reported association between environmental chemical exposures, primarily PCBs, and NAFLD (Clair et al. 2018, Wahlang et al. 2016, Wahlang et al. 2014). However, to date there is still limited epidemiologic information on associations between organochlorine insecticide/pesticide exposures and NAFLD.

Although ALT is a commonly used biomarker for liver disease in epidemiologic and clinical studies, its reported sensitivity for fatty liver diseases is somewhat ambiguous, especially at higher cutoff values (Hadizadeh et al. 2017). In fact, in some patients with biopsy-proven NAFLD, ALT may be normal, and ALT levels do not always correlate with disease severity (Wong et al. 2009). Furthermore, in fatty liver related to industrial chemicals, ALT sensitivity could be limited (Brautbar & Williams 2002, Cave et al. 2010b). We therefore hypothesized that the higher ALT cut-offs used previously may have reduced the number of cases of suspected NAFLD, and this may, in turn, have reduced the power to detect industrial chemicals possibly associated with suspected NAFLD. Thus, the objective of the current study is to analyze associations between environmental chemical exposures and suspected NAFLD in adult NHANES 2003–2004 participants, using the lower ALT cut-offs proposed by Prati *et al.*, in an effort to detect additional chemicals dose-dependently associated with increased odds ratios for ALT elevation and suspected NAFLD.

Materials and Methods

Study Design and Participants

Adult participants from NHANES 2003–2004 were evaluated in this cross-sectional study which was approved by the University of Louisville Institutional Review Board. The following exclusion criteria were utilized: age <18 years, positive serum hepatitis B surface antigen, positive serum hepatitis C antibody, elevated transferrin saturation (>60% for males and >50% for females), and alcohol consumption 20 grams/day for males and 10 grams/day for females. The maximum final sample size was 4,582 and has been described previously.

Pollutants

All pollutant data posted by the National Center for Health Statistics (NCHS) prior to December 2008, were accessed and downloaded. This data included 196 pollutants from 17 subclasses. Subclasses comprised of serum or blood organochlorine insecticides; metals including lead, cadmium and mercury (total and inorganic); volatile organic compounds (VOCs), non-dioxin like PCBs; dioxin-like compounds including polychlorinated dibenzo-p-dioxins (PCDDs), polychlorinated dibenzofurans (PCDFs) and coplanar PCBs; perfluorinated chemicals (PFCs); polybrominated diphenyl ethers (PBDEs); and cotinine. In

addition, urinary total (elemental plus inorganic) mercury; heavy metals; total arsenic and speciated arsenics; polycyclic aromatic hydrocarbons; phthalates; organophosphate insecticides; perchlorates; environmental phenols; and iodine were also included. The full list of chemicals belonging to each of these 17 subclasses, as well as the laboratories conducting these measurements and their methods with lower limit of detection (LLOD) have been described previously (Cave et al. 2010a). Only pollutants with a 60% or greater detection rate (111 of 196 pollutants) were evaluated to avoid bias in estimation among those below the limit of detection. Importantly, not all pollutants were measured in every NHANES subject. For most pollutants subclasses, data were collected from a random one-third subsample of subjects (Cave et al. 2010a). Lipid (cholesterol and triglyceride)-and creatinine adjustment were appropriately made for pollutants as previously described (Cave et al. 2010a).

Outcome Variables and Statistical Methods

In the current study, elevated ALT was defined by the cutoffs proposed by Prati *et al.* (>30 IU/L for men and >19 IU/L for women) (Prati et al. 2002). The prevalence of ALT elevation was determined in 4,582 subjects from various demographic groups based on sex, age, race, and body mass index (BMI). Statistical significance was determined by the chi-square test.

Individuals are likely to be exposed to multiple pollutants within a subclass, thus the cumulative subclass measure was obtained by summing the ranks according to the magnitude of detectable levels of each pollutant within that subclass (Cave et al. 2010a). For each pollutant subclass, subjects were stratified into quartiles by their cumulative exposure rank with the first quartile representing subjects with the lowest levels. Multivariate-adjusted odds ratios for ALT elevation were then determined across increasing quartiles of chemical exposure using the 1st quartile as the reference group by using logistic regression models. Multiple pollutants have previously been associated with obesity and insulin resistance in NHANES, so the analysis was conducted with adjustments for both BMI and homeostasis model assessment of insulin resistance (HOMA-IR) (Lee et al. 2007a, Lee et al. 2007b). Fasting glucose and insulin were measured in only a subset of NHANES participants, and adjusting for obesity and HOMA- IR further reduces the sample size to 2,211 subjects. Because most of the pollutant subclasses were measured only in a subset of this sample, the maximum sample size used for determination of associations was further reduced (Cave et al. 2010a). Adjustments were also made for age, sex, race, and poverty income ratio (PIR). Estimates of the main results were calculated accounting for stratification and clustering, and adjusting for age, race, and PIR using a previously described logistic model (Cave et al. 2010a). *P*-values were determined both with ($p_{\text{trend-adj}}$) and without (p_{trend}) adjustment for multiple comparisons.

For subclasses yielding significant results, these analyses were repeated for the individual chemicals within that subclass. Subjects with detectable levels of individual pollutants were ranked and placed into quartiles, and compared with a reference group which consisted of either i) subjects with levels below the LLOD or ii) individuals in the first quartile of exposure level.

Finally, chi-squared tests were used to determine whether demographic characteristics (sex, race, age, and BMI) were associated with high levels of exposure to selected subclasses of compounds associated with elevated ALT. All statistical analyses were performed using the Survey procedures, SURVEYFREQ and SURVEYLOGISTIC, from SAS 9.1 (SAS Institute Inc., Cary, NC, USA). A p-value of 0.05 or less was used to determine statistical significance.

Results

Demographic Information

After applying the exclusion criteria, a total of 4,582 adult subjects remained, and their demographic information is presented in Table 1. The percentage of females was slightly higher than the percentage of males (52.2% vs. 47.8%, respectively). The mean age was 47.2 ± 21.2 years with a range from 18–85 years. Non-Hispanic Whites accounted for 72.3% of the population. Body weights, as defined by NIH guidelines, were fairly evenly distributed between normal weight (31.5%), overweight (34.3%), and obese (32.5%) with very few subjects being underweight (1.7%).

Prevalence of Unexplained ALT Elevation

Out of the 4,582 remaining adult subjects, 1561 subjects had unexplained ALT elevation or suspected NAFLD (Table 1). After incorporating the NHANES sampling weights for these individuals, this corresponded to 37.6% of the adult United States population, or 68.9 million people. As shown in Table 1, ALT elevation was slightly more common in females than males (42.9% vs. 31.8%, $p=0.002$). Compared to non-Hispanic Whites, ALT elevation was more common in Hispanics, while non-Hispanic Blacks had a lower prevalence of ALT elevation (37.8% vs. 47.7% vs. 28.4%, $p<0.001$). Moreover, ALT elevation was most prevalent during the 5th and 6th decades ($p=0.022$). With regards to body weight, compared to normal weight participants, ALT elevation was more prevalent in overweight and obese participants (25.3% vs. 38.9% vs. 49.0%, respectively, $p<0.001$).

Pollutant Subclass Results

Measures of association were based on a maximum of the 2,211 participants with HOMA-IR scores as described previously (Cave et al. 2010a). Out of the 17 NHANES categorized pollutant subclasses investigated, only two were associated with significant dose-dependent increased adjusted odds ratios for ALT elevation (Table 2). These subclasses included (i) serum lipid-adjusted organochlorine insecticides/pesticides and (ii) blood lead, mercury, and cadmium. For these classes, the adjusted odds ratios for the highest exposure quartile, compared to the lowest, were 3.5 (95% CI 2.3–5.5, $p_{\text{trend-adj}}<0.001$) and 1.6 (95% CI 1.2–2.3, $p_{\text{trend-adj}}=0.015$) respectively.

Individual Pollutant Results

Eleven individual pollutants from these two subclasses were subsequently analyzed (Table 3). Whole blood lead (99.6%) and total mercury (92.5%) had extremely high detection rates and also demonstrated increased adjusted odds ratios for ALT elevation (lead: $OR_{\text{adj}}=3.3$ comparing 4th quartile vs. unexposed, 95% CI 0.3–34.2, $p_{\text{trend-adj}}=0.026$; and mercury:

OR_{adj}=2.2 comparing 4th quartile vs. unexposed, 95% CI 1.4–3.3, $p_{\text{trend-adj}} < 0.001$). Cadmium, the third member of this subclass, was not associated with increased odds ratios for ALT elevation.

Eight organochlorine insecticides (or their metabolites) were present at detection rates ranging from 61.7%–99.9%, and three of these were associated with increased adjusted odds ratios for ALT elevation (Table 3). These included the bio-accumulated chlordane constituents, heptachlor epoxide and trans-nonachlor, as well as the related insecticide dieldrin. The adjusted odds ratios for the highest exposure quartile compared to unexposed subjects were 3.1 (95% CI 1.3–5.0, $p_{\text{trend}}=0.001$), 1.6 (95% CI 0.6–3.8, $p_{\text{trend}}=0.05$), and 2.6 (95% CI 1.3–7.2, $p_{\text{trend}}=0.007$), respectively, for these chemicals. After adjusting for multiple comparisons, trans-nonachlor was no longer statistically significant ($p_{\text{trend-adj}}=0.093$). All five of the other insecticides analyzed from this class showed a trend toward increased odds ratios for ALT elevation across quartiles ($p_{\text{trend-adj}} < 0.095$). These included the chlordane metabolite, oxychlordane (OR_{adj}=2.7 comparing 4th quartile vs. unexposed, 95% CI 0.9–8.2, $p_{\text{trend}}=0.053$); dichlorodiphenyldichloroethylene (DDE) (OR_{adj}=1.7 comparing 4th quartile vs. unexposed, 95% CI 0.9–2.9, $p_{\text{trend}}=0.062$); hexachlorobenzene (OR_{adj}=1.4 comparing 4th quartile vs. unexposed, 95% CI 1.0–2.0, $p_{\text{trend}}=0.075$); hexachlorocyclohexane (OR_{adj}=1.7 comparing 4th quartile vs. unexposed, 95% CI 0.9–3.5, $p_{\text{trend}}=0.082$), and dichlorodiphenyltrichloroethane (DDT) (OR_{adj}=1.2 comparing 4th quartile vs. unexposed, 95% CI 0.7–1.9, $p_{\text{trend}}=0.095$). Of the eight analyzed insecticides, the DDT metabolite, DDE, had the highest median lipid-adjusted serum concentration (1535 ng/g) in the 4th quartile.

Next, demographic risk factors for organochlorine insecticide and metal exposures were determined. Both older age and higher BMI, but neither race/ethnicity nor sex, were associated with increased odds for having lipid-adjusted organochlorine insecticide levels in the highest quartile (Table 4). Age had the most pronounced effect, with 58.4% of subjects aged 70 years and older having insecticide levels in the highest quartile, compared to only 1.0% of subjects age <30 years old ($p < 0.001$). A greater percentage of obese subjects (25.7%) were in the highest insecticide exposure quartile compared to normal weight subjects (11.5%) ($p=0.002$). Demographic factors associated with higher whole blood metal levels included older age, having a lower BMI, and being a non-Hispanic Black male (Table 5). Again, age seemed to have the most pronounced effect, with only 8.7% of subjects <30 years old falling in the highest blood metal concentration quartile, compared to 40.1% of subjects aged 70 years and older. However, in contrast to organochlorine insecticide exposure, the percentage of subjects in the highest quartile of blood metal concentration were subjects with lower BMI (30.3% for lean subjects compared to 19.4% for obese subjects, $p=0.004$).

Discussion

Using the ALT reference range proposed by Prati *et al.*, the prevalence of suspected NAFLD was 37.6% in adult NHANES 2003–2004, compared to 10.6% in our prior study (Cave *et al.* 2010a). Although this prevalence value is much higher, it is however in line with current estimates for the prevalence of suspected NAFLD in western countries and worldwide (20–

30%), as well as reported prevalence in the U.S. population (30–46%) (Le et al. 2017, Perumpail et al. 2017, Ruhl & Everhart 2015, Williams et al. 2011). With regards to demographics, suspected NAFLD was more common with females (sex); Hispanics (race/ethnicity); middle age; and with increasing BMI. These results were concordant with our previous study. However, the race/ethnicity differences were lower in the current study when compared to our previous study which used higher ALT cut-offs. In the current study, when compared to Non-Hispanic White participants, Non-Hispanic Blacks and Other groups had 16% and 12% lower levels of NAFLD prevalence, respectively, while Hispanics had 24% higher levels of NAFLD prevalence. In the previous study, Non-Hispanic Blacks and Other populations had 44% and 1% lower levels of NAFLD respectively and Hispanics had 86% higher levels of NAFLD when compared to Non-Hispanic White participants. This observation is broadly consistent with the distribution of loss of function alleles of the NASH gene, namely, papatin-like phospholipase A 3 (PNPLA3) within groups that identified with Hispanic ethnicity; and supported the concept that PNPLA3 gene dysfunction can worsen NAFLD and NASH (Hernaes et al. 2013, Rotman et al. 2010).

With regards to environmental pollutants, the present analysis demonstrated that lead and mercury exposures were associated with dose-dependent increased adjusted odds ratios for ALT elevation which was consistent with our earlier reports (Cave et al. 2010a). The biologic plausibility that these metals could play a causal role in the development of liver diseases including NAFLD have also been reported by other groups (Kang et al. 2013, Lin et al. 2014, Yorita Christensen et al. 2013). On the contrary, the present study identified that organochlorine insecticides were associated with significantly increased odds ratios for ALT elevation which was not observed when using the higher ALT cut-offs. Nonetheless, in the previous study, the odds ratio for ALT elevation was higher in the third quartile of organochlorine insecticide exposure compared to unexposed subjects, but this trend did not reach statistical significance (Cave et al. 2010a). However, due to the low number of cases of ALT elevation noted (63/587), the first analysis could have been underpowered to detect a significant difference. In comparison, 202 cases of ALT elevation occurred in the present analysis in the organochlorine insecticide group, and this may have potentially increased statistical power.

Organochlorine insecticides are categorized as persistent organic pollutants due to their intact chemical integrity and resistance to degradation and metabolism. These lipid-soluble and poorly metabolized chemicals tend to bioaccumulate in the adipose tissue and other organs of living organisms including marine wildlife and humans (Kim et al. 2014, Pedro et al. 2017). Indeed, the current study demonstrated that obese subjects, who are at risk for NAFLD, also have the highest levels of serum lipid-adjusted organochlorine insecticides. Organochlorine insecticides include DDT and its metabolite, DDE; cyclodienes such as technical chlordane - a synthetic mixture comprising of over 50 chemicals with major components (60–85%) including *cis*- and *trans*-chlordane, hexachlor, heptachlor and trans-nonachlor, and their bioaccumulated metabolites (heptachlor epoxide and oxychlordane). Additional compounds in this class include dieldrin, aldrin and endrin as well as caged structures such as mirex and chlordane. Because of the toxic impact of DDT on the ecosystem, particularly wildlife endangerment (Carson & Darling 1962); the chlorinated cyclodienes were produced commercially as alternatives to DDT. These chemicals were

officially banned in 2001 at the Stockholm Convention on Persistent Environmental Pollutants (Porta & Zumeta 2002). Importantly, even though these insecticides have been banned from commercial production and use, they continue to persist in the environment (Turusov et al. 2002). For example, although DDT was banned from use in the US in 1972, 99.7% of NHANES subjects had detectable levels of the DDT metabolite, DDE. Furthermore, the organochlorine insecticides analyzed in this study continue to contaminate the US food supply, and US DDT per capita daily consumption was estimated to be greater than 250 ng/day (Schechter et al. 2010). Also, there are still considerable serum levels of chlordane metabolites including oxychlordane and trans-nonachlor in the US population according to the *Fourth National Report on Human Exposure to Environmental Chemicals* published by the Centers for Disease Control and Prevention (CDC), utilizing the NHANES datasets (1999–2004 and 2005–2010) (Anonymous 2018). Notably, in the current study, increased pesticide body burden was associated with increased age, implicating that bioaccumulation occurred over time.

An appreciable number of studies have reported associations of organochloride insecticide exposures with neurotoxicity, cancer, as well as negative effects on endocrine and metabolic health (De Coster & van Larebeke 2012, Kamel et al. 2007, Lee et al. 2007a, Lee et al. 2007b, Wahlang 2018). Organochlorine insecticides also have well-documented hepatotoxicity, especially at high doses (Al-Eryani et al. 2015, Freire et al. 2015, Ogata & Izushi 1991, Reuber 1978b, a). Therefore, serum organochlorine insecticides could be previously unrecognized mediators of liver disease in the general US adult population. While high-level exposures (e.g. acute poisonings) to these chemicals are sufficient to cause liver disease, the current data implicated that chronic low-level exposures, particularly occurring on a background of obesity or genetic susceptibility, may also contribute to the genesis and progression of liver diseases including NAFLD. Indeed, epidemiologic findings from human observational studies and toxicological studies using animal models have demonstrated that organochlorines, particularly chlordane and its metabolites are associated with metabolic disorders that encompass NAFLD (Ji et al. 2016, Liu et al. 2017b, Rosenbaum et al. 2017, Wang et al. 2017).

One of the most compelling data on the hepatic effects of organochlorine insecticide exposures in human subjects stemmed from an extremely high chlordecone exposure in 32 plant workers in Virginia (Guzelian et al. 1980). Although liver enzymes were repeatedly normal in these workers, many had hepatomegaly which eventually led to liver biopsy in 12 cases. These biopsies revealed mild steatosis, mild portal inflammation and fibrosis, glycogenated nuclei and lipofuscin accumulation. Additionally, organochlorine insecticides have also been associated with metabolic dysfunction including hyperglycemia, hypertriglyceridemia, diabetes and the metabolic syndrome in epidemiologic studies (Kim et al. 2014, Lee et al. 2007a, Lee et al. 2007b). Even though mechanisms of hepatotoxicity are not clearly defined in animal models, organochlorine insecticides have long been recognized as endocrine-disrupting chemicals through their interaction with steroid hormones receptors such as estrogen, androgen and thyroid receptors (De Coster & van Larebeke 2012). Organochlorine insecticides also induce hepatic expression of cytochrome P450s by activating xenobiotic receptors that can also influence hepatic energy metabolism such as the pregnane-xenobiotic receptor, constitutive androstane receptor and aryl hydrocarbon

receptor (De Coster & van Larebeke 2012, Kiyosawa et al. 2008). Emerging new hypotheses for organochlorine insecticide' mechanistic actions contributing to NAFLD development or progression include inhibition of the epidermal growth factor receptor (EGFR) signaling, similar to PCBs (Hardesty et al. 2018). It was recently demonstrated that chlordane and one of its most bioaccumulated component, trans-nonachlor, are potent EGFR inhibitors that diminished EGFR phosphorylation, a crucial step in maintenance of normal signaling processes (Hardesty et al. 2018). Another recently proposed mechanism of action for organochlorine insecticides is their ability to influence the gut-liver axis by modulating intestinal microbiota and altering bile acid metabolism (Liu et al. 2017a).

The current study is not without limitations and potential problems are inherent to the study design. Firstly, ALT levels in the general population may not rise to cross diagnostic thresholds in NAFLD, particularly in populations exposed to industrial chemicals. This is because the rise may either be transient and occurred early in the process or the entire exposed population may have lower ALT expression than unexposed individuals. Secondly, some of these chemicals have been proposed to be diet-induced obesogens that act in conjunction with high fat diet to mechanistically cause NAFLD (Wahlang et al. 2019), and adjusting for BMI may reduce the significance of association with ALT elevations. In the current cross-sectional analysis, higher BMI was associated with both suspected NAFLD and increased pesticide levels; and perhaps unadjusted BMI analysis may yield some insight into the potential obesogenic effect of these chemicals. However, this will need further investigation such as cohort studies to establish temporal trends between obesogen exposure and NAFLD/obesity incidences. Another limitation is that the 'true' normal range for ALT levels in adult NHANES is largely unknown, and no causal inferences can be obtained from the current study. Furthermore, age was not taken into consideration as a factor that could drive both increased chemical exposures as well as increased ALT elevation. It is well known that persistent chemicals bioaccumulate over time, and age often dictates length or duration of chronic chemical exposure. Additionally, age is known to influence NAFLD development and progression (Estes et al. 2018), and increased age could possibly subject the liver to higher risks of environment insults; therefore future studies should take this into consideration. However, a major strength of the study is that it utilized the lower ALT reference range suggested by Prati *et al.* which has been rigorously validated, albeit in a different population cohort.

Conclusion

In summary, the current study demonstrated that organochlorine insecticides/pesticides, lead, and mercury were present in the serum of nearly all US adults in 2003–2004. These common pollutants were associated with significant dose-dependent increased adjusted odds ratios for ALT elevation in subjects whose ALT elevations were not explained by viral hepatitis, hemochromatosis, or alcoholism. The results suggested a possible interaction between low-level environmental pollutant exposures and the development of liver disease and suspected NAFLD in the general US adult population, dependent on age and race. Importantly, the data from the current study contributes to the body of literature on the associations between insecticide exposures and liver disease and implicate that non-occupational, low-level exposures may pose an even bigger problem for liver disease in the

general US population. Future studies employing animal exposure models at doses relevant to human exposures are needed to confirm the causal role of these environmental pollutants, primarily organochlorine insecticides, in NAFLD initiation, development and progression.

Funding Support:

This research was supported in part by the National Institute of Environmental Health Sciences [R35ES028373, P42ES023716, T32ES011564]; the National Institute of General Medical Sciences [P20GM113226]; and the National Institute on Alcohol Abuse and Alcoholism [P50AA024337].

Abbreviations:

ALT	alanine aminotransferase
BMI	body mass index
CI	confidence intervals
DDE	dichlorodiphenyldichloroethylene
DDT	dichlorodiphenyltrichloroethane
EGFR	epidermal growth factor receptor
HOMA-IR	homeostasis model assessment of insulin resistance
LLOD	lower limit of detection
MeHg	methylmercury
NAFLD	nonalcoholic fatty liver disease
NASH	non-alcoholic steatohepatitis
NCHS	National Center for Health Statistics
NHANES	National Health and Nutrition Examination Survey
OR	odds ratio
PBDEs	polybrominated diphenyl ethers
PCBs	polychlorinated biphenyls
PCDDs	polychlorinated dibenzo-p-dioxins
PCDFs	polychlorinated dibenzofurans
PFCs	perfluorinated compounds
PIR	poverty income ratio
TASH	toxicant associated steatohepatitis

References

- Aguilera-Mendez A (2019): Nonalcoholic hepatic steatosis: a silent disease. *Rev Med Inst Mex Seguro Soc* 56, 544–549 [PubMed: 30889343]
- Al-Eryani L, Wahlang B, Falkner KC, Guardiola JJ, Clair HB, Prough RA, Cave M (2015): Identification of Environmental Chemicals Associated with the Development of Toxicant-associated Fatty Liver Disease in Rodents. *Toxicol Pathol* 43, 482–97. 10.1177/0192623314549960 [PubMed: 25326588]
- Anonymous (2018): Fourth National Report on Human Exposure to Environmental Chemicals Updated Tables, Volume Two. Centers for Disease Control and Prevention (CDC). U.S. Department of Health and Human Services
- Bassler J, Ducatman A, Elliott M, Wen S, Wahlang B, Barnett J, Cave MC (2019): Environmental perfluoroalkyl acid exposures are associated with liver disease characterized by apoptosis and altered serum adipocytokines. *Environ Pollut* 247, 1055–1063. 10.1016/j.envpol.2019.01.064 [PubMed: 30823334]
- Brautbar N, Williams J 2nd (2002): Industrial solvents and liver toxicity: risk assessment, risk factors and mechanisms. *Int J Hyg Environ Health* 205, 479–91. 10.1078/1438-4639-00175 [PubMed: 12455270]
- Carson R, Darling L (1962): *Silent spring*. Houghton Mifflin; Riverside Press, Boston; Cambridge, Mass.
- Cave M, Appana S, Patel M, Falkner KC, McClain CJ, Brock G (2010a): Polychlorinated biphenyls, lead, and mercury are associated with liver disease in American adults: NHANES 2003–2004. *Environ Health Perspect* 118, 1735–42. 10.1289/ehp.1002720 [PubMed: 21126940]
- Cave M, Falkner KC, Ray M, Joshi-Barve S, Brock G, Khan R, Bon Homme M, McClain CJ (2010b): Toxicant-associated steatohepatitis in vinyl chloride workers. *Hepatology* 51, 474–81. 10.1002/hep.23321 [PubMed: 19902480]
- Cave M, Falkner KC, Henry L, Costello B, Gregory B, McClain CJ (2011): Serum cytokeratin 18 and cytokine elevations suggest a high prevalence of occupational liver disease in highly exposed elastomer/polymer workers. *J Occup Environ Med* 53, 1128–33. 10.1097/JOM.0b013e31822cfd68 [PubMed: 21915069]
- Clair HB, Pinkston CM, Rai SN, Pavuk M, Dutton ND, Brock GN, Prough RA, Falkner KC, McClain CJ, Cave MC (2018): Liver Disease in a Residential Cohort With Elevated Polychlorinated Biphenyl Exposures. *Toxicol Sci* 164, 39–49. 10.1093/toxsci/kfy076 [PubMed: 29684222]
- Clark JM, Brancati FL, Diehl AM (2003): The prevalence and etiology of elevated aminotransferase levels in the United States. *Am J Gastroenterol* 98, 960–7. 10.1111/j.1572-0241.2003.07486.x [PubMed: 12809815]
- Clark JM (2006): The epidemiology of nonalcoholic fatty liver disease in adults. *J Clin Gastroenterol* 40 Suppl 1, S5–10. 10.1097/01.mcg.0000168638.84840.ff [PubMed: 16540768]
- De Coster S, van Larebeke N (2012): Endocrine-disrupting chemicals: associated disorders and mechanisms of action. *J Environ Public Health* 2012, 713696 10.1155/2012/713696 [PubMed: 22991565]
- Dunn W, Xu R, Wingard DL, Rogers C, Angulo P, Younossi ZM, Schwimmer JB (2008): Suspected nonalcoholic fatty liver disease and mortality risk in a population-based cohort study. *Am J Gastroenterol* 103, 2263–71. 10.1111/j.1572-0241.2008.02034.x [PubMed: 18684196]
- Estes C, Razavi H, Loomba R, Younossi Z, Sanyal AJ (2018): Modeling the epidemic of nonalcoholic fatty liver disease demonstrates an exponential increase in burden of disease. *Hepatology* 67, 123–133. 10.1002/hep.29466 [PubMed: 28802062]
- Freire C, Koifman RJ, Koifman S (2015): Hematological and hepatic alterations in Brazilian population heavily exposed to organochlorine pesticides. *J Toxicol Environ Health A* 78, 534–48. 10.1080/15287394.2014.999396 [PubMed: 25849770]
- Guardiola JJ, Beier JI, Falkner KC, Wheeler B, McClain CJ, Cave M (2016): Occupational exposures at a polyvinyl chloride production facility are associated with significant changes to the plasma metabolome. *Toxicol Appl Pharmacol* 313, 47–56. 10.1016/j.taap.2016.10.001 [PubMed: 27765658]

- Guzelian PS, Vranian G, Boylan JJ, Cohn WJ, Blanke RV (1980): Liver structure and function in patients poisoned with chlordecone (Kepone). *Gastroenterology* 78, 206–13. [PubMed: 6153073]
- Hadizadeh F, Faghihimani E, Adibi P (2017): Nonalcoholic fatty liver disease: Diagnostic biomarkers. *World J Gastrointest Pathophysiol* 8, 11–26. 10.4291/wjgp.v8.i2.11 [PubMed: 28573064]
- Hardesty JE, Al-Eryani L, Wahlang B, Falkner KC, Shi H, Jin J, Vivace BJ, Ceresa BP, Prough RA, Cave MC (2018): Epidermal Growth Factor Receptor Signaling Disruption by Endocrine and Metabolic Disrupting Chemicals. *Toxicol Sci* 162, 622–634. 10.1093/toxsci/kfy004 [PubMed: 29329451]
- Hernaez R, McLean J, Lazo M, Brancati FL, Hirschhorn JN, Borecki IB, Harris TB, Genetics of Obesity-Related Liver Disease C, Nguyen T, Kamel IR, Bonekamp S, Eberhardt MS, Clark JM, Kao WH, Speliotes EK (2013): Association between variants in or near PNPLA3, GCKR, and PPP1R3B with ultrasound-defined steatosis based on data from the third National Health and Nutrition Examination Survey. *Clin Gastroenterol Hepatol* 11, 1183–1190 e2. 10.1016/j.cgh.2013.02.011 [PubMed: 23416328]
- Ji G, Xu C, Sun H, Liu Q, Hu H, Gu A, Jiang ZY (2016): Organochloride pesticides induced hepatic ABCG5/G8 expression and lipogenesis in Chinese patients with gallstone disease. *Oncotarget* 7, 33689–702. 10.18632/oncotarget.9399 [PubMed: 27203212]
- Kamel F, Engel LS, Gladen BC, Hoppin JA, Alavanja MC, Sandler DP (2007): Neurologic symptoms in licensed pesticide applicators in the Agricultural Health Study. *Hum Exp Toxicol* 26, 243–50. 10.1177/0960327107070582 [PubMed: 17439927]
- Kang MY, Cho SH, Lim YH, Seo JC, Hong YC (2013): Effects of environmental cadmium exposure on liver function in adults. *Occup Environ Med* 70, 268–73. 10.1136/oemed-2012-101063 [PubMed: 23322921]
- Kim KS, Lee YM, Kim SG, Lee IK, Lee HJ, Kim JH, Kim J, Moon HB, Jacobs DR Jr., Lee DH (2014): Associations of organochlorine pesticides and polychlorinated biphenyls in visceral vs. subcutaneous adipose tissue with type 2 diabetes and insulin resistance. *Chemosphere* 94, 151–7. 10.1016/j.chemosphere.2013.09.066 [PubMed: 24161582]
- Kiyosawa N, Kwekel JC, Burgoon LD, Dere E, Williams KJ, Tashiro C, Chittim B, Zacharewski TR (2008): Species-specific regulation of PXR/CAR/ER-target genes in the mouse and rat liver elicited by o, p'-DDT. *BMC Genomics* 9, 487 10.1186/1471-2164-9-487 [PubMed: 18925944]
- Kwo PY, Cohen SM, Lim JK (2017): ACG Clinical Guideline: Evaluation of Abnormal Liver Chemistries. *Am J Gastroenterol* 112, 18–35. 10.1038/ajg.2016.517 [PubMed: 27995906]
- Lazo M, Selvin E, Clark JM (2008): Brief communication: clinical implications of short-term variability in liver function test results. *Ann Intern Med* 148, 348–52. [PubMed: 18316753]
- Le MH, Devaki P, Ha NB, Jun DW, Te HS, Cheung RC, Nguyen MH (2017): Prevalence of non-alcoholic fatty liver disease and risk factors for advanced fibrosis and mortality in the United States. *PLoS One* 12, e0173499 10.1371/journal.pone.0173499 [PubMed: 28346543]
- Lee DH, Lee IK, Jin SH, Steffes M, Jacobs DR Jr. (2007a): Association between serum concentrations of persistent organic pollutants and insulin resistance among nondiabetic adults: results from the National Health and Nutrition Examination Survey 1999–2002. *Diabetes Care* 30, 622–8. 10.2337/dc06-2190 [PubMed: 17327331]
- Lee DH, Lee IK, Porta M, Steffes M, Jacobs DR Jr. (2007b): Relationship between serum concentrations of persistent organic pollutants and the prevalence of metabolic syndrome among non-diabetic adults: results from the National Health and Nutrition Examination Survey 1999–2002. *Diabetologia* 50, 1841–1851. 10.1007/s00125-007-0755-4 [PubMed: 17624515]
- Liangpunsakul S, Chalasani N (2004): Relationship between unexplained elevations in alanine aminotransferase and serum leptin in U.S. adults: results from the Third National Health and Nutrition Examination Survey (NHANES III). *J Clin Gastroenterol* 38, 891–7. [PubMed: 15492608]
- Liangpunsakul S, Chalasani N (2005): Unexplained elevations in alanine aminotransferase in individuals with the metabolic syndrome: results from the third National Health and Nutrition Survey (NHANES III). *Am J Med Sci* 329, 111–6. [PubMed: 15767815]

- Lin YS, Ginsberg G, Caffrey JL, Xue J, Vulimiri SV, Nath RG, Sonawane B (2014): Association of body burden of mercury with liver function test status in the U.S. population. *Environ Int* 70, 88–94. 10.1016/j.envint.2014.05.010 [PubMed: 24908642]
- Liu Q, Shao W, Zhang C, Xu C, Wang Q, Liu H, Sun H, Jiang Z, Gu A (2017a): Organochloride pesticides modulated gut microbiota and influenced bile acid metabolism in mice. *Environ Pollut* 226, 268–276. 10.1016/j.envpol.2017.03.068 [PubMed: 28392238]
- Liu Q, Wang Q, Xu C, Shao W, Zhang C, Liu H, Jiang Z, Gu A (2017b): Organochloride pesticides impaired mitochondrial function in hepatocytes and aggravated disorders of fatty acid metabolism. *Sci Rep* 7, 46339 10.1038/srep46339 [PubMed: 28397872]
- Martin-Rodriguez JL, Gonzalez-Cantero J, Gonzalez-Cantero A, Arrebola JP, Gonzalez-Calvin JL (2017): Diagnostic accuracy of serum alanine aminotransferase as biomarker for nonalcoholic fatty liver disease and insulin resistance in healthy subjects, using 3T MR spectroscopy. *Medicine (Baltimore)* 96, e6770 10.1097/MD.0000000000006770 [PubMed: 28445310]
- Ogata M, Izushi F (1991): Effects of chlordane on parameters of liver and muscle toxicity in man and experimental animals. *Toxicol Lett* 56, 327–37. [PubMed: 2035178]
- Pedro S, Boba C, Dietz R, Sonne C, Rosing-Asvid A, Hansen M, Provatas A, McKinney MA (2017): Blubber-depth distribution and bioaccumulation of PCBs and organochlorine pesticides in Arctic-invasive killer whales. *Sci Total Environ* 601–602, 237–246. 10.1016/j.scitotenv.2017.05.193
- Perumpail BJ, Khan MA, Yoo ER, Cholankeril G, Kim D, Ahmed A (2017): Clinical epidemiology and disease burden of nonalcoholic fatty liver disease. *World J Gastroenterol* 23, 8263–8276. 10.3748/wjg.v23.i47.8263 [PubMed: 29307986]
- Porta M, Zumeta E (2002): Implementing the Stockholm Treaty on Persistent Organic Pollutants. *Occup Environ Med* 59, 651–2. [PubMed: 12356922]
- Prati D, Taioli E, Zanella A, Della Torre E, Butelli S, Del Vecchio E, Vianello L, Zanuso F, Mozzi F, Milani S, Conte D, Colombo M, Sirchia G (2002): Updated definitions of healthy ranges for serum alanine aminotransferase levels. *Ann Intern Med* 137, 1–10. [PubMed: 12093239]
- Reuber MD (1978a): Carcinomas and other lesions of the liver in mice ingesting organochlorine pesticides. *Clin Toxicol* 13, 231–56. 10.3109/15563657808988235 [PubMed: 216523]
- Reuber MD (1978b): Carcinogenicity testing of chemicals with particular reference to organochlorine pesticides. *Sci Total Environ* 10, 105–15. [PubMed: 212831]
- Rosenbaum PF, Weinstock RS, Silverstone AE, Sjodin A, Pavuk M (2017): Metabolic syndrome is associated with exposure to organochlorine pesticides in Anniston, AL, United States. *Environ Int* 108, 11–21. 10.1016/j.envint.2017.07.017 [PubMed: 28779625]
- Rotman Y, Koh C, Zmuda JM, Kleiner DE, Liang TJ, Nash CRN (2010): The association of genetic variability in patatin-like phospholipase domain-containing protein 3 (PNPLA3) with histological severity of nonalcoholic fatty liver disease. *Hepatology* 52, 894–903. 10.1002/hep.23759 [PubMed: 20684021]
- Ruhl CE, Everhart JE (2015): Fatty liver indices in the multiethnic United States National Health and Nutrition Examination Survey. *Aliment Pharmacol Ther* 41, 65–76. 10.1111/apt.13012 [PubMed: 25376360]
- Schechter A, Colacino J, Haffner D, Patel K, Opel M, Papke O, Birnbaum L (2010): Perfluorinated compounds, polychlorinated biphenyls, and organochlorine pesticide contamination in composite food samples from Dallas, Texas, USA. *Environ Health Perspect* 118, 796–802. 10.1289/ehp.0901347 [PubMed: 20146964]
- Takyar V, Nath A, Beri A, Gharib AM, Rotman Y (2017): How healthy are the “Healthy volunteers”? Penetrance of NAFLD in the biomedical research volunteer pool. *Hepatology* 66, 825–833. 10.1002/hep.29247 [PubMed: 28470683]
- Turusov V, Rakitsky V, Tomatis L (2002): Dichlorodiphenyltrichloroethane (DDT): ubiquity, persistence, and risks. *Environ Health Perspect* 110, 125–8. 10.1289/ehp.02110125 [PubMed: 11836138]
- Wahlang B, Beier JJ, Clair HB, Bellis-Jones HJ, Falkner KC, McClain CJ, Cave MC (2013): Toxicant-associated steatohepatitis. *Toxicol Pathol* 41, 343–60. 10.1177/0192623312468517 [PubMed: 23262638]

- Wahlang B, Song M, Beier JI, Cameron Falkner K, Al-Eryani L, Clair HB, Prough RA, Osborne TS, Malarkey DE, Christopher States J, Cave MC (2014): Evaluation of Aroclor 1260 exposure in a mouse model of diet-induced obesity and non-alcoholic fatty liver disease. *Toxicol Appl Pharmacol* 279, 380–90. 10.1016/j.taap.2014.06.019 [PubMed: 24998970]
- Wahlang B, Prough RA, Falkner KC, Hardesty JE, Song M, Clair HB, Clark BJ, States JC, Arteel GE, Cave MC (2016): Polychlorinated Biphenyl-Xenobiotic Nuclear Receptor Interactions Regulate Energy Metabolism, Behavior, and Inflammation in Non-alcoholic-Steatohepatitis. *Toxicol Sci* 149, 396–410. 10.1093/toxsci/kfv250 [PubMed: 26612838]
- Wahlang B (2018): Exposure to persistent organic pollutants: impact on women's health. *Rev Environ Health* 33, 331–348. 10.1515/reveh-2018-0018 [PubMed: 30110273]
- Wahlang B, Jin J, Beier JI, Hardesty JE, Daly EF, Schnegelberger RD, Falkner KC, Prough RA, Kirpich IA, Cave MC (2019): Mechanisms of Environmental Contributions to Fatty Liver Disease. *Curr Environ Health Rep* 6, 80–94. 10.1007/s40572-019-00232-w [PubMed: 31134516]
- Wang D, Wang X, Zhang P, Wang Y, Zhang R, Yan J, Zhou Z, Zhu W (2017): The fate of technical-grade chlordane in mice fed a high-fat diet and its roles as a candidate obesogen. *Environ Pollut* 222, 532–542. 10.1016/j.envpol.2016.11.028 [PubMed: 28041837]
- Williams CD, Stengel J, Asike MI, Torres DM, Shaw J, Contreras M, Landt CL, Harrison SA (2011): Prevalence of nonalcoholic fatty liver disease and nonalcoholic steatohepatitis among a largely middle-aged population utilizing ultrasound and liver biopsy: a prospective study. *Gastroenterology* 140, 124–31. 10.1053/j.gastro.2010.09.038 [PubMed: 20858492]
- Wong VW, Wong GL, Tsang SW, Hui AY, Chan AW, Choi PC, Chim AM, Chu S, Chan FK, Sung JJ, Chan HL (2009): Metabolic and histological features of non-alcoholic fatty liver disease patients with different serum alanine aminotransferase levels. *Aliment Pharmacol Ther* 29, 387–96. 10.1111/j.1365-2036.2008.03896.x [PubMed: 19035982]
- Yorita Christensen KL, Carrico CK, Sanyal AJ, Gennings C (2013): Multiple classes of environmental chemicals are associated with liver disease: NHANES 2003–2004. *Int J Hyg Environ Health* 216, 703–9. 10.1016/j.ijheh.2013.01.005 [PubMed: 23491026]
- Zhou JH, Cai JJ, She ZG, Li HL (2019): Noninvasive evaluation of nonalcoholic fatty liver disease: Current evidence and practice. *World J Gastroenterol* 25, 1307–1326. 10.3748/wjg.v25.i11.1307 [PubMed: 30918425]

Table 1.
Prevalence of Unexplained ALT Elevation in Adult NHANES 2003–2004 by Demographic Groups.

BMI - body mass index; SE - standard error of the mean. p -values were calculated by χ^2 tests.

Demographic Variable	Population Distribution (%)	Prevalence of Unexplained ALT Elevation (%)	SE	p -value
Sex				0.002
Male	47.8	31.9	1.6	
Female	52.2	42.9	1.5	
Race				<.001
Non-Hispanic White	72.3	37.8	0.8	
Non-Hispanic Black	10.8	28.4	1.9	
Hispanic	11.7	46.7	1.7	
Other Race	5.1	33.3	3.4	
Age (years)				0.022
< 30	21.5	30.1	1.9	
30–40	19.5	38.4	1.9	
40–50	20.4	44.0	2.4	
50–60	16.0	43.5	1.9	
60–70	10.5	41.4	2.5	
>= 70	12.1	28.1	2.4	
BMI (kg/m²)				<.001
< 18.5	1.7	23.0	5.7	
18.5–24.9	31.5	25.3	1.6	
25–29.9	34.3	38.9	1.1	
>= 30	32.5	49.0	1.5	

Table 2.
Adjusted* Odds Ratios (95% CI) for ALT Elevation by Exposure Quartile (With Number of Cases/Total Number) for 17 Pollutant Subclasses in Adult NHANES 2003–2004.

1st quartile: 25th percentile, 2nd quartile: 25th - 50th percentile, 3rd quartile: 50th - 75th percentile, 4th quartile: >75th percentile. Detectable values of each pollutant were individually ranked, and the rank orders of the individual pollutants in each subclass were summed to arrive at the subclass value. All non-detectable values were ranked as one. The summary values were categorized by cut-off points of 25th, 50th, and 75th values of the sum of ranks. CI - confidence interval; PCB - polychlorinated biphenyl.

Pollutant Subclass	Pollutants Screened/ Analyzed ^a					<i>P</i> _{trend}	<i>P</i> _{trend-adj} [†]
		1 st	2 nd	3 rd	4 th		
Organochlorine insecticides (serum) (n=587)	13/8	38/136 ^b	39/140	55/155	70/156	<.001	<.001
		Referent	1.1 (0.62,0) ^c	1.9 (1.3–2.7)	3.5 (2.35,5)		
Lead, cadmium, mercury (blood) (n=2051)	4/3	165/520	161/505	169/518	185/508	0.002	0.013
		Referent	1.1 (0.8–1.4)	1.3 (1.0–1.6)	1.6 (1.2–2.3)		
Heavy metals (urine) (n=709)	12/9	46/184	58/197	47/179	62/149	0.082	0.399
		Referent	1.0 (0.6–1.9)	0.8 (0.5–1.3)	1.8 (1.0–3.1)		
Cotinine (serum) (n=2050)	1/1	210/527	168/515	169/504	131/504	0.094	0.399
		Referent	0.8 (0.6–1.0)	0.9 (0.7–1.1)	0.7 (0.5–1.0)		
Total & speciated arsenics (urine) (n=710)	8/3	49/188	53/187	56/170	55/165	0.144	0.488
		Referent	1.0 (0.6–1.6)	1.2 (0.9–1.6)	1.2 (0.8–1.9)		
Total mercury (urine) (n=707)	1/1	44/180	47/167	52/169	68/191	0.202	0.571
		Referent	1.1 (0.6–2.1)	1.1 (0.7–1.7)	1.4 (0.9–2.3)		
Phthalates (urine) (n=655)	13/9	62/177	55/158	48/150	61/170	0.237	0.576
		Referent	0.9 (0.5–1.6)	0.7 (0.5–1.1)	0.8 (0.6–1.2)		
Organophosphate Insecticides (urine) (n=631)	6/4	65/162	55/159	47/159	58/151	0.429	0.686
		Referent	0.7 (0.4–1.3)	0.5 (0.3–1.0)	0.9 (0.5–1.5)		
Environmental phenols (urine) (n=643)	4/3	52/145	61/162	59/165	56/171	0.495	0.899
		Referent	1.1 (0.7–1.9)	0.9 (0.5–1.7)	0.9 (0.5–1.5)		
Volatile organic compounds (blood) (n=433)	33/6	37/108	43/97	45/111	32/117	0.529	0.899
		Referent	1.7 (0.83,6)	1.4 (0.7–2.9)	0.8 (0.4–1.6)		
Non-dioxin-like PCBs (serum) (n=532)	26/25	49/125	46/145	50/131	47/131	0.605	0.903
		Referent	1.0 (0.5–1.8)	1.2 (0.8–1.8)	1.2 (0.6–2.3)		
Polyaromatic hydrocarbons (urine) (n=563)	21/10	47/147	48/139	48/142	42/135	0.747	0.903
		Referent	1.0 (0.7–1.6)	0.9 (0.6–1.6)	0.9 (0.6–1.6)		
Iodine (urine) (n=645)	1/1	47/158	61/160	65/180	55/147	0.801	0.903
		Referent	1.2 (0.9–1.7)	1.0 (0.6–1.5)	1.0 (0.7–1.5)		
Perchlorate (urine) (n=638)	1/1	45/161	59/163	66/166	56/148	0.807	0.903
		Referent	1.3 (0.8–2.0)	1.4 (0.8–2.3)	1.1 (0.6–2.0)		

Pollutant Subclass	Pollutants Screened/ Analyzed ^a					P_{trend}	$P_{\text{trend-adj}}^{\ddagger}$
		1 st	2 nd	3 rd	4 th		
Perfluorinated compounds (serum) (n=694)	12/4	57/189	56/169	65/167	32/169	0.816	0.903
		Referent	1.2 (0.7–2.0)	1.9 (1.0–3.6)	0.7 (0.3–1.7)		
Polybrominated diphenyl ethers (serum) (n=614)	11/6	57/169	63/149	44/149	52/147	0.850	0.903
		Referent	1.4 (0.9–2.0)	0.8 (0.5–1.2)	1.1 (0.7–1.9)		
Dioxins, furans, coplanar PCBs (serum) (n=535)	29/17	49/138	44/136	49/135	48/126	0.962	0.962
		Referent	1.0 (0.6–1.6)	0.9 (0.4–2.5)	1.0 (0.3–2.8)		

* ORs were adjusted for age, sex, race, poverty income ratio, HOMA-IR, and BMI.

[‡] Additionally adjusted for multiple comparisons.

^a In any given pollutant subclass, only chemicals present with at least a 60% detection rate were included in the analysis.

^b Number of cases / total number.

^c Adjusted odds ratios with 95% confidence intervals.

Table 3.
Adjusted* Odds Ratios (95% CI) for ALT Elevation by Exposure Quartile (With Median Concentration Levels and Number of Cases/Total Number) for Pollutant Subclasses Lead, Cadmium, and Mercury and Organochlorine Insecticides in Adult NHANES 2003–2004.

1st quartile: 25th percentile, 2nd quartile: 25th - 50th percentile, 3rd quartile: 50th - 75th percentile, 4th quartile: >75th percentile. Lead, cadmium, and mercury were measured in whole blood. Organochlorine insecticides were measured in serum and are reported as lipid-adjusted values. For each pollutant, the reference group consisted of those individuals with values below the limit of detection for that pollutant, unless otherwise noted. CI - confidence interval.

Pollutant	Detection Rate (%)	Not Detectable	Detectable				P _{trend}	P _{trend-adj} [†]
			1 st	2 nd	3 rd	4 th		
Lead (ug/dL)	99.6	--	0.80 ^a	1.30	1.90	3.30	0.017	0.026
		1/6	176/579 ^b	162/494	205/498	136/474		
		Referent	2.5(0.2–24.4) ^c	3.3 (0.3–33.7)	5.1 (0.5–52.5)	3.3 (0.3–34.2)		
Cadmium (ug/L)	82.8	--	0.30	0.40	0.60	1.10	0.510	0.510
		117/345	241/672	82/257	129/406	111/371		
		Referent	1.1 (0.9–1.3)	0.9 (0.6–1.3)	0.9 (0.6–1.4)	0.9 (0.6–1.4)		
Mercury, total (ug/L)	92.5	--	0.40	0.80	1.40	3.10	<.001	<.001
		38/158	145/500	178/540	148/395	171/458		
		Referent	1.3 (0.9–1.9)	1.7 (1.1–2.6)	1.9 (1.3–3.0)	2.2 (1.4–3.3)		
Dieldrin Lipid Adj (ng/g)	88.5	--	3.90	5.90	8.20	14.15	0.007	0.027
		13/62	38/129	38/119	53/141	67/155		
		Referent	1.6 (0.7–3.5)	1.8 (0.9–3.6)	2.2 (1.1–4.4)	3.1 (1.3–7.2)		
Heptachlor Epoxide Lipid Adj (ng/g)	61.7	--	3.60	5.60	8.90	16.70	0.001	0.009
		56/214	29/82	35/112	44/106	48/97		
		Referent	1.4 (0.8–2.4)	1.3 (0.7–2.2)	1.9 (1.1–3.2)	2.6 (1.3–5.0)		
Trans-nonachlor Lipid Adj (ng/g)	93.9	--	5.90	12.85	27.50	57.50	0.050	0.093
		11/37	35/139	54/138	54/137	54/153		
		Referent	0.7 (0.4–1.3)	1.6 (0.8–3.2)	1.7 (0.6–4.6)	1.6 (0.6–3.8)		
B-hexachlorocyclohexane Lipid Adj (ng/g)	75.4	--	4.30	8.80	19.10	52.40	0.082	0.093
		36/146	33/97	56/133	43/117	43/116		
		Referent	1.6 (1.0–2.8)	2.3 (1.4–3.8)	1.8 (1.0–3.4)	1.7 (0.9–3.5)		
Hexachloroben zene Lipid Adj (ng/g)	99.9	--	9.90	13.75	17.60	24.60	0.075	0.093
		44/148	47/142	60/167	59/150			
		Referent	1.1 (0.8–1.6)	1.2 (0.9–1.6)	1.4 (1.0–2.0)			

Pollutant	Detection Rate (%)	Not Detectable	Detectable				P _{trend}	P _{trend-adj} [†]
			1 st	2 nd	3 rd	4 th		
Oxychlorane Lipid Adj (ng/g)	83.8	--	5.00	10.50	19.30	35.50	0.053	0.093
		25/106	40/117	42/123	61/142	44/125		
		Referent	1.8 (0.9–3.7)	2.1 (0.8–5.7)	3.5 (1.4–8.4)	2.7 (0.9–8.2)		
p,p'-DDE Lipid Adj (ng/g)	99.7	--	82.40	183.00	464.00	1535.0	0.062	0.093
		0/1	43/155	52/153	57/148	57/149		
		Referent	1.4 (0.8–2.4)	1.7 (1.0–2.8)	1.7 (0.9–2.9)			
p,p'-DDT Lipid Adj (ng/g)	79.0	--	3.50	5.10	7.70	19.30	0.095	0.095
		35/135	35/112	42/111	54/121	42/126		
		Referent	1.3 (0.7–2.4)	1.7 (0.8–3.8)	2.2 (1.5–3.3)	1.2 (0.7–1.9)		

* ORs were adjusted for age, sex, race, poverty income ratio, HOMA-IR, and BMI.

^a Median concentration levels.

^b Number of cases / total number.

^c Adjusted odds ratios with 95% confidence intervals.

[†] Additionally adjusted for multiple comparisons.

Table 4.
Demographic Variables Associated with Having Lipid-adjusted Serum Organochlorine Insecticide Levels in the Highest Exposure Quartile.

BMI - body mass index. *p*-values were calculated by χ^2 tests.

Demographic Variable	Population Distribution (%)	Insecticides > 75%-ile (%)	SE	<i>p</i> -value
Sex				0.406
Male	47.1	18.1	2.5	
Female	52.9	20.1	1.7	
Race				0.342
Non-Hispanic White	72.6	19.0	2.1	
Non-Hispanic Black	10.3	23.9	3.2	
Hispanic	11.4	15.8	3.6	
Other Race	5.7	19.0	3.6	
Age (years)				<.001
< 30	21.2	1.0	0.7	
30–40	20.5	3.5	1.6	
40–50	20.8	12.1	3.1	
50–60	15.4	30.1	5.5	
60–70	10.6	41.1	4.9	
>= 70	11.5	58.4	5.1	
BMI (kg/m²)				<.001
< 18.5	2.0	2.7	2.5	
18.5–24.9	31.7	11.5	2.3	
25–29.9	32.0	20.7	2.1	
>= 30	34.3	25.7	2.6	

Table 5.
Demographic Variables Associated with Having Whole Blood Lead, Cadmium, and Mercury Levels in the Highest Exposure Quartile.

BMI - body mass index. *p*-values were calculated by χ^2 tests.

Demographic Variable	Population Distribution (%)	Metals > 75%-ile (%)	SE	<i>p</i> -value
Sex				0.056
Male	47.5	25.9	1.5	
Female	52.5	21.7	1.8	
Race				<.001
Non-Hispanic White	72.4	22.0	1.7	
Non-Hispanic Black	10.9	31.1	2.9	
Hispanic	11.7	18.6	1.9	
Other Race	5.1	43.3	4.1	
Age (years)				<.001
< 30	21.6	8.7	1.4	
30–40	19.4	17.0	1.4	
40–50	20.2	22.4	1.9	
50–60	16.1	33.5	3.5	
60–70	10.4	35.5	2.9	
>= 70	12.2	40.1	3.2	
BMI (kg/m²)				0.004
< 18.5	1.7	30.3	6.1	
18.5–24.9	31.5	24.5	1.5	
25–29.9	34.2	26.9	1.5	
>= 30	32.6	19.4	1.7	