



Published in final edited form as:

*Bioorg Med Chem.* 2020 February 15; 28(4): 115237. doi:10.1016/j.bmc.2019.115237.

## Identification of Potent Pyrazole Based APELIN Receptor (APJ) Agonists

Sanju Narayanan<sup>†</sup>, Vineetha Vasukuttan<sup>†</sup>, Sudarshan Rajagopal<sup>‡</sup>, Rangan Maitra<sup>†</sup>, Scott P. Runyon<sup>\*,†</sup>

<sup>†</sup>Center for Drug Discovery, RTI International, Research Triangle Park, NC 27709, United States

<sup>‡</sup>Center for Pulmonary Vascular Disease, Duke University Medical Center, Durham, NC 27710, United States

### Abstract

The apelinergic system comprises the apelin receptor and its cognate apelin and elabela peptide ligands of various lengths. This system has become an increasingly attractive target for pulmonary and cardiometabolic diseases. Small molecule regulators of this receptor with good drug-like properties are needed. Recently, we discovered a novel pyrazole based small molecule agonist **8** of APJ ( $EC_{50} = 21.5 \mu\text{M}$ ,  $K_i = 5.2 \mu\text{M}$ ) through focused screening which was further optimized to initial lead **9** ( $EC_{50} = 0.800 \mu\text{M}$ ,  $K_i = 1.3 \mu\text{M}$ ). In our efforts to synthesize more potent agonists and to explore the structural features important for apelin receptor agonism, we carried out structural modifications at N1 of the pyrazole core as well as the amino acid side-chain of **9**. Systematic modifications at these two positions provided potent small molecule agonists exhibiting  $EC_{50}$  values of  $< 100 \text{ nM}$ . Recruitment of  $\beta$ -arrestin as a measure of desensitization potential of select compounds was also investigated. Functional selectivity was a feature of several compounds with a bias towards calcium mobilization over  $\beta$ -arrestin recruitment. These compounds may be suitable as tools for in vivo studies of APJ function.

### Graphical Abstract

\*Corresponding Author srnyon@rti.org. Tel: 919-316-3842. Fax: 919 541 8868.


#### X-RAY CRYSTALLOGRAPHIC DATA

Atomic coordinates for **43** and **44** have been deposited with the Cambridge Crystallographic Data Centre (deposition numbers 1946460, 1946461). Copies of the data can be obtained, free of charge, on application to CCDC, 12 Union Road, Cambridge, CB2 1EZ, UK [fax: +44(0)-1223-336033 or e-mail: deposit@ccdc.cam.ac.uk].

#### Declaration of interests

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

**Publisher's Disclaimer:** This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



## Keywords

Apelin; APJ; agonist; pyrazole; AGTRL1; APLNR

## Introduction

The apelinergic system comprises the endogenous apelin peptides and its cognate G-protein coupled receptor, APJ, encoded by the gene AGTRL1 (angiotensin receptor like 1).<sup>1</sup> The apelin receptor is considered physiologically important due to its presence across many species including human, monkey, pig, rodents, frog and zebrafish.<sup>2</sup> The apelin receptor is a 380 amino acid protein having ~33% sequence identity to the AT1 receptor.<sup>1</sup> Apelin-36 was the first endogenous peptide identified along with other smaller peptides (apelin-17, apelin-13, apelin-12, pyr-apelin-13) that were derived from the cleavage of paired basic residues (Arg-Arg and Arg-Lys) of the 77 amino acid precursor protein.<sup>3–7</sup> Known apelin peptides are readily hydrolyzed *in vivo* (half-life ~5 min).<sup>8</sup> These proteins belong to an understudied component of the renin-angiotensin system with angiotensin-converting-enzyme II (ACE II) being involved in degradation of apelin *in vivo*.<sup>8,9</sup> However, none of the known angiotensin receptor ligands or peptides activate apelin receptor.<sup>1</sup> Recently, Elabela/Toddler was identified as the second endogenous ligand of this receptor.<sup>10–12</sup> Identification of these two sets of endogenous ligands facilitated molecular characterization of apelin receptor. Apelin receptors are primarily coupled to inhibitory G<sub>αi</sub> proteins and their activation is associated with inhibition of forskolin-stimulated cAMP production.<sup>3</sup> Additionally, apelin receptor can also activate PLC mediated calcium flux through G<sub>αq</sub> coupling<sup>13,14</sup> and recruit β-arrestin that desensitizes the receptor utilizing clathrin mediated endocytosis.<sup>15</sup> Apelin and apelin receptor are widely expressed across tissues in both human and rodents. In rat CNS, apelin mRNA is expressed within the hypothalamus where it is co-expressed with vasopressin mRNA in paraventricular (PVN) and supraoptic nuclei (SON).<sup>16,17</sup> Peripherally, apelin receptor is expressed in many organs including kidney, lungs, adipose tissue, liver and heart.<sup>18–20</sup> Activation of the apelinergic system has many beneficial effects including protection against hypertensive disorders, excitotoxic neuronal damage, pulmonary arterial hypertension, metabolic syndrome and heart failure.<sup>21,22</sup> The receptor also plays an important role during embryonal development through regulation of vasculogenesis.<sup>23</sup>

Efforts are underway to produce small molecule agonists of apelin receptor with improved drug-like properties because the endogenous peptide ligands have short half-lives and limited utility as therapeutics. A number of small molecules have been reported. Iturrioz and colleagues reported a non-peptide functional apelin receptor agonist **E339-3D6** (Figure 1) that exhibited an  $EC_{50}$  of 90 nM (FRET) and  $K_i$  value of ~400 nM in both rat and human apelin receptor.<sup>24</sup>

The **ML233** core (Figure 1) ((E)-2-cyclohexyl-5-methyl-4-(phenylsulfonyloxyimino)cyclohexa-2,5-dienone) was identified through a high throughput screen of ~330 600 compounds from the NIH small molecule library.<sup>25</sup> **ML233** is a full agonist and exhibits an  $EC_{50}$  of 3.7  $\mu$ M ( $\beta$ -arrestin), but it displayed poor solubility and stability in microsomal fractions. In addition, **ML233** had significant off-target binding at several other receptors including the 5-HT<sub>1A</sub>,  $\alpha$ 2C adrenergic, mu-opioid, benzylpiperazine receptors and norepinephrine transporter. A number of patent disclosures have also emerged. Sanofi Aventis,<sup>26</sup> and Sanford-Burnham Medical Research Institute,<sup>27</sup> have disclosed small molecule agonists based on the substituted benzimidazole (US20140094450 A1) and triazole core (WO 2015/184011 A2) respectively (Figure 1). In addition, Bristol-Myers Squibb and Amgen have disclosed small molecules based on dihydropyrimidine-4-one core (WO 2017/096130A1)<sup>28</sup> and triazole core (WO 2017/192485A1)<sup>29</sup> respectively. A small molecule benzimidazole derivative related to the Sanofi Aventis scaffold **CMF-019** (Figure 1) was reported to be G protein biased small molecule apelin receptor agonist. **CMF-019** was ~400 fold bias towards G $\alpha$ i signaling over  $\beta$ -arrestin recruitment and ~6000 fold bias over receptor internalization. Studies using these biased agonists that preferentially stimulate G-protein activation over  $\beta$ -arrestin recruitment have demonstrated improved in vivo efficacy (vasodilation and inotropic actions) without  $\beta$ -arrestin dependent cardiac hypertrophy.<sup>30-32</sup>

Our laboratory has previously reported a novel pyrazole scaffold **8** ( $EC_{50}$  = 21.5  $\mu$ M (calcium),  $K_i$  = 5.2  $\mu$ M) identified through focused screening, which was further optimized to an initial lead **9** ( $EC_{50}$  = 0.800  $\mu$ M (calcium),  $K_i$  = 1.3  $\mu$ M) (Figure 2).<sup>33</sup> In our efforts to identify more potent agonists as tool compounds and to explore the structural features important for apelin receptor agonism, we carried out modifications at N1 position of the pyrazole core as well as the amino acid side-chain of the lead compound **9**. Synthesized compounds were characterized using calcium mobilization, inhibition of forskolin-induced cAMP accumulation and  $\beta$ -arrestin recruitment assays. Considering the potential importance of selectivity over the  $\beta$ -arrestin pathway,<sup>30,31</sup> bias factors for the most promising compounds were calculated.

## Chemistry

Scheme 1 depicts the synthetic route employed to prepare target compounds **9** – **38**. Synthesis of analogs **25** and **37** are shown as representative examples. 2, 6-Dimethoxy acetophenone **39** was condensed with diethyl oxalate to afford the sodium salt of diketone **40** in quantitative yield. Reaction of **40** with isobutylhydrazine or cyclopentylhydrazine trifluoroacetate in refluxing ethanol provided 1, 5-pyrazoles **41a**, **42a** and 1,3-pyrazoles **41b**, **42b** in a ratio of 4:1. Transformation of **41a** and **42a** to acids **43** and **44** was achieved using

alkaline hydrolysis with lithium hydroxide monohydrate. Coupling of carboxylic acids **43** and **44** with (*S*)-*tert*-butyl 3-amino-5-cyclohexylpentanoate<sup>33,34</sup> in the presence of benzotriazole-1-yl-oxy-tris(dimethylamino)-phosphonium hexafluorophosphate (BOP) afforded **45** and **46**. *Tert*-butyl esters **45** and **46** were deprotected using trifluoroacetic acid to provide acids **12** and **13**. Finally, carboxylic acids **12** and **13** were coupled to cyclobutylamine using BOP to obtain **25** and **37** respectively.

As shown in Scheme 1 the synthesis of compounds **41** and **42** provides a mixture of isomers at the N1 and N2 pyrazole positions in ratio 4:1 that are readily separable using normal phase chromatography. The structures of hydrolyzed carboxylic acids **43** and **44** were not previously described but were confirmed using X-Ray crystallography (Figures 3 & 4). The NMR shift assignments for **41** and **42** were used as a guide to identify the additional N1 vs N2 alkyl substituted pyrazoles synthesized in this study.

## Results and Discussion

Our laboratory has previously reported the novel pyrazole based small molecule agonist hit **8** ( $\text{Ca}^{2+}$   $\text{EC}_{50}$  = 21.5  $\mu\text{M}$ ,  $\text{K}_i$  = 5.2  $\mu\text{M}$ ) identified through focused screening that was further optimized to an initial lead **9** ( $\text{Ca}^{2+}$   $\text{EC}_{50}$  = 0.800  $\mu\text{M}$ , cAMP  $\text{EC}_{50}$  = 844 nM,  $\text{K}_i$  = 1.3  $\mu\text{M}$ ). In addition to improving potency and affinity, compound **9** exhibited significantly reduced off-target NTR2 activity and did not activate or inhibit the AT1 receptor (Table 1).<sup>33</sup> Considering the structural similarity with the Sanofi scaffold **7**, we hypothesized that modification of **9** at N1 position could potentially enhance apelin receptor potency. Substitution of 4-fluorophenyl in **9** with various alkyl or cycloalkyl groups were initially considered as these were envisioned to provide similar hydrophobicity and steric bulk (Table 1) as phenyl. In addition, these analogs would also help determine whether aromatic substitutions were optimal at the N1 position for apelin receptor potency with this scaffold. The agonist activities of synthesized analogs were evaluated using both calcium mobilization and cAMP generation functional assays. In addition to potency and efficacy, recruitment of  $\beta$ -arrestin was also evaluated.

### A. Modification at N1 pyrazole core of compound with cyclohexyl ethyl amino acid side chain:

The N1 methyl substituted analog **10** ( $\text{Ca}^{2+}$   $\text{EC}_{50}$  = 1.46  $\mu\text{M}$ , cAMP  $\text{EC}_{50}$  = 2.55  $\mu\text{M}$ ) exhibited approximately 2-fold reduced potency compared to the aromatic N1 phenyl substituent **9** ( $\text{Ca}^{2+}$   $\text{EC}_{50}$  = 0.800  $\mu\text{M}$ ). Increasing the chain length to three carbon propyl **11**, enhanced potency by 2-fold ( $\text{Ca}^{2+}$   $\text{EC}_{50}$  = 0.416  $\mu\text{M}$ ) compared to **9**. Adding bulk with addition of branched alkyl substitution (isobutyl) **12**, however, did not enhance potency ( $\text{Ca}^{2+}$   $\text{EC}_{50}$  = 0.649  $\mu\text{M}$ ). Further, increasing the hydrophobic volume to five carbon cyclopentyl **13** ( $\text{Ca}^{2+}$   $\text{EC}_{50}$  = 0.424  $\mu\text{M}$ ) and six carbon cyclohexyl **14** ( $\text{Ca}^{2+}$   $\text{EC}_{50}$  = 0.380  $\mu\text{M}$ ) retained the 2-fold enhanced potency. Extending the cyclohexyl ring by one methylene **15** ( $\text{Ca}^{2+}$   $\text{EC}_{50}$  >10  $\mu\text{M}$ ) significantly reduced the potency suggesting tolerance for alkyl/cycloalkyl moiety with optimum hydrophobic volume of 3–6 carbons. This was further confirmed by replacing 4-fluorophenyl with larger hydrophobic cyclooctyl moiety **16** ( $\text{Ca}^{2+}$   $\text{EC}_{50}$  >10  $\mu\text{M}$ ) that proved detrimental for activity. The agonist activities of these

compounds were also evaluated using cAMP assay to monitor the influence of these modifications on functional selectivity. Compounds **11** (cAMP EC<sub>50</sub> = 0.281 μM), **13** (cAMP EC<sub>50</sub> = 0.238 μM) and **14** (cAMP EC<sub>50</sub> = 0.518 μM) exhibited 2 to 3-fold enhanced potency compared to the N1 phenyl substituted lead compound **9** (cAMP EC<sub>50</sub> = 0.844 μM). By contrast, **12** had similar potency to **9** in the cAMP assay. Compounds **15** and **16** with bulky cyclohexyl methyl and cyclooctyl group were completely inactive. Recruitment of β-arrestin was also investigated as a potential measure of desensitization. Compounds **11**, **12**, **13**, **14** exhibited > 2 μM potencies in β-arrestin assay suggesting the possibility of reduced receptor internalization and desensitization with these analogs. These studies suggested that an aryl ring was not required at N1 position of the pyrazole core to retain APJ receptor activity. Compounds substituted with alkyl/cycloalkyl moieties and with optimum hydrophobic volumes (n-propyl, cyclopentyl, cyclohexyl) exhibited enhanced potency and preference towards calcium/cAMP signaling over β-arrestin recruitment.

## B. Modification of carboxylic acid moiety of **12**:

Next, we focused on optimizing the carboxyl acid of **12**. Simple conversion of carboxylic acid to primary amide **17** (Table 2) was detrimental for activity (Ca<sup>2+</sup> EC<sub>50</sub> > 5.75 μM). This suggested that either an acidic group or bulky amide substitution was necessary to retain activity. The first substituted amide compound **18** with glycine methyl ester displayed ~2-fold enhanced potency (Ca<sup>2+</sup> EC<sub>50</sub> = 0.288 μM) compared to the carboxylic acid derivative **12** (Ca<sup>2+</sup> EC<sub>50</sub> = 0.649 μM) suggesting substituted amides versus acids could also retain or enhance potency. The N-methyl analog **19** was synthesized to probe the importance of amide proton for receptor interaction. Compound **19** exhibited similar potency (Ca<sup>2+</sup> EC<sub>50</sub> = 0.237 μM) compared to the unsubstituted amide, suggesting the amide proton is not critical for activity through the calcium signaling pathway. In contrast, compound **19** exhibited 3-fold reduced potency (cAMP EC<sub>50</sub> = 1.85 μM) in cAMP assay confirming the importance of the amide N-H to engage this pathway. Similarly, conversion of the glycine ester **18** (Ca<sup>2+</sup> EC<sub>50</sub> = 0.288 μM) to glycine acid **20** displayed 6-fold reduced potency (Ca<sup>2+</sup> EC<sub>50</sub> = 1.85 μM) signifying a preference for the hydrophobic ester over the polar carboxylic acid group. Since esters are metabolically labile and easily hydrolyzed in vivo, we synthesized relatively stable analogs **21** and **23** with 2-butanol and 2-butanone substitution respectively. Interestingly, compound **21** and **23** displayed 14-fold (Ca<sup>2+</sup> EC<sub>50</sub> = 0.046 μM) and 4-fold (Ca<sup>2+</sup> EC<sub>50</sub> = 0.144 μM) enhanced potency compared to the carboxylic acid derivative **12** (Ca<sup>2+</sup> EC<sub>50</sub> = 0.649 μM). In addition, compound **25** with cyclobutyl amide substitution also exhibited ~8-fold enhanced potency (Ca<sup>2+</sup> EC<sub>50</sub> = 81 nM). Taken together, these data suggest the hydrophobic amide substitution is important to enhance agonist potency. However, replacing glycine methyl ester with the more stable ethyl ether **24** exhibited reduced potency (Ca<sup>2+</sup> EC<sub>50</sub> = 1.70 μM). Compounds **17-25** were also evaluated for their ability to inhibit forskolin stimulated cAMP accumulation. Compounds **21** (cAMP EC<sub>50</sub> = 0.118 μM), **22** (cAMP EC<sub>50</sub> = 0.157 μM), **23** (cAMP EC<sub>50</sub> = 0.246 μM) and **25** (cAMP EC<sub>50</sub> = 0.283 μM) exhibited 2 to 3-fold enhanced potency compared to the carboxylic acid analog **12** (cAMP EC<sub>50</sub> = 0.915 μM). Compounds **18**, **21**, **22**, **23**, **24**, **25** were also potent in β-arrestin recruitment and exhibited an EC<sub>50</sub> value < 0.5 μM. Thus, conversion of carboxylic acid to alkyl and cycloalkyl amides retained or enhanced potency for β-arrestin recruitment indicating a

preference for hydrophobic amides is favorable for activating this signaling pathway. Binding affinities of select compounds were also evaluated using [<sup>125</sup>I]-apelin-13 in radioligand displacement assay. High binding affinities of compounds **21** ( $K_i = 0.036 \mu\text{M}$ ), **22** ( $K_i = 0.038 \mu\text{M}$ ), **23** ( $K_i = 0.172 \mu\text{M}$ ), **25** ( $K_i = 0.054 \mu\text{M}$ ) indicated that these compounds competitively displaced [<sup>125</sup>I]-apelin-13 and were bound to the orthosteric site of apelin receptor (Table 2). Compounds **21**, **22**, **23**, **25** also did not activate AT1 receptor ( $\text{EC}_{50} > 10 \mu\text{M}$ ), exhibiting selectivity of cyclohexyl ethyl series over the off-target AT1 receptor.

### C. Modification of carboxylic acid moiety of **26**:

Early on in our program we had evaluated if phenyl ethyl could replace the cyclohexyl ethyl sidechain because structural modifications of an aryl ring appeared more synthetically feasible. Since we had identified amide substituents that greatly enhanced agonist potency such as cyclobutyl amide (**25**), we evaluated if potency for (**26**,  $\text{Ca}^{2+} \text{EC}_{50} > 10 \mu\text{M}$ ) could be rescued with a similar set of modifications of the carboxylic acid moiety as shown in Table 2. All of the new amides with the phenylethyl sidechain (Table 3) had significantly enhanced potency over the parent acid **26**, in keeping with the observations for the cyclohexyl ethyl analogs in Table 2. Compound **27** with glycine methyl ester exhibited > 400fold ( $\text{Ca}^{2+} \text{EC}_{50} = 0.021 \mu\text{M}$ ) enhanced potency compared to the carboxylic acid derivative **26** ( $\text{Ca}^{2+} \text{EC}_{50} > 10 \mu\text{M}$ ). In addition, compound **27** displayed > 100-fold functional selectivity toward calcium signaling over cAMP ( $\text{cAMP EC}_{50} = 3.09 \mu\text{M}$ ) and  $\beta$ -arrestin pathway ( $\beta\text{-arr EC}_{50} = 2.33 \mu\text{M}$ ). However, functional selectivity was lost when the amide proton was replaced with methyl in analog **28** ( $\text{Ca}^{2+} \text{EC}_{50} = 0.027 \mu\text{M}$ ,  $\text{cAMP EC}_{50} = 0.266 \mu\text{M}$ ). Similarly, analogs **29** ( $\text{Ca}^{2+} \text{EC}_{50} = 0.411 \mu\text{M}$ ,  $\text{cAMP EC}_{50} > 10 \mu\text{M}$ ,  $\beta\text{-arr EC}_{50} > 10 \mu\text{M}$ ), **30** ( $\text{Ca}^{2+} \text{EC}_{50} = 0.313 \mu\text{M}$ ,  $\text{cAMP EC}_{50} > 10 \mu\text{M}$ ,  $\beta\text{-arr EC}_{50} > 10 \mu\text{M}$ ), **31** ( $\text{Ca}^{2+} \text{EC}_{50} = 0.162 \mu\text{M}$ ,  $\text{cAMP EC}_{50} > 10 \mu\text{M}$ ,  $\beta\text{-arr EC}_{50} = 8.67 \mu\text{M}$ ) with ethyl ether, 2-butanol and 2-butanone substitution were also selectively potent in the calcium assay. The cyclobutyl amide compound **32**, however, had moderate potency ( $\text{Ca}^{2+} \text{EC}_{50} = 1.26 \mu\text{M}$ ,  $\text{cAMP EC}_{50} > 3.48 \mu\text{M}$ ) and did not demonstrate signaling bias. Interestingly, the cyclobutyl amide  $\beta$ -amino acid analog **34** displayed enhanced potency ( $\text{Ca}^{2+} \text{EC}_{50} = 0.083 \mu\text{M}$ ) and was also selective towards calcium signaling over cAMP pathway. Thus, elongation of the carboxylic acid by one methylene in **34** demonstrated signaling bias for calcium over cAMP compared to **32**. However, no signaling bias was observed in the cyclohexyl series with  $\beta$ -amino acid derivative **25**. Binding affinities of select compounds were also evaluated. Compounds **27** ( $K_i = 0.187 \mu\text{M}$ ), **28** ( $K_i = 0.134 \mu\text{M}$ ), **31** ( $K_i = 0.571 \mu\text{M}$ ), **34** ( $K_i = 0.449 \mu\text{M}$ ) competitively displaced [<sup>125</sup>I]-apelin-13. Compounds **27**, **28**, **31**, **34** were also selective over the off-target AT1 receptor (Table 3).

### D. Modification at N1 position of pyrazole of the compound with cyclohexyl ethyl carboxylic acid side chain:

As noted earlier in Table 1, modification at the N1 position of pyrazole core had been critical in enhancing potency of carboxylic acid scaffold. To monitor the effects of similar N1 substituents on cyclobutylamide scaffold, compounds **35** – **38** (Table 4) were synthesized. The cyclobutyl carboxamide scaffold was chosen for modification since it showed better



potency (Table 2) and was considered more stable than the ester or other alkyl moieties. At first, substitution at N1 with aromatic 4-fluorophenyl moiety **35** displayed moderate potency ( $\text{Ca}^{2+}$   $\text{EC}_{50}$  = 0.425  $\mu\text{M}$ ). However, alkyl and cycloalkyl substituents **36**, **25**, **37**, **38** significantly enhanced the potency of compounds compared to their carboxylic acid derivatives **11-14**. Specifically, compound **36** ( $\text{Ca}^{2+}$   $\text{EC}_{50}$  = 0.049  $\mu\text{M}$ ) and **37** ( $\text{Ca}^{2+}$   $\text{EC}_{50}$  = 0.070  $\mu\text{M}$ ) with propyl and cyclopentyl substitution displayed 8-fold and 6-fold enhanced potency compared to the corresponding carboxylic acid derivative **11** ( $\text{Ca}^{2+}$   $\text{EC}_{50}$  = 0.416  $\mu\text{M}$ ) and **13** ( $\text{Ca}^{2+}$   $\text{EC}_{50}$  = 0.424  $\mu\text{M}$ ). Thus, modification at N1 position resulted in potent compounds **25**, **36**, **37**, **38** that exhibited  $\text{EC}_{50}$  values  $\leq$  100 nM. However, these compounds were not functionally selective. Select potent compounds were also evaluated in radioligand binding assay. Compounds **25** ( $K_i$  = 0.054  $\mu\text{M}$ ), **36** ( $K_i$  = 0.060  $\mu\text{M}$ ), **37** ( $K_i$  = 0.059  $\mu\text{M}$ ), **38** ( $K_i$  = 0.147  $\mu\text{M}$ ) exhibited high binding affinities and competitively displaced [ $^{125}\text{I}$ ]-apelin-13.

**Human liver microsomal stability studies**—Our long-term goal is to develop potent, metabolically stable APJ biased agonists for chronic *in vivo* studies to determine the importance of individual signaling cascades. Early *in vitro* assessment of metabolic stability of compounds is necessary to determine possible *in vivo* metabolic liabilities. Liver microsomal assays that includes cytochrome P450 system are the primary assays used to evaluate the stability of compounds. A small set of potent compounds (Table 5) were therefore evaluated for their stability in human liver microsomes. Our lead compound **9** demonstrated excellent stability ( $t_{1/2}$  = 229 min,  $\text{CL}_{\text{INT}}$  = 5 mL/min/kg). All the other analogs tested except **31** ( $t_{1/2}$  = 34.71 min,  $\text{CL}_{\text{INT}}$  = 36 mL/min/kg) with 2-butanone substitution are having very poor stability as indicated by high clearance values. This may be because all the other analogs contain alkyl/cycloalkyl substitution at N1 that possibly undergo oxidative metabolism at  $\alpha$ -carbon attached to the N1 nitrogen. Excellent stability of compound **9** is possibly due to the presence of stable aromatic 4-fluorophenyl moiety at N1 pyrazole.

**Effects of Modifications on Ligand Bias**—As noted earlier, small molecule apelin receptor agonists have been generated that are G protein biased and result in increased cardiac contractility *in vivo*.<sup>30</sup> We tested the effect of these site modifications on bias relative to the reference endogenous agonist Pyr-Apelin-13 by calculating bias factors<sup>35,36</sup> between assays for G protein signaling (calcium and cAMP) and  $\beta$ -arrestin recruitment (Table 5). In general, two patterns were observed (Figure 5): compounds that displayed bias towards calcium signaling compared to cAMP inhibition and  $\beta$ -arrestin recruitment (Figure 5A) and compounds that did not display significant bias ( $<$  10-fold) between pathways (Figure 5B). For those compounds that displayed significant bias between calcium signaling and cAMP or  $\beta$ -arrestin, bias factors generally ranged between 2 and 3 on a log scale (corresponding to 100- to 1000-fold bias towards calcium signaling).

## Conclusions

Systematic modifications at N1 position of the pyrazole core as well as the amino acid side chain resulted in potent small molecule agonists exhibiting  $\text{EC}_{50}$  values of  $\leq$  100 nM.

Functional selectivity was a feature of several compounds with a bias towards calcium mobilization over  $\beta$ -arrestin recruitment. Modification at N1 position of the pyrazole core resulted in compounds **11**, **13**, **14** (n-propyl, cyclopentyl, cyclohexyl substitutions) that exhibited enhanced potency and functional selectivity towards calcium/cAMP over  $\beta$ -arrestin signaling. Conversion of the carboxylic acid to alkyl and cycloalkyl amides were well tolerated and enhanced potency. Compounds **21**, **22**, **25** with cyclohexyl ethyl side chain exhibited potency 100 nM in calcium assay were equally potent towards other signaling cascades (cAMP and  $\beta$ -arrestin). Modification at carboxylic acid portion of compounds having a phenylethyl side chain resulted in compounds **29**, **30**, **31** that displayed significant bias toward calcium signaling over  $\beta$ -arrestin cascade (bias factors between 2 (100-fold) and 3 (1000-fold) on a log scale). Similarly, compounds **27–31**, **34** exhibited significant bias toward calcium signaling over cAMP cascade. Among the set of biased compounds, compound **31** displayed better metabolic stability ( $t_{1/2}$  = 34.71 min,  $CL_{INT}$  = 36 mL/min/kg) and also did not activate the off-target AT1 receptor. Interestingly, elongation of the carboxylic acid by one methylene ( $\beta$ -amino acid) in **34** demonstrated signaling bias for calcium over cAMP compared to the one methylene short carboxylic acid derivative **32**. However, no signaling bias was observed in cyclohexyl series with  $\beta$ -amino acid **25**. Modification at N1 position of the pyrazole resulted in compounds **25**, **36**, **37**, **38** that exhibited high potency across all the three signaling cascade. Specifically, compound **37** with N1 cyclopentyl substitution exhibited the highest potency (cAMP  $EC_{50}$  = 0.097  $\mu$ M) inhibiting forskolin-induced cAMP accumulation. Compound **37** (cAMP  $EC_{50}$  = 0.097  $\mu$ M) also exhibited ~10-fold enhanced potency compared to the starting lead **9** (cAMP  $EC_{50}$  = 0.844  $\mu$ M). Lipinski's "Rule of 5" have been useful guides for predicting oral absorption of compounds. The calculated properties of **37** (clogP 5.5, 2 donors, 5 acceptors, TPSA of 94.5 and a MW of 550.7) suggested that this compound might potentially be orally bioavailable. In conclusion, we have identified potent small molecule apelin receptor agonist **37** including several others that can be suitable as tool compounds for in vivo studies of apelin receptor upon additional characterization.

## Experimental Section:

Reagents and starting materials were obtained from commercial suppliers and were used without purification. Reactions were conducted under  $N_2$  atmosphere using oven-dried glassware. All solvents and chemicals used were reagent grade. Anhydrous tetrahydrofuran (THF), dichloromethane (DCM), and *N,N*-dimethylformamide (DMF) were purchased from Fisher Scientific and used as such. Flash column chromatography was carried out using a Teledyne ISCO Combiflash Rf system and Rediseq Rf gold pre-packed HP silica columns. Purity and characterization of compounds were established by combination of HPLC, TLC, LC-MS, and NMR analytical technique described below.  $^1H$  NMR spectra were recorded on a Bruker Avance DPX300 (300 MHz) spectrometer using  $CHCl_3-d$ ,  $MeOH-d_4$ ,  $DMSO-d_6$  with tetramethylsilane (TMS) (0.00 ppm) as the internal reference. Chemical shifts are reported in ppm relative to the solvent signal and coupling constants (*J*) values are reported in Hertz (Hz). Thin-layer chromatography (TLC) was performed on precoated silica gel GF Uniplates from Analtech and spots were visualized with UV light or  $I_2$  detection or phosphomolybdic acid stain. Low-resolution mass spectra were obtained using a Waters



Alliance HT/Micromass ZQ system (ESI). Optical rotations were measured on an Auto Pol III polarimeter at the sodium D line. Analytical and preparative HPLC was performed on an automated Varian ProStar HPLC system equipped with a Xterra® C<sub>18</sub> RP18 (4.6 × 100 mm i. d., 3.5 μm) column, with a flow rate of 0.1 mL/min.; λ<sub>max</sub> = 254 nm; mobile phase A: H<sub>2</sub>O and mobile phase B: CH<sub>3</sub>CN. Purity of the target compounds was determined to be 95% by HPLC. Chemical names were generated using ChemDraw Ultra (CambridgeSoft, version 10.0).

### Procedure: Scheme 1

**Ethyl 4-(2,6-dimethoxyphenyl)-2,4-dioxobutanoate (40):** To a solution of sodium ethoxide (21% in EtOH) (5.4 mL, 14.37 mmol) was added dropwise a mixture of diethyl oxalate (1.85 mL, 13.690 mmol) and 2,6-dimethoxy acetophenone (2.45 g, 13.69 mmol) in anhydrous ethanol (15 mL). The resultant mixture was stirred at room temperature for 30 minutes, upon which yellow suspension formed. The reaction mixture was heated to reflux for 4 h. The reaction was cooled to room temperature. Ethanol was evaporated in vacuo. The resultant residue was triturated with diethyl ether (30 mL) and filtered to obtain sodium salt of ethyl 4-(2,6-dimethoxyphenyl)-2,4-dioxobutanoate as yellow solid (4.0 g, 97 %). MS (ESI) *m/z*: Calcd. for C<sub>14</sub>H<sub>16</sub>O<sub>6</sub> 280.09 [M]<sup>+</sup>, found 279.3 [M-H]<sup>-</sup>.

### **Ethyl 5-(2,6-dimethoxyphenyl)-1-isobutyl-1H-pyrazole-3-carboxylate (41a)**

**Step 1: Preparation of Isobutylhydrazine trifluoroacetate:** Isobutyraldehyde (1g, 13.87 mmol) and *tert*-butyl carbazate (1.8 g, 13.87 mmol) in methanol (25 mL) was stirred at room temperature for 1 h. The solvent was evaporated and the resulting solid was dried under in vacuo to give white solid of (*E*)-*tert*-butyl 2-(2-methylpropylidene)hydrazinecarboxylate in quantitative yield. Sodium cyanoborohydride (1.2 g, 20.13 mmol) was added portionwise to a mixture of the (*E*)-*tert*-butyl 2-(2-methylpropylidene)hydrazinecarboxylate (2.5 g, 13.42 mmol) in 75 % of aqueous acetic acid (25 mL) at room temperature. The resultant solution was stirred for 3 h at room temperature. The reaction mixture was neutralized with 1N NaOH, extracted with CH<sub>2</sub>Cl<sub>2</sub> (3 × 25 mL), washed with saturated NaHCO<sub>3</sub>, dried with Na<sub>2</sub>SO<sub>4</sub>, filtered, and evaporated to give *tert*-butyl 2-isobutylhydrazinecarboxylate as oil (2.40 g, 95 %). <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz) δ 0.93 (d, *J*=6.78 Hz, 6 H), 1.46 (s, 9 H), 1.64 – 1.82 (m, 1 H), 2.43 (br. s., 1 H), 2.67 (d, *J*=6.78 Hz, 2 H). MS (ESI) *m/z*: Calcd. for C<sub>9</sub>H<sub>20</sub>N<sub>2</sub>O<sub>2</sub> 188.15 [M]<sup>+</sup>, found 189.3 [M+H]<sup>+</sup>.

Trifluoroacetic acid (12 mL) was added dropwise to a solution of the *tert*-butyl 2-isobutylhydrazinecarboxylate (2.4 g, 12.75 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (12 mL). The reaction mixture was stirred at room temperature for 1.5 h. The solvent was evaporated to give the trifluoroacetate salt of the title compound as colorless oil in quantitative yield. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz) δ 1.04 (dd, *J*=9.04, 6.78 Hz, 6 H), 2.04 – 2.25 (m, 1 H), 3.02 (dd, *J*=6.97, 3.96 Hz, 2 H). MS (ESI) *m/z*: Calcd. for C<sub>4</sub>H<sub>12</sub>N<sub>2</sub> 88.10 [M]<sup>+</sup>, found 89.4 [M+H]<sup>+</sup>.

**Step 2: Preparation of ethyl 5-(2,6-dimethoxyphenyl)-1-isobutyl-1H-pyrazole-3-carboxylate (41a):** Sodium salt of ethyl 4-(2,6-dimethoxyphenyl)-2,4-dioxobutanoate (40) (1.2 g, 3.96 mmol) and isobutylhydrazine trifluoroacetate (0.962 g, 4.76 mmol) was mixed with glacial acetic acid (25 mL) and conc. HCl (0.6 mL). The reaction mixture was heated to

reflux for 3.5 h. After cooling, reaction mixture was poured into water (25 mL). The aqueous layer was extracted with CH<sub>2</sub>Cl<sub>2</sub> (3 × 30 mL) and the combined CH<sub>2</sub>Cl<sub>2</sub> layer was washed with saturated aqueous NaHCO<sub>3</sub>. The organic layer was then washed with saturated brine, dried over Na<sub>2</sub>SO<sub>4</sub>, followed by filtration. The solvent was evaporated in vacuo. The residue was purified by silica gel flash chromatography (0–40% EtOAc:Hex) to give major N<sub>1</sub>-substituted title compound as colorless oil (535 mg, 40 %) and less polar N<sub>2</sub>-substituted, 2-isobutyl-5-(2,6-dimethoxyphenyl)-1*H*-pyrazole-3-carboxylate as colorless oil (190 mg, 14 %).

**Ethyl 5-(2,6-dimethoxyphenyl)-1-isobutyl-1*H*-pyrazole-3-carboxylate (41a)**—<sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz) δ 0.72 (d, *J*=6.78 Hz, 6 H), 1.39 (t, *J*=7.15 Hz, 3 H), 2.10–2.24 (m, 1 H), 3.72 (d, *J*=6.0 Hz, 2 H), 3.74 (s, 6 H), 4.41 (q, *J*=7.16 Hz, 2 H), 6.62 (d, *J*=8.67 Hz, 2 H), 6.73 (s, 1 H), 7.38 (t, *J*=8.48 Hz, 1 H). MS (ESI) *m/z*. Calcd. for C<sub>18</sub>H<sub>24</sub>N<sub>2</sub>O<sub>4</sub> 332.17 [M]<sup>+</sup>, found 333.4 [M+H]<sup>+</sup>.

**Ethyl 5-(2,6-dimethoxyphenyl)-2-isobutyl-1*H*-pyrazole-3-carboxylate (41b)**—<sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz) δ 0.94 (d, *J*=6.78 Hz, 6 H), 1.37 (t, *J*=7.16 Hz, 3 H), 2.26 (m, 1 H), 3.76 (s, 6 H), 4.33 (q, *J*=7.16 Hz, 2 H), 4.44 (d, *J*=7.16 Hz, 2 H), 6.61 (d, *J*=8.67 Hz, 2 H), 6.92 (s, 1 H), 7.28 (t, *J*=8.48 Hz, 1 H). MS (ESI) *m/z*. Calcd. for C<sub>18</sub>H<sub>24</sub>N<sub>2</sub>O<sub>4</sub> 332.17 [M]<sup>+</sup>, found 333.3 [M+H]<sup>+</sup>.

**Ethyl 1-cyclopentyl-5-(2,6-dimethoxyphenyl)-1*H*-pyrazole-3-carboxylate (42a)**

**Step 1: Preparation of Cyclopentylhydrazine trifluoroacetate:** Cyclopentanone (2.0 g, 23.77 mmol) and *tert*-butyl carbazate (3.1 g, 23.77 mmol) in methanol (20 mL) was stirred at room temperature for 3 h. The solvent was evaporated and the resulting solid was dried under in vacuo to give white solid of *Tert*-butyl 2-cyclopentylidenehydrazinecarboxylate (4.6 g, 98 %). Sodium cyanoborohydride (2.1 g, 34.06 mmol) was added portionwise to a mixture of *tert*-butyl 2-cyclopentylidenehydrazinecarboxylate (4.5 g, 22.697 mmol) in 75 % of aqueous acetic acid (25 mL) at room temperature. The resultant solution was stirred for 3 h at room temperature. The reaction mixture was neutralized with 1N NaOH, extracted with CH<sub>2</sub>Cl<sub>2</sub> (3 × 30 mL), washed with saturated NaHCO<sub>3</sub>, dried with Na<sub>2</sub>SO<sub>4</sub>, filtered, and evaporated to give *tert*-butyl 2-cyclopentylhydrazinecarboxylate as oil (4.4 g, 97 %). <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz) δ 1.40–1.55 (m, 4 H), 1.46 (s, 9 H), 1.63–1.75 (m, 4 H), 3.45–3.58 (m, 1 H). MS (ESI) *m/z*. Calcd. for C<sub>10</sub>H<sub>20</sub>N<sub>2</sub>O<sub>2</sub> 200.15 [M]<sup>+</sup>, found 223.3 [M+Na]<sup>+</sup>.

Trifluoroacetic acid (24 mL) was added dropwise to a solution of the *tert*-butyl 2-cyclopentylhydrazinecarboxylate (4.4 g, 21.97 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (24 mL). The reaction mixture was stirred at room temperature for 2.5 h. The solvent was evaporated to give the trifluoroacetate salt of the title compound as colorless oil in quantitative yield. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz) δ 1.58 – 1.70 (m, 2 H), 1.72– 1.92 (m, 4 H), 1.95–2.09 (m, 2 H), 3.60–3.78 (m, 1 H). MS (ESI) *m/z*. Calcd. for C<sub>5</sub>H<sub>12</sub>N<sub>2</sub> 100.10 [M]<sup>+</sup>, found 101.2 [M+H]<sup>+</sup>.

**Step 2: Preparation of ethyl 1-cyclopentyl-5-(2,6-dimethoxyphenyl)-1*H*-pyrazole-3-carboxylate (42a):** Sodium salt of ethyl 4-(2,6-dimethoxyphenyl)-2,4-dioxobutanoate (40) (1.2 g, 3.96 mmol) and cyclopentylhydrazine trifluoroacetate (1.3 g, 5.95 mmol) was mixed

with glacial acetic acid (25 mL) and conc. HCl (0.6 mL). The reaction mixture was heated to reflux for 3 h. After cooling, reaction mixture was poured into water (50 mL). The aqueous layer was extracted with DCM (3 × 30 mL). The combined DCM layer was washed with saturated aqueous NaHCO<sub>3</sub>. The organic layer was then washed with saturated brine, and then dried over Na<sub>2</sub>SO<sub>4</sub>, followed by filtration. The solvent was evaporated in vacuo. The residue was purified by silica gel flash chromatography (0–40% EtOAc:Hex) to give N<sub>1</sub>-substituted title compound (1.0 g, 73 %, major product) as light yellow solid and less polar N<sub>2</sub>-substituted, 2-cyclopentyl-5-(2,6-dimethoxyphenyl)-1*H*-pyrazole-3-carboxylate as colorless oil (50 mg, 5 %).

**1-cyclopentyl-5-(2,6-dimethoxyphenyl)-1*H*-pyrazole-3-carboxylate (42a):** <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz) δ 1.38 (t, *J*=7.2 Hz, 3 H), 1.44 – 1.57 (m, 2 H), 1.84 – 1.98 (m, 4 H), 2.08 – 2.26 (m, 2 H), 3.74 (s, 6 H), 4.21–4.31 (m, 1 H), 4.39 (q, *J*=6.9 Hz, 2 H), 6.63 (d, *J*=8.3 Hz, 2 H), 6.69 (s, 1 H), 7.38 (t, *J*=8.48 Hz, 1 H). MS (ESI) *m/z*: Calcd. for C<sub>19</sub>H<sub>24</sub>N<sub>2</sub>O<sub>4</sub> 344.41 [M]<sup>+</sup>, found 345.0 [M+H]<sup>+</sup>.

**2-cyclopentyl-5-(2,6-dimethoxyphenyl)-1*H*-pyrazole-3-carboxylate (42b):** <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz) δ 1.36 (t, *J*=7.2 Hz, 3 H), 1.62 – 1.74 (m, 2 H), 1.85 – 1.98 (m, 2 H), 2.08 – 2.26 (m, 4 H) 3.77 (s, 6 H), 4.33 (q, *J*=7.16 Hz, 2 H), 5.68 (quin, *J*=7.49 Hz, 1 H), 6.62 (d, *J*=8.48 Hz, 2 H), 6.91 (s, 1 H), 7.27 (t, *J*=8.48 Hz, 1 H). MS (ESI) *m/z*: Calcd. for C<sub>19</sub>H<sub>24</sub>N<sub>2</sub>O<sub>4</sub> 344.41 [M]<sup>+</sup>, found 345.3 [M+H]<sup>+</sup>.

**5-(2,6-Dimethoxyphenyl)-1-isobutyl-1*H*-pyrazole-3-carboxylic acid (43):** Lithium hydroxide monohydrate (189 mg, 4.51 mmol) in 1 mL of water was added to a solution of ethyl 5-(2,6-dimethoxyphenyl)-1-isobutyl-1*H*-pyrazole-3-carboxylate (500 mg, 1.50 mmol) in MeOH (11 mL) and THF (2 mL). The mixture was stirred at room temperature for 18 h. The reaction mixture was concentrated to about half the volume and then extracted with ether (2 × 15 mL). The aqueous layer was acidified with 1 N HCl and extracted with CH<sub>2</sub>Cl<sub>2</sub> (3 × 25 mL). The combined organic layers were washed with water, brine and then dried with Na<sub>2</sub>SO<sub>4</sub>. The solvent was evaporated in vacuo to give the title compound as white solid (440 mg, 96 %). <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz) δ 0.74 (d, *J*=6.40 Hz, 6 H), 2.10–2.24 (m, 1 H), 3.72 (d, *J*=7.54 Hz, 2 H), 3.75 (s, 6 H), 6.63 (d, *J*=9.0 Hz, 2 H), 6.79 (s, 1 H), 7.40 (t, *J*=8.48 Hz, 1 H). MS (ESI) *m/z*: Calcd. for C<sub>16</sub>H<sub>20</sub>N<sub>2</sub>O<sub>4</sub> 304.14 [M]<sup>+</sup>, found 303.3 [M-H]<sup>-</sup>.

**1-Cyclopentyl-5-(2,6-dimethoxyphenyl)-1*H*-pyrazole-3-carboxylic acid (44):** Following the procedure described in **43**, compound **44** was obtained from ethyl 1-cyclopentyl-5-(2,6-dimethoxyphenyl)-1*H*-pyrazole-3-carboxylate. 98 % yield; white solid; <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz) δ 1.48 – 1.64 (m, 2 H), 1.84 – 2.02 (m, 4 H), 2.03 – 2.20 (m, 2 H), 3.75 (s, 6 H), 4.30 (quin, *J*=7.4 Hz, 1 H), 6.64 (d, *J*=8.3 Hz, 2 H), 6.76 (s, 1 H), 7.40 (t, *J*=8.3 Hz, 1 H). MS (ESI) *m/z*: Calcd. for C<sub>17</sub>H<sub>20</sub>N<sub>2</sub>O<sub>4</sub> 316.14 [M]<sup>+</sup>, found 317.2 [M+H]<sup>+</sup>, 315.4 [M-H]<sup>-</sup>.

**(*S*)-*Tert*-butyl 5-cyclohexyl-3-(5-(2,6-dimethoxyphenyl)-1-isobutyl-1*H*-pyrazole-3-carboxamido)pentanoate (45):** 5-(2,6-Dimethoxyphenyl)-1-isobutyl-1*H*-pyrazole-3-carboxylic acid (**43**) (40 mg, 0.13 mmol) was dissolved in THF (2 mL). To the solution was added BOP (58 mg, 0.13 mmol) and triethylamine (0.055 mL, 0.39 mmol). The resulting mixture was stirred at room temperature for 15 minutes. (*S*)-*Tert*-butyl 3-amino-5-

cyclohexylpentanoate (37 mg, 0.14 mmol) was added and stirred at room temperature for 3 h. THF was evaporated in vacuo, water was added to the residue and the aqueous layer was extracted with CH<sub>2</sub>Cl<sub>2</sub> (3 × 30 mL). The combined organic layers were washed with water, brine and then dried with Na<sub>2</sub>SO<sub>4</sub>. The solvent was evaporated in vacuo to give the crude residue. The residue was purified by silica gel flash chromatography (0–50 % EtOAc:Hex) to give (*S*)-*tert*-butyl 5-cyclohexyl-3-(5-(2,6-dimethoxyphenyl)-1-isobutyl-1*H*-pyrazole-3-carboxamido)pentanoate (**45**) as foam (65 mg, 91%). <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz) δ 0.75 (d, *J*=5.65 Hz, 6 H), 0.81–0.96 (m, 2 H), 1.10 – 1.36 (m, 6 H), 1.47 (s, 8 H), 1.61 – 1.73 (m, 8 H), 2.12 (dt, *J*=13.75, 7.06 Hz, 1 H), 2.54 (d, *J*=5.65 Hz, 2 H), 3.58 – 3.67 (m, 2 H), 3.72 (s, 3 H), 3.74 (s, 3 H), 4.31 – 4.42 (m, 1 H), 6.61 (d, *J*=8.67 Hz, 2 H), 6.69 (s, 1 H), 7.23 (s, 1 H), 7.37 (t, *J*=8.48 Hz, 1 H). MS (ESI) *m/z*. Calcd. for C<sub>31</sub>H<sub>47</sub>N<sub>3</sub>O<sub>5</sub> 541.35 [M]<sup>+</sup>, found 542.6 [M+H]<sup>+</sup>.

**(S)-Tert-butyl 5-cyclohexyl-3-(1-cyclopentyl-5-(2,6-dimethoxyphenyl)-1H-pyrazole-3-carboxamido)pentanoate (46):**

Following the procedure described in **45**, compound **46** was obtained using 1-cyclopentyl-5-(2,6-dimethoxyphenyl)-1*H*-pyrazole-3-carboxylic acid (**44**) and (*S*)-*tert*-butyl 3-amino-5-cyclohexylpentanoate. 87% yield; yellow solid; <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz) δ 0.81 – 0.96 (m, 2 H), 1.10 – 1.38 (m, 8 H), 1.48 (s, 9 H), 1.59 – 1.77 (m, 8 H), 1.81 – 1.96 (m, 4 H), 2.00 – 2.16 (m, 2 H), 2.55 (d, *J*=5.46 Hz, 2 H), 3.73 (s, 3 H), 3.74 (s, 3 H), 4.21 – 4.30 (m, 1 H), 4.30 – 4.43 (m, 1 H), 6.62 (d, *J*=8.48 Hz, 2 H), 6.67 (s, 1 H), 7.25 (br. s., 1 H), 7.37 (t, *J*=8.38 Hz, 1 H). MS (ESI) *m/z*. Calcd. for C<sub>32</sub>H<sub>47</sub>N<sub>3</sub>O<sub>5</sub> 553.35 [M]<sup>+</sup>, found 555.0 [M+H]<sup>+</sup>

**(S)-5-Cyclohexyl-3-(5-(2,6-dimethoxyphenyl)-1-isobutyl-1H-pyrazole-3-carboxamido)pentanoic acid (12):**

Trifluoroacetic acid (0.6 mL) was added dropwise to a solution of the (*S*)-*tert*-butyl 5-cyclohexyl-3-(5-(2,6-dimethoxyphenyl)-1-isobutyl-1*H*-pyrazole-3-carboxamido)pentanoate (**45**) (60 mg, 0.11 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (1.5 mL). The reaction mixture was stirred at room temperature for 2.5 h. The solvent was evaporated to give the trifluoroacetate salt of the title compound. 54 % yield; white solid; <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz) δ 0.74 (d, *J*=6.78 Hz, 6 H), 0.81–0.98 (m, 3 H), 1.11–1.40 (m, 6 H), 1.60 – 1.78 (m, 8 H), 2.02–2.17 (m, 1 H), 3.66 (d, *J*=7.54 Hz, 2 H), 3.74 (s, 6 H), 4.29–4.44 (m, 1 H), 6.62 (d, *J*=8.67 Hz, 2 H), 6.72 (s, 1 H), 7.38 (t, *J*=8.48 Hz, 1 H), 7.37–7.51 (m, 1 H). MS (ESI) *m/z*. Calcd. for C<sub>27</sub>H<sub>39</sub>N<sub>3</sub>O<sub>5</sub> 485.29 [M]<sup>+</sup>, found 486.5 [M+H]<sup>+</sup>.

**(S)-5-Cyclohexyl-3-(1-cyclopentyl-5-(2,6-dimethoxyphenyl)-1H-pyrazole-3-**

**carboxamido)pentanoic acid (13):** Following the procedure described in **12**, compound **13** was obtained using (*S*)-*tert*-butyl 5-cyclohexyl-3-(1-cyclopentyl-5-(2,6-dimethoxyphenyl)-1*H*-pyrazole-3-carboxamido)pentanoate (**46**). 94% yield; white solid; <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz) δ ppm 0.78 – 1.00 (m, 3 H), 1.09 – 1.40 (m, 6 H), 1.49 – 1.77 (m, 4 H), 1.71 (t, *J*=7.44 Hz, 4 H), 1.821.98 (m, 4 H), 1.99 – 2.13 (m, 2 H), 2.65 – 2.80 (m, 2 H), 3.73 (s, 3 H), 3.74 (s, 3 H), 4.21–4.33 (m, 2 H), 6.63 (d, *J*=8.48 Hz, 2 H), 6.68 (s, 1 H), 7.29 (s, 1 H), 7.38 (t, *J*=8.38 Hz, 1 H). MS (ESI) *m/z*. Calcd. for C<sub>28</sub>H<sub>39</sub>N<sub>3</sub>O<sub>5</sub> 497.29 [M]<sup>+</sup>, found 496.7[M-H]<sup>-</sup>

**(S)-N-(1-(Cyclobutylamino)-5-cyclohexyl-1-oxopentan-3-yl)-5-(2,6-dimethoxyphenyl)-isobutyl-1H-pyrazole-3-carboxamide (25)**: Following the procedure described in **45**, compound **25** was synthesized using (*S*)-5-cyclohexyl-3-(5-(2,6-dimethoxyphenyl)-1-isobutyl-1*H*-pyrazole-3-carboxamido)pentanoic acid (**12**) and cyclobutylamine. 61 % yield; white solid; <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz) δ 0.74 (d, *J*=6.40 Hz, 6 H), 0.80 – 0.98 (m, 3 H), 1.10 – 1.40 (m, 8 H), 1.61–1.72 (m, 5 H), 1.77 – 1.93 (m, 2 H), 2.05 – 2.17 (m, 1 H), 2.20–2.34 (m, 2 H), 2.52 (d, *J*=6.03 Hz, 2 H), 3.63 (d, *J*=7.16 Hz, 2 H), 3.73 (s, 3 H), 3.74 (s, 3 H), 4.19 – 4.29 (m, 1 H), 4.32 – 4.43 (m, 1 H), 6.62 (d, *J*=8.67 Hz, 2 H), 6.67–6.72 (s, 1 H), 6.69 (s, 1 H), 7.09 (d, *J*=8.67 Hz, 1 H), 7.37 (t, *J*=8.67 Hz, 1 H). MS (ESI) *m/z*: Calcd. for C<sub>31</sub>H<sub>46</sub>N<sub>4</sub>O<sub>4</sub> 538.35 [M]<sup>+</sup>, found 539.6 [M+H]<sup>+</sup>.

**(S)-N-(1-(Cyclobutylamino)-5-cyclohexyl-1-oxopentan-3-yl)-1-cyclopentyl-5-(2,6dimethoxyphenyl)-1H-pyrazole-3-carboxamide (37)**: Following the procedure described in **45**, compound **37** was synthesized using (*S*)-5-cyclohexyl-3-(1-cyclopentyl-5-(2,6dimethoxyphenyl)-1*H*-pyrazole-3-carboxamido)pentanoic acid (**13**) and cyclobutylamine. 75 % yield; white solid; <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz) δ 0.88 (d, *J*=11.30 Hz, 2 H), 1.12 – 1.39 (m, 6 H), 1.60 – 1.77 (m, 9 H), 1.80 – 1.97 (m, 6 H), 2.00–2.13 (m, 2 H), 2.18–2.32 (m, 2 H), 2.24 (d, *J*=7.16 Hz, 2 H), 2.53 (d, *J*=6.40 Hz, 2 H), 3.74 (s, 3 H), 3.73 (s, 3 H), 4.15 – 4.46 (m, 3 H), 6.63 (d, *J*=8.29 Hz, 2 H), 6.67 (s, 1 H), 6.75 (d, *J*=8.29 Hz, 1 H), 7.06 (d, *J*=9.04 Hz, 1 H), 7.38 (t, *J*=8.38 Hz, 1 H). MS (ESI) *m/z*: Calcd. for C<sub>32</sub>H<sub>46</sub>N<sub>4</sub>O<sub>4</sub> 550.35 [M]<sup>+</sup>, found 551.6 [M+H]<sup>+</sup>.

#### Preparation of Analogs:

##### **(S)-5-cyclohexyl-3-(5-(2,6-dimethoxyphenyl)-1-methyl-1H-pyrazole-3-carboxamido)pentanoic acid (10)**

**Step 1: Preparation of ethyl 5-(2,6-dimethoxyphenyl)-1-methyl-1H-pyrazole-3-carboxylate**: Sodium salt of ethyl 4-(2,6-dimethoxyphenyl)-2,4-dioxobutanoate (**40**) (1.5 g, 4.96 mmol) and commercially available, methylhydrazine (0.26 ml, 4.96 mmol), was mixed with glacial acetic acid (30 mL) and conc. HCl (0.6 mL). The reaction mixture was heated to reflux for 4 h. After cooling, reaction mixture was poured into water (30 mL). The aqueous layer was extracted with CH<sub>2</sub>Cl<sub>2</sub> (3 × 30 mL) and the combined CH<sub>2</sub>Cl<sub>2</sub> layer was washed with saturated aqueous NaHCO<sub>3</sub>. The organic layer was then washed with saturated brine, dried over Na<sub>2</sub>SO<sub>4</sub>, followed by filtration. The solvent was evaporated in vacuo and the residue was purified by silica gel flash chromatography (0–40 % EtOAc:Hex) to give the title compound as oil (160 mg, 11 %). <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz) δ 1.40 (t, *J*=7.0 Hz, 3 H), 3.72 (s, 3 H), 3.76 (s, 6 H), 4.42 (q, *J*=7.2 Hz, 2 H), 6.64 (d, *J*=8.3 Hz, 2 H), 6.79 (s, 1 H), 7.39 (t, *J*=8.3 Hz, 1 H). MS (ESI) *m/z*: Calcd. for C<sub>15</sub>H<sub>18</sub>N<sub>2</sub>O<sub>4</sub> 290.13 [M]<sup>+</sup>, found 291.2 [M+H]<sup>+</sup>.

**Step 2: Preparation of 5-(2,6-dimethoxyphenyl)-1-methyl-1H-pyrazole-3-carboxylic acid**: Following the procedure as described in **45**, compound 5-(2,6-dimethoxyphenyl)-1-methyl-1*H*-pyrazole-3-carboxylic acid was obtained using ethyl 5-(2,6-dimethoxyphenyl)-1-methyl-1*H*-pyrazole-3-carboxylate. 87 % yield; white solid; <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz) δ 3.73 (s, 3 H), 3.78 (s, 6 H), 6.65 (d, *J*=8.5 Hz, 2 H), 6.84 (s, 1 H), 7.40 (t, *J*=8.48 Hz, 1 H). MS (ESI) *m/z*: Calcd. for C<sub>13</sub>H<sub>14</sub>N<sub>2</sub>O<sub>4</sub> 262.10 [M]<sup>+</sup>, found 263.3 [M+H]<sup>+</sup>, 261.3 [M-H]<sup>-</sup>.

**Step 3: Preparation of (*S*)-*tert*-butyl 5-cyclohexyl-3-(5-(2,6-dimethoxyphenyl)-1-methyl-1*H*-pyrazole-3-carboxamido)pentanoate:** Following the procedure described in **45**, compound (*S*)-*tert*-butyl 5-cyclohexyl-3-(5-(2,6-dimethoxyphenyl)-1-methyl-1*H*-pyrazole-3-carboxamido)pentanoate was obtained using 5-(2,6-dimethoxyphenyl)-1-methyl-1*H*-pyrazole-3-carboxylic acid and (*S*)-*tert*-butyl 3-amino-5-cyclohexylpentanoate. 87% yield; yellow solid; <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz) δ 0.78–0.95 (m, 2 H), 1.08–1.32 (m, 7 H), 1.48 (s, 9 H), 1.52 – 1.78 (m, 8 H), 2.53 (dd, *J*=5.46, 3.01 Hz, 1 H), 3.65 (s, 3 H), 3.76 (s, 6 H), 6.63 (d, *J*=8.48 Hz, 2 H), 6.76 (s, 1 H), 7.31–7.33 (br. s., 1 H), 7.37 (t, *J*=8.38 Hz, 1 H). MS (ESI) *m/z*: Calcd. for C<sub>28</sub>H<sub>41</sub>N<sub>3</sub>O<sub>5</sub> 499.30 [M]<sup>+</sup>, found 501.0 [M+H]<sup>+</sup>.

**Step 4: Preparation of (*S*)-5-cyclohexyl-3-(5-(2,6-dimethoxyphenyl)-1-methyl-1*H*-pyrazole-3-carboxamido)pentanoic acid (**10**):** Using the procedure described in **12**, compound (*S*)-5-cyclohexyl-3-(5-(2,6-dimethoxyphenyl)-1-methyl-1*H*-pyrazole-3-carboxamido)pentanoic acid was obtained from (*S*)-*tert*-butyl 5-cyclohexyl-3-(5-(2,6-dimethoxyphenyl)-1-methyl-1*H*-pyrazole-3-carboxamido)pentanoate. 51 % yield; white solid; <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz) δ 0.80 – 0.97 (m, 2 H), 1.11 – 1.37 (m, 6 H), 1.59–1.78 (m, 7 H), 2.71 (d, *J*=5.09 Hz, 2 H), 3.66 (s, 3 H), 3.76 (s, 6 H), 4.29–4.42 (m, 1 H), 6.63 (d, *J*=8.48 Hz, 2 H), 6.78 (s, 1 H), 7.38 (t, *J*=8.48 Hz, 1 H), 7.39–7.43 (m, 1 H). MS (ESI) *m/z*: Calcd. for C<sub>24</sub>H<sub>33</sub>N<sub>3</sub>O<sub>5</sub> 443.24 [M]<sup>+</sup>, found 442.6 [M-H]<sup>-</sup>.

### Preparation of (*S*)-5-cyclohexyl-3-(5-(2,6-dimethoxyphenyl)-1-propyl-1*H*-pyrazole-3-carboxamido)pentanoic acid (**11**)

#### Step 1: Preparation of ethyl 5-(2,6-dimethoxyphenyl)-1-propyl-1*H*-pyrazole-3-carboxylate

**Step 1.1: Preparation of propylhydrazine trifluoroacetate :** Propionaldehyde (1g, 17.22 mmol) and *tert*-butyl carbazate (2.3 g, 17.22 mmol) in methanol (20 mL) was stirred at room temperature for 2 h. The solvent was evaporated and the resulting solid was dried under in vacuo to give white solid of (*E*)-*tert*-butyl 2-propyldiazene-carboxylate in quantitative yield. Sodium cyanoborohydride (1.5 g, 24.39 mmol) was added portionwise to a mixture of the (*E*)-*tert*-butyl 2-propyldiazene-carboxylate (2.8 g, 16.26 mmol) in 75 % of aqueous acetic acid (20 mL) at room temperature. The resultant solution was stirred for 3 h at room temperature. The reaction mixture was neutralized with 1N NaOH, extracted with CH<sub>2</sub>Cl<sub>2</sub> (3 × 25 mL), washed with saturated NaHCO<sub>3</sub>, dried with Na<sub>2</sub>SO<sub>4</sub>, filtered, and evaporated to give title compound as oil in quantitative yield. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz) δ 0.92 (t, *J*=9.0 Hz, 3 H), 1.46 (s, 9 H), 1.44–1.55 (m, 2 H), 2.62 (bs, 2 H).

Trifluoroacetic acid (11 mL) was added dropwise to a solution of the *tert*-butyl 2-propylhydrazine-carboxylate (2.3 g, 13.20 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (11 mL). The reaction mixture was stirred at room temperature for 2.5 h. The solvent was evaporated to give the trifluoroacetate salt of the title compound as colorless oil in quantitative yield. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz) δ 1.01 (t, *J*=9.0 Hz, 3 H), 1.71–1.87 (m, 2 H), 3.10 (t, *J*=7.0 Hz, 2 H). MS (ESI) *m/z*: Calcd. for C<sub>3</sub>H<sub>10</sub>N<sub>2</sub> 74.08 [M]<sup>+</sup>, found 75.2 [M+H]<sup>+</sup>.

**Step 1.2: Preparation of ethyl 5-(2,6-dimethoxyphenyl)-1-propyl-1*H*-pyrazole-3-carboxylate:** Sodium salt of ethyl 4-(2,6-dimethoxyphenyl)-2,4-dioxobutanoate (1.2 g, 3.96



mmol) and propylhydrazine trifluoroacetate (1.12 g, 5.95 mmol) was mixed with glacial acetic acid (30 mL) and conc. HCl (0.6 mL). The reaction mixture was heated to reflux for 3 h. After cooling, reaction mixture was poured into water (30 mL). The aqueous layer was extracted with CH<sub>2</sub>Cl<sub>2</sub> (3 × 30 mL) and the combined CH<sub>2</sub>Cl<sub>2</sub> layer was washed with saturated aqueous NaHCO<sub>3</sub>. The organic layer was then washed with saturated brine, dried over Na<sub>2</sub>SO<sub>4</sub>, followed by filtration. The solvent was evaporated in vacuo and the residue was purified by silica gel flash chromatography (0–40% EtOAc:Hex) to give the title light yellow foam (158 mg, 15 %). <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz) δ 0.76 (t, *J*=7.3 Hz, 3 H), 1.39 (t, *J*=7.2 Hz, 3 H), 1.69–1.82 (m, 2 H), 3.74 (s, 6 H), 3.88 (t, *J*=7.5 Hz, 2 H), 4.41 (q, *J*=7.2 Hz, 2 H), 6.63 (d, *J*=8.3 Hz, 2 H), 6.73 (s, 1 H), 7.39 (t, *J*=8.5 Hz, 1 H). MS (ESI) *m/z*: Calcd. for C<sub>17</sub>H<sub>22</sub>N<sub>2</sub>O<sub>4</sub> 318.16 [M]<sup>+</sup>, found 319.3 [M+H]<sup>+</sup>.

**Step 2: Preparation of 5-(2,6-dimethoxyphenyl)-1-propyl-1*H*-pyrazole-3-carboxylic acid:**

Following the procedure as described in **45**, compound 5-(2,6-dimethoxyphenyl)-1-propyl-1*H*-pyrazole-3-carboxylic acid was obtained using ethyl 5-(2,6-dimethoxyphenyl)-1-propyl-1*H*-pyrazole-3-carboxylate. 88 % yield; light yellow solid; <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz) δ 0.78 (t, *J*=7.5 Hz, 3 H), 1.72 – 1.84 (m, 2 H), 3.76 (s, 6 H), 3.87 (t, *J*=7.5 Hz, 2 H), 6.64 (d, *J*=8.3 Hz, 2 H), 6.78 (s, 1 H), 7.40 (t, *J*=8.3 Hz, 1 H). MS (ESI) *m/z*: Calcd. for C<sub>15</sub>H<sub>18</sub>N<sub>2</sub>O<sub>4</sub> 290.13 [M]<sup>+</sup>, found 289.2 [M-H]<sup>-</sup>.

**Step 3: Preparation of (*S*)-*tert*-butyl 5-cyclohexyl-3-(5-(2,6-dimethoxyphenyl)-1-propyl-1*H*-pyrazole-3-carboxamido)pentanoate:**

Following the procedure described in **46**, compound (*S*)-*tert*-butyl 5-cyclohexyl-3-(5-(2,6-dimethoxyphenyl)-1-propyl-1*H*-pyrazole-3-carboxamido)pentanoate was obtained using 5-(2,6-dimethoxyphenyl)-1-methyl-1*H*-pyrazole-3-carboxylic acid and (*S*)-*tert*-butyl 3-amino-5-cyclohexylpentanoate. 84% yield; white solid; <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz) δ 0.77 (t, *J*=7.44 Hz, 3 H), 0.82 – 0.96 (m, 2 H), 1.10 – 1.36 (m, 7 H), 1.47 (s, 9 H), 1.61 – 1.80 (m, 7 H), 2.54 (d, *J*=5.65 Hz, 2 H), 3.73 (s, 3 H), 3.74 (s, 3 H), 3.79 (t, *J*=9.0 Hz, 2 H), 4.30 – 4.47 (m, 1 H), 6.62 (d, *J*=8.29 Hz, 2 H), 6.70 (s, 1 H), 7.25 (br. s., 1 H), 7.37 (t, *J*=8.38 Hz, 1 H). MS (ESI) *m/z*: Calcd. for C<sub>30</sub>H<sub>45</sub>N<sub>3</sub>O<sub>5</sub> 527.34 [M]<sup>+</sup>, found 529.1 [M+H]<sup>+</sup>.

**Step 4: Preparation of (*S*)-5-cyclohexyl-3-(5-(2,6-dimethoxyphenyl)-1-propyl-1*H*-pyrazole-3-carboxamido)pentanoic acid (**11**):**

Following the procedure described in **12**, compound **11** was obtained from deprotection of (*S*)-*tert*-butyl 5-cyclohexyl-3-(5-(2,6-dimethoxyphenyl)-1-propyl-1*H*-pyrazole-3-carboxamido)pentanoate. 84 % yield; white solid; <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz) δ ppm 0.78 (t, *J*=7.44 Hz, 3 H), 0.84 – 0.99 (m, 2 H), 1.09 – 1.37 (m, 6 H), 1.60 – 1.80 (m, 9 H), 2.65 – 2.78 (m, 2 H), 3.74 (s, 3 H), 3.75 (s, 3 H), 3.81 (t, *J*=6.0 Hz, 3 H), 4.28 – 4.39 (m, 1 H), 6.62 (d, *J*=8.48 Hz, 2 H), 6.72 (s, 1 H), 7.32 – 7.41 (m, 2 H). MS (ESI) *m/z*: Calcd. for C<sub>29</sub>H<sub>37</sub>N<sub>3</sub>O<sub>5</sub> 471.27 [M]<sup>+</sup>, found 470.6 [M-H]<sup>-</sup>.

**Preparation of (*S*)-5-Cyclohexyl-3-(1-cyclohexyl-5-(2,6-dimethoxyphenyl)-1*H*-pyrazole-3-carboxamido)pentanoic acid (**14**)**

**Step 1: Ethyl 1-cyclohexyl-5-(2,6-dimethoxyphenyl)-1*H*-pyrazole-3-**

**carboxylate:** Sodium salt of methyl 4-(2,6-dimethoxyphenyl)-2,4-dioxobutanoate (**40**) (1.5 g, 3.31 mmol) and commercially available cyclohexylhydrazine hydrochloride (0.75 g, 3.31

mmol) was mixed with glacial acetic acid (30 mL) and conc. HCl (0.6 mL). The reaction mixture was heated to reflux for 4 h. After cooling, reaction mixture was poured into water (50 mL). The aqueous layer was extracted with CH<sub>2</sub>Cl<sub>2</sub> (3 × 30 mL) and the combined CH<sub>2</sub>Cl<sub>2</sub> layer was washed with saturated aqueous NaHCO<sub>3</sub>. The organic layer was then washed with saturated brine, dried over Na<sub>2</sub>SO<sub>4</sub>, followed by filtration. The solvent was evaporated in vacuo to give crude residue. The residue was purified by silica gel flash chromatography (0–40 % EtOAc:Hex) to give the title compound as white solid (820 mg, 46 %). <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz) δ 1.09 – 1.31 (m, 5 H), 1.37 (t, *J*=6.97 Hz, 3 H), 1.74 – 1.92 (m, 5 H), 1.98 – 2.11 (m, 3 H), 3.63 – 3.72 (m, 1 H), 3.74 (s, 3 H), 4.40 (q, *J*=7.16 Hz, 2 H), 6.64 (d, *J*=8.29 Hz, 2 H), 6.69 (s, 1 H), 7.39 (t, *J*=9.0 Hz, 1 H). MS (ESI) *m/z*: Calcd. for C<sub>20</sub>H<sub>26</sub>N<sub>2</sub>O<sub>4</sub> 358.19 [M]<sup>+</sup>, found 359.5 [M-H]<sup>-</sup>.

**Step 2: 1-Cyclohexyl-5-(2,6-dimethoxyphenyl)-1H-pyrazole-3-carboxylic**

**acid:** Following the procedure as described in **45**, compound 5-(2,6-dimethoxyphenyl)-1-propyl-1H-pyrazole-3-carboxylic acid was obtained using ethyl 1-cyclohexyl-5-(2,6-dimethoxyphenyl)-1H-pyrazole-3-carboxylate. 96 % yield; white solid; <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz) δ 1.11 – 1.29 (m, 4 H), 1.57 – 1.67 (m, 1 H), 1.76– 2.06 (m, 6 H), 3.66 – 3.73 (m, 1 H), 3.75 (s, 6H), 6.65 (d, *J*=8.29 Hz, 2 H), 6.75 (s, 1 H), 7.40 (t, *J*=8.48 Hz, 1 H). MS (ESI) *m/z*: Calcd. for C<sub>18</sub>H<sub>22</sub>N<sub>2</sub>O<sub>4</sub> 330.38 [M]<sup>+</sup>, found 331.4 [M+H]<sup>+</sup>, 329.4 [M-H]<sup>-</sup>.

**Step 3: (S)-Tert-butyl 5-cyclohexyl-3-(1-cyclohexyl-5-(2,6-dimethoxyphenyl)-1H-pyrazole-3-carboxamido)pentanoate:**

Following the procedure described in **45**, compound (*S*)-*tert*-butyl 5-cyclohexyl-3-(1-cyclohexyl-5-(2,6-dimethoxyphenyl)-1H-pyrazole-3-carboxamido)pentanoate was obtained using 1-cyclohexyl-5-(2,6-dimethoxyphenyl)-1H-pyrazole-3-carboxylic acid and (*S*)-*tert*-butyl 3-amino-5-cyclohexylpentanoate. 93% yield; white solid; <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz) δ 0.80 – 0.96 (m, 3 H), 1.10 – 1.35 (m, 10 H), 1.49 (s, 9 H), 1.61– 1.74 (m, 7 H), 1.77–1.95 (m, 5 H), 2.55 (d, *J*=5.27 Hz, 2 H), 3.58 – 3.70 (m, 1 H), 3.72 (s, 3 H), 3.74 (s, 3 H), 4.28 – 4.46 (m, 1 H), 6.63 (d, *J*=8.29 Hz, 2 H), 6.66 (s, 1 H), 7.29 (s, 1 H), 7.37 (t, *J*=8.48 Hz, 1 H). MS (ESI) *m/z*: Calcd. for C<sub>33</sub>H<sub>49</sub>N<sub>3</sub>O<sub>5</sub> 567.37 [M]<sup>+</sup>, found 568.7 [M+H]<sup>+</sup>.

**Step 4: (S)-5-Cyclohexyl-3-(1-cyclohexyl-5-(2,6-dimethoxyphenyl)-1H-pyrazole-3-carboxamido)pentanoic acid (14):**

Following the procedure described in **12**, compound **14** was obtained from deprotection of (*S*)-*tert*-butyl 5-cyclohexyl-3-(1-cyclohexyl-5-(2,6-dimethoxyphenyl)-1H-pyrazole-3-carboxamido)pentanoate. 94 % yield; white solid; <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz) δ ppm 0.79 – 0.99 (m, 2 H), 1.10 – 1.37 (m, 10 H), 1.51 – 1.77 (m, 11 H), 1.78–1.94 (m, 2 H), 2.65 – 2.80 (m, 2 H), 3.60 – 3.71 (m, 1 H), 3.74 (s, 6 H), 4.20 – 4.33 (m, 1 H), 6.64 (d, *J*=8.48 Hz, 2 H), 6.67 (s, 1 H), 7.24 (br. s., 1 H), 7.38 (t, *J*=8.48 Hz, 1 H). MS (ESI) *m/z*: Calcd. for C<sub>29</sub>H<sub>41</sub>N<sub>3</sub>O<sub>5</sub> 511.30 [M]<sup>+</sup>, found 510.6 [M-H]<sup>-</sup>.

**(S)-5-Cyclohexyl-3-(1-(cyclohexylmethyl)-5-(2,6-dimethoxyphenyl)-1H-pyrazole-3-carboxamido)pentanoic acid (15):**

**Step 1: Preparation of ethyl 1-(cyclohexylmethyl)-5-(2,6-dimethoxyphenyl)-1H-pyrazole-3-carboxylate**

**Step 1.1: Preparation of (Cyclohexylmethyl)hydrazine trifluoroacetate :** Commercially available cyclohexanecarbaldehyde (1g, 8.91 mmol) and *tert*-butyl carbazate (1.2 g, 8.91 mmol) in methanol (25 mL) was stirred at room temperature for 1 h. The solvent was evaporated and the resulting solid was dried under in vacuo to give white solid of (*E*)-*tert*-butyl 2-(cyclohexylmethylene)hydrazinecarboxylate in quantitative yield. Sodium cyanoborohydride (0.82 g, 13.12 mmol) was added portionwise to a mixture of the (*E*)-*tert*-butyl 2-(cyclohexylmethylene)hydrazinecarboxylate (2.0 g, 8.75 mmol) in 50 % of aqueous acetic acid (25 mL) at room temperature. The resultant solution was stirred for 2.5 h at room temperature. The reaction mixture was neutralized with 1N NaOH, extracted with CH<sub>2</sub>Cl<sub>2</sub> (3 × 30 mL), washed with saturated NaHCO<sub>3</sub>, dried with Na<sub>2</sub>SO<sub>4</sub>, filtered, and evaporated to give title compound as oil (1.85 g, 92 %). <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz) δ 0.82 – 1.05 (m, 2 H), 1.13 – 1.33 (m, 4 H), 1.46 (s, 9 H), 1.59 – 1.84 (m, 5 H), 2.69 (d, *J*=6.78 Hz, 2 H). MS (ESI) *m/z*: Calcd. for C<sub>12</sub>H<sub>24</sub>N<sub>2</sub>O<sub>2</sub> 228.18 [M]<sup>+</sup>, found 229.4 [M+H]<sup>+</sup>.

Trifluoroacetic acid (8 mL) was added dropwise to a solution of the *tert*-butyl 2-isobutylhydrazinecarboxylate (2.2 g, 9.63 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (8 mL). The reaction mixture was stirred at room temperature for 1.5 h. The solvent was evaporated to give the trifluoroacetate salt of the title compound as colorless oil in quantitative yield. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz) δ 0.20 – 0.85 (m, 4 H), 0.86 – 1.34 (m, 6 H), 1.62–1.69 (m, 1 H), 1.70 – 1.82 (m, 2 H). MS (ESI) *m/z*: Calcd. for C<sub>7</sub>H<sub>16</sub>N<sub>2</sub> 128.13 [M]<sup>+</sup>, found 129.3 [M+H]<sup>+</sup>.

**Step 1.2: Preparation of ethyl 1-(cyclohexylmethyl)-5-(2,6-dimethoxyphenyl)-1H-pyrazole-3-carboxylate:** Sodium salt of ethyl 4-(2,6-dimethoxyphenyl)-2,4-dioxobutanoate (1.2 g, 3.96 mmol) and (cyclohexylmethyl)hydrazine trifluoroacetate (0.960 g, 3.965 mmol) was mixed with glacial acetic acid (25 mL) and conc. HCl (0.6 mL). The reaction mixture was heated to reflux for 4 h. After cooling, reaction mixture was poured into water (25 mL). The aqueous layer was extracted with DCM (3 × 30 mL). The combined DCM layer was washed with saturated aqueous NaHCO<sub>3</sub>. The organic layer was then washed with saturated brine, and then dried over Na<sub>2</sub>SO<sub>4</sub>, followed by filtration. The solvent was evaporated in vacuo. The residue was purified by silica gel flash chromatography (0–50% EtOAc:Hex) to give the title compound as oil (0.510 g, 35 %) and 2-methyl cyclohexyl substituted isomer as oil (0.215 g, 15 %). <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz) δ 0.61 – 0.78 (m, 2 H), 0.97 – 1.19 (m, 2 H), 1.39 (t, *J*=8.48 Hz, 3 H), 1.45 – 1.66 (m, 6 H), 1.78–1.93 (m, 1 H), 3.74 (s, 6 H), 3.74 (d, *J*=6.78 Hz, 2 H), 4.41 (q, *J*=7.16 Hz, 2 H), 6.63 (d, *J*=8.67 Hz, 2 H), 6.72 (s, 1 H), 7.39 (t, *J*=8.48 Hz, 1 H). MS (ESI) *m/z*: Calcd. for C<sub>21</sub>H<sub>28</sub>N<sub>2</sub>O<sub>4</sub> 372.20 [M]<sup>+</sup>, found 373.5 [M+H]<sup>+</sup>.

**Step 2: Preparation of 1-(cyclohexylmethyl)-5-(2,6-dimethoxyphenyl)-1H-pyrazole-3-carboxylic acid:** Following the procedure as described in **45**, compound 1-(cyclohexylmethyl)-5-(2,6-dimethoxyphenyl)-1H-pyrazole-3-carboxylic acid was obtained using ethyl 1-(cyclohexylmethyl)-5-(2,6-dimethoxyphenyl)-1H-pyrazole-3-carboxylate. 95 % yield; white solid; <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz) δ 0.62 – 0.81 (m, 2 H), 0.98 – 1.22 (m, 3

H), 1.43 – 1.66 (m, 5 H), 1.76–1.93 (m, 1 H), 3.73 (d,  $J=6.78$  Hz, 2 H), 3.75 (s, 6 H), 6.63 (d,  $J=8.29$  Hz, 2 H), 6.78 (s, 1 H), 7.40 (t,  $J=8.48$  Hz, 1 H). MS (ESI)  $m/z$ . Calcd. for  $C_{19}H_{24}N_2O_4$  344.17  $[M]^+$ , found 343.4  $[M-H]^-$

**Step 3: (*S*)-*Tert*-butyl 5-cyclohexyl-3-(1-(cyclohexylmethyl)-5-(2,6-**

**dimethoxyphenyl)-1*H*-pyrazole-3-carboxamido)pentanoate:** Following the procedure described in **45**, compound (*S*)-*tert*-butyl 5-cyclohexyl-3-(1-(cyclohexylmethyl)-5-(2,6-dimethoxyphenyl)-1*H*-pyrazole-3-carboxamido)pentanoate was obtained using 1-(cyclohexylmethyl)-5-(2,6-dimethoxyphenyl)-1*H*-pyrazole-3-carboxylic acid and (*S*)-*tert*-butyl 3-amino-5-cyclohexylpentanoate. 66% yield; clear foam;  $^1H$  NMR ( $CDCl_3$ , 300 MHz)  $\delta$  0.63 – 0.78 (m, 2 H), 0.80 – 0.98 (m, 3 H), 1.02 – 1.36 (m, 9 H), 1.48 (s, 9 H), 1.53 – 1.85 (m, 12 H), 2.54 (d,  $J=5.46$  Hz, 2 H), 3.62 – 3.69 (m, 2 H), 3.72 (s, 3 H), 3.74 (s, 3 H), 4.30 – 4.42 (m, 1 H), 6.62 (d,  $J=8.48$  Hz, 2 H), 6.68 (s, 1 H), 7.22 (s, 1 H), 7.37 (t,  $J=8.29$  Hz, 1 H). MS (ESI)  $m/z$ . Calcd. for  $C_{34}H_{51}N_3O_5$  581.38  $[M]^+$ , found 582.8  $[M+H]^+$ .

**Step 4: Preparation of (*S*)-5-cyclohexyl-3-(1-(cyclohexylmethyl)-5-(2,6-**

**dimethoxyphenyl)1*H*-pyrazole-3-carboxamido)pentanoic acid (15):** Following the procedure described in **12**, compound **15** was obtained from deprotection of (*S*)-*tert*-butyl 5-cyclohexyl-3-(1-(cyclohexylmethyl)-5-(2,6-dimethoxyphenyl)-1*H*-pyrazole-3-carboxamido)pentanoate. 84 % yield; white solid;  $^1H$  NMR ( $CDCl_3$ , 300 MHz)  $\delta$  0.62–0.98 (m, 3 H), 1.02–1.87(m, 10 H), 1.421.84 (m, 13 H), 2.64–2.81 (m, 2 H), 3.62–3.72 (m, 2 H), 3.74 (s, 6 H), 4.24–4.39 (m, 1 H), 6.62 (d,  $J=8.10$  Hz, 2 H), 6.71 (s, 1 H), 7.20–7.26 (m, 1 H), 7.28 – 7.41 (m, 1 H). MS (ESI)  $m/z$ . Calcd. for  $C_{30}H_{43}N_3O_5$  525.32  $[M]^+$ , found 524.8  $[M-H]^-$ .

**(*S*)-5-Cyclohexyl-3-(1-cyclooctyl-5-(2,6-dimethoxyphenyl)-1*H*-pyrazole-3-carboxamido)pentanoic acid (16)**

**Step 1: Ethyl 1-cyclooctyl-5-(2,6-dimethoxyphenyl)-1*H*-pyrazole-3-carboxylate**

**Step 1.1: Preparation of *tert*-butyl 2-cyclooctylhydrazine carboxylate:** Octanone (1.43 g, 11.34 mmol) and *tert*-butyl carbazate (1.5 g, 11.34 mmol) in methanol (25 mL) was stirred at room temperature for 2 h. The solvent was evaporated and the resulting solid was dried under in vacuo to give white solid of (*E*)-*tert*-butyl 2-cyclooctyldiazene-carboxylate in quantitative yield. Sodium cyanoborohydride (0.980 g, 15.60 mmol) was added portionwise to a mixture of the (*E*)-*tert*-butyl 2-cyclooctyldiazene-carboxylate (2.50 g, 10.40 mmol) in 75 % of aqueous acetic acid (35 mL) at room temperature. The resultant solution was stirred for 3 h at room temperature. The reaction mixture was neutralized with 1N NaOH, extracted with  $CH_2Cl_2$  ( $3 \times 30$  mL), washed with saturated  $NaHCO_3$ , dried with  $Na_2SO_4$ , filtered, and evaporated to give title compound as oil (2.40 g, 95 %).  $^1H$  NMR ( $CDCl_3$ , 300 MHz)  $\delta$  1.33–1.59 (m, 4 H), 1.46 (s, 9 H), 1.63–1.79 (m, 11 H). MS (ESI)  $m/z$ . Calcd. for  $C_{13}H_{26}N_2O_2$  242.20  $[M]^+$ , found 243.4  $[M+H]^+$ .

Trifluoroacetic acid (10 mL) was added dropwise to a solution of the *tert*-butyl 2-(naphthalen-2-ylmethyl)hydrazine-carboxylate (3.0 g, 12.38 mmol) in  $CH_2Cl_2$  (10 mL). The reaction mixture was stirred at room temperature for 1.5 h. The solvent was evaporated to give the trifluoroacetate salt of the title compound as colorless oil in quantitative yield.  $^1H$

NMR (CDCl<sub>3</sub>, 300 MHz)  $\delta$  1.36 – 2.04 (m, 14 H), 3.22– 3.47 (m, 1 H). MS (ESI)  $m/z$ : Calcd. for C<sub>8</sub>H<sub>18</sub>N<sub>2</sub> 142.15 [M]<sup>+</sup>, found 143.3 [M+H]<sup>+</sup>.

**Step 1.2: Preparation of ethyl 1-cyclooctyl-5-(2,6-dimethoxyphenyl)-1H-pyrazole-3-carboxylate:** Sodium salt of ethyl 4-(2,6-dimethoxyphenyl)-2,4-dioxobutanoate (1.2 g, 3.97 mmol) and cyclooctylhydrazine trifluoroacetate (1.2 g, 4.76 mmol) was mixed with glacial acetic acid (25 mL) and conc. HCl (0.6 mL). The reaction mixture was heated to reflux for 3.5 h. After cooling, reaction mixture was poured into water (30 mL). The aqueous layer was extracted with DCM (3 × 30 mL). The combined DCM layer was washed with saturated aqueous NaHCO<sub>3</sub>. The organic layer was then washed with saturated brine, and then dried over Na<sub>2</sub>SO<sub>4</sub>, followed by filtration. The solvent was evaporated in vacuo. The residue was purified by silica gel flash chromatography (0–30% EtOAc:Hex) to give the title compound as gum (600 mg, 39 %). <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz)  $\delta$  1.18 – 1.29 (m, 2 H), 1.32 – 1.43 (m, 6 H), 1.37 (t,  $J=7.15$  Hz, 3 H), 1.67 – 1.86 (m, 4 H), 2.15 – 2.29 (m, 2 H), 3.74 (s, 6 H), 3.98 – 4.09 (m, 1 H), 4.39 (q,  $J=7.03$  Hz, 2 H), 6.64 (d,  $J=8.29$  Hz, 2 H), 6.69 (s, 1 H), 7.39 (t,  $J=8.48$  Hz, 1 H). MS (ESI)  $m/z$ : Calcd. for C<sub>22</sub>H<sub>30</sub>N<sub>2</sub>O<sub>4</sub> 386.22 [M]<sup>+</sup>, found 387.4 [M+H]<sup>+</sup>.

**Step 2: Preparation of 1-cyclooctyl-5-(2,6-dimethoxyphenyl)-1H-pyrazole-3-carboxylic acid:** Following the procedure as described in 46, compound 1-cyclooctyl-5-(2,6-dimethoxyphenyl)-1H-pyrazole-3-carboxylic acid was obtained from hydrolysis of ethyl 1-cyclooctyl-5-(2,6-dimethoxyphenyl)-1H-pyrazole-3-carboxylate. 77 % yield; white solid; <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz)  $\delta$  1.21 – 1.65 (m, 8 H), 1.68 – 1.89 (m, 4 H), 2.10 – 2.33 (m, 2 H), 3.75 (s, 6 H), 3.98 – 4.09 (m, 1 H), 6.65 (d,  $J=8.29$  Hz, 2 H), 6.74 (s, 1 H), 7.41 (t,  $J=8.48$  Hz, 1 H). MS (ESI)  $m/z$ : Calcd. for C<sub>20</sub>H<sub>26</sub>N<sub>2</sub>O<sub>4</sub> 358.19 [M]<sup>+</sup>, found 357.5 [M-H]<sup>-</sup>.

**Step 3: Preparation of (S)-tert-butyl 5-cyclohexyl-3-(1-cyclooctyl-5-(2,6-dimethoxyphenyl)-1H-pyrazole-3-carboxamido)pentanoate:** Following the procedure described in 46, compound (S)-tert-butyl 5-cyclohexyl-3-(5-(2,6-dimethoxyphenyl)-1-propyl-1H-pyrazole-3-carboxamido)pentanoate was obtained using 5-(2,6-dimethoxyphenyl)-1-methyl-1H-pyrazole-3-carboxylic acid and (S)-tert-butyl 3-amino-5-cyclohexylpentanoate. 73% yield; clear foam; <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz)  $\delta$  0.80 – 0.97 (m, 2 H), 1.13 – 1.45 (m, 13 H), 1.48 (s, 9 H), 1.62 – 1.86 (m, 13 H), 2.02 – 2.18 (m, 2 H), 2.55 (d,  $J=5.46$  Hz, 2 H), 3.73 (s, 3 H), 3.74 (s, 3 H), 3.92–4.04 s, 1 H), 4.31 – 4.42 (m, 1 H), 6.62 (s, 1 H), 6.65 (d,  $J=4.33$  Hz, 1 H), 7.25 – 7.28 (m, 1 H), 7.38 (t,  $J=8.38$  Hz, 1 H). MS (ESI)  $m/z$ : Calcd. for C<sub>35</sub>H<sub>53</sub>N<sub>3</sub>O<sub>5</sub> 595.40 [M]<sup>+</sup>, found 596.9 [M+H]<sup>+</sup>.

**Step 4: Preparation of (S)-5-cyclohexyl-3-(1-cyclooctyl-5-(2,6-dimethoxyphenyl)-1H-pyrazole-3-carboxamido)pentanoic acid (16):** Following the procedure described in 12, compound 16 was obtained from deprotection of (S)-tert-butyl 5-cyclohexyl-3-(1-cyclooctyl-5-(2,6-dimethoxyphenyl)-1H-pyrazole-3-carboxamido)pentanoate. 86 % yield; white solid; <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz)  $\delta$  0.81 – 0.99 (m, 2 H), 1.11 – 1.47 (m, 13 H), 1.49–1.88 (m, 14 H), 2.03–2.18 (m, 2 H), 2.66 – 2.81 (m, 2 H), 3.74 (s, 5 H), 3.94 – 4.05 (m, 1 H), 4.21 – 4.32 (m, 1 H), 6.62 (s, 1 H), 6.66 (d,  $J=6.59$  Hz, 2 H), 7.28 (br. s., 1 H), 7.39 (t,  $J=8.38$  Hz, 1 H). MS (ESI)  $m/z$ : Calcd. for C<sub>31</sub>H<sub>45</sub>N<sub>3</sub>O<sub>5</sub> 539.34 [M]<sup>+</sup>, found 538.7 [M-H]<sup>-</sup>.

**(S)-N-(1-amino-5-cyclohexyl-1-oxopentan-3-yl)-5-(2,6-dimethoxyphenyl)-1-isobutyl-1H-pyrazole-3-carboxamide (17)**: Following the procedure described in 45, compound 17 was synthesized using (S)-5-cyclohexyl-3-(5-(2,6-dimethoxyphenyl)-1-isobutyl-1H-pyrazole-3-carboxamido)pentanoic acid (12) and 2 M NH<sub>3</sub>. 72 % yield; white solid; <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz) δ 0.75 (dd, *J*=6.59, 2.83 Hz, 6 H), 0.81 – 1.01 (m, 3 H), 1.11 – 1.36 (m, 6 H), 1.61 – 1.78 (m, 6 H), 2.06 – 2.17 (m, 1 H), 2.23 – 2.36 (m, 1 H), 2.58 (d, *J*=6.03 Hz, 2 H), 3.64 (d, *J*=7.16 Hz, 2 H), 3.74 (d, *J*=2.26 Hz, 6 H), 4.26 – 4.39 (m, 1 H), 6.62 (d, *J*=8.29 Hz, 2 H), 6.68 (s, 1 H), 6.81 (br. s., 1 H), 7.09 (d, *J*=9.42 Hz, 1 H), 7.38 (t, *J*=8.48 Hz, 1 H). MS (ESI) *m/z*: Calcd. for C<sub>27</sub>H<sub>40</sub>N<sub>4</sub>O<sub>4</sub> 484.30 [M]<sup>+</sup>, found 485.6 [M+H]<sup>+</sup>.

**(S)-Methyl 2-(5-cyclohexyl-3-(5-(2,6-dimethoxyphenyl)-1-isobutyl-1H-pyrazole-3-carboxamido)pentanamido)acetate (18)**: Following the procedure described in 45, compound 18 was synthesized using (S)-5-cyclohexyl-3-(5-(2,6-dimethoxyphenyl)-1-isobutyl-1H-pyrazole-3-carboxamido)pentanoic acid (12) and glycine methyl ester hydrochloride. 21 % yield; white solid; <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz) δ 0.74 (dd, *J*=6.78, 1.51 Hz, 6 H), 0.82 – 0.99 (m, 2 H), 1.10 – 1.36 (m, 7 H), 1.63 – 1.77 (m, 6 H), 2.11 (td, *J*=14.22, 7.35 Hz, 1 H), 2.62 (d, *J*=5.65 Hz, 2 H), 3.63 (d, *J*=7.16 Hz, 2 H), 3.68 (s, 3 H), 3.73 (s, 3 H), 3.74 (s, 3 H), 4.04 (d, *J*=5.65 Hz, 2 H), 4.27 – 4.41 (m, 1 H), 6.62 (d, *J*=8.29 Hz, 2 H), 6.68 (s, 1 H), 6.91 – 7.00 (m, 1 H), 7.18 (d, *J*=9.04 Hz, 1 H), 7.37 (t, *J*=8.48 Hz, 1 H). MS (ESI) *m/z*: Calcd. for C<sub>30</sub>H<sub>44</sub>N<sub>4</sub>O<sub>6</sub> 556.33 [M]<sup>+</sup>, found 557.8 [M+H]<sup>+</sup>.

**(S)-Methyl 2-(5-cyclohexyl-3-(5-(2,6-dimethoxyphenyl)-1-isobutyl-1H-pyrazole-3-carboxamido)-N-methylpentanamido)acetate (19)**: Following the procedure described in 45, compound 19 was synthesized using (S)-5-cyclohexyl-3-(5-(2,6-dimethoxyphenyl)-1-isobutyl-1H-pyrazole-3-carboxamido)pentanoic acid (12) and sarcosine methyl ester hydrochloride. 38 % yield; white solid; <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz) δ 0.74 (d, *J*=6.78 Hz, 6 H), 0.81 – 1.01 (m, 3 H), 1.10 – 1.40 (m, 6 H), 1.54 – 1.83 (m, 5 H), 2.06 – 2.18 (m, 1 H), 2.62 (dd, *J*=15.45, 6.78 Hz, 1 H), 2.84 – 2.96 (m, 2 H), 3.14 (s, 3 H), 3.63 (d, *J*=7.16 Hz, 2 H), 3.71 (s, 3 H), 3.72 (s, 3 H), 3.73 (s, 3 H), 4.04 – 4.21 (m, 2 H), 4.27–4.39 (m, 1 H), 6.61 (d, *J*=8.67 Hz, 2 H), 6.67 (s, 1 H), 7.30 (br. s., 1 H), 7.37 (t, *J*=8.29 Hz, 1 H). MS (ESI) *m/z*: Calcd. for C<sub>31</sub>H<sub>46</sub>N<sub>4</sub>O<sub>6</sub> 570.34 [M]<sup>+</sup>, found 572.0 [M+H]<sup>+</sup>.

**(S)-2-(5-Cyclohexyl-3-(5-(2,6-dimethoxyphenyl)-1-isobutyl-1H-pyrazole-3-carboxamido)pentanamido)acetic acid (20)**: Following the procedure as described in 45, compound (S)-2-(5-Cyclohexyl-3-(5-(2,6-dimethoxyphenyl)-1-isobutyl-1H-pyrazole-3-carboxamido)pentanamido)acetic acid (20) was obtained using (S)-methyl 2-(5-cyclohexyl-3-(5-(2,6-dimethoxyphenyl)-1-isobutyl-1H-pyrazole-3-carboxamido)pentanamido)acetate (18). 52 % yield; white solid; <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz) δ 0.74 (d, *J*=6.40 Hz, 6 H), 0.85–0.95 (m, 2 H), 1.10 – 1.39 (m, 5 H), 1.55–1.87 (m, 8 H), 2.06 – 2.15 (m, 2 H), 2.50–2.82 (m, 2 H), 3.64 (d, *J*=7.54 Hz, 2 H), 3.74 (s, 6 H), 4.01 – 4.17 (m, 2 H), 4.26–4.39 (m, 1 H), 6.62 (d, *J*=8.29 Hz, 2 H), 6.67 (s, 1 H), 7.16 (d, *J*=8.67 Hz, 1 H), 7.38 (t, *J*=8.48 Hz, 1 H), 7.42 (br. s., 1 H). MS (ESI) *m/z*: Calcd. for C<sub>29</sub>H<sub>42</sub>N<sub>4</sub>O<sub>6</sub> 542.31 [M]<sup>+</sup>, found 543.7 [M+H]<sup>+</sup>.



***N*-((3*S*)-5-Cyclohexyl-1-((2-hydroxybutyl)amino)-1-oxopentan-3-yl)-5-(2,6-dimethoxyphenyl)-1-isobutyl-1*H*-pyrazole-3-carboxamide (21):** Following the procedure described in **45**, compound **21** was synthesized using (*S*)-5-cyclohexyl-3-(5-(2,6-dimethoxyphenyl)-1-isobutyl-1*H*-pyrazole-3-carboxamido)pentanoic acid (**12**) and 1-amino-2-butanol. 65 % yield; white solid; <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz) δ 0.74 (d, *J*=6.00 Hz, 6 H), 0.90 (t, *J*=7.35 Hz, 3 H), 0.81–0.96 (m, 2 H), 1.10 – 1.46 (m, 7 H), 1.62 – 1.76 (m, 5 H), 2.04–2.19 (m, 1 H), 2.45 – 2.69 (m, 2 H), 2.88 – 3.03 (m, 1 H), 3.18 – 3.29 (m, 1 H), 3.35 – 3.52 (m, 3 H), 3.57 (br. s., 2 H), 3.64 (d, *J*=7.16 Hz, 2 H), 3.74 (m, 6 H), 4.25 – 4.46 (m, 1 H), 6.62 (d, *J*=8.29 Hz, 2 H), 6.68 (s, 1 H), 6.88–6.98 (d, *J*=5.65 Hz, 1 H), 7.04 (t, *J*=9.61 Hz, 1 H), 7.38 (t, *J*=8.48 Hz, 1 H). MS (ESI) *m/z*: Calcd. for C<sub>31</sub>H<sub>48</sub>N<sub>4</sub>O<sub>5</sub> 556.36 [M]<sup>+</sup>, found 558.1 [M+H]<sup>+</sup>.

***N*-((*S*)-5-Cyclohexyl-1-(((*R*)-2-hydroxybutyl)amino)-1-oxopentan-3-yl)-5-(2,6-dimethoxyphenyl)-1-isobutyl-1*H*-pyrazole-3-carboxamide (22):** Following the procedure described in **45**, compound **22** was synthesized using (*S*)-5-cyclohexyl-3-(5-(2,6-dimethoxyphenyl)-1-isobutyl-1*H*-pyrazole-3-carboxamido)pentanoic acid (**12**) and (*2R*)-1-amino-2-butanol. 63 % yield; white solid; <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz) δ 0.74 (dd, *J*=6.69, 1.41 Hz, 6 H), 0.79 – 0.98 (m, 2 H), 0.90 (t, *J*=6.0 Hz, 3 H), 1.08 – 1.47 (m, 8 H), 1.58 – 1.75 (m, 6 H), 1.89 (br. s., 1 H), 2.11 (m, 1 H), 2.50 – 2.64 (m, 2 H), 3.17 – 3.26 (m, 1 H), 3.37 (dd, *J*=5.65, 2.64 Hz, 1 H), 3.45–3.51 (m, 1 H), 3.54–3.61 (m, 1 H), 3.64 (d, *J*=8.48 Hz, 2 H), 3.73 (s, 3 H), 3.74 (s, 3 H), 4.25 – 4.37 (m, 1 H), 6.62 (d, *J*=8.48 Hz, 2 H), 6.68 (s, 1 H), 6.97 (t, *J*=5.46 Hz, 1 H), 7.09 (d, *J*=9.23 Hz, 1 H), 7.38 (t, *J*=9.23 Hz, 1 H). MS (ESI) *m/z*: Calcd. for C<sub>31</sub>H<sub>48</sub>N<sub>4</sub>O<sub>5</sub> 556.36 [M]<sup>+</sup>, found 557.8 [M+H]<sup>+</sup>.

***(S)*-*N*-((5-Cyclohexyl-1-oxo-1-((2-oxobutyl)amino)pentan-3-yl)-5-(2,6-dimethoxyphenyl)-1-isobutyl-1*H*-pyrazole-3-carboxamide (23):** To a solution of *N*-((*S*)-5-cyclohexyl-1-(((*R*)-2-hydroxybutyl)amino)-1-oxopentan-3-yl)-5-(2,6-dimethoxyphenyl)-1-isobutyl-1*H*-pyrazole-3-carboxamide (**21**) (22 mg, 0.39 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (5 mL) was added PCC (15 mg, 0.06 mmol), Celite 545 (10 mg) and stirred at room temperature for 16 h. The reaction mixture was diluted with ethyl ether (1 mL), stirred at rt for 1 h, before it was filtered through celite and silica gel (1:1) pad. The filtrate was concentrated to give crude residue. Crude product was purified by silica gel flash chromatography (0–50% EtOAc/hexanes) to give the title compound. 27 % yield; light brown solid; <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz) δ 0.74 (d, *J*=6.03 Hz, 6 H), 0.81 – 1.01 (m, 2 H), 1.08 (t, *J*=7.16 Hz, 4 H), 1.13 – 1.39 (m, 7 H), 1.63–1.78 (m, 7 H), 2.46 (q, *J*=7.16 Hz, 2 H), 2.60 (br. s., 2 H), 3.63 (d, *J*=6.40 Hz, 2 H), 3.74 (s, 6 H), 4.12 (m, 2 H), 6.62 (d, *J*=8.29 Hz, 2 H), 6.68 (br. s., 1 H), 6.85 (br. s., 1 H), 7.24–7.31 (m, 1 H), 7.37 (t, *J*=9.00 Hz, 1 H). MS (ESI) *m/z*: Calcd. for C<sub>31</sub>H<sub>46</sub>N<sub>4</sub>O<sub>5</sub> 554.35 [M]<sup>+</sup>, found 556.2 [M+H]<sup>+</sup>.

***(S)*-*N*-((5-Cyclohexyl-1-((2-methoxyethyl)amino)-1-oxopentan-3-yl)-5-(2,6-dimethoxyphenyl)-1-isobutyl-1*H*-pyrazole-3-carboxamide (24):** Following the procedure described in **45**, compound **24** was synthesized using (*S*)-5-cyclohexyl-3-(5-(2,6-dimethoxyphenyl)-1-isobutyl-1*H*-pyrazole-3-carboxamido)pentanoic acid (**12**) and 2-methoxy ethylamine. 91 % yield; white solid; <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz) δ 0.74 (d, *J*=6.78 Hz, 6 H), 0.82 – 1.01 (m, 2 H), 1.12 – 1.40 (m, 7 H), 1.63 – 1.77 (m, 7 H),

2.55 (d,  $J=6.03$  Hz, 2 H), 3.27 – 3.31 (m, 3 H), 3.39 – 3.48 (m, 4 H), 3.63 (d,  $J=7.16$  Hz, 2 H), 3.72 (s, 3 H), 3.74 (s, 3 H), 4.23–4.36 (m, 1 H), 6.51 (br. s., 1 H), 6.62 (d,  $J=8.67$  Hz, 2 H), 6.68 (s, 1 H), 7.21 (d,  $J=8.29$  Hz, 1 H), 7.37 (t,  $J=8.29$  Hz, 1 H). MS (ESI)  $m/z$ . Calcd. for  $C_{30}H_{46}N_4O_5$  542.35  $[M]^+$ , found 544.1  $[M+H]^+$ .

**(S)-2-(5-(2,6-Dimethoxyphenyl)-1-isobutyl-1H-pyrazole-3-carboxamido)-4-phenylbutanoic acid (26):** Following the procedure described in **12**, compound **26** was obtained from deprotection of (*S*)-*tert*-butyl 2-(5-(2,6-dimethoxyphenyl)-1-isobutyl-1H-pyrazole-3-carboxamido)-4-phenylbutanoate. 80 % yield; white solid;  $^1H$  NMR ( $CDCl_3$ , 300 MHz)  $\delta$  0.75 (dd,  $J=6.59$ , 2.45 Hz, 6 H), 0.81–0.95 (m, 2 H), 2.08 – 2.26 (m, 2 H), 2.31 – 2.46 (m, 1 H), 2.82 (t,  $J=8.10$  Hz, 2 H), 3.67 (d,  $J=7.35$  Hz, 2 H), 3.73 (s, 3 H), 3.74 (s, 3 H), 4.63 – 4.74 (m, 1 H), 6.62 (d,  $J=8.29$  Hz, 2 H), 6.73 (s, 1 H), 7.12 – 7.25 (m, 3 H), 7.27 – 7.32 (m, 1 H), 7.29 – 7.44 (m, 2 H). MS (ESI)  $m/z$ . Calcd. for  $C_{26}H_{31}N_3O_5$  465.23  $[M]^+$ , found 464.8  $[M-H]^-$ .

**(S)-Methyl 2-(2-(5-(2,6-dimethoxyphenyl)-1-isobutyl-1H-pyrazole-3-carboxamido)-4-phenylbutanamido)acetate (27):** Following the procedure described in **45**, compound **27** was synthesized using 5-(2,6-dimethoxyphenyl)-1-isobutyl-1H-pyrazole-3-carboxylic acid (**43**) and (*S*)-methyl 2-(2-amino-4-phenylbutanamido)acetate trifluoroacetate. 50 % yield; white solid;  $^1H$  NMR ( $CDCl_3$ , 300 MHz)  $\delta$  0.76 (d,  $J=6.78$  Hz, 3 H), 0.74 (d,  $J=6.78$  Hz, 3 H), 2.07–2.21 (m, 2 H), 2.32 – 2.44 (m, 1 H), 2.79 (t,  $J=7.91$  Hz, 2 H), 3.64 (d,  $J=7.16$  Hz, 2 H), 3.73 (s, 6 H), 3.75 (s, 3 H), 4.05 (dd,  $J=5.65$ , 2.26 Hz, 2 H), 4.64 (td,  $J=8.10$ , 6.03 Hz, 1 H), 6.62 (d,  $J=8.29$  Hz, 2 H), 6.71 (s, 1 H), 6.85 – 6.95 (m, 1 H), 7.14 – 7.27 (m, 4 H), 7.28 – 7.33 (m, 2 H), 7.38 (t,  $J=8.48$  Hz, 1 H). MS (ESI)  $m/z$ . Calcd. for  $C_{29}H_{36}N_4O_6$  536.26  $[M]^+$ , found 537.5  $[M+H]^+$ .

**(S)-Methyl 2-(2-(5-(2,6-dimethoxyphenyl)-1-isobutyl-1H-pyrazole-3-carboxamido)-N-methyl-4-phenylbutanamido)acetate (28):** Following the procedure described in **45**, compound **28** was synthesized using 5-(2,6-dimethoxyphenyl)-1-isobutyl-1H-pyrazole-3-carboxylic acid (**43**) and (*S*)-methyl 2-(2-amino-*N*-methyl-4-phenylbutanamido)acetate trifluoroacetate. 50 % yield; white solid;  $^1H$  NMR ( $CDCl_3$ , 300 MHz)  $\delta$  0.76 (d,  $J=4.52$  Hz, 3 H), 0.73 (d,  $J=4.52$  Hz, 3 H), 2.03 – 2.23 (m, 4 H), 2.70 – 2.84 (m, 2 H), 3.05 (s, 3 H), 3.63 (d,  $J=9.00$  Hz, 3 H), 3.72 (s, 3 H), 3.74 (s, 6 H), 4.44–4.52 (m, 1 H), 5.20–5.28 (m, 1 H), 6.62 (dd,  $J=8.67$ , 1.51 Hz, 2 H), 6.69 (s, 1 H), 7.14 – 7.23 (m, 1 H), 7.29 (m, 3 H), 7.37 (t,  $J=8.29$  Hz, 1 H), 7.65 (d,  $J=8.67$  Hz, 1 H). MS (ESI)  $m/z$ . Calcd. for  $C_{30}H_{38}N_4O_6$  550.28  $[M]^+$ , found 551.7  $[M+H]^+$ .

**(S)-5-(2,6-Dimethoxyphenyl)-1-isobutyl-N-(1-((2-methoxyethyl)amino)-1-oxo-4-phenylbutan-2-yl)-1H-pyrazole-3-carboxamide (29):** Following the procedure described in **45**, compound **29** was synthesized using 5-(2,6-dimethoxyphenyl)-1-isobutyl-1H-pyrazole-3-carboxylic acid (**43**) and (*S*)-2-amino-*N*-(2-methoxyethyl)-4-phenylbutanamide trifluoroacetate. 73 % yield; white solid;  $^1H$  NMR ( $CDCl_3$ , 300 MHz)  $\delta$  0.76 (d,  $J=6.78$  Hz, 3 H), 0.74 (d,  $J=6.78$  Hz, 3 H), 2.02 – 2.20 (m, 2 H), 2.25 – 2.39 (m, 1 H), 2.76 (t,  $J=7.91$  Hz, 2 H), 3.34 (s, 3 H), 3.39 – 3.53 (m, 4 H), 3.64 (d,  $J=7.16$  Hz, 2 H), 3.74 (s, 3 H), 3.75 (s, 3 H), 4.56–4.63 (m, 1 H), 6.53 (br. s., 1 H), 6.62 (d,  $J=8.29$  Hz, 2 H),

6.70 (s, 1 H), 7.15 – 7.25 (m, 3 H), 7.28 – 7.33 (m, 3 H), 7.38 (t,  $J=9.00$  Hz, 1 H). MS (ESI)  $m/z$ : Calcd. for  $C_{29}H_{38}N_4O_5$  522.28  $[M]^+$ , found 523.9  $[M+H]^+$ .

**5-(2,6-Dimethoxyphenyl)-*N*-((2*S*)-1-((2-hydroxybutyl)amino)-1-oxo-4-phenylbutan-2-yl)-1-isobutyl-1*H*-pyrazole-3-carboxamide (30)**: Following the procedure described in **45**, compound **30** was synthesized using 5-(2,6-dimethoxyphenyl)-1-isobutyl-1*H*-pyrazole-3-carboxylic acid (**43**) and (2*S*)-2-amino-*N*-(2-hydroxybutyl)-4-phenylbutanamide trifluoroacetate. 90 % yield; white solid;  $^1H$  NMR ( $CDCl_3$ , 300 MHz)  $\delta$  0.76 (d,  $J=3.39$  Hz, 3 H), 0.74 (d,  $J=3.77$  Hz, 3 H), 0.95 (t,  $J=6.00$  Hz, 3 H), 1.38–1.52 (m, 2 H), 2.06 – 2.18 (m, 2 H), 2.30 – 2.44 (m, 1 H), 2.75 – 2.85 (m, 2 H), 3.09 – 3.19 (m, 1 H), 3.40 – 3.54 (m, 1 H), 3.64 (d,  $J=7.54$  Hz, 3 H), 3.74 (m, 6 H), 4.48 – 4.58 (m, 1 H), 6.63 (d,  $J=9.00$  Hz, 2 H), 6.69 (s, 1 H), 6.81 (br. s., 1 H), 7.15 – 7.24 (m, 2 H), 7.28 – 7.32 (m, 4 H), 7.38 (t,  $J=8.29$  Hz, 1 H). MS (ESI)  $m/z$ : Calcd. for  $C_{30}H_{40}N_4O_5$  536.30  $[M]^+$ , found 535.7  $[M-H]^-$ .

**(*S*)-5-(2,6-Dimethoxyphenyl)-1-isobutyl-*N*-(1-oxo-1-((2-oxobutyl)amino)-4-phenylbutan-2-yl)-1*H*-pyrazole-3-carboxamide (31)**: Following the procedure described in **23**, compound **31** was synthesized using 5-(2,6-dimethoxyphenyl)-*N*-((2*S*)-1-(2-hydroxybutyl)amino)-1-oxo-4-phenylbutan-2-yl)-1-isobutyl-1*H*-pyrazole-3-carboxamide (**31**). 45 % yield; light brown solid;  $^1H$  NMR ( $CDCl_3$ , 300 MHz)  $\delta$  0.75 (dd,  $J=6.78, 3.39$  Hz, 6 H), 1.10 (t,  $J=7.16$  Hz, 3 H), 2.04 – 2.20 (m, 2 H), 2.28–2.41 (m, 1 H), 2.47 (q,  $J=7.16$  Hz, 2 H), 2.78 (t,  $J=7.72$  Hz, 2 H), 3.64 (d,  $J=7.54$  Hz, 2 H), 3.74 (s, 6 H), 4.09–4.18 (m, 2 H), 4.60–4.71 (m, 1 H), 6.62 (d,  $J=8.29$  Hz, 2 H), 6.70 (s, 1 H), 6.94 (br. s., 1 H), 7.13 – 7.25 (m, 3 H), 7.27 – 7.34 (m, 3 H), 7.38 (t,  $J=8.48$  Hz, 1 H). MS (ESI)  $m/z$ : Calcd. for  $C_{30}H_{38}N_4O_5$  534.28  $[M]^+$ , found 535.6  $[M+H]^+$ .

**(*S*)-*N*-(1-(Cyclobutylamino)-1-oxo-4-phenylbutan-2-yl)-5-(2,6-dimethoxyphenyl)-1-isobutyl-1*H*-pyrazole-3-carboxamide (32)**: Following the procedure described in **45**, compound **32** was synthesized using 5-(2,6-dimethoxyphenyl)-1-isobutyl-1*H*-pyrazole-3-carboxylic acid (**43**) and (*S*)-2-amino-*N*-cyclobutyl-4-phenylbutanamide trifluoroacetate. 85 % yield; white solid;  $^1H$  NMR ( $CDCl_3$ , 300 MHz)  $\delta$  0.75 (dd,  $J=6.78, 2.26$  Hz, 6 H), 1.66 – 1.74 (m, 2 H), 1.81 – 1.96 (m, 2 H), 2.01 – 2.18 (m, 2 H), 2.26 – 2.38 (m, 3 H), 2.75 (t,  $J=7.91$  Hz, 2 H), 3.64 (d,  $J=7.16$  Hz, 2 H), 3.75 (s, 6 H), 4.29 – 4.41 (m, 1 H), 4.47 – 4.56 (m, 1 H), 6.50 (d,  $J=7.91$  Hz, 1 H), 6.63 (d,  $J=8.29$  Hz, 2 H), 6.71 (m, 1 H), 7.14 – 7.24 (m, 3 H), 7.28 – 7.33 (m, 3 H), 7.38 (t,  $J=8.48$  Hz, 1 H). MS (ESI)  $m/z$ : Calcd. for  $C_{30}H_{38}N_4O_4$  518.29  $[M]^+$ , found 519.9  $[M+H]^+$ .

**(*S*)-3-(5-(2,6-Dimethoxyphenyl)-1-isobutyl-1*H*-pyrazole-3-carboxamido)-5-phenylpentanoic acid (33)**: Following the procedure described in **12**, compound **33** was obtained from deprotection of (*S*)-*tert*-butyl 3-(5-(2,6-dimethoxyphenyl)-1-isobutyl-1*H*-pyrazole-3-carboxamido)-5-phenylpentanoate. 69 % yield; white solid;  $^1H$  NMR ( $CDCl_3$ , 300 MHz)  $\delta$  0.74 (dd,  $J=6.59, 1.88$  Hz, 6 H), 2.00–2.17 (m, 4 H), 2.72–2.81 (m, 2 H), 2.76 (d,  $J=6.00$  Hz, 2 H), 3.66 (d,  $J=7.54$  Hz, 2 H), 3.73 (s, 3 H), 3.74 (s, 3 H), 4.38 – 4.49 (m, 1 H), 6.62 (d,  $J=8.29$  Hz, 2 H), 6.72 (s, 1 H), 7.15 – 7.23 (m, 3 H), 7.26 – 7.34 (m, 2 H), 7.38 (t,  $J=8.48$  Hz, 1 H), 7.44 (m, 1 H). MS (ESI)  $m/z$ : Calcd. for  $C_{27}H_{33}N_3O_5$  479.24  $[M]^+$ , found 478.7  $[M-H]^-$ .

**(S)-N-(1-(Cyclobutylamino)-1-oxo-5-phenylpentan-3-yl)-5-(2,6-dimethoxyphenyl)-1-isobutyl-1H-pyrazole-3-carboxamide (34)**: Following the procedure described in **45**, compound **34** was synthesized using (*S*)-3-(5-(2,6-dimethoxyphenyl)-1-isobutyl-1*H*-pyrazole-3-carboxamido)-5-phenylpentanoic acid (**33**) and cyclobutylamine. 93 % yield; white solid; <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz) δ 0.74 (d, *J*=6.78 Hz, 6 H), 1.61 – 1.72 (m, 2 H), 1.73 – 1.87 (m, 2 H), 1.92 – 2.18 (m, 3 H), 2.19 – 2.32 (m, 2 H), 2.55 (d, *J*=6.22 Hz, 2 H), 2.68 – 2.79 (m, 2 H), 3.64 (d, *J*=7.35 Hz, 2 H), 3.73 (m, 3 H), 3.74 (m, 3 H), 4.28 – 4.41 (m, 2 H), 6.62 (d, *J*=8.29 Hz, 2 H), 6.70 (s, 1 H), 7.15 – 7.26 (m, 5 H), 7.28 – 7.41 (m, 3 H). MS (ESI) *m/z*: Calcd. for C<sub>31</sub>H<sub>40</sub>N<sub>4</sub>O<sub>4</sub> 532.30 [M]<sup>+</sup>, found 533.7 [M+H]<sup>+</sup>.

**(S)-N-(1-(Cyclobutylamino)-5-cyclohexyl-1-oxopentan-3-yl)-5-(2,6-dimethoxyphenyl)-1-(4-fluorophenyl)-1H-pyrazole-3-carboxamide (35)**: Following the procedure described in **45**, compound **35** was synthesized using (*S*)-5-cyclohexyl-3-(5-(2,6-dimethoxyphenyl)-1-(4-fluorophenyl)-1*H*-pyrazole-3-carboxamido)pentanoic acid (**9**) and cyclobutylamine. 91 % yield; white solid; <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz) δ 0.78 – 0.97 (m, 2 H), 1.10 – 1.40 (m, 7 H), 1.57 – 1.75 (m, 8 H), 1.79 – 1.91 (m, 2 H), 2.21 – 2.35 (m, 2 H), 2.49 – 2.54 (m, 2 H), 3.55 (s, 3 H), 3.60 (s, 3 H), 4.27 – 4.42 (m, 2 H), 6.47 – 6.60 (m, 3 H), 6.94 (s, 1 H), 6.97 (d, *J*=8.29 Hz, 2 H), 7.19 – 7.25 (m, 2 H), 7.27 – 7.34 (m, 2 H). MS (ESI) *m/z*: Calcd. for C<sub>33</sub>H<sub>41</sub>FN<sub>4</sub>O<sub>4</sub> 576.31 [M]<sup>+</sup>, found 577.5 [M+H]<sup>+</sup>.

**(S)-N-(1-(Cyclobutylamino)-5-cyclohexyl-1-oxopentan-3-yl)-5-(2,6-dimethoxyphenyl)-1-propyl-1H-pyrazole-3-carboxamide (36)**: Following the procedure described in **45**, compound **36** was synthesized using (*S*)-5-cyclohexyl-3-(5-(2,6-dimethoxyphenyl)-1-propyl-1*H*-pyrazole-3-carboxamido)pentanoic acid (**11**) and cyclobutylamine. 75 % yield; white solid; <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz) δ ppm 0.78 (t, *J*=7.44 Hz, 3 H), 0.82 – 0.98 (m, 2 H), 1.12 – 1.35 (m, 7 H), 1.62 – 1.95 (m, 12 H), 2.18 – 2.35 (m, 2 H), 2.52 (d, *J*=9.0 Hz, 2 H), 3.73 (s, 3 H), 3.74 (s, 3 H), 3.79 (t, *J*=9.0 Hz, 2 H), 4.20 – 4.30 (m, 1 H), 4.31 – 4.44 (m, 1 H), 6.64 (d, *J*=9.0 Hz, 2 H), 6.65–6.68 (m, 1 H), 6.69 (s, 1 H), 7.10 (d, *J*=8.67 Hz, 1 H), 7.38 (t, *J*=8.38 Hz, 1 H). MS (ESI) *m/z*: Calcd. for C<sub>30</sub>H<sub>44</sub>N<sub>4</sub>O<sub>4</sub> 524.34 [M]<sup>+</sup>, found 526.0 [M+H]<sup>+</sup>

**(S)-N-(1-(Cyclobutylamino)-5-cyclohexyl-1-oxopentan-3-yl)-1-cyclohexyl-5-(2,6-dimethoxyphenyl)-1H-pyrazole-3-carboxamide (38)**: Following the procedure described in **45**, compound **38** was synthesized using (*S*)-5-cyclohexyl-3-(1-cyclohexyl-5-(2,6-dimethoxyphenyl)-1*H*-pyrazole-3-carboxamido)pentanoic acid (**14**) and cyclobutylamine. 61 % yield; white solid; <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz) δ ppm 0.80 – 0.97 (m, 2 H), 1.09 – 1.38 (m, 9 H), 1.58 – 1.76 (m, 10 H), 1.77 – 2.00 (m, 8 H), 2.18 – 2.33 (m, 2 H), 2.53 (d, *J*=6.40 Hz, 2 H), 3.58 – 3.69 (m, 1 H), 3.74 (s, 3 H), 3.73 (s, 3 H), 4.19 – 4.30 (m, 1 H), 4.31 – 4.44 (m, 1 H), 6.63 (d, *J*=9.0 Hz, 2 H), 6.66 (s, 1 H), 6.74 (d, *J*=7.54 Hz, 1 H), 7.08 (d, *J*=9.23 Hz, 1 H), 7.38 (t, *J*=8.38 Hz, 1 H). MS (ESI) *m/z*: Calcd. for C<sub>33</sub>H<sub>48</sub>N<sub>4</sub>O<sub>4</sub> 564.37 [M]<sup>+</sup>, found 565.7 [M+H]<sup>+</sup>.

### Bioassays

**Lance<sup>™</sup> cAMP accumulation assay**: Stimulation buffer containing 1X Hank's Balanced Salt Solution (HBSS), 5 mM HEPES, 0.1% BSA stabilizer, and 0.5 mM final IBMX was

prepared and titrated to pH 7.4 at room temperature. Serial dilutions of the test compounds and 1  $\mu$ M forskolin, both prepared at 4x the desired final concentration in stimulation buffer, were added to a 96-well white  $\frac{1}{2}$  area microplate (PerkinElmer). A cAMP standard curve prepared at 4x the desired final concentration in stimulation buffer was added to the assay plate. The total assay volume for 96-well plate was 40  $\mu$ L. The final DMSO concentration in all wells was 1.5%. Stable hAPJ-CHO cells were lifted with a non-enzymatic solution (Cell-stripper, Mediatech Inc., Orlando, FL) and spun at 270xg for 10 min. The cell pellet was resuspended in stimulation buffer and 4000 cells were added to each well except wells containing the cAMP standard curve. After incubating for 30 min at RT, Eu-cAMP tracer and uLIGHT-anti-cAMP working solutions were added per the manufacturer's instructions. After incubation at RT for 1 h, the TR-FRET signal (ex 337 nm, em 620 and 650 nm) was read on a CLARIOstar multimode plate reader (BMG Biotech, Cary, NC). For the screen, the same procedure was followed except that instead of adding serial dilutions of the test compounds, a single concentration (1  $\mu$ M, prepared at 4x final) of each test compound was added to the assay plates. Data were analyzed using Prism software (GraphPad, La Jolla, CA). Nonlinear regression analysis was performed to fit data and obtain maximum response (Emax), EC<sub>50</sub>, correlation coefficient ( $r^2$ ), and other parameters. All experiments were performed in duplicate at least two times to ensure reproducibility and data are reported as mean  $\pm$  SEM, unless noted otherwise.

**Calcium mobilization assay:** CHO cells stably expressing either the G $\alpha_{q16}$  protein (CHO-RD-HGA16; Molecular Devices) plus the human apelin receptor (CHO hAPJ) or human angiotensin II receptor, type I (CHO hAGTR1; DiscoverX) were removed from flasks using Cell-stripper and quenched with DMEM/F12, 10 % FBS and 1X Penicillin/Streptomycin, centrifuged, and re-suspended in the serum-containing media. Cells were counted with a hemocytometer and 30,000 cells of CHO hAPJ and 20,000 cells of CHO hAGTR1 were transferred to each well of a black Costar 96-well optical bottom plate (Corning Corporation, Corning, NY). Each plate was incubated at 37°C for 24 h. The culture media was removed from the plates, cells were washed with DPBS and were subsequently loaded with a fluorescent calcium probe (Calcium 5 dye; Molecular Devices, Sunnyvale, CA) in an HBSS-based buffer containing 20 mM HEPES, 1% BSA and 10  $\mu$ M probenecid (Sigma) in a total volume of 225  $\mu$ L. Cells were incubated at 37°C for 1h and then stimulated with test compounds or pyrapelin-13 (Anaspec, Fremont, CA) at various concentrations using a FLIPR Tetra plate-reader. Agonist-mediated change in fluorescence (ex 488 nm, em 525 nm) was monitored in each well at 1 sec intervals for 90 sec. Data were collected using Softmax version 4.8 (MDS Analytical Technologies) and analyzed using Prism software. Nonlinear regression analysis was performed to fit data and obtain maximum response (Emax), EC<sub>50</sub>, correlation coefficient ( $r^2$ ), and other parameters. All experiments were performed in duplicate at least two times to ensure reproducibility and data are reported as mean  $\pm$  SEM, unless noted otherwise.

**PathHunter  $\beta$ -arrestin recruitment assay:** The stable CHO-K1 human AGTRL1  $\beta$ -arrestin-2 cell line (DiscoverX) was maintained in TC-treated flasks with growth media containing DMEM/F12 (Corning Cellgro, Manassas, VA), 10% FBS, 1X Penicillin-streptomycin L-glutamine (Gibco, Carlsbad, CA). The selection reagents used were



geneticin (800 µg/mL) and hygromycin (300 µg/mL). The cells were removed from the flasks using Cell stripper. Cells were resuspended in complete media and spun at 300xg for 5 min. The supernatant was discarded and cells were resuspended in Assay Complete Cell plating 2 Reagent (DiscoverX). Cells were counted using hemocytometer and seeded in white clear bottom TC-treated 96-well plate (DiscoverX) at 10,000 cells per well. The plates were incubated at 37°C and 5% CO<sub>2</sub> for 24 hours. Compound dilutions were prepared at 10X final in DPBS (1X) buffer with 10% DMSO and 1% BSA. Compound dilutions were added to the 80–90% confluent assay plate and the plates were incubated at 37°C and 5% CO<sub>2</sub> for 90 min, followed by the addition of PathHunter Detection Reagent (DiscoverX), prepared as per the manufacturer's instructions. The assays were run at 1% DMSO and 0.1% BSA final. The culture plates were incubated another 60 min at room temperature. The luminescent signal was detected on an EnSpire multimode plate reader (PerkinElmer CT, USA). Data were collected using EnSpire Software 4.1 and analyzed using GraphPad Prism. Nonlinear regression analysis was performed to fit data and obtain maximum response (E<sub>max</sub>), EC<sub>50</sub>, correlation coefficient (r<sup>2</sup>), and other parameters. All experiments were performed in duplicate at least two times to ensure reproducibility and data reported as mean ± SEM, unless noted otherwise.

**Radioligand displacement assay**—Competitive inhibition binding was performed using [<sup>125</sup>I]-apelin-13 (Perkin Elmer, Akron, OH). Eleven concentrations (ranging from 10<sup>-11</sup> M to 3.16X10<sup>-5</sup> M) of compound were incubated in assay buffer (0.025 M HEPES, (pH=7.4), 0.01 M MgCl<sub>2</sub>, 0.001 M CaCl<sub>2</sub>, 0.5% fatty acid-free BSA) with 7.5X10<sup>-11</sup> M [<sup>125</sup>I]-apelin-13 (Perkin-Elmer) in the presence of 0.125 µg of CHO-K1 cell membrane homogenates expressing the human apelin receptor (Perkin Elmer). Non-specific binding (NSB) was determined in the presence of 10<sup>-6</sup> M unlabeled apelin-13 (Anaspec). The assays were run in 1.4 mL polypropylene tubes (Thermo Scientific, Waltham, MA) in a 96-well format. All assays were incubated at room temperature on a shaking platform for two hours. Incubation was terminated and bound radioligand was separated from free radioligand by rapid vacuum filtration over 0.5% PEI treated GF/C filter plates using ice-cold 50 mM Tris-HCl (pH=7.4) in a Brandel Scientific (Gaithersburg, MD) 96-well harvester. Filter plates were allowed to dry and Microscint 20 scintillation cocktail (Perkin-Elmer) was added to each well. Bound radioactivity was determined with a TopCount 12-detector instrument (Packard Instruments) using standard scintillation counting techniques. The binding data from each assay plate were normalized to NSB and total counts per minute (CPM) bound values prior to analysis. The K<sub>i</sub> values were calculated from a three-parameter logistic curve fit to the data with Prism (version 6.0, GraphPad Software, Inc., San Diego, CA) using 2.1X10<sup>-10</sup> M as the K<sub>d</sub> for [<sup>125</sup>I]-apelin-13. All experiments were performed in duplicate at least two times to ensure reproducibility and data reported as mean ± SEM, unless noted otherwise.

**Hepatic Liver Microsome Stability Studies**—Human liver microsomal (HLM) stability assays were performed as described previously.<sup>37</sup> Briefly, test compounds were incubated at a 1 µM final concentration with 0.5 mg/ml pooled human liver microsomes from 200 unidentified donors (Xenotech, LLC, Lenexa, KS) in a 100 mM phosphate buffer (pH 7.4) containing 3 mM MgCl<sub>2</sub>, 1 mM nicotinamide adenine dinucleotide phosphate



(NADPH), 5 mM uridine diphosphate glucuronic acid (UDPGA), and 50 µg/ml alamethicin. Triplicate samples were incubated for up to 120 min. Samples were removed at regular intervals. Reactions were terminated by addition of 3 volumes of methanol and processed for LC-MS by centrifugation. Standard curves were prepared in blank matrix for each compound for quantitative assessment. Intrinsic clearance rate was calculated for each compound using the formula:  $Cl_{int} (\mu\text{l}/\text{min}/\text{mg}) = 0.693/(t_{1/2} \times \text{microsomal protein concentration})$ . Data reported are average values from 3 measurements.

**Bias Factor Calculation**—Data from technical replicates on the same day were baseline-corrected and normalized relative to the reference agonist Pyr-Apelin-13. Data from multiple days were then combined and fit to logistic functions with Hill coefficient equal to 1 to determine  $E_{max}$ ,  $EC_{50}$  and their standard errors. Bias factors, which quantify the degree of signaling through pathway 1 compared to pathway 2 for a ligand relative to a reference agonist on a logarithmic scale, were calculated as previously described.<sup>35,36</sup>

$$Bias\ Factor = \log\left(\left(\frac{E_{max,1}}{EC_{50,1}} \frac{EC_{50,2}}{E_{max,2}}\right)_{lig} * \left(\frac{E_{max,2}}{EC_{50,2}} \frac{EC_{50,1}}{E_{max,1}}\right)_{ref}\right)$$

## Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

## ACKNOWLEDGEMENTS

Authors wish to thank Dr. Elaine Gay, Dr. Ann Decker for help with various assays and Dr. Thomas Prisinzano, Victor Day, and the University of Kansas X-Ray Crystallography Laboratory for the crystal structures.

Funding Sources

This work was supported by NIH grants: 1R01HD079547-01A1 and 1R01DK103625-01A1

## ABBREVIATIONS

<b>AT1</b>	Angiotensin 1
<b>NPS</b>	Neuropeptide S receptor
<b>NTR-1</b>	Neurotensin receptor 1
<b>NTR-2</b>	Neurotensin receptor 2
<b>BOP</b>	Benzotriazole-1-yl-oxy-tris-(dimethylamino)-phosphonium hexafluorophosphate
<b>LiOH.H<sub>2</sub>O</b>	Lithium hydroxide monohydrate
<b>TFA</b>	Trifluoroacetic acid
<b>HCl</b>	Hydrochloric acid
<b>Et3N</b>	Triethylamine


<b>DCM</b>	Dichloromethane
<b>THF</b>	Tetrahydrofuran
<b>MeOH</b>	Methanol
<b>SAR</b>	Structure Activity Relationship
<b>TLC</b>	thin layer chromatography

## REFERENCES


- O'Dowd BF; Heiber M; Chan A; Heng HH; Tsui LC; Kennedy JL; Shi X; Petronis A; George SR; Nguyen T A human gene that shows identity with the gene encoding the angiotensin receptor is located on chromosome 11. *Gene* 1993; 136, 355–360. [PubMed: 8294032]
- Tatemoto K; Hosoya M; Habata Y; Fujii R; Kakegawa T; Zou MX; Kawamata Y; Fukusumi S; Hinuma S; Kitada C; Kurokawa T; Onda H; Fujino M Isolation and characterization of a novel endogenous peptide ligand for the human APJ receptor. *Biochem Biophys Res Commun* 1998; 251, 471–476. [PubMed: 9792798]
- Hosoya M; Kawamata Y; Fukusumi S; Fujii R; Habata Y; Hinuma S; Kitada C; Honda S; Kurokawa T; Onda H; Nishimura O; Fujino M Molecular and functional characteristics of APJ. Tissue distribution of mRNA and interaction with the endogenous ligand apelin. *J Biol Chem* 2000; 275, 21061–21067. [PubMed: 10777510]
- Maguire JJ; Kleinz MJ; Pitkin SL; Davenport AP [Pyr(1)]Apelin-13 Identified as the Predominant Apelin Isoform in the Human Heart Vasoactive Mechanisms and Inotropic Action in Disease. *Hypertension* 2009; 54, 598–U296. [PubMed: 19597036]
- Langelan DN; Bebbington EM; Reddy T; Rainey JK Structural insight into G-protein coupled receptor binding by apelin. *Biochemistry* 2009; 48, 537–548. [PubMed: 19123778]
- De Mota N; Goazigo ARL; El Messari S; Chartrel N; Roesch D; Dujardin C; Kordon C; Vaudry H; Moos FO; Llorens-Cortes C Apelin, a potent diuretic neuropeptide counteracting vasopressin actions through inhibition of vasopressin neuron activity and vasopressin release. *P Natl Acad Sci USA* 2004; 101, 10464–10469.
- Azizi M; Iturriz X; Blanchard A; Peyrard S; De Mota N; Chartrel N; Vaudry H; Corvol P; Llorens-Cortes C Reciprocal regulation of plasma apelin and vasopressin by osmotic stimuli. *J Am Soc Nephrol* 2008; 19, 1015–1024. [PubMed: 18272843]
- Murza A; Parent A; Besserer-Offroy E; Tremblay H; Karadereye F; Beaudet N; Leduc R; Sarret P; Marsault E Elucidation of the structure-activity relationships of apelin: influence of unnatural amino acids on binding, signaling, and plasma stability. *Chemmedchem* 2012; 7, 318–325. [PubMed: 22170700]
- Murza A; Belleville K; Longpre JM; Sarret P; Marsault E Stability and Degradation Patterns of Chemically Modified Analogs of Apelin-13 in Plasma and Cerebrospinal Fluid. *Biopolymers* 2014; 102, 297–303. [PubMed: 24728860]
- Pauli A; Norris ML; Valen E; Chew GL; Gagnon JA; Zimmerman S; Mitchell A; Ma J; Dubrulle J; Reyon D; Tsai SQ; Joung JK; Saghatelian A; Schier AF Toddler: An Embryonic Signal That Promotes Cell Movement via Apelin Receptors. *Science* 2014; 343, 746–+.
- Chng SC; Ho LN; Tian J; Reversade B ELABELA: A Hormone Essential for Heart Development Signals via the Apelin Receptor. *Dev Cell* 2013; 27, 672–680. [PubMed: 24316148]
- Laflamme B ELABELA, a peptide hormone for heart development. *Nat Genet* 2014; 46, 7.
- Szokodi I; Tavi P; Foldes G; Voutilainen-Myllyla S; Ilves M; Tokola H; Pikkariainen S; Piihola J; Rysa J; Toth M; Ruskoaho H Apelin, the novel endogenous ligand of the orphan receptor APJ, regulates cardiac contractility. *Circ Res* 2002; 91, 434–440. [PubMed: 12215493]
- Lee DK; Jeong JH; Oh S; Jo YH Apelin-13 Enhances Arcuate POMC Neuron Activity via Inhibiting M-Current. *PLoS One* 2015; 10, e0119457. [PubMed: 25782002]

15. El Messari S; Iturrioz X; Fassot C; De Mota N; Roesch D; Llorens-Cortes C Functional dissociation of apelin receptor signaling and endocytosis: implications for the effects of apelin on arterial blood pressure. *J Neurochem* 2004; 90, 1290–1301. [PubMed: 15341513]
16. De Mota N; Lenkei Z; Llorens-Cortes C Cloning, pharmacological characterization and brain distribution of the rat apelin receptor. *Neuroendocrinology* 2000; 72, 400–407. [PubMed: 11146423]
17. Reaux-Le Goazigo A; Morinville A; Burllet A; Llorens-Cortes C; Beaudet A Dehydration-induced cross-regulation of apelin and vasopressin immunoreactivity levels in magnocellular hypothalamic neurons. *Endocrinology* 2004; 145, 4392–4400. [PubMed: 15166125]
18. Medhurst AD; Jennings CA; Robbins MJ; Davis RP; Ellis C; Winborn KY; Lawrie KW; Hervieu G; Riley G; Bolaky JE; Herrity NC; Murdock P; Darker JG Pharmacological and immunohistochemical characterization of the APJ receptor and its endogenous ligand apelin. *J Neurochem* 2003; 84, 1162–1172. [PubMed: 12603839]
19. Reaux-Le Goazigo A; Alvear-Perez R; Zizzari P; Epelbaum J; Bluet-Pajot MT; Llorens-Cortes C Cellular localization of apelin and its receptor in the anterior pituitary: evidence for a direct stimulatory action of apelin on ACTH release. *Am J Physiol Endocrinol Metab* 2007; 292, E7–15. [PubMed: 16896162]
20. Yokomori H; Oda M; Yoshimura K; Machida S; Kaneko F; Hibi T Overexpression of apelin receptor (APJ/AGTRL1) on hepatic stellate cells and sinusoidal angiogenesis in human cirrhotic liver. *J Gastroenterol* 2011; 46, 222–231. [PubMed: 20725750]
21. Narayanan S; Harris DL; Maitra R; Runyon SP Regulation of the Apelinergic System and Its Potential in Cardiovascular Disease: Peptides and Small Molecules as Tools for Discovery. *J Med Chem* 2015; 58, 7913–7927. [PubMed: 26102594]
22. Wysocka MB; Pietraszek-Gremplewicz K; Nowak D The Role of Apelin in Cardiovascular Diseases, Obesity and Cancer. *Front Physiol* 2018; 9, 557. [PubMed: 29875677]
23. Papangeli I; Kim J; Maier I; Park S; Lee A; Kang Y; Tanaka K; Khan OF; Ju H; Kojima Y; Red-Horse K; Anderson DG; Siekmann AF; Chun HJ MicroRNA 139–5p coordinates APLNR-CXCR4 crosstalk during vascular maturation. *Nature communications* 2016; 7, 11268.
24. Iturrioz X; Alvear-Perez R; De Mota N; Franchet C; Guillier F; Leroux V; Dabire H; Le Jouan M; Chabane H; Gerbier R; Bonnet D; Berdeaux A; Maigret B; Galzi JL; Hibert M; Llorens-Cortes C Identification and pharmacological properties of E339–3D6, the first nonpeptidic apelin receptor agonist *Faseb J* 2010; 24, 1506–1517. [PubMed: 20040517]
25. Khan P; Maloney PR; Hedrick M; Gosalia P; Milewski M; Li L; Roth GP; Sergienko E; Suyama E; Sugarman E; Nguyen K; Mehta A; Vasile S; Su Y; Shi S; Stonich D; Nguyen H; Zeng FY; Novo AM; Vicchiarelli M; Diwan J; Chung TDY; Pinkerton AB; Smith LH In Probe Reports from the NIH Molecular Libraries Program Bethesda (MD), 2010.
26. Hachtel SW, Wolfart P; Weston J; Mueller M; Defossa E; Mertsch K; Weng JH; Binnie RA; Abdul-Latif F; Bock WJ; Walser A; ; Benzimidazole-carboxylic acid amide derivatives as APJ Receptor Modulators.; US20140094450 A1; Sanofi[Fr]: 2014.
27. Pinkerton ABS, L. H.; Agonists of the Apelin Receptor and Methods of Use Thereof.; WO2015/184011 A2; Sanford-Burnham Medical Research Institute: 2015.
28. Meng WC, H. J.; Finlay H; Lawrence RM; Myers MC Apelin Receptor Agonists and Methods of Use. WO2017096130A1 2017, Bristo-Myers Squibb Company.
29. Chen XC, Y.; Cheng AC; Connors RV; Debenedetti MV; Dransfield PJ; Fu Z; Harvey JS; Heath JA; Hedley SJ; Houze J; Judd TC; Khakoo AY; Kopecky DJ; Lai SJ; Ma Z; Nishimura N; Olson SH; Pattaropong V; Swaminath G; Wang X; Yeh WC; Heterocyclic Triazole Compounds as Agonists of the APJ Receptor. WO2017192485A1 2017, Amgen Inc.
30. Read C; Fitzpatrick CM; Yang P; Kuc RE; Maguire JJ; Glen RC; Foster RE; Davenport AP Cardiac action of the first G protein biased small molecule apelin agonist. *Biochemical pharmacology* 2016; 116, 63–72. [PubMed: 27475715]
31. Brame AL; Maguire JJ; Yang P; Dyson A; Torella R; Cheriyan J; Singer M; Glen RC; Wilkinson IB; Davenport AP Design, characterization, and first-in-human study of the vascular actions of a novel biased apelin receptor agonist. *Hypertension* 2015; 65, 834–840. [PubMed: 25712721]

32. Trifonov L; Afri M; Palczewski K; Korshin EE; Gruzman A An Expedient Synthesis of CMF-019: (S)-5-Methyl-3-[1-(pentan-3-yl)-2-(thiophen-2-ylmethyl)-1H-benzo[d]imidazole-5-carboxamido]hexanoic Acid, a Potent Apelin Receptor (APJ) Agonist. *Med Chem* 2018; 14, 688–694. [PubMed: 29651942]
33. Narayanan S; Maitra R; Deschamps JR; Bortoff K; Thomas JB; Zhang Y; Warner K; Vasukuttan V; Decker A; Runyon SP Discovery of a novel small molecule agonist scaffold for the APJ receptor. *Bioorg Med Chem* 2016; 24, 3758–3770. [PubMed: 27369451]
34. Davies SG; Mulvaney AW; Russell AJ; Smith AD Parallel synthesis of homochiral  $\alpha$ -amino acids. *Tetrahedron: Asymmetry* 2007; 18, 1554–1566.
35. Rajagopal S; Ahn S; Rominger DH; Gowen-MacDonald W; Lam CM; DeWire SM; Violin JD; Lefkowitz RJ Quantifying Ligand Bias at Seven-Transmembrane Receptors. *Molecular Pharmacology* 2011; 80, 367–377. [PubMed: 21610196]
36. Onaran HO; Ambrosio C; Ugur O; Koncz EM; Gro MC; Vezzi V; Rajagopal S; Costa T Systematic errors in detecting biased agonism: Analysis of current methods and development of a new model-free approach. *Sci Rep-Uk* 2017; 7.
37. Fulp A; Bortoff K; Seltzman H; Zhang YA; Mathews J; Snyder R; Fennell T; Maitra R Design and Synthesis of Cannabinoid Receptor 1 Antagonists for Peripheral Selectivity. *J Med Chem* 2012; 55, 2820–2834. [PubMed: 22372835]




**Figure 1.**  
 Non-peptide apelin receptor agonists reported in the literature




**Figure 2.**  
 Chemical structure of initial hit **8**, identified through focused screen and optimized to lead compound **9**




**Scheme 1:**Synthesis of compound **25** and **37**


*Reagents and conditions:* a) Diethyl oxalate, EtONa, EtOH, reflux, 6 h, 97%; b) Isobutylhydrazine trifluoroacetate/cyclopentylhydrazine trifluoroacetate, EtOH, reflux, 16 h; 40/73% c) LiOH, MeOH/THF/H<sub>2</sub>O, rt, 18 h, 96/98%; d) (*S*)-*tert*-butyl 3-amino-5-cyclohexylpentanoate, BOP, Et<sub>3</sub>N, THF, rt, 1.5 h, 91/87%; e) TFA, DCM, rt, 1.5 h, 54/94%; f) cyclobutylamine, BOP, Et<sub>3</sub>N, THF, rt, 3 h, 61/75%.



**Figure 3.**  
Crystal structure of compound **43**




**Figure 4.**  
Crystal structure of compound **44**






**Figure 5.** Radar plots of bias factors for (A) compounds that display bias towards calcium signaling and (B) compounds that have minimal (< 10-fold) bias towards any signaling pathways compared to the reference agonist Pyr-Apelin-13. Bias factors are shown on a logarithmic scale.

Table 1.

Modification at N1 pyrazole core of compound with cyclohexyl ethyl amino acid side chain



Compd.	R	APJ Calcium EC <sub>50</sub> * μM±SEM (%E <sub>max</sub> )	APJ cAMP EC <sub>50</sub> * μM±SEM (%E <sub>max</sub> )	APJ β-arrestin EC <sub>50</sub> * μM±SEM (%E <sub>max</sub> )	APJ Binding K <sub>i</sub> * (μM±SEM)	AT1 Calcium EC <sub>50</sub> * (μM±SEM)
Pyr-Apelin-13	-	0.002 ± 0.0003	0.0003 ± 0.0001	0.001 ± 0.0005	0.001 ± 0.0005	ND*
9.		0.800 ± 0.1 (91)	0.844 ± 0.2 (103)	6.22 ± 2.0 (86)	1.3 ± 0.3	>10
10.	CH <sub>3</sub>	1.46 ± 0.4 (65)	2.55 ± 1.6 (106)	9.94 ± 1.4 (75)	ND	ND
11.	CH <sub>2</sub> -CH <sub>2</sub> -CH <sub>3</sub>	0.416 ± 0.02 (89)	0.281 ± 0.09 (109)	3.28 ± 0.4 (67)	ND	ND
12.		0.649 ± 0.1 (101)	0.915 ± 0.5 (108)	5.59 ± 0.06 (81)	ND	ND
13.		0.424 ± 0.07 (96)	0.238 ± 0.05 (103)	2.47 ± 0.4 (77)	ND	ND


Compd.	R	APJ Calcium EC <sub>50</sub> <sup>*</sup> μM±SEM (%E <sub>max</sub> )	APJ cAMP EC <sub>50</sub> <sup>*</sup> μM±SEM (%E <sub>max</sub> )	APJ β-arrestin EC <sub>50</sub> <sup>*</sup> μM±SEM (%E <sub>max</sub> )	APJ Binding K <sub>i</sub> <sup>*</sup> (μM±SEM)	AT1 Calcium EC <sub>50</sub> <sup>*</sup> (μM±SEM)
14.		0.380 ± 0.03 (91)	0.518 ± 0.08 (113)	3.26 ± 0.08 (83)	ND	ND
15.		>10	>10	>10	ND	ND
16.		>10	>10	>10	ND	ND

\* EC<sub>50</sub> and K<sub>i</sub> values are averages of multiple experiments performed in duplicate unless otherwise stated ± standard error of the mean.






\* ND = Not Done



Table 2.

Modification of carboxylic acid moiety of **12**

Compd.	R	APJ Calcium EC <sub>50</sub> <sup>*</sup> μM±SEM (%E <sub>max</sub> )	APJ cAMP EC <sub>50</sub> <sup>*</sup> μM±SEM (%E <sub>max</sub> )	APJ β-arrestin EC <sub>50</sub> <sup>*</sup> μM±SEM (%E <sub>max</sub> )	APJ Binding K <sub>i</sub> <sup>*</sup> (μM±SEM)	AT1 Calcium EC <sub>50</sub> <sup>*</sup> (μM±SEM)
17.	-NH <sub>2</sub>	5.75 ± 1.1 (94)	>10	7.24 ± 0.9 (133)	ND <sup>*</sup>	ND
18.		0.288 ± 0.03 (92)	0.684 ± 0.1 (105)	0.280 ± 0.001 (89)	ND	ND
19.		0.237 ± 0.04 (91)	1.85 ± 0.2 (106)	0.453 ± 0.014 (87)	ND	ND
20.		1.85 ± 0.5 (77)	4.82 ± 1.1 (129)	8.87 ± 2.7 (87)	ND	ND


Compd.	R	APJ Calcium EC <sub>50</sub> <sup>*</sup> μM±SEM (%E <sub>max</sub> )	APJ cAMP EC <sub>50</sub> <sup>*</sup> μM±SEM (%E <sub>max</sub> )	APJ β-arrestin EC <sub>50</sub> <sup>*</sup> μM±SEM (%E <sub>max</sub> )	APJ Binding K <sub>i</sub> <sup>*</sup> (μM±SEM)	AT1 Calcium EC <sub>50</sub> <sup>*</sup> (μM±SEM)
21.		0.046 ± 0.008 (88)	0.118 ± 0.02 (104)	0.113 ± 0.02 (89)	0.036 ± 0.006	>10
22.		0.069 ± 0.02 (139)	0.157 ± 0.08 (104)	0.114 ± 0.005 (90)	0.038 ± 0.01	>10
23.		0.144 ± 0.03 (102)	0.246 ± 0.02 (106)	0.182 ± 0.045 (87)	0.172 ± 0.1	>10
24.		1.70 ± 0.04 (94)	3.17 ± 0.8 (134)	0.313 ± 0.04 (81)	ND	ND
25.		0.081 ± 0.02 (91)	0.283 ± 0.08 (97)	0.144 ± 0.008 (90)	0.054 ± 0.01	>10

\* EC<sub>50</sub> and K<sub>i</sub> values are averages of multiple experiments performed in duplicate unless otherwise stated ± standard error of the mean.





\* ND = Not Done

Table 3.

Modification of carboxylic acid moiety of 26.



Compd.	R	n	APJ Calcium EC <sub>50</sub> <sup>*</sup> μM±SEM (%E <sub>max</sub> )	APJ cAMP EC <sub>50</sub> <sup>*</sup> μM±SEM (%E <sub>max</sub> )	APJ β-arrestin EC <sub>50</sub> <sup>*</sup> μM±SEM (%E <sub>max</sub> )	APJ Binding K <sub>i</sub> <sup>*</sup> (μM±SEM) (%E <sub>max</sub> )	AT1 Calcium EC <sub>50</sub> <sup>*</sup> (μM±SEM)
26.	-OH	0	>10	>10	>10	ND <sup>*</sup>	ND
27.		0	0.021±0.007 (131)	3.09 ± 3.0 (124)	2.33 ± 0.9 (119)	0.187± 0.06	ND
28.		0	0.027 ± 0.02 (106)	0.266 ± 0.057 (106)	0.207 ± 0.002 (99)	0.134± 0.09	>10
29.		0	0.411± 0.09 (97)	>10	>10	ND	ND


Compd.	R	n	APJ Calcium EC <sub>50</sub> * μM±SEM (%E <sub>max</sub> )	APJ cAMP EC <sub>50</sub> * μM±SEM (%E <sub>max</sub> )	APJ β-arrestin EC <sub>50</sub> * μM±SEM (%E <sub>max</sub> )	APJ Binding K <sub>i</sub> * (μM±SEM) (%E <sub>max</sub> )	AT1 Calcium EC <sub>50</sub> * (μM±SEM)
30.		0	0.313± 0.01 (93)	>10	>10	ND	ND
31.		0	0.162± 0.001 (98)	>10	8.67± 0.1 (54)	0.571± 0.3	>10
32.		0	1.26 ± 0.8 (81)	3.48± 0.1 (13)	>10	ND	ND
33.	-OH	1	>10	>10	>10	ND	ND
34.		1	0.083± 0.01 (96)	>10	3.11 ± 0.2 (80)	0.449± 0.08	>10

\* EC<sub>50</sub> and K<sub>i</sub> values are averages of multiple experiments performed in duplicate ± standard error of the mean.


\* ND = Not Done

Table 4.

Modification at N1 position of pyrazole core



Compd.	R	APJ Calcium EC <sub>50</sub> <sup>*</sup> μM±SEM (%E <sub>max</sub> )	APJ cAMP EC <sub>50</sub> <sup>*</sup> μM±SEM (%E <sub>max</sub> )	APJ β-arrestin EC <sub>50</sub> <sup>*</sup> μM±SEM (%E <sub>max</sub> )	APJ Binding K <sub>i</sub> <sup>*</sup> μM±SEM, (%E <sub>max</sub> )	AT1 Calcium EC <sub>50</sub> <sup>*</sup> (μM)
35.		0.425 ± 0.06 (45)	0.344 ± 0.004 (91)	0.422 ± 0.008 (75)	ND <sup>*</sup>	ND
36.		0.049 ± 0.004 (91)	0.107 ± 0.046 (110)	0.078 ± 0.02 (77)	0.060 ± 0.03	>10
25.		0.081 ± 0.025 (91)	0.282 ± 0.08 (97)	0.144 ± 0.008 (90)	0.054 ± 0.01	>10
37.		0.070 ± 0.009 (94)	0.097 ± 0.032 (106)	0.063 ± 0.004 (89)	0.059 ± 0.02	>10

Compd.	R	APJ Calcium EC <sub>50</sub> <sup>*</sup> μM±SEM (%E <sub>max</sub> )	APJ cAMP EC <sub>50</sub> <sup>*</sup> μM±SEM (%E <sub>max</sub> )	APJ β-arrestin EC <sub>50</sub> <sup>*</sup> μM±SEM (%E <sub>max</sub> )	APJ Binding K <sub>i</sub> <sup>*</sup> μM±SEM), (%E <sub>max</sub> )	AT1 Calcium EC <sub>50</sub> <sup>*</sup> (μM)
38.		0.101 ± 0.001 (87)	0.151 ± 0.04 (101)	0.092 ± 0.007 (78)	0.147 ± 0.06	>10

\*EC<sub>50</sub> and K<sub>i</sub> values are averages of multiple experiments performed in duplicate unless otherwise stated ± standard error of the mean.

\*ND = Not Done

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript



**Table 5:**

Microsomal stability of synthesized compounds

Compd.	Half-life, $t_{1/2}$ (min)	Clearance, $CL_{INT}$ (mL/min/kg)
7-ethoxy-4-methylcoumarin	3.84	325
9.	229.59	5
21.	3.21	389
22.	3.10	402
23.	3.55	351
25.	3.26	383
28.	8.18	153
31.	34.71	36
34.	3.59	347
36.	2.90	431
37.	6.25	199
38.	6.28	199

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

**Table 6:**

Bias factors for a series of APJ agonists (shown in dimensionless units on a logarithmic scale  $\pm$  standard error of the mean). Positive values denote bias towards the first pathway listed and negative values denote bias towards the second pathway listed (pathway 1: pathway 2).

Compd.	Bias Ca:cAMP	Bias Ca: $\beta$ arr	Bias cAMP: $\beta$ arr
<b>Pyr-Apelin-13</b>	0 $\pm$ 0.136	0 $\pm$ 0.194	0 $\pm$ 0.180
<b>9.</b>	0.793 $\pm$ 0.172	0.89 $\pm$ 0.239	0.097 $\pm$ 0.243
<b>10.</b>	0.743 $\pm$ 0.229	0.952 $\pm$ 0.189	0.209 $\pm$ 0.23
<b>11.</b>	0.549 $\pm$ 0.133	1.18 $\pm$ 0.153	0.634 $\pm$ 0.157
<b>12.</b>	0.947 $\pm$ 0.173	1.13 $\pm$ 0.17	0.182 $\pm$ 0.178
<b>13.</b>	0.606 $\pm$ 0.163	1.06 $\pm$ 0.157	0.454 $\pm$ 0.189
<b>14.</b>	0.895 $\pm$ 0.132	1.14 $\pm$ 0.164	0.241 $\pm$ 0.167
<b>19.</b>	1.75 $\pm$ 0.169	0.509 $\pm$ 0.185	-1.24 $\pm$ 0.199
<b>27.</b>	2.45 $\pm$ 0.149	1.95 $\pm$ 0.215	-0.498 $\pm$ 0.231
<b>28.</b>	2.27 $\pm$ 0.217	1.56 $\pm$ 0.392	-0.709 $\pm$ 0.344
<b>29.</b>	2.73 $\pm$ 0.39	2.82 $\pm$ 0.805	0.095 $\pm$ 0.882
<b>30.</b>	2.75 $\pm$ 0.376	2.46 $\pm$ 0.32	-0.288 $\pm$ 0.479
<b>31.</b>	2.72 $\pm$ 0.161	2.14 $\pm$ 0.219	-0.584 $\pm$ 0.243
<b>34.</b>	2.45 $\pm$ 0.17	1.8 $\pm$ 0.193	-0.65 $\pm$ 0.186