



Sexually dimorphic role of oxytocin in medaka mate choice

Saori Yokoi^{a,b,1}, Kiyoshi Naruse^b, Yasuhiro Kamei^b, Satoshi Ansai^{b,c}, Masato Kinoshita^d, Mari Mito^e, Shintaro Iwasaki^{e,f}, Shuntaro Inoue^a, Teruhiro Okuyama^g, Shinichi Nakagawa^a, Larry J. Young^{h,i,j}, and Hideaki Takeuchi^{c,k,1}

^aFaculty of Pharmacological Sciences, Hokkaido University, 060-0812 Hokkaido, Japan; ^bNational Institute for Basic Biology, 444-8585 Aichi, Japan; ^cGraduate School of Life Sciences, Tohoku University, 980-8577 Miyagi, Japan; ^dDivision of Applied Biosciences, Graduate school of Agriculture, Kyoto University, 606-8502 Kyoto, Japan; ^eRNA Systems Biochemistry Laboratory, RIKEN Cluster for Pioneering Research, 351-0198 Wako, Saitama, Japan; ^fDepartment of Computational Biology and Medical Sciences, Graduate School of Frontier Sciences, The University of Tokyo, 277-8561 Kashiwa, Chiba, Japan; ^gInstitute for Quantitative Biosciences, The University of Tokyo, 113-0032 Tokyo, Japan; ^hCenter for Social Neural Networks, University of Tsukuba, 305-8577 Tsukuba, Japan; ⁱCenter for Translational Social Neuroscience, Department of Psychiatry and Behavioral Sciences, Emory University School of Medicine, Atlanta, GA 30322; ^jSilvio O. Conte Center for Oxytocin and Social Cognition, Yerkes National Primate Research Center, Emory University, Atlanta, GA 30329; and ^kGraduate School of Natural Science and Technology, Okayama University, 700-8530 Okayama, Japan

Edited by Gene E. Robinson, University of Illinois at Urbana–Champaign, Urbana, IL, and approved January 15, 2020 (received for review December 9, 2019)

Oxytocin is a central neuromodulator required for facilitating mate preferences for familiar individuals in a monogamous rodent (prairie vole), irrespective of sex. While the role of oxytocin in mate choice is only understood in a few monogamous species, its function in nonmonogamous species, comprising the vast majority of vertebrate species, remains unclear. To address this issue, we evaluated the involvement of an oxytocin homolog (isotocin, referred herein as *oxt*) in mate choice in medaka fish (*Oryzias latipes*). Female medaka prefer to choose familiar mates, whereas male medaka court indiscriminately, irrespective of familiarity. We generated mutants of the *oxt* ligand (*oxt*) and receptor genes (*oxtr1* and *oxtr2*) and revealed that the *oxt-oxtr1* signaling pathway was essential for eliciting female mate preference for familiar males. This pathway was also required for unrestricted and indiscriminate mating strategy in males. That is, either *oxt* or *oxtr1* mutation in males decreased the number of courtship displays toward novel females, but not toward familiar females. Further, males with these mutations exhibited enhanced mate-guarding behaviors toward familiar females, but not toward novel females. In addition, RNA-sequencing (seq) analysis revealed that the transcription of genes involved in gamma-amino butyric acid metabolism as well as those encoding ion-transport ATPase are up-regulated in both *oxt* and *oxtr1* mutants only in female medaka, potentially explaining the sex difference of the mutant phenotype. Our findings provide genetic evidence that *oxt-oxtr1* signaling plays a role in the mate choice for familiar individuals in a sex-specific manner in medaka fish.

sexual preference | mate guarding | social recognition | genome editing

Oxytocin is a neuropeptide considered to contribute to the biological basis of prosociality for familiar individuals. For example, the oxytocin system facilitates maternal behaviors and mother-kin bonding (1, 2). In some monogamous rodents (e.g., prairie voles), oxytocin has an essential role in both sexes to enhance the preference for familiar individuals (3–5). Little attention, however, has been paid to the function of oxytocin in mate choice in nonmonogamous species. Mate choice based on familiarity recognition is divergent and variable across species and, in most nonmonogamous species, females and males exhibit different mating strategies (3, 6–9). Thus, studies on the sexual difference in nonmonogamous species offering a comparative perspective will broaden our understanding of the role of oxytocin in mate choice. Medaka is a fascinating model for studying the molecular underpinnings of sexual dimorphism in mate choice because the male and female mating strategies are distinct. Under laboratory conditions, we previously demonstrated that female medaka prefer to mate with visually familiar males (10, 11); familiarization could facilitate the selection of dominant males that

prominently exhibit mate-guarding behavior in triadic relationships (two males and one female) (12, 13). We also found that male medaka did not exhibit any preference based on familiarity. Here, we performed a functional dissection of the oxytocin system using a series of medaka mutants and demonstrated a sexually dimorphic role of the oxytocin signaling pathway (*oxt-oxtr1*) in mate choice.

Results

Generation of Medaka with Oxytocin and Oxytocin Receptor Gene Mutations. Among teleost fish, the homolog of the nonapeptide oxytocin is isotocin, which has two amino acid substitutions at positions 4 (Gln to Ser) and 8 (Leu to Ile) in the mammalian OXT (*SI Appendix, Fig. S1B*). A series of medaka-carrying mutations of genes encoding *oxt* and its receptors were generated using TILLING (Targeting Induced Local Lesions In Genomes) (14, 15), TALEN (transcription activator-like effector nucleases) (16), and CRISPR/Cas9 (Clustered Regularly Interspaced Short Palindromic Repeat), which is the most widely used technology for genome editing using RNA-guided endonuclease (Cas) (17)

Significance

In some monogamous species, the oxytocin system facilitates mating preference for a familiar individual, irrespective of sex. In contrast, in nonmonogamous species, the mating strategy mostly differs between males and females; males tend to compete with rival males and court indiscriminately, whereas females choose their mates selectively. Here, we studied the role of oxytocin in mate choice in medaka fish, a nonmonogamous species. Analysis of oxytocin system mutants in medaka revealed that the oxytocin system regulates mate preference in a sex-specific manner, indicating that the oxytocin system is required for mate choice in both sexes, regardless of the differences in the mating strategy between monogamous and nonmonogamous species.

Author contributions: S.Y., K.N., S.N., and H.T. designed research; S.Y., S.A., M.M., and S. Inoue performed research; K.N., Y.K., S.A., M.K., M.M., S. Iwasaki, and T.O. contributed new reagents/analytic tools; S.Y. analyzed data; and S.Y., L.J.Y., and H.T. wrote the paper.

The authors declare no competing interest.

This article is a PNAS Direct Submission.

Published under the PNAS license.

Data deposition: RNA-seq data used in this study have been deposited in the Gene Expression Omnibus (GEO) database, <https://www.ncbi.nlm.nih.gov/geo> (accession no. GSE139158).

¹To whom correspondence may be addressed. Email: yokois@pharm.hokudai.ac.jp or hideaki.takeuchi.a8@tohoku.ac.jp.

This article contains supporting information online at <https://www.pnas.org/lookup/suppl/doi:10.1073/pnas.1921446117/-DCSupplemental>.

First published February 18, 2020.

(SI Appendix, Table S1). Using TILLING, we first generated *oxt* mutant fish (*oxt*^{I22F/I22F}), in which a conserved isoleucine residue was changed to phenylalanine (SI Appendix, Fig. S1). A previous study (18) indicated that this single amino acid substitution decreases the affinity for the receptor and diminishes the biological efficacy of oxytocin. We also generated another *oxt* mutant fish (*oxt*^{-/-}) using CRISPR/Cas9. A 4-bp deletion in the first exon in the mutant resulted in a frameshift upstream of *oxt* (SI Appendix, Fig. S2).

In addition, we created medaka mutants of genes for *oxt* receptors. Like in cichlid (19), we found two paralogous genes encoding *oxtr* receptors (*oxtr1* [Ensembl gene ID: ENSORLG00000000719] and *oxtr2* [Ensembl gene ID: ENSORLG00000003297], respectively) in the medaka genome. As neither *oxtr1* nor *oxtr2* expression has been reported in medaka fish, we confirmed the expression of these genes in the brain using RT-PCR (SI Appendix, Fig. S3A). Both *oxtr1* and *oxtr2* were expressed in the adult brain in both sexes. To evaluate the role of the *oxtr*, we generated *oxtr1* and *oxtr2* mutants: The *oxtr1* mutants (*oxtr1*^{-/-}, generated by TALEN) and *oxtr2* mutants (*oxtr2*^{-/-}, generated by CRISPR/Cas9) have 7-bp and 8-bp deletions in the first exon, respectively (SI Appendix, Fig. S3 B–D), and both have mutated transcripts encoding C-terminal-deleted proteins that lack at least six of the seven transmembrane domains. Considering that *oxtr1* and *oxtr2* are seven-transmembrane receptors and the topology of the deletions along the domain structures (SI Appendix, Fig. S3B), these deletions should lead to a loss of function.


***oxt* and *oxtr1* Mutant Females Exhibited a Loss of Mate Preference for Visually Familiarized Males.** Wild-type (WT) medaka females can recognize and select visually familiarized males as their mating partners (10, 11). We quantified the effect of visual familiarization (VF; allowing a pair of fish can see each other through a transparent wall for a short-time: 1 d) on the degree of female receptivity in a series of mutants. This short period of VF decreased the latency to mate on the basis of the shorter time interval between the first male courtship behavior and the first mating in WT females (Mann–Whitney *U* test: *Z* = -2.608, *P* = 0.0091; Fig. 1A). The shorter latency correlated with higher female receptivity. The *oxt*^{I22F/I22F} and *oxtr1*^{-/-} females, however, had a short latency to mate even without VF and exhibited high receptivity for any male (Mann–Whitney *U* test: *Z* = -0.340, *P* = 0.733; *Z* = 0.340, *P* = 0.733, respectively, Fig. 1A and B), suggesting a loss of mate preference. However, *oxt*^{+I22F} and *oxtr2*^{-/-} females, like WT fish, exhibited a mate preference for familiar males (Mann–Whitney *U* test: *Z* = -2.494, *P* = 0.012; *Z* = -2.532, *P* = 0.011, respectively, Fig. 1A and B). We confirmed that the other *oxt* mutant female (deletion mutant: *oxt*^{-/-}) exhibited the same behavioral phenotype as the *oxt*^{I22F/I22F} mutant (Mann–Whitney *U* test: *Z* = -0.265, *P* = 0.791; SI Appendix, Fig. S4A),

indicating that this phenotype was due to loss of function. Moreover, we confirmed that none of these mutations affected visual responses or locomotion in the males (Kruskal–Wallis χ^2 [3, *n* = 16] = 5.801, *P* = 0.12; SI Appendix, Fig. S5). Taken together, these findings indicated that *oxt* and *oxtr1* are required for mate preference in medaka females.

***oxt* and *oxtr1* Males Preferred to Mate with Socially Familiarized Females.** We previously reported that short-time VF did not influence male mating preference (10). Consistent with this observation, even long-time social familiarization (SF; group rearing in a single tank) had no effect on mate preference in WT males (Fig. 2A). Thus, WT males have no mate preference based on familiarity. To examine the effect of the *oxt* mutation on male courtship activity toward socially unfamiliar or familiar females, we performed a courtship behavior assay using females whose genotype was the same as that of the males (SF group) and females whose genotype differed from that of the males (non-SF group). In the experiments, individuals with the same genotype were group reared from birth, i.e., socially familiarized for a long period of time. Whereas in WT and *oxt*^{+I22F} males, neither SF nor the female genotype significantly affected the number of courtship behaviors (Steel–Dwass’s *P* > 0.999; Fig. 2A), the number of courtship behaviors of homozygous *oxt*^{I22F/I22F} males in the non-SF group was significantly lower than that of WT males in the non-SF group (Kruskal–Wallis χ^2 [5, *n* = 66] = 19.602, *P* = 0.0014, WT male, *oxt*^{I22F/I22F} female, non-SF vs. *oxt*^{I22F/I22F} male, WT female, non-SF; Steel–Dwass’s *P* = 0.013; Fig. 2A).

There are two possible interpretations of these results. The WT females could react *oxt*^{I22F/I22F} males differently, which might decrease male courtship behaviors. Alternatively, the *oxt*^{I22F/I22F} males could recognize socially familiarized females and had a mate preference for socially familiarized females, as the courtship behaviors of *oxt*^{I22F/I22F} males in the SF group were not significantly different from that of WT (WT male, WT female, SF vs. *oxt*^{I22F/I22F} male, *oxt*^{I22F/I22F} female, SF; Steel–Dwass’s *P* = 0.989, WT male, *oxt*^{I22F/I22F} female, non-SF vs. *oxt*^{I22F/I22F} male, *oxt*^{I22F/I22F} female, non-SF; Steel–Dwass’s *P* = 0.969; Fig. 2A). To evaluate these possibilities, we examined whether SF enhance the courtship activities in *oxt*^{I22F/I22F} males toward unfamiliar WT females that had been reared in other tanks. We group reared *oxt*^{I22F/I22F} males and WT females in the same tank for a certain period of time (1–20 d) and examined the effect of SF on the number of courtship behaviors (Fig. 2B). Without SF, *oxt*^{I22F/I22F} males exhibited courtship behaviors approximately three times within 5 min. In contrast, after SF for 19–20 d, the number of courtship behaviors significantly increased (Kruskal–Wallis χ^2 [3, *n* = 40] = 14.624, *P* = 0.0021, 0 d after familiarization vs. 19–20 d after familiarization; Steel’s *P* = 0.048; Fig. 2C). The other *oxt* mutant males (*oxt*^{-/-}) also performed significantly more courtship behaviors toward WT females after SF for 19–20 d (Mann–Whitney *U* test: *Z* = -2.425, *P* = 0.015; SI Appendix, Fig. S4B). We then investigated behavioral changes in WT and *oxt*^{+I22F} males following SF for 19–20 d and revealed no effect in WT (Cab) or *oxt*^{+I22F} males (Mann–Whitney *U* test: *Z* = -0.427, *P* = 0.670; *Z* = 0, *P* > 0.999, respectively; Fig. 2D), consistent with previous observations that WT males exhibit courtship behaviors irrespective of SF. In addition, SF for 19–20 d facilitated the courtship activities of *oxt*^{I22F/I22F} males to a similar level as WT (~8 times within 5 min; Fig. 2C and D), implying that 19–20 d was sufficient for SF. Next, we examined the possible involvement of *oxtrs* in the emergence of mate preference. SF influenced mating preference for females in *oxtr1*^{-/-} males like *oxt* mutant males, whereas it did not influence mating preference in *oxtr2*^{-/-} males (Mann–Whitney *U* test: *Z* = -2.281, *P* = 0.023; *Z* = -0.491, *P* = 0.624, respectively; Fig. 2E).

In addition, we compared social motivation of nonfocal females toward WT and mutant males. We performed simple behavioral tests to evaluate approaching behavior of a single fish






Fig. 2. Courtship behavior of *oxt* and *oxtrs* mutant males toward unfamiliar or familiar females. (A) Males and females with the same genotype were group reared from birth, i.e., socially familiarized for a long period of time. Neither female genotype nor SF influenced the number of courtship behaviors in WT and *oxt*^{+122F} males (pink). In contrast, *oxt*^{122F/122F} males (dark pink) exhibited courtship behaviors toward WT females less frequently than toward *oxt*^{122F/122F} females. Mean \pm SEM, $n = 11$ per group, Kruskal–Wallis: Steel–Dwass’s post hoc $^{***}P < 0.01$, $^{*}P < 0.05$. (B) Procedure for assessing the effect of social familiarization on courtship behavior in *oxt*^{122F/122F} males. *oxt*^{122F/122F} males and WT females were introduced for the first time during the courtship behavior test (day 0) and then two *oxt*^{122F/122F} males and two WT females were kept in the same tank (familiarization) for 20 d. (C) The number of courtship behaviors of *oxt*^{122F/122F} males toward unfamiliar WT females significantly increased following 19–20 d of SF. Mean \pm SEM, $n = 10$ per group, Kruskal–Wallis: Steel’s post hoc: $^{*}P < 0.05$ vs. “0 day post familiarization”. (D) WT and *oxt*^{+122F} (pink) males exhibited courtship behavior toward familiarized WT females to the same extent as toward unfamiliar WT females. Mean \pm SEM, $n = 11$ per group, Mann–Whitney *U* test. (E) WT and *oxtr2*^{-/-} (brown) males exhibited courtship behavior toward familiarized WT females to the same extent as toward unfamiliar WT females. However, the number of courtship behaviors of *oxtr1*^{-/-} males (orange) toward unfamiliar WT females was significantly lower than that toward familiarized females. Mean \pm SEM, $n = 12$ per group, Mann–Whitney *U* test: $^{*}P < 0.05$; n.s., not significant. ♀, female; ♂, male.

toward the target fish (ref. 20 and *SI Appendix*, Fig. S6A). Indeed, WT females exhibited approaching behavior toward males, and this trend was not changed by the genotype of the male (Mann–Whitney *U* test: WT $Z = -3.926$, $P < 0.0001$; *oxt*^{122F/122F} $Z = -3.580$, $P = 0.0003$, *oxt*^{-/-} $Z = -3.291$, $P = 0.0009$; *oxtr1*^{-/-} $Z = -2.338$, $P = 0.019$, respectively; *SI Appendix*, Fig. S6B). This finding suggested that females did not react differently to mutant and WT males. Taken together, our data indicated that the activation of *oxtr1*, but not *oxtr2*, by *oxt* ligands blocks the emergence of mate preference toward socially familiarized females.

***oxt* and *oxtr1* Mutant Males Exhibit a Loss of Mate-Guarding Behaviors toward Unfamiliar Individuals.** To further investigate whether *oxt* and *oxtr1* mutant males lack sexual motivation toward

unfamiliar females, we used another behavioral test to assess male mate-guarding behaviors. In triadic relationships (two males and one female), the two males compete and the male nearest the female dominantly exhibits mate-guarding behavior (occupying a dominant position near the female while interfering with the other male’s approach toward the female) (12, 13). Our previous study in medaka demonstrated that male sexual motivation toward a target female is essential to drive mate-guarding in triadic relationships (12, 13). Thus, we could indirectly test male sexual motivation toward a target female using this behavioral paradigm. We placed two males with the same allele and one unfamiliar female in the tank and quantified the guarding index of the dominant male. The guarding index correlates with the frequency of the male fish locating in the area between the rival male and female. If the guarding index is higher than that of the merged control groups (negative control, *Materials and Methods*), it is considered that mate-guarding emerged in the triadic relationship. Indeed, the guarding indices of the WT (*oxt*^{+122F}), and *oxtr2*^{-/-} males were significantly higher than those of the merged control groups (Mann–Whitney *U* test: $Z = 2.165$, $P = 0.030$; $Z = 2.800$, $P = 0.0051$; $Z = 2.656$, $P = 0.0074$, respectively; Fig. 3A and B), indicating that these males exhibited mate-guarding behavior toward unfamiliar females. However, *oxt*^{122F/122F} and *oxtr1*^{-/-} males did not exhibit mate-guarding behavior toward unfamiliar females (Mann–Whitney *U* test: $Z = -0.953$, $P = 0.341$; $Z = 0.693$, $P = 0.489$, respectively; Fig. 3A and B). A 20-d period of SF facilitated the mate-guarding behavior in these mutants (Mann–Whitney *U* test: $Z = 2.858$, $P = 0.0043$; $Z = 2.540$, $P = 0.011$, respectively; Fig. 3A and B). The other *oxt* mutant (*oxt*^{-/-}) males exhibited the same behavioral phenotype (Mann–Whitney *U* test: before SF $Z = 0.265$, $P = 0.791$; after SF $Z = 3.023$, $P = 0.0025$, respectively; *SI Appendix*, Fig. S7A). Taken together, these findings suggested that *oxt* and *oxtr1* mutant males might lack sexual motivation toward unfamiliar females, which could lead to decreased motivation for mate guarding.

***oxt* and *oxtr1* Mutant Males Exhibit Dominant Mate-Guarding Behaviors toward Socially Familiarized Females.** Next, we examined whether familiarity with target females could affect the dominance of mutant males in the mate-guarding behaviors. To compare the dominance of *oxt* homozygous mutant males with that of heterozygous mutants, we performed the dominance test (*Materials and Methods*). We allowed two males with different alleles to compete for one female and examined the dominance of individual males in mate-guarding behavior. When we allowed one *oxt*^{+122F} male and one *oxt*^{122F/122F} male to compete for an unfamiliar female, the guarding index of the *oxt*^{+122F} males was significantly higher than that of the *oxt*^{122F/122F} males (Wilcoxon signed ranks: $Z = -2.863$, $P = 0.0042$; *SI Appendix*, Fig. S7B), indicating that *oxt*^{+122F} males were dominant toward unfamiliar females (*Movie S1*). This observation was consistent with previous findings that *oxt*^{122F/122F} males did not exhibit mate-guarding behavior toward unfamiliar females (Fig. 3A). Surprisingly, when we used long-time socially familiarized females instead of unfamiliar females, the tendency toward dominance was completely reversed (Wilcoxon signed ranks: $Z = -2.628$, $P = 0.0086$; Fig. 3C). The *oxt*^{122F/122F} males exhibited dominant mate-guarding behavior toward socially familiarized females (*Movie S2*). In addition, *oxt*^{122F/122F} males exhibited dominance when competing with WT males for socially familiarized females (Wilcoxon signed ranks: $Z = -2.981$, $P = 0.0029$; *SI Appendix*, Fig. S7C). We confirmed that other *oxt* mutant (*oxt*^{-/-}) males had the same behavioral phenotype (Wilcoxon signed ranks: before SF $Z = -2.089$, $P = 0.037$; *SI Appendix*, Fig. S7D, after SF $Z = -2.701$, $P = 0.0069$; *SI Appendix*, Fig. S7E). Therefore, *oxt* mutations enhanced mate-guarding behavior toward familiar females, but not toward unfamiliar females. We next performed the dominance test using *oxtr* mutants and showed that *oxtr1*^{-/-} males, but not *oxtr2*^{-/-} males,

exhibited dominant mate-guarding behavior toward socially familiarized females, similar to *oxtr1* mutants (Wilcoxon signed ranks: *oxtr1* $Z = -2.980$, $P = 0.0029$, *oxtr2* $Z = -0.178$, $P = 0.859$; Fig. 3D, *oxtr1* $Z = -2.275$, $P = 0.023$, *oxtr2* $Z = -0.275$, $P = 0.784$; SI Appendix, Fig. S7F). These findings indicated that *oxtr1* signaling mediated the suppression of dominant mate-guarding behavior toward socially familiarized females.

Signaling Pathways Influenced by *oxtr* System Defects. The sex differences in mating behavior induced by the *oxtr* system led us to investigate the molecular basis of these differences. In the medaka brain, *oxtr* is expressed mainly in the preoptic area, and there are no significant sex differences (21), but whether the *oxtr1* expression pattern in medaka differs between sexes was unknown. To identify *oxtr1*-expressing neurons, we performed in situ hybridization and showed that *oxtr1* was expressed in the telencephalon, preoptic area, habenula, optic tectum, hypothalamus, and corpus glomerulosum. We found no significant sex differences in the distribution pattern (SI Appendix, Fig. S8).

Next, to identify signaling pathways that could be affected by defects in the *oxtr-oxtr1* pathway and clarify the molecular basis underlying the sex differences, we performed RNA-sequencing (seq) (22). We used the whole adult brains, as *oxtrs* are widely distributed throughout the medaka brain, similar to what has been reported in the brains of other teleost fish and mammals (23). We compared gene expression profiles among six groups: 1) WT female, 2) WT male, 3) *oxtr*^{I22F/I22F} female, 4) *oxtr*^{I22F/I22F} male, 5) *oxtr1*^{-/-} female, and 6) *oxtr1*^{-/-} male. The *oxtr* or *oxtr1* mutations significantly changed the transcription levels of 1,389 and 1,181 genes, respectively, in females (SI Appendix, Fig. S9A and Dataset S1), and in 1,052 and 813 genes, respectively, in males (SI Appendix, Fig. S9B and Dataset S1). As

oxtr and *oxtr1* mutants exhibited similar behavioral phenotypes, we focused on genes whose transcription levels significantly increased or decreased with both *oxtr* and *oxtr1* mutations. There were no genes regulated in an opposite manner between females and males (SI Appendix, Fig. S10 and Dataset S2). At first, we searched for genes associated with the *oxtr* system in both sexes. Gene ontology (GO) analysis and pathway analysis revealed no enrichments of up-regulated genes (SI Appendix, Fig. S11A), whereas down-regulated gene clusters included the three genes for all of the components of C1q (Fig. 4A and B and SI Appendix, Fig. S11B). Functional C1q is a heterotrimer complex consisting of the products of three genes: *c1qa*, *c1qb*, and *c1qc*. Next, as mutations in the *oxtr-oxtr1* system have different effects on mate preference between the sexes, we reasoned that the *oxtr* system regulated signaling pathway in a sex-specific manner. GO analysis showed an enrichment of genes involved in cellular sodium ion homeostasis in a female-selective manner. The GO term (12 genes) included 4 genes encoding ion-transporting ATPase subunits (*atp1a3a*, *atp1a3b*, *atp1b1b*, and *atp6v1ab*), which were up-regulated (Fig. 4C and SI Appendix, Fig. S12A). Pathway analysis revealed an enrichment of a gene set involved in gamma-aminobutyric acid (GABA) metabolism, suggesting that the mutation prominently up-regulated the transcription of these genes (*slc32a1*, *slc6a1b*, *stxbp1a*, and *dnajc5aa*) in a female-selective manner (Fig. 4C and SI Appendix, Fig. S12A). There was no enrichment of female-specific down-regulated genes for any terms (SI Appendix, Fig. S12B).

We determined 20 male-specific up-regulated genes and 39 male-specific down-regulated genes (SI Appendix, Fig. S13 and Dataset S2). Pathway analysis revealed the enrichment of extracellular matrix-related genes up-regulated in a male-specific manner that are reported to regulate the outgrowth of neuronal dendrites (24). In contrast, these analyses showed no enrichment of male-specific down-regulated genes for any terms (SI Appendix, Fig. S13B).

Discussion

The findings of the present study demonstrated that the *oxtr* system facilitates mate choice in a sexually dimorphic manner in a nonmonogamous species (medaka fish) in which males and females have different mating strategies. Here, we showed that the *oxtr-oxtr1* pathway is required in females for mate preference for familiar mates, and in males for an unrestricted and indiscriminate mating strategy (Fig. 5). Our findings also revealed that mutant males have the ability to discriminate between SF and non-SF females. Previously, we demonstrated that medaka females primarily use visual cues from “faces (head part)” to discriminate visually familiarized males from unfamiliar males (11, 25). However, SF (group rearing) provides exposure to both visual and olfactory cues. Reproductive hormones, such as 17 α ,20 β -dihydroxy-4-pregnen-3-one and prostaglandin F2 α , which act as pheromones released by females, modulate male behaviors in some fishes (26, 27). These pheromones also might be candidate substances that affect male mating preference in medaka fish. Future studies should investigate if, like females, mutant males use visual cues and/or chemical substances to recognize SF females.

The Different *oxtr* Functions Between Sexes in Medaka Fish. Among monogamous rodents, the *oxtr* system facilitates sexual preference for a familiar mate, irrespective of sex (1–5). Generally, sexual dimorphism of the mating strategy in monogamous species is reduced, providing advantages for shared parenting, protection of resources, and social support (28). Our findings suggest that the *oxtr* system influences mate choice in both sexes regardless of the mating strategy, and that sexual dimorphism in the mating strategy might relate to functional differences in the *oxtr* system between sexes (Fig. 5). Here, transcriptome analysis showed that an *oxtr/oxtr1* mutation led to up-regulation of gene




Fig. 3. Effect of social familiarization on mate-guarding behavior. (A) Although *oxtr*^{I22F/I22F} males did not exhibit mate-guarding behavior toward unfamiliar WT females (dark pink), social familiarization (SF) enhanced the mate-guarding behavior of these mutant males toward familiarized mates (purple). (B) *oxtr2*^{-/-} males exhibited mate-guarding toward unfamiliar females (brown), whereas *oxtr1*^{-/-} males did not (orange). *oxtr1*^{-/-} males exhibited mate-guarding toward socially familiarized females (beige). (C) *oxtr*^{I22F/I22F} males (dark pink) tended to be dominant in the dominance test using familiar females. (D) *oxtr1*^{-/-} males (orange) tended to be dominant in the dominance test using familiar females. (A–D) Mean \pm SEM, $n = 12$ per group, Mann-Whitney U test: * $P < 0.05$; ** $P < 0.01$; n.s., not significant. ♀, female; ♂, male.

expression associated with GABA metabolism/ion-transport ATPase in a female-selective manner and down-regulation of genes encoding complement component C1q in both sexes. In human studies, abnormalities in glutamate/GABA signaling were recently hypothesized to underlie a neurodevelopmental syndrome, autism spectrum disorder (29). As *atp1a3* are prominently expressed in GABAergic interneurons (30) in the mammalian brain, the enhanced expression of the ATPases might be associated with up-regulation of GABA metabolism. In addition, rodent and human studies suggest the involvement of complement

component C1q in fundamental neurodevelopmental pathways (axon pruning) and the pathogenesis of autism spectrum disorders (31). Therefore, it might be that *oxt/oxtr1* mutation in medaka fish impairs the neural mechanism underlying such a neurodevelopmental disorder. One possibility is that the *oxt/oxtr1* mutation causes a developmental defect in TN-GnRH3 neurons, because TN-GnRH3 neurons are required for the suppression of female receptivity toward unfamiliar males. Two mutants (*cxc4* and *cxc7*), which have an abnormal TN-GnRH3 neuron morphology, exhibit the same behavioral phenotype as *oxt/oxtr1*




Fig. 4. Volcano plots of up-regulated/down-regulated genes in *oxt* and *oxtr1* mutants. Log2foldChange of gene transcription levels in female mutants (A) and in male mutants (B) compared with those in WT fish. (C) Magnified views of the graph outlined by the dotted line in A and B. Each dot represents one gene. Gray dots indicate genes whose transcription levels were not significantly changed in mutants compared with those in WT (FDR > 0.05). Coral and light blue dots represent significantly up-regulated/down-regulated genes in mutants, respectively. Dark blue and blue dots indicate genes up-regulated in female mutants and involved in the GABA system and sodium ion homeostasis, respectively. Green dots represent genes down-regulated in both sexes of mutants and involved in the complement activation system. Horizontal dotted line represents a padj of 0.05.

mutant females (10). Further studies might clarify the involvement of the oxt system in the development of TN-GnRH3 neurons.

oxt Function in Mate Preference in the Ray-Finned Fish. In the monogamous convict cichlid (*Amatitlania nigrofasciata*), administration of an arginine vasotocin (homolog of vasopressin)/oxt pathway antagonist into the brain impaired both affiliative behavior toward a potential mate and aggression toward neighbors (32). The oxt-selective function, however, has not been elucidated by pharmacological methods because of the cross-talk between the oxt and vasopressin systems (33). Thus, evaluation of the possible involvement of the oxt system on neurodevelopment cannot be performed by pharmacological studies (antagonist and agonist administration into the adult brain). Furthermore, most teleost fish have two paralogue genes for oxtrs, and it would be impossible to functionally dissect oxtrs without genetic techniques. The present genetic dissection using medaka fish showed their respective functions. In the African cichlid fish, the presence of rival males increases *oxtr1* (*istr2*) expression in the preoptic area of the male brain (34), suggesting the possible involvement of *oxtr1* (*istr2*) in social habituation. In contrast, among African cichlids, the expression pattern of two paralogue genes in the brain exhibits considerable species specificity, irrespective of sociality (19). Therefore, we cannot exclude the possibility that the function of oxtrs in medaka fish is species specific.

Our findings provide genetic evidence that oxt function is required for mate choice based on familiarity recognition in ray-finned fish. In medaka fish, *oxt/oxtr1* mutant females lost their mate preference for familiar males, which may be due to the loss of an aversion for unfamiliar males rather than an impaired affiliation toward familiar males. The interpretation that OXT mediates its effect via the modulation of a negative response (rejection) rather than a positive response differs from the typical interpretation that OXT systems promote social affiliation in mammals. Therefore, our findings could suggest a divergence in oxt system function across species. Various ray-finned fish such as the African cichlid (*Neolamprologus pulcher*) (35), the Trinidadian guppy (*Poecilia reticulata*) (1), bluegill sunfish (*Lepomis macrochirus*) (36), and threespine stickleback (*Gasterosteus aculeatus*) (37) are able to recognize familiar mates, an ability that may be important for selecting sexual and social partners. Further studies of oxt function in other ray-finned fish could shed light on the evolutionary origin of oxt central brain function

in mate choice, dating back 400 million years with the emergence of Actinopterygii on earth.

Materials and Methods

Ethics Statement. The experiments described herein were conducted using protocols approved by the Animal Care and Use Committee of the Hokkaido University (permit no. 17-0130). Surgeries were performed under anesthesia using MS-222, and all efforts were made to minimize suffering following the NIH *Guide for the Care and Use of Laboratory Animals* (38).

Fish and Breeding Conditions. Medaka fish (*Oryzias latipes*; drR strain, cab strain and mutants) were maintained in groups in plastic aquariums (13 cm × 19 cm × 12 cm [height]). All fish were hatched and bred in our laboratory. WT fish and each mutant fish type were bred in different tanks (only *oxt* heterozygous and *oxt* homozygous mutants were bred in the same tanks because they were born to the same parents). Sexually mature (4–5 mo of age) male (2.8–3.5 cm) and female (3.0–3.5 cm) medaka producing fertilized eggs every morning were used. The water temperature was ~28 °C, and light was provided by standard fluorescent lamps for 14 h per day (0800–2200).

Female Mating Receptivity Test. To quantify the motivation of a female to mate with a male of interest, a female mating receptivity assay was performed as previously described (10). Please see *SI Appendix, Methods*.

Courtship Behavior Assay. This procedure was performed as previously described (13). Males and females were separated in the evening (1800–1900) the day before the assay. The mating pair was then placed together in a single tank the next morning, and mating behavior was recorded for 5 min. We counted the number of courtship displays.

Social Familiarization. Four fish (two focal males and two WT females) were reared in a single tank for SF. The focal males and WT females were unfamiliar to each other before this procedure. For this behavioral test, we used the same fish before and after the SF procedure (20 d) as “before SF” fish and “after SF” fish, respectively.

Mate-Guarding Test. To determine whether males exhibited mate guarding, a mate-guarding test was performed as previously described (13). One female and two males (all male pairs were size-matched) were placed in the tank, and their behavior was recorded from the bottom of the tank in the morning (1000–1200). As a negative control group (merged group), we performed the same experiment using virtually merged trios, recording one female and two males one by one, each placed in a separate aquarium. For details, please see *SI Appendix, Methods*.

Dominance Test. We used one *genotype A* male and one *genotype B* male and allowed them to compete for the female to evaluate their dominance in mate-guarding behavior as previously described (13). All male pairs were size matched. The relative locations of the three fish were measured and guarding indices of *genotype A* and *genotype B* were calculated. Their guarding indices were compared with a higher guarding index indicating higher dominance in the mate-guarding behavior compared with the other male.

RNA-Seq and Data Analyses. Total RNA was extracted from the whole brains of two adult females or males for each sample (*n* = 3 per group). RNA-seq libraries were sequenced on HiSeq 4000 (Illumina). Significantly changed transcripts were defined as false discovery rate (FDR) < 0.05. For GO and pathway analysis, the transcript ID was converted into zebrafish homologs and then run on GO-MWU and PANTHER, respectively. See *SI Appendix, Methods* for details.

Statistical Analysis. In the female mating preference test, the male courtship test and the male mate-guarding test, each behavioral index was compared using Mann–Whitney *U* test implemented in Statcel (OMS Ltd.). In the dominance test, we compared the guarding index of two males using the Wilcoxon signed ranks test implemented in Statcel (OMS Ltd.). To compare the behavioral index of multiple groups, we used Kruskal–Wallis and Steel–Dwass’s post hoc test or Steel’s post hoc test implemented in the R system. All statistical tests are two-tailed, and *P* values < 0.05 were considered significant. See *SI Appendix, Methods* for details.




Fig. 5. Requirement of the oxt system for the mate preference for familiar mates. WT medaka females, but not males, have a sexual preference for familiar mates (red and blue arrows). This tendency was reversed, however, in *oxt* and *oxtr1* mutant fish. Females lost mate preference (dark red arrows) and the mutant males gained mate preference for familiar mates (dark blue arrow). ♀, female; ♂, male.

Data Availability. RNA-seq data used in this study have been deposited in the Gene Expression Omnibus (GEO) database, <https://www.ncbi.nlm.nih.gov/geo> (accession no. GSE139158).

ACKNOWLEDGMENTS. We thank Profs. T. Kubo, T. Kikusui, Y. Taniguchi, T. Miyatake, and M. Kawata for constructive discussion; the National BioResource Project Medaka for supplying the medaka strains (<https://shigen.nig.ac.jp/medaka>); Drs. T. Sakuma and T. Yamamoto for providing the pFUS_A2A and pFUS_A2B vectors; I. Hara for technical assistance in TILLING methods; T. Yamazaki for breeding the mutant fish; and M. Takei and M. Endo for

developing the optomotor response (OMR) apparatus. This work was supported by the National Institute for Basic Biology Priority Collaborative Research Project 10-104 and Cooperative Research Project 19-347; Joint Research Grant 01111904 by the National Institutes of Natural Sciences; Japan Society for the Promotion of Science (JSPS) KAKENHI Grants 16K18369 (to S.Y.), 19K16247 (to S.Y.), 16H06524 (to S.Y.), 18H02479 (to H.T.), 16H01276 (to H.T.); Grant-in-Aid for JSPS fellows (S.Y.); and Sumitomo Foundation (S.Y.). The contribution by L.J.Y. was supported by NIH Grant P50MH100023 (to L.J.Y.) and Yerkes National Primate Research Center (YNPRC) Base Grant P51OD11132 to YNPRC.

1. M. Numan, L. J. Young, Neural mechanisms of mother-infant bonding and pair bonding: Similarities, differences, and broader implications. *Horm. Behav.* **77**, 98–112 (2016).
2. J. K. Rilling, L. J. Young, The biology of mammalian parenting and its effect on offspring social development. *Science* **345**, 771–776 (2014).
3. H. Walum, L. J. Young, The neural mechanisms and circuitry of the pair bond. *Nat. Rev. Neurosci.* **19**, 643–654 (2018).
4. L. A. McGraw, L. J. Young, The prairie vole: An emerging model organism for understanding the social brain. *Trends Neurosci.* **33**, 103–109 (2010).
5. Z. V. Johnson *et al.*, Central oxytocin receptors mediate mating-induced partner preferences and enhance correlated activation across forebrain nuclei in male prairie voles. *Horm. Behav.* **79**, 8–17 (2016).
6. K. A. Hughes, L. Du, F. H. Rodd, D. N. Reznick, Familiarity leads to female mate preference for novel males in the guppy, *Poecilia reticulata*. *Anim. Behav.* **58**, 907–916 (1999).
7. C. M. Bull, Monogamy in lizards. *Behav. Processes* **51**, 7–20 (2000).
8. J. D. Klatt, J. L. Goodson, Oxytocin-like receptors mediate pair bonding in a socially monogamous songbird. *Proc. Biol. Sci.* **280**, 20122396 (2013).
9. S. A. Cheetham, M. D. Thom, R. J. Beynon, J. L. Hurst, “The effect of familiarity on mate choice” in *Chemical Signals in Vertebrates 11*, J. L. Hurst, R. J. Beynon, S. C. Roberts, T. D. Wyatt, Eds. (Springer New York, New York, NY, 2008), pp. 271–280.
10. T. Okuyama *et al.*, A neural mechanism underlying mating preferences for familiar individuals in medaka fish. *Science* **343**, 91–94 (2014).
11. M. Y. Wang, H. Takeuchi, Individual recognition and the ‘face inversion effect’ in medaka fish (*Oryzias latipes*). *eLife* **6**, e24728 (2017).
12. S. Yokoi *et al.*, Mate-guarding behavior enhances male reproductive success via familiarization with mating partners in medaka fish. *Front. Zool.* **13**, 21 (2016).
13. S. Yokoi *et al.*, An essential role of the arginine vasotocin system in mate-guarding behaviors in triadic relationships of medaka fish (*Oryzias latipes*). *PLoS Genet.* **11**, e1005009 (2015).
14. Y. Taniguchi *et al.*, Generation of medaka gene knockout models by target-selected mutagenesis. *Genome Biol.* **7**, R116 (2006).
15. T. Ishikawa *et al.*, High-resolution melting curve analysis for rapid detection of mutations in a Medaka TILLING library. *BMC Mol. Biol.* **11**, 70 (2010).
16. S. Ansai *et al.*, Efficient targeted mutagenesis in medaka using custom-designed transcription activator-like effector nucleases. *Genetics* **193**, 739–749 (2013).
17. S. Ansai, M. Kinoshita, Targeted mutagenesis using CRISPR/Cas system in medaka. *Biol. Open* **3**, 362–371 (2014).
18. R. Walter, C. W. Smith, J. Roy, Importance of the third amino acid residue of oxytocin for its action on isolated rat uterus: Study of relationship between hormone conformation and activity. *Proc. Natl. Acad. Sci. U.S.A.* **73**, 3054–3058 (1976).
19. C. M. O’Connor, S. E. Marsh-Rollo, S. C. Ghio, S. Balshine, N. Aubin-Horth, Is there convergence in the molecular pathways underlying the repeated evolution of sociality in African cichlids? *Horm. Behav.* **75**, 160–168 (2015).
20. Y. Isoe, Y. Konagaya, S. Yokoi, T. Kubo, H. Takeuchi, Ontogeny and sexual differences in swimming proximity to conspecifics in response to visual cues in medaka fish. *Zool. Sci.* **33**, 246–254 (2016).
21. Y. Kawabata, T. Hiraki, A. Takeuchi, K. Okubo, Sex differences in the expression of vasotocin/isotocin, gonadotropin-releasing hormone, and tyrosine and tryptophan hydroxylase family genes in the medaka brain. *Neuroscience* **218**, 65–77 (2012).
22. S. Yokoi, M. Mito, S. Iwasaki, S. Nakagawa, H. Takeuchi, The gene expression of the brain/pituitary from oxt/oxr1 homozygous mutants and WT medaka in both sexes. GEO. <https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE139158>. Deposited 21 October 2019.
23. L. S. Huffman *et al.*, Distribution of nonapeptide systems in the forebrain of an African cichlid fish, *Astatotilapia burtoni*. *J. Chem. Neuroanat.* **44**, 86–97 (2012).
24. U. Novak, A. H. Kaye, Extracellular matrix and the brain: Components and function. *J. Clin. Neurosci.* **7**, 280–290 (2000).
25. M. Y. Wang, H. Takeuchi Individual recognition and the ‘face inversion effect’ in medaka fish (*Oryzias latipes*). *eLife* **6**, e24728 (2017).
26. M. Kobayashi, P. W. Sorensen, N. E. Stacey, Hormonal and pheromonal control of spawning behavior in the gold fish. *Fish Physiol. Biochem.* **26**, 71–84 (2002).
27. Y. Yabuki *et al.*, Olfactory receptor for prostaglandin F2 α mediates male fish courtship behavior. *Nat. Neurosci.* **19**, 897–904 (2016).
28. R. L. Trivers, “Parental investment and sexual selection” in *Sexual Selection and the Descent of Man*, B. Campbell, Ed. (Heinemann, 1972), pp. 136–179.
29. E. Kang *et al.*, Interplay between a mental disorder risk gene and developmental polarity switch of GABA action leads to excitation-inhibition imbalance. *Cell Rep.* **28**, 1419–1428.e3 (2019).
30. A. R. Paciorkowski *et al.*, Novel mutations in ATP1A3 associated with catastrophic early life epilepsy, episodic prolonged apnea, and postnatal microcephaly. *Epilepsia* **56**, 422–430 (2015).
31. C. Fang, C1q as a regulator of brain development: Implications for autism spectrum disorders. *Brain Disord. Ther.* **4**, 1 (2015).
32. R. G. Oldfield, H. A. Hofmann, Neuropeptide regulation of social behavior in a monogamous cichlid fish. *Physiol. Behav.* **102**, 296–303 (2011).
33. Z. Song, H. E. Albers, Cross-talk among oxytocin and arginine-vasopressin receptors: Relevance for basic and clinical studies of the brain and periphery. *Front. Neuroendocrinol.* **51**, 14–24 (2018).
34. C. A. Weitekamp *et al.*, A role for oxytocin-like receptor in social habituation in a teleost. *Brain Behav. Evol.* **89**, 153–161 (2017).
35. S. Balshine-Earn, A. Lotem, Individual recognition in a cooperatively breeding cichlid: Evidence from video playback experiments. *Behav* **135**, 369–386 (1998).
36. J. A. Brown, P. W. Colgan, Individual and species recognition in centrarchid fishes: Evidence and hypotheses. *Behav. Ecol. Sociobiol.* **19**, 373–379 (1986).
37. G. J. FitzGerald, J. Morrisette, Kin recognition and choice of shoal mates by threespine sticklebacks. *Ethol. Ecol. Evol.* **4**, 273–283 (1992).
38. National Research Council, *Guide for the Care and Use of Laboratory Animals* (National Academies Press, Washington, DC, ed. 8, 2011).