



Published in final edited form as:

*Annu Rev Physiol.* 2020 February 10; 82: 21–43. doi:10.1146/annurev-physiol-021119-034453.

## Cardiac Pacemaker Activity and Aging

Colin H. Peters<sup>1</sup>, Emily J. Sharpe<sup>1</sup>, Catherine Proenza<sup>1,2</sup>

<sup>1</sup>Department of Physiology and Biophysics, University of Colorado Anschutz Medical Campus, Aurora, Colorado 80045, USA;

<sup>2</sup>Department of Cardiology, University of Colorado Anschutz Medical Campus, Aurora, Colorado 80045, USA

### Abstract

A progressive decline in maximum heart rate (mHR) is a fundamental aspect of aging in humans and other mammals. This decrease in mHR is independent of gender, fitness, and lifestyle, affecting in equal measure women and men, athletes and couch potatoes, spinach eaters and fast food enthusiasts. Importantly, the decline in mHR is the major determinant of the age-dependent decline in aerobic capacity that ultimately limits functional independence for many older individuals. The gradual reduction in mHR with age reflects a slowing of the intrinsic pacemaker activity of the sinoatrial node of the heart, which results from electrical remodeling of individual pacemaker cells along with structural remodeling and a blunted  $\beta$ -adrenergic response. In this review, we summarize current evidence about the tissue, cellular, and molecular mechanisms that underlie the reduction in pacemaker activity with age and highlight key areas for future work.

### Keywords

sinoatrial node; maximum heart rate; intrinsic heart rate; aging; cardiac pacemaking

## INTRODUCTION

Aging has long been associated with declines in physical fitness and aerobic capacity. In Homer's *Iliad*, Nestor laments that his "limbs are gnarled now, the old power's gone" (1, p. 318). This proclamation from an old warrior highlights that while some of the effects of aging can be counteracted by exercise training, they cannot be fully mitigated.

One of the unalterable consequences of aging is a progressive decline in maximum heart rate (mHR) (Figure 1). The age-dependent decline in mHR was first published in a cross-sectional study by Sid Robinson in 1938 (2), who followed up with a longitudinal study in 1975 (3). Importantly, the decline in mHR is an inherent property of cardiac aging. Unlike other measures of cardiac fitness, such as left ventricular mass, that can be increased by exercise training (4), mHR declines with age at approximately the same rate for all

---

colin.peters@cuanschutz.edu.

### DISCLOSURE STATEMENT

The authors are not aware of any affiliations, memberships, funding, or financial holdings that might be perceived as affecting the objectivity of this review.

individuals without regard for gender, diet, or physical fitness (5, 6). The rate of decline of mHR is commonly approximated by the formula  $mHR \approx 220 - \text{age}$  (7), or about 1 beat per minute (bpm)/year, although the original study by Robinson and more recent studies suggest that the actual rate is 0.6–0.8 bpm/year (2, 5, 8). A similar age-dependent decline in mHR has been observed in many other species, including mice, rats, rabbits, dogs, and cats (9–14).

The decline in mHR begets an inevitable decline in the maximum rate of oxygen consumption ( $V_{O2max}$ ) and, consequently, physical performance, with age. Indeed, decreased mHR is the primary determinant of the age-dependent decline in  $V_{O2max}$  and the main factor that limits exercise performance with age (15, 16). Moreover, of the factors that determine  $V_{O2max}$ , mHR alone cannot be improved by exercise training. While mHR and  $V_{O2max}$  are not important considerations for the majority of young and middle-aged individuals who do not require or utilize their full aerobic capacity, low  $V_{O2max}$  is a leading factor that limits functional independence for many otherwise-healthy older individuals (17–19). The age-dependent decline in mHR is thus of paramount concern as the population ages because it diminishes quality of life for individuals and increases health care costs for society.

It has long been appreciated that the age-dependent decline in mHR is caused by a nearly parallel age-dependent decline in intrinsic heart rate (iHR) (20–22). iHR is the heart rate measured under autonomic blockade. It thus serves as a systemic readout of the function of the primary cardiac pacemaker, the sinoatrial node (SAN). The age-dependent decline in iHR indicates, by definition, that pacemaker activity of the SAN is compromised with age. Despite the decline in iHR, resting heart rate (rHR) is fairly constant with age because it is maintained by autonomic nervous system activity. An important corollary is that sympathetic tone must increase with age in order to maintain rHR (Figure 1). In this review, we focus on the mechanisms that cause the age-dependent loss of cardiac pacemaker function.

## SINOATRIAL NODE OVERVIEW

The SAN is the pacemaker of the heart. Its eventual discovery in 1907 built on observations of numerous early scientists over the course of several centuries (for review see 23). In the eighteenth century Albrecht von Haller provided a convincing argument that the heartbeat originated within the heart itself, without requiring input from the nervous system that others had proposed (24). In the late nineteenth century, Walter Gaskell noted that the electrical signal began in the sinus auricle and was then conducted through the atrium and on to the ventricles (24). Subsequently, in 1907 Keith & Flack (25) published an extensive study detailing a special tissue within the sinus auricle—the SAN—as the site from which the heart rhythm arises.

In denervated human hearts, the SAN generates a spontaneous rhythmic electrical depolarization at a rate of approximately 100 bpm (26, 27). This depolarization propagates through the cardiac conduction system and throughout the heart to initiate each heartbeat (Figure 2a). The automaticity of the SAN is the basis of the iHR that can be measured in vivo in humans and animals under autonomic blockade (20).

The SAN is highly innervated by the autonomic nervous system, which profoundly regulates heart rate on a beat-to-beat basis to closely match the cardiac output to ever-changing metabolic demand. Efferents from the vagus nerve provide parasympathetic innervation, first synapsing in fat pad ganglia on the heart, and then projecting to the SAN where they decrease heart rate by slowing the intrinsic firing rate of the SAN. Conversely, thoracic sympathetic nerves from T1–T4 first synapse in sympathetic chain ganglia adjacent to the spinal cord and then project to the SAN where they elicit the fight-or-flight increase in heart rate. At rest, humans below the age of about 85 have a predominant parasympathetic tone that inhibits iHR to produce an average rHR of ~60–100 bpm (Figure 1) (28). During exercise or stress, sympathetic nervous system activity increases SAN firing, which in turn increases heart rate.

Within the SAN, sinoatrial node myocytes (SAMs) serve as the pacemaker cells. As noted by Keith & Flack (25), SAMs are surrounded by fibrous tissue and are in close proximity to processes from the vagus and sympathetic nerves (Figure 2c) (25). SAMs are highly specialized cells that differ considerably from the working myocytes of the atrial and ventricular myocardium in terms of function, morphology and gene expression. Compared to atrial and ventricular myocytes, SAMs are smaller in size and appear unstriated (Figure 2b), owing to fewer and poorly organized sarcomeres (29–31). SAMs are also characterized by numerous caveolae that increase the membrane surface area and likely serve to organize signaling microdomains (31). Isolated SAMs are also quite heterogeneous, which is thought to reflect a functional gradient of cellular subtypes from the center to periphery of the SAN, at least in some species (32, 33).

In the intact SAN, SAMs are electrically coupled by low-conductance (<40 pS) gap junctions formed by connexins Cx45, Cx40, and Cx30.2 (34–36). In the rest of the heart, high-conductance (>70 pS) gap junctions are formed primarily by Cx43 or Cx40 (37). The presence of lower-conductance connexins in the SAN suggests greater electrical resistance between cells (38). There is also evidence to suggest that gap junctions in the SAN are particularly sparse, which further increases electrical resistance (29, 39). This electrical isolation is thought to be necessary to protect the SAN from the hyperpolarizing influence of the much larger atrial myocardium. Computational models predict that higher electrical resistance in the central SAN coupled with a gradual decrease in resistance toward the atrial myocardium allows for the relatively small SAN to pace the atria (40). Indeed, both low-conduction velocity in the SAN and an increase in velocity at the boundary with the atria have been measured experimentally (41). A zone of transition in connexin expression between the SAN and atria has also been observed (33, 35). It is thought that these transitional areas form complex block zones and exit pathways that optimize excitation of the atria while minimizing reentry (33, 42, 43). Additionally, expression of low-conductance connexins allows for the possible formation of asymmetric gap junctions with highly rectified conductances that would allow for nearly unidirectional flow between cells (44, 45). SAMs are further isolated from the surrounding atrial myocardium by the nonconducting fibrous tissue that ensheathes them (25, 43).

Age-dependent slowing of SAN conduction is well characterized, but as is the case for many aspects of age-dependent changes in the SAN, data on changes to connexin protein

expression with aging are limited (43, 46, 47). One study in guinea pigs shows a progressive loss in Cx43 protein with no change in Cx40 or Cx45 (46). The loss of Cx43 was accompanied by slowed conduction velocity through the SAN, although the underlying mechanism is not clear, given the low expression of Cx43 in SAN. In rats, the abundance of Cx30.2 transcripts in the SAN decreases with age, but whether this effect is translated to the protein level is unknown (48).

### Sinoatrial Action Potentials and Aging

SAMs function as cardiac pacemaker cells by firing spontaneous action potentials (APs) that underlie the autorhythmicity of the heart. Spontaneous APs in SAMs have a characteristic waveform (Figure 3*a*). Compared to APs in ventricular myocytes, the upstroke of sinoatrial APs is considerably slower, and the amplitude is smaller. Sinoatrial APs also lack the transient repolarization phase, and the plateau phase is considerably less pronounced. Most importantly, sinoatrial APs do not have a stable resting membrane potential. Instead, a spontaneous depolarization during diastole brings the membrane potential to threshold to trigger the subsequent AP.

Age-dependent slowing of mHR is caused, at least in part, by slower firing of individual SAMs. Using transmembrane recordings of aged rabbit and cat pacemaker tissue, Alings & Bouman (10) showed that spontaneous AP firing rate decreases with age due to both increased AP duration and slowing of the diastolic depolarization rate (DDR). This was consistent with earlier work showing an age-dependent slowing of the firing rate and prolongation of AP duration in rat atrial cells within tissue preparations (9, 49). Our group and others have observed a decrease in the basal and maximal AP firing rates of isolated SAMs and SAN tissue from aged mice that mirrors the slowing of iHR and mHR (11, 13) (Figure 4*a*). In isolated SAMs, we found that the slower intrinsic firing rate was caused by a more negative maximum diastolic potential (MDP) and a slower DDR, while the AP duration trended toward prolongation with age (13), consistent with the previous studies (10, 11) (Figure 4*b*). Indeed, reanalysis of our data set from aged versus young SAMs (13) using refined parameter definitions and automated AP waveform analysis (50) revealed a significant AP prolongation in the aged cells. An exception to this trend is a study in mice that showed a slower DDR in aging, but in contrast to other studies, a shortened AP duration (47).

### IONIC REMODELING OF THE SINOATRIAL NODE WITH AGE

It is beyond the scope of this review to exhaustively describe the long and ever-growing list of ionic currents that are important for pacemaking in SAMs. For more details, the reader is referred to many excellent review articles (51–53). Here, we provide a brief overview of the major ionic currents in SAMs, followed by a more-detailed discussion of the currents that have so far been experimentally assayed in young versus aged SAMs. The sinoatrial AP waveform parameters that change with age (DDR, MDP, and AP duration) (10, 11, 13) constrain the possible underlying ionic mechanisms, indicating that the balance of currents active during these phases is altered by the natural aging process. It should be noted that the focus on ionic currents as the end effectors of changes in membrane potential does not

impugn the clear roles of intracellular  $\text{Ca}^{2+}$  dynamics and second messengers in regulating these transmembrane conductances.

### Overview of Age-Dependent Changes in Ionic Currents in Sinoatrial Node Myocytes

A unique complement of ionic currents is critical for the production of spontaneous APs in SAMs. Studies through many decades have focused primarily on the identity of the currents that cause the diastolic depolarization; comparatively less is known about currents active during other phases of the AP. The precise identities and relative amplitudes of currents active in any given cell depend on the species, the location of the cell within the SAN, and the physiological context (including short-term status such as sympathetic stimulation and longer-term processes such as aging or disease). Many inward currents contribute to the diastolic depolarization, including but not limited to the funny current ( $I_f$ ), the  $\text{Na}^+$ - $\text{Ca}^{2+}$  exchange current ( $I_{\text{NCX}}$ ), L- and T-type  $\text{Ca}^{2+}$  currents ( $I_{\text{Ca,L}}$  and  $I_{\text{Ca,T}}$ ), and possibly voltage-gated  $\text{Na}^+$  currents ( $I_{\text{Na}}$ ). Following the diastolic depolarization,  $I_{\text{Ca,L}}$  and  $I_{\text{Ca,T}}$  are thought to be primarily responsible for the upstroke of the AP. Major outward currents in SAMs include voltage-gated  $\text{K}^+$  currents ( $I_{\text{Kr}}$ ,  $I_{\text{Ks}}$ , and  $I_{\text{to}}$ ),  $\text{Ca}^{2+}$ -activated  $\text{K}^+$  currents ( $I_{\text{KCa}}$ ), inward rectifiers ( $I_{\text{KACh}}$ ,  $I_{\text{KATP}}$ , and varying amounts of  $I_{\text{K1}}$ ), and possibly  $I_f$  (see below).

**Funny current.**—The  $I_f$  was first discovered 40 years ago as an adrenaline-sensitive current activated by hyperpolarization in rabbit SAN tissue (54).  $I_f$  is produced by hyperpolarization-activated, cyclic-nucleotide sensitive (HCN) channels. There are four HCN channel isoforms (HCN1–4), of which HCN4 is the predominant isoform in the SAN of all mammals (55, 56). HCN1 and HCN2 are expressed at lower levels in the SAN in a species-dependent manner (57–61).

As the name implies, HCN channels are activated by membrane hyperpolarization. Consistent with the adrenaline sensitivity of  $I_f$ , the open probability of HCN4 channels is modulated by binding of cyclic nucleotides to a conserved C-terminal cyclic nucleotide-binding domain (62). Sympathetic stimulation increases cAMP in SAMs, and binding of cAMP to HCN4 channels shifts pore opening to more depolarized membrane potentials and slows channel closing. The resulting increase in inward current contributes to the sympathetic nervous system-induced increase in AP firing rate of SAMs and, consequently, heart rate (63).

Although  $I_f$  is best known for conducting inward current during diastole, less appreciated is the potential role of  $I_f$  during repolarization. HCN channels are permeable to both  $\text{Na}^+$  and  $\text{K}^+$ , with a net reversal potential of approximately  $-30$  mV. Computational models and our preliminary data suggest that the channels also pass an outward, repolarizing current during systole that may shape the AP waveform and modulate firing rate (64, 65).

Strong evidence supports a role for  $I_f$  in age-dependent declines in SAM firing rate. In isolated SAMs from aged mice, the voltage dependence of  $I_f$  is shifted to more hyperpolarized potentials, thereby reducing current available during the AP (13, 66) (Figure 4*d*). This hyperpolarized voltage dependence persists in excised inside-out membrane patches, thereby precluding a simple decrease in intracellular cAMP as the causative mechanism (66). Rather, the data suggest that the HCN4 channel itself or an unknown,

channel-associated modulator must be altered in aging (66, 67). In addition to the shifted voltage dependence,  $I_f$  density is decreased in aged mouse SAMs, although this decrease is fully explained by an increase in membrane capacitance (13, 68). While a decrease in  $I_f$  is consistent with experiments showing reduced effects of HCN blockers in aged SAN (58, 69), there is no consensus about whether aging changes HCN channel expression: In rodents, HCN2 and HCN4 transcripts have been reported to either decrease (20, 21) or not change with age (47, 48), whereas HCN2 and HCN4 transcripts and protein were found to be upregulated in right atrial appendage from human patients (70). Regardless of mechanism, the functional reduction in  $I_f$  is predicted to slow pacemaking, in a manner similar to that of  $I_f$  blockers and *HCN4* mutations (71–73).

**Intracellular  $Ca^{2+}$  release and  $I_{NCX}$ .**—A role for sarcoplasmic reticulum (SR)  $Ca^{2+}$  release in heart rate regulation dates back to the early 1900s. In 1912, Pilcher (74) found that applying small amounts of caffeine—unknown to him as an activator of ryanodine receptors—to dog hearts increased the heart rate. In the 1970s, oscillations in  $Ca^{2+}$ , cAMP, and conductance were proposed to contribute to spontaneous activity in neurons and cardiac pacemaker cells (75, 76). Nearly 80 years after Pilcher (74), Rubenstein & Lipsius (77) showed in feline secondary pacemaker tissue the presence of a ryanodine-sensitive current during the late diastolic depolarization. A large body of work from many groups has established that this current is mediated by the plasma membrane  $Na^+$ - $Ca^{2+}$  exchanger (NCX).  $I_{NCX}$  in SAMs reflects the concerted activity of  $Ca^{2+}$  channels and transporters on the plasma and SR membranes—the same molecular players that mediate excitation-contraction coupling in other cardiac myocytes.

During the latter part of the diastolic depolarization, type 2 ryanodine receptors (RYR2) on the SR membrane release  $Ca^{2+}$  into the cytosol (78). This release is augmented by  $Ca^{2+}$  entry through voltage-gated  $Ca^{2+}$  channels on the plasma membrane that open in response to the diastolic depolarization (79). The ryanodine-sensitive current arises due to the subsequent extrusion of  $Ca^{2+}$  from the cell by NCX1, which exchanges three  $Na^+$  ions for each  $Ca^{2+}$  ion and thus generates a net inward current that contributes to the diastolic depolarization (80).

Block of RYR2 decreases the AP firing rate in SAMs, presumably through a decrease in  $I_{NCX}$  (78, 80, 81). Firing rate is also reduced by block of SERCA2A, the SR  $Ca^{2+}$ -ATPase responsible for pumping cytosolic  $Ca^{2+}$  back into the SR (82, 83). Block of SERCA2A prevents the replenishment of SR  $Ca^{2+}$  stores, which in turn decreases the  $Ca^{2+}$  release through RYR2 and presumably limits  $I_{NCX}$  (83).

Age-dependent changes have been observed in a number of these  $Ca^{2+}$ -handling proteins. At both transcript and protein levels, RYR2 is decreased in rat SAN with aging, whereas SERCA2A, RYR2, and NCX1 protein are all decreased in aged mouse SAN (11, 48) (Figure 4*f*). In aged mouse SAMs, Liu et al. (11) suggested that SR  $Ca^{2+}$  load is reduced; the amplitude and rate of rise of the  $Ca^{2+}$  transient were decreased and the duration of the  $Ca^{2+}$  transient was increased. Although the direct impact of aging on  $I_{NCX}$  in SAMs has yet to be measured experimentally, decreased SR  $Ca^{2+}$  release would be expected to contribute to slower AP firing rate by reducing  $I_{NCX}$ .

**Voltage-gated Ca<sup>2+</sup> currents.**—In SAMs, I<sub>Ca,L</sub> and I<sub>Ca,T</sub> contribute to the diastolic depolarization and are largely responsible for the AP upstroke. Larson et al. (13) found that the I<sub>Ca,L</sub> and I<sub>Ca,T</sub> densities are lower in aged SAMs (13) (Figure 4e). As is the case for I<sub>f</sub>, this decrease was fully accounted for by cellular hypertrophy without a concurrent upregulation of channel expression. In accordance, age-dependent decreases in I<sub>Ca,L</sub> and Ca<sup>2+</sup> transients have been observed in isolated human atrial tissue, canine atrial tissue, and rat ventricular tissue (11, 84–86). However, there is some variability in reported effects of aging on voltage-gated Ca<sup>2+</sup> channel expression. While Jones et al. (86) demonstrated an age-dependent decrease in Ca<sub>v</sub>1.2 protein expression in guinea pig SAN, experiments in rats showed an increase in Ca<sub>v</sub>1.2 mRNA expression (48, 87). Despite the differences in expression results, the functional decreases observed in I<sub>Ca,L</sub> and I<sub>Ca,T</sub> densities would be predicted to contribute to slower AP firing rates in aged SAMs, both directly by reducing inward Ca<sup>2+</sup> current and indirectly by reducing Ca<sup>2+</sup>-induced Ca<sup>2+</sup> release and I<sub>NCX</sub>, as discussed above.

**Na<sup>+</sup> currents.**—Unfortunately, few data are available on age-dependent changes to Na<sup>+</sup> channels in the SAN and their functional consequences. In fact, the role of voltage-gated Na<sup>+</sup> channels is often overlooked in the generation of spontaneous APs in pacemaker cells: The Na<sup>+</sup> channel blocker tetrodotoxin (TTX) slows the AP upstroke of subsidiary pacemaker cells from guinea pig (88) and cultured SAMs from rabbit (89) and slows the spontaneous beating rate of isolated mouse hearts (90). The cardiac Na<sub>v</sub>1.5 channel isoform is the site of multiple mutants that have been linked to inherited sinoatrial node dysfunction (SND) (91). And in mouse models, mutant Na<sub>v</sub>1.5 channels decrease iHR and cause SND (91).

Experiments in rats suggest an increase in Na<sub>v</sub>1.5 transcript expression with age (48), whereas in canine SAMs, I<sub>Na</sub> density decreases between newborns and young dogs and again between young and adult dogs (92). However, as with I<sub>Ca,L</sub>, I<sub>Ca,T</sub>, and I<sub>f</sub>, the decrease in I<sub>Na</sub> density between young and adult canine SAMs can be accounted for by the increase in membrane capacitance (92). Yanni et al. (68) showed that in aged mice there is an increase in the area of the SAN that lacks the relatively TTX-resistant Na<sub>v</sub>1.5 sodium channel isoform and yet an increased effect of 2 μM TTX on firing rate, possibly reflecting an upregulation of TTX-sensitive Na<sup>+</sup> channel isoforms.

**K<sup>+</sup> currents.**—There are clear roles for voltage-gated, Ca<sup>2+</sup>-activated, and inward rectifier K<sup>+</sup> channels in the repolarization and diastolic depolarization phases of the SAN AP (93–98). The delayed rectifier K<sup>+</sup> channels K<sub>v</sub>7.1 (*KCNQ1*), which underlies I<sub>Ks</sub>, and K<sub>v</sub>11.1 (*ERG*), which produces I<sub>Kr</sub>, are thought to be primarily responsible for repolarization of the sinoatrial AP. While mutations in these channels are most commonly associated with long QT syndrome, the same mutations are also associated with episodic bradycardia (98–100). Furthermore, pharmacological blockade of *ERG* causes bradycardia in both dogs and mice (101, 102).

Recently, a mutant in the G protein-gated inwardly rectifying K<sup>+</sup> (GIRK) channel Kir3.4, which along with Kir3.1 passes the I<sub>KACH</sub> current, was found to cause SND (103). Activation of I<sub>KACH</sub> contributes to slowed sinoatrial pacemaking in response to parasympathetic stimulation by slowing the DDR. Even in mice, which typically have

predominant sympathetic drive at rest, knockout of Kir3.1 causes resting tachycardia (93), while knockout of either Kir3.1 or Kir3.4 attenuates the decreases in heart rate seen with application of parasympathetic agonists (93, 104). Similarly, Ca<sup>2+</sup>-activated K<sup>+</sup> channels also play a role in determining the DDR and firing rate of SAMs. Both big conductance and intermediate conductance Ca<sup>2+</sup>-activated K<sup>+</sup> channels (BK and SK4, respectively) have been found to regulate pacemaking (94–96).

Despite their importance, there is a distinct lack of functional data on K<sup>+</sup> channels in aging SAMs. This lack of data is particularly significant given that the increase in AP duration and slowed DDR with age predict that there are likely to be changes in outward currents (10, 11). At the transcript level in aged rat SAN, some changes in delayed rectifier K<sup>+</sup> channels have been reported: K<sub>v</sub>7.1 is increased and K<sub>v</sub>1.5, the channel that underlies I<sub>Kur</sub>, is decreased (48). There is also an age-dependent decrease in Kir3.4 transcript and an increase in transcripts encoding K<sub>v</sub>1.2 and K<sub>v</sub>4.2, which produce I<sub>to</sub>, (48). Given that I<sub>to</sub> is increased in right atrial tissue in aged dogs, this may translate to increased sinoatrial I<sub>to</sub> with age; however, this has yet to be confirmed experimentally (83). Unfortunately, there is no protein or current expression data yet available from aged SAMs to elucidate the functional significance of any of these transcriptional changes.

### Conclusions About Electrical Remodeling of Sinoatrial Node Myocytes in Aging

Although it is clear that aging changes the expression and function of several ion channels in SAMs, a comprehensive understanding of how aging changes the ionic basis of pacemaking is not possible due to a paucity of functional data. In many cases the only existing studies are those of mRNA expression, which is often poorly correlated with protein expression (105). Many additional functional studies of age-dependent changes in ionic currents, including K<sup>+</sup> currents, are necessary to fully understand electrical remodeling in aged SAMs.

## STRUCTURAL REMODELING

As early as 1907, the SAN was noted to have a relatively high amount of fibrotic and connective tissue relative to the rest of the heart, and a further increase in fibrosis was observed in the SAN from diseased hearts (25). In an extensive review, Csepe et al. (43) note that the fibrotic tissue in the SAN may serve as an electrical insulator that, in addition to the SAN-specific gap junctions mentioned above, may protect the SAN from the larger and more hyperpolarized atrial myocardium.

A natural age-dependent increase in fibrotic tissue in the SAN has been consistently noted in humans and animal models and is strongly associated with reduced pacemaker function (106–108) (Figure 4c). Increased interstitial fibrosis and altered expression of matrix metalloproteinases correlate with SAN dysfunction and frailty in aged mice (47). In human patients, age-dependent atrial remodeling and slowed conduction velocity are associated with impaired SAN function (109) and are exacerbated in individuals with clinical SND (110), a disorder that is most prevalent in older individuals (111). Hao et al. (112) found that increased collagen, fibrosis, and TGFβ1 in the SAN were particularly pronounced in aged heterozygous SCN5A knockout mice, in association with decreased SAN function. And caldesmon 2-null mice have increased SAN fibrosis and sinus bradycardia (113).



However, in these latter cases, the genetic knockouts might also decrease SAN function by changing ionic currents; blockade of sodium channels can decrease SAN function (87–89) and calsequestrin 2 knockout also alters SR  $\text{Ca}^{2+}$  transients, diastolic  $\text{Ca}^{2+}$  levels, and spontaneous  $\text{Ca}^{2+}$  release patterns (113).

Despite the strong correlations, increased fibrosis in the aged SAN is not always associated with decreased pacemaker activity; severe fibrosis can occur in aged hearts with normal sinus rhythm (107). Thus, while fibrosis is associated with pacemaker anomalies, the extent to which it slows iHR in aging is unknown, in part due to the difficulty of experimentally manipulating fibrosis to reveal causal relationships.

Histological studies also show that the volume of the SAN and the number of SAMs decrease with age (106–108). Multiple labs have noted cellular hypertrophy and increased membrane capacitance of SAMs with aging (13, 68, 114). Along with the reduced volume of the SAN, this is consistent with a decrease in the number of cells. The relatively small number of SAMs makes the SAN vulnerable to loss of cells; a critical mass of SAMs is required to excite the atria. Indeed, animal models with apoptosis of SAMs have slower iHRs (115), and two-dimensional models of the SAN suggest that slower atrial rates are a direct consequence of a reduced number of SAMs (115, 116). However, it is not clear whether cellular hypertrophy of SAMs may compensate to some degree for the loss of cells. Moreover, normal sinus rhythm is maintained in some patients with only 10% of their SAMs remaining (107). Thus, as for the increase in fibrosis, a decrease in number of SAMs can be correlated with a decrease in heart rate, but the extent to which it contributes to the reduced SAN function in aging has not yet been resolved.

## **$\beta$ -ADRENERGIC RESPONSIVITY**

In addition to electrical and structural remodeling, aging also decreases mHR in part by blunting the heart rate response to sympathetic stimulation. The observation of a decreased inotropic response to catecholamines in heart muscle from aged rats (117) led to the hypothesis that aging reduces mHR by similarly decreasing the response of the SAN to  $\beta$ -adrenergic receptor ( $\beta$ AR) stimulation (14). Indeed, isoproterenol injection or maximal exercise produces smaller increases in heart rate above rHR in older human subjects (4, 118), and the  $\beta$ AR blocker, propranolol, causes a correspondingly smaller decrease in mHR in older individuals (119). However, as discussed above, the constant rHR with age in the context of the reduction in iHR necessitates an increase in sympathetic tone at rest that naturally decreases the capacity of the heart rate to increase above resting levels (Figure 1) (20, 120–122).

In spite of the confounds of the sympathetic response from a base of iHR versus rHR, a reduction in the absolute response to  $\beta$ AR stimulation with age is revealed by other measurements. For example, the dose of isoproterenol needed to increase heart rate by 25 bpm increases with age (123). Decreased  $\beta$ AR sensitivity is also suggested by differences in the rates of decline in iHR and mHR with age (Figure 1); while early work, citing parallel declines in iHR and mHR, concluded that there was no evidence for an age-related decrease in the mechanisms which increased heart rate above its intrinsic level (20, 21), more recent

studies have revealed slight differences in the rates of decline in iHR and mHR. In humans, iHR decreases by ~0.5 bpm/year (20, 27, 124), whereas mHR decreases by ~0.7 bpm/year (2, 5). Similarly, in mice the declines in iHR and mHR differ between 3 and 32 months (~120 vs 170 bpm, respectively) (13). Overall, these data suggest that while decreases in iHR remain the primary driver of age-dependent declines in mHR, a modest role for declines in sympathetic nervous function also exists. Accordingly, Christou & Seals (22) found that the reduction in the heart rate response to isoproterenol accounts for ~17% of the decrease in mHR with age.

### Molecular Basis of the Reduced $\beta$ -Adrenergic Receptor Response with Age

Stimulation of  $\beta$ ARs on SAMs activates adenylyl cyclases to increase intracellular cAMP, which activates the cAMP-dependent protein kinase (PKA). PKA phosphorylates a host of molecules in SAMs that contribute to the fight-or-flight increase in AP firing rate. As discussed above,  $\beta$ AR-stimulated cAMP also speeds SAM firing by binding directly to HCN channels to increase  $I_f$  (125).

Molecules involved in intracellular  $Ca^{2+}$  cycling are among the best-known targets of PKA. PKA-mediated phosphorylation of L-type  $Ca^{2+}$  channels increases  $I_{Ca,L}$  (126), which in turn increases  $Ca^{2+}$  release from the SR through RYR2 (127, 128). SR release is further augmented by PKA phosphorylation of RYR2 itself, which increases its open probability (129, 130). The resulting increase in cytoplasmic  $Ca^{2+}$  in turn increases extrusion by NCX (131, 132). The increase in both  $I_{Ca,L}$  and  $I_{NCX}$  is thought to contribute to the fight-or-flight increase in AP firing rate. HCN4 channels are also a site for PKA phosphorylation, which may contribute to the shift in voltage dependence of  $I_f$  upon  $\beta$ AR stimulation and thereby to the increase in SAM firing rate (133). Finally, PKA phosphorylation of the SERCA inhibitory protein, phospholamban, relieves its inhibition of SERCA to increase the rate of  $Ca^{2+}$  reuptake into the SR (82, 134). Continued beating of the heart requires that the increased  $Ca^{2+}$  influx through L-type channels must be balanced by increased extrusion by NCX, while increased SR  $Ca^{2+}$  release through RYR2 must in turn be balanced by increased  $Ca^{2+}$  reuptake through SERCA.

As mentioned above, aging changes a number of these PKA-sensitive molecules: RYR2 transcript and protein levels are decreased in aged rat SAN, whereas SERCA2A, RYR2, and NCX1 proteins are all decreased in aged mouse SAN (11, 48). Although aging decreases adenylyl cyclase activity in the ventricular myocardium (135), its activity has not been measured in aged SAN. Several groups have reported a lack of change of  $\beta$ AR transcripts with age in rodents (48, 136). To our knowledge, no studies have examined age-dependent protein expression of either  $\beta$ ARs or adenylyl cyclases in isolated SAN tissue.

Studies from our lab have characterized cellular responses to stimulation of components of the  $\beta$ AR signaling cascade. In SAMs, stimulation of  $\beta$ ARs with isoproterenol, stimulation of adenylyl cyclase with forskolin, and/or inhibition of phosphodiesterases with IBMX are all insufficient to restore firing rate in aged cells; firing rates increase by similar amounts in young and old SAMs in response to these manipulations but fail to overcome the slower intrinsic firing rate of aged SAMs (13, 66). At face value, these data could indicate a fundamental limit to the firing rate of aged SAMs. However, application of a high

concentration of cAMP to the cytosol of SAMs restores AP firing rate in aged SAMs to youthful levels (66). These data establish that aged SAMs are capable of firing at the same rates as younger SAMs and suggest that the endogenous  $\beta$ AR machinery is insufficient to surmount the reduced intrinsic firing rate to maximally stimulate the cells.

Interestingly, a high concentration of IBMX was observed to restore maximum beating rate to young levels in SAN tissue strips from aged mice (11). This difference from the response of single cells could reflect an emergent tissue-level property, perhaps akin to the mechanism responsible for higher beating rates of SAN tissue compared to isolated cells from rabbits and mice (13, 137). Alternatively, it could result from IBMX activation of other cell types in the tissue or off-target effects of IBMX. Indeed, when data from aged SAMs were incorporated into an existing SAM AP model, the simulated APs reproduced the inability of  $\beta$ AR stimulation to rescue firing rate in aged cells; further tailoring of parameters was required to replicate the tissue-level effects (138). Ultimately, a great deal of additional experimental data will be required to formulate strong mathematical models of the effects of aging on sinoatrial cells and tissue.

## CELLULAR-LEVEL CHANGES ARE THE PRIMARY DETERMINANT OF AGE-DEPENDENT DECLINES IN MAXIMUM HEART RATE

Surprisingly, age-dependent effects at the tissue and cellular levels are not additive. In both rabbits and mice, the firing rate of intact SAN tissue is higher than the firing rate of individual SAMs, indicating that there is an emergent, tissue-level property that effectively increases heart rate, with the simplest mechanism being that the fastest cell paces all others (13, 137). Age-dependent deficits in this tissue-level compensation, due to fibrosis and/or decreases in cell number, might be expected to compound with those of individual SAMs to yield a slower beating rate than that caused by either factor alone. However, this has not been observed. In fact, the changes in firing rate of isolated SAMs prior to and after isoproterenol perfusion directly mirror the change in iHR and mHR in these animals. In mice, the intrinsic AP firing rate slows by  $\sim 100$  AP/min between 2 and 24 months, while iHR and the beating rate of whole nodes slow by  $\sim 60$ – $90$  bpm (11, 13, 66). The same pattern is present for the maximum AP firing rate and mHR; aging slows maximum firing rate by  $\sim 150$  AP/min and mHR by  $\sim 120$  bpm (13). If anything, for both intrinsic and maximum rates, there is less slowing at the tissue level than the cell level, not more as would be predicted for additive effects.

Taken together, studies of human and animal heart rates, isolated SAN tissue, and isolated SAMs suggest that age-dependent declines in mHR are driven primarily by electrical remodeling at the cellular level with smaller declines attributable to structural remodeling and reduced  $\beta$ AR responsiveness.

## CONSEQUENCES FOR HUMAN HEALTH

The age-dependent decline in cardiac pacemaker function and ensuing declines in mHR and  $V_{O2max}$  have profound negative consequences for health span, quality of life, athletic performance, and societal health care costs. Physical exercise mitigates some effects of

aging and increases health span by slowing the decline in  $V_{O_{2max}}$  (4, 15); however, an inevitable decline in  $V_{O_{2max}}$  is imposed by the age-dependent decline in mHR (2–4).

$V_{O_{2max}}$  is a key determinant of functional independence and quality of life. Loss of functional independence in old age is correlated with  $V_{O_{2max}}$  below a threshold of 15–18 mL/(kg·min) (18). As the percentage of the population over the age of 65 continues to increase—20% of the US population will be over the age of 65 by 2035 (139)—the costs associated with this loss of independence will also increase. Age-dependent declines in mHR and  $V_{O_{2max}}$  also limit peak physical performance for elite athletes and weekend warriors alike. For the ever-growing and increasingly older population of recreational athletes, the age-dependent decline in  $V_{O_{2max}}$  limits not only performance but also the ability to participate at all.

While the age-dependent decline in pacemaker function reduces physical capacity for all people, some individuals experience more severe age-related pacemaker deficits. Clinically symptomatic SND—marked by bradyarrhythmias, alternating bradycardia and tachycardia, sinoatrial block, and dementia due to cerebral hypofusion—affects 1 in 600 people over the age of 65 (140). SND accounts for the majority of the 225,000 electronic pacemakers that are implanted each year in the United States at an annual cost of \$8.3 billion (141). In addition to the increased health care costs of treating SND as the population ages, surgically implanted pacemakers also carry risks for individuals, including infection and cardiac perforation (141).

The human and societal costs of age-dependent pacemaker deficits are exacerbated by a limited number of drugs to control heart rate. One FDA-approved drug, ivabradine, is available to slow heart rate by blocking HCN channels; however, it is not specific for the sinoatrial node HCN4 channel isoform and has some off-target effects (142–144). Similarly, there are few drugs to safely increase heart rate for treatment of bradycardia. Atropine is sometimes effective at increasing heart rate, but some elderly patients with bradyarrhythmia are resistant to atropine (145, 146), consistent with the loss of parasympathetic tone at rest in the elderly (120–122).

The ideal pharmaceutical to treat SND and enhance  $V_{O_{2max}}$  in elderly individuals would be one that acts specifically on the SAN to increase iHR, and therefore mHR, without affecting sympathetic tone or cardiac contractility. Of the factors that contribute to the age-related decrease in mHR, cellular mechanisms that drive the loss of pacemaker function with age appear to be the first-line drug targets. Structural remodeling would be challenging to target and hard to dose in response to continued changes as the individual ages. Meanwhile, targeting the decrease in  $\beta$ AR responsivity poses obvious challenges for sinoatrial specificity, as most targets would also impact other cardiovascular and systemic processes. Among cellular targets, HCN4 channels are particularly attractive owing to their relatively specific expression in the SAN and the proven effectiveness of  $I_f$  as a target for control of heart rate (71). While ion channel blockers often act on the structurally conserved pore domain and so can be prone to off-target effects, agonists can be targeted to less-conserved allosteric domains, making them alluring targets for high-specificity drugs (147).

## LIMITATIONS, GAPS IN KNOWLEDGE, AND FUTURE DIRECTIONS

The primary limitation to understanding the age-dependent slowing of mHR is a simple lack of data. Although the effects of aging on human heart rate, and even the firing rate of the SAN, are well characterized, few mechanistic data exist. Moreover, the conclusions that can be drawn from existing data sets are limited by difficulties in integrating information from different species and across different levels (molecular, cellular, tissue).

Gaps in knowledge about pacemaking exist across all levels of SAN and SAM function. At the cellular level, one of the key limitations to progress is the field's disproportionate focus on identifying a single most important mechanism for the diastolic depolarization in SAMs (52), which has left large gaps in knowledge about currents active during other phases of the AP. The relative lack of information about  $K^+$  currents in SAMs is particularly important for aging because the AP prolongation along with reduced  $Ca^{2+}$  current density predicts a decrease in  $K^+$  currents in aged SAMs (10, 11, 13). Although, as discussed above,  $K^+$  currents play an important role in determining cycle length through modulation of multiple AP waveform parameters and are implicated in SND (95, 97–100, 103), there is no information to our knowledge about age-dependent changes in  $K^+$  currents in SAMs.

There is also a lack of functional data corresponding to the more numerous, and sometimes conflicting, transcript studies. For example, although expression is reduced in aged SAN for a number of  $Ca^{2+}$ -handling proteins (11, 48, 86), there has not yet been a comparison of  $I_{NCX}$  in young versus old SAMs (in part due to difficulties inherent to experimental isolation of  $I_{NCX}$ ). Even in the case of currents for which age-dependent changes have been reported, the underlying mechanisms are unknown. For example, our data suggest that a novel HCN4-associated factor may be responsible for the age-dependent changes in  $I_f$  in SAMs (66, 67), but this factor has yet to be identified.

The lack of data on age-dependent ionic remodeling in single cells is compounded at the tissue level. Little is known about innervation and cell–cell coupling in general, much less how they may change with age. And while studies suggest that gap junction expression is altered with aging (46), cell–cell coupling between SAMs remains understudied.

The dearth of mechanistic data at the cellular and tissue levels in turn hinders the generation of validated mathematical models of APs in aged SAMs and precludes scaling of cellular data to the tissue level at this time. Modeling is a particularly important tool for understanding the complex and interrelated mechanisms that drive pacemaking in part because experimental perturbations often cause the entire dynamical system to fail. Acquisition of robust data sets under standardized conditions is a priority for future work in order to refine and improve mathematical models of the SAN.

Another important area for future work is a description of gender-dependent differences in the aging SAN. Such differences seem likely to exist—and to have profound implications for treatments—based on gender differences in aging of other regions of the heart (148). Indeed, while the rate of decline of mHR and iHR is indistinguishable between men and women (20), some studies suggest that women have higher rHRs and iHRs than men (149). Women are also more likely to have an electronic pacemaker implanted for SND or atrial fibrillation

with bradycardia (150), and numerous cardiovascular drugs have different effects in men and women (151).

## ACKNOWLEDGMENTS

This work was supported by grants from the US National Institutes of Health (R01-HL088427, T32-AG000279, F31-HL132408) and the American Heart Association (19POST34380777). The content is solely the responsibility of the authors and does not represent the official views of the National Institutes of Health or the American Heart Association.

## Glossary

|                          |  |
|--------------------------|--|
| <b>mHR</b>               | maximum heart rate                                   |
| <b>V<sub>O2max</sub></b> | maximum rate of oxygen consumption                   |
| <b>iHR</b>               | intrinsic heart rate                                 |
| <b>SAN</b>               | sinoatrial node                                      |
| <b>rHR</b>               | resting heart rate                                   |
| <b>SAM</b>               | sinoatrial node myocyte                              |
| <b>AP</b>                | action potential                                     |
| <b>DDR</b>               | diastolic depolarization rate                        |
| <b>MDP</b>               | maximum diastolic potential                          |
| <b>I<sub>f</sub></b>     | funny current  |
| <b>SR</b>                | sarcoplasmic reticulum                               |
| <b>NCX</b>               | Na <sup>+</sup> -Ca <sup>2+</sup> exchanger          |
| <b>RYR</b>               | ryanodine receptor                                   |
| <b>SERCA</b>             | sarco-/endoplasmic reticulum Ca <sup>2+</sup> ATPase |
| <b>SND</b>               | sinoatrial node dysfunction                          |
| <b>βAR</b>               | β-adrenergic receptor                                |

## LITERATURE CITED

1. Homer. 1990 Iliad. London: Penguin
2. Robinson S. 1938 Experimental studies of physical fitness in relation to age. *Eur. J. Appl. Physiol* 10(3):251–323 Seminal study showing age-dependent declines in mHR.
3. Robinson S, Dill DB, Tzankoff SP, Wagner JA, Robinson RD. 1975 Longitudinal studies of aging in 37 men. *J. Appl. Physiol* 38(2):263–67 [PubMed: 1120749]
4. Heath GW, Hagberg JM, Ehsani AA, Holloszy JO. 1981 A physiological comparison of young and older endurance athletes. *J. Appl. Physiol* 51(3):634–40 [PubMed: 7327965]
5. Tanaka H, Monahan KD, Seals DR. 2001 Age-predicted maximal heart rate revisited. *J. Am. Coll. Cardiol* 37(1):153–56 [PubMed: 11153730]

6. Tanaka H, Desouza CA, Jones PP, Stevenson ET, Davy KP, Seals DR. 1997 Greater rate of decline in maximal aerobic capacity with age in physically active versus sedentary healthy women. *J. Appl. Physiol* 83(6):1947–53 [PubMed: 9390967]
7. Fox SM, Naughton JP, Haskell WL. 1971 Physical activity and the prevention of coronary heart disease. *Ann. Clin. Res* 3(6):404–32 [PubMed: 4945367]
8. Nes BM, Janszky I, Wisløff U, Støylen A, Karlsen T. 2013 Age-predicted maximal heart rate in healthy subjects: the HUNT fitness study. *Scand. J. Med. Sci. Sports* 23(6):697–704 [PubMed: 22376273]
9. Di Gennaro M, Bernabei R, Sgadari A, Carosella L, Carbonin PU. 1987 Age-related differences in isolated rat sinus node function. *Basic Res. Cardiol* 82(6):530–36 [PubMed: 3435399]
10. Alings AM, Bouman LN. 1993 Electrophysiology of the ageing rabbit and cat sinoatrial node—a comparative study. *Eur. Heart J* 14(9):1278–88 [PubMed: 8223742]
11. Liu J, Sirenko S, Juhaszova M, Sollott SJ, Shukla S, et al. 2014 Age-associated abnormalities of intrinsic automaticity of sinoatrial nodal cells are linked to deficient cAMP-PKA-Ca<sup>2+</sup> signaling. *Am. J. Physiol. Heart Circ. Physiol* 306(10):H1385–97 [PubMed: 24633551]
12. Yaniv Y, Ahmet I, Tsutsui K, Behar J, Moen JM, et al. 2016 Deterioration of autonomic neuronal receptor signaling and mechanisms intrinsic to heart pacemaker cells contribute to age-associated alterations in heart rate variability in vivo. *Aging Cell* 15(4):716–24 [PubMed: 27168363]
13. Larson ED, St Clair JR, Sumner WA, Bannister RA, Proenza C. 2013 Depressed pacemaker activity of sinoatrial node myocytes contributes to the age-dependent decline in maximum heart rate. *PNAS* 110(44):18011–16 [PubMed: 24128759] Description of age-dependent declines in pacemaking and membrane currents of isolated sinoatrial myocytes.
14. Yin FC, Spurgeon HA, Greene HL, Lakatta EG, Weisfeldt ML. 1979 Age-associated decrease in heart rate response to isoproterenol in dogs. *Mech. Ageing Dev* 10(1–2):17–25 [PubMed: 449422]
15. Hagberg JM, Allen WK, Seals DR, Hurley BF, Ehsani AA, Holloszy JO. 1985 A hemodynamic comparison of young and older endurance athletes during exercise. *J. Appl. Physiol* 58(6):2041–46 [PubMed: 4008419]
16. Tanaka H, Seals DR. 2008 Endurance exercise performance in Masters athletes: age-associated changes and underlying physiological mechanisms. *J. Physiol* 586(1):55–63 [PubMed: 17717011]
17. Paterson DH, Govindasamy D, Vidmar M, Cunningham DA, Koval JJ. 2004 Longitudinal study of determinants of dependence in an elderly population. *J. Am. Geriatr. Soc* 52(10):1632–38 [PubMed: 15450038]
18. Shephard RJ. 2009 Maximal oxygen intake and independence in old age. *Br. J. Sports Med* 43(5):342–46 [PubMed: 18403414]
19. Hawkins S, Wiswell R. 2003 Rate and mechanism of maximal oxygen consumption decline with aging: implications for exercise training. *Sports Med.* 33(12):877–88 [PubMed: 12974656]
20. Jose AD, Collison D. 1970 The normal range and determinants of the intrinsic heart rate in man. *Cardiovasc. Res* 4(2):160–67 [PubMed: 4192616] Seminal study on age-dependent declines in iHR.
21. Jose AD, Stitt F, Collison D. 1970 The effects of exercise and changes in body temperature on the intrinsic heart rate in man. *Am. Heart J* 79(4):488–98 [PubMed: 5418022]
22. Christou DD, Seals DR. 2008 Decreased maximal heart rate with aging is related to reduced  $\beta$ -adrenergic responsiveness but is largely explained by a reduction in intrinsic heart rate. *J. Appl. Physiol* 105(1):24–29 [PubMed: 18483165] Study measuring the contribution of decreased iHR and  $\beta$ AR agonist sensitivity to declines in mHR.
23. Silverman ME, Hollman A. 2007 Discovery of the sinus node by Keith and Flack: on the centennial of their 1907 publication. *Heart* 93(10):1184–87 [PubMed: 17890694]
24. Fye WB. 1987 The origin of the heart beat: a tale of frogs, jellyfish, and turtles. *Circulation* 76(3):493–500 [PubMed: 3304703]
25. Keith A, Flack M. 1907 The form and nature of the muscular connections between the primary divisions of the vertebrate heart. *J. Anat. Physiol* 41(Part 3):172–89 [PubMed: 17232727] Description of SAN anatomy in multiple species.
26. Bexton RS, Nathan AW, Hellestrand KJ, Cory-Pearce R, Spurrell RA, et al. 1984 Sinoatrial function after cardiac transplantation. *J. Am. Coll. Cardiol* 3(3):712–23 [PubMed: 6363487]

27. Strobel JS, Epstein AE, Bourge RC, Kirklin JK, Kay GN. 1999 Nonpharmacologic validation of the intrinsic heart rate in cardiac transplant recipients. *J. Interv. Card. Electrophysiol. Int. J. Arrhythm. Pacing* 3(1):15–18
28. Robinson BF, Epstein SE, Beiser GD, Braunwald E. 1966 Control of heart rate by the autonomic nervous system. Studies in man on the interrelation between baroreceptor mechanisms and exercise. *Circ. Res* 19(2):400–11 [PubMed: 5914852]
29. Masson-Pévet M, Bleeker WK, Gros D. 1979 The plasma membrane of leading pacemaker cells in the rabbit sinus node. A qualitative and quantitative ultrastructural analysis. *Circ. Res* 45(5):621–29 [PubMed: 487525]
30. James TN, Sherf L, Fine G, Morales AR. 1966 Comparative ultrastructure of the sinus node in man and dog. *Circulation* 34(1):139–63 [PubMed: 5942665]
31. Ophhof T, de Jonge B, Jongsma HJ, Bouman LN. 1987 Functional morphology of the mammalian sinoatrial node. *Eur. Heart J* 8(11):1249–59 [PubMed: 3691562]
32. Boyett MR, Honjo H, Kodama I. 2000 The sinoatrial node, a heterogeneous pacemaker structure. *Cardiovasc. Res* 47(4):658–87 [PubMed: 10974216]
33. Csepe TA, Zhao J, Hansen BJ, Li N, Sul LV, et al. 2016 Human sinoatrial node structure: 3D microanatomy of sinoatrial conduction pathways. *Prog. Biophys. Mol. Biol* 120(1–3):164–78 [PubMed: 26743207]
34. Kwong KF, Schuessler RB, Green KG, Laing JG, Beyer EC, et al. 1998 Differential expression of gap junction proteins in the canine sinus node. *Circ. Res* 82(5):604–12 [PubMed: 9529165]
35. Coppén SR, Kodama I, Boyett MR, Dobrzynski H, Takagishi Y, et al. 1999 Connexin45, a major connexin of the rabbit sinoatrial node, is co-expressed with connexin43 in a restricted zone at the nodal-crista terminalis border. *J. Histochem. Cytochem* 47(7):907–18 [PubMed: 10375379]
36. Kreuzberg MM, Söhl G, Kim J-S, Verselis VK, Willecke K, Bukauskas FF. 2005 Functional properties of mouse connexin30.2 expressed in the conduction system of the heart. *Circ. Res* 96(11):1169–77 [PubMed: 15879306]
37. Haubrich S, Schwarz HJ, Bukauskas F, Lichtenberg-Fraté H, Traub O, et al. 1996 Incompatibility of connexin 40 and 43 Hemichannels in gap junctions between mammalian cells is determined by intracellular domains. *Mol. Biol. Cell* 7(12):1995–2006 [PubMed: 8970160]
38. Kreuzberg MM, Willecke K, Bukauskas FF. 2006 Connexin-mediated cardiac impulse propagation: connexin 30.2 slows atrioventricular conduction in mouse heart. *Trends Cardiovasc. Med* 16(8):266–72 [PubMed: 17055382]
39. Anumonwo JM, Wang HZ, Trabka-Janik E, Dunham B, Veenstra RD, et al. 1992 Gap junctional channels in adult mammalian sinus nodal cells. Immunolocalization and electrophysiology. *Circ. Res* 71(2):229–39 [PubMed: 1378358]
40. Joyner RW, van Capelle FJ. 1986 Propagation through electrically coupled cells. How a small SA node drives a large atrium. *Biophys. J* 50(6):1157–64 [PubMed: 3801575]
41. Yamamoto M, Honjo H, Niwa R, Kodama I. 1998 Low-frequency extracellular potentials recorded from the sinoatrial node. *Cardiovasc. Res* 39(2):360–72 [PubMed: 9798521]
42. Fedorov VV, Glukhov AV, Chang R, Kosteki G, Aferol H, et al. 2010 Optical mapping of the isolated coronary-perfused human sinus node. *J. Am. Coll. Cardiol* 56(17):1386–94 [PubMed: 20946995]
43. Csepe TA, Kalyanasundaram A, Hansen BJ, Zhao J, Fedorov VV. 2015 Fibrosis: a structural modulator of sinoatrial node physiology and dysfunction. *Front. Physiol* 6:37 [PubMed: 25729366] Review of fibrosis in the dysfunction of aging SAN.
44. Valiunas V, Weingart R, Brink PR. 2000 Formation of heterotypic gap junction channels by connexins 40 and 43. *Circ. Res* 86(2):E42–49 [PubMed: 10666425]
45. Bukauskas FF, Angele AB, Verselis VK, Bennett MVL. 2002 Coupling asymmetry of heterotypic connexin 45/connexin 43-EGFP gap junctions: properties of fast and slow gating mechanisms. *PNAS* 99(10):7113–18 [PubMed: 12011467]
46. Jones SA, Lancaster MK, Boyett MR. 2004 Ageing-related changes of connexins and conduction within the sinoatrial node. *J. Physiol* 560(Part 2):429–37 [PubMed: 15308686]



47. Moghtadaei M, Jansen HJ, Mackasey M, Rafferty SA, Bogachev O, et al. 2016 The impacts of age and frailty on heart rate and sinoatrial node function. *J. Physiol* 594(23):7105–26 [PubMed: 27598221]
48. Tellez JO, M czewski M, Yanni J, Sutyagin P, Mackiewicz U, et al. 2011 Ageing-dependent remodelling of ion channel and Ca<sup>2+</sup> clock genes underlying sino-atrial node pacemaking. *Exp. Physiol* 96(11):1163–78 [PubMed: 21724736] Review of fibrosis in the dysfunction of aging SAN.
49. Cavoto FV, Kelliher GJ, Roberts J. 1974 Electrophysiological changes in the rat atrium with age. *Am. J. Physiol* 226(6):1293–97 [PubMed: 4833987]
50. Rickert C, Proenza C. 2017 ParamAP: standardized parameterization of sinoatrial node myocyte action potentials. *Biophys. J* 113(4):765–69 [PubMed: 28834713]
51. Mangoni ME, Nargeot J. 2008 Genesis and regulation of the heart automaticity. *Physiol. Rev* 88(3):919–82 [PubMed: 18626064]
52. Lakatta EG, DiFrancesco D. 2009 JMCC point-counterpoint. *J. Mol. Cell. Cardiol* 47(2):157–70 [PubMed: 19361514]
53. Monfredi O, Dobrzynski H, Mondal T, Boyett MR, Morris GM. 2010 The anatomy and physiology of the sinoatrial node—a contemporary review. *PACE Pacing Clin. Electrophysiol* 33(11):1392–406 [PubMed: 20946278]
54. Brown H, DiFrancesco D, Noble S. 1979 Cardiac pacemaker oscillation and its modulation by autonomic transmitters. *J. Exp. Biol* 81:175–204 [PubMed: 315983]
55. Moosmang S, Biel M, Hofmann F, Ludwig A. 1999 Differential distribution of four hyperpolarization-activated cation channels in mouse brain. *Biol. Chem* 380(7–8):975–80 [PubMed: 10494850]
56. Moosmang S, Stieber J, Zong X, Biel M, Hofmann F, Ludwig A. 2001 Cellular expression and functional characterization of four hyperpolarization-activated pacemaker channels in cardiac and neuronal tissues. *Eur. J. Biochem* 268(6):1646–52 [PubMed: 11248683]
57. Li N, Csepe TA, Hansen BJ, Dobrzynski H, Higgins RSD, et al. 2015 Molecular mapping of sinoatrial node HCN channel expression in the human heart. *Circ. Arrhythm. Electrophysiol* 8(5):1219–27 [PubMed: 26304511]
58. Huang X, Yang P, Yang Z, Zhang H, Ma A. 2016 Age-associated expression of HCN channel isoforms in rat sinoatrial node. *Exp. Biol. Med* 241(3):331–39
59. Herrmann S, Layh B, Ludwig A. 2011 Novel insights into the distribution of cardiac HCN channels: an expression study in the mouse heart. *J. Mol. Cell. Cardiol* 51(6):997–1006 [PubMed: 21945247]
60. Moroni A, Gorza L, Beltrame M, Gravante B, Vaccari T, et al. 2001 Hyperpolarization-activated cyclic nucleotide-gated channel 1 is a molecular determinant of the cardiac pacemaker current *I<sub>f</sub>*. *J. Biol. Chem* 276(31):29233–41 [PubMed: 11328811]
61. Ludwig A, Budde T, Stieber J, Moosmang S, Wahl C, et al. 2003 Absence epilepsy and sinus dysrhythmia in mice lacking the pacemaker channel HCN2. *EMBO J.* 22(2):216–24 [PubMed: 12514127]
62. Flynn GE, Zagotta WN. 2018 Insights into the molecular mechanism for hyperpolarization-dependent activation of HCN channels. *PNAS* 115(34):E8086–95 [PubMed: 30076228]
63. Bucchi A, Barbuti A, DiFrancesco D, Baruscotti M. 2012 Funny current and cardiac rhythm: insights from HCN knockout and transgenic mouse models. *Front. Physiol* 3:240 [PubMed: 22783204]
64. Verkerk AO, Wilders R. 2013 Hyperpolarization-activated current, *I<sub>f</sub>* in mathematical models of rabbit sinoatrial node pacemaker cells. *BioMed. Res. Int* 2013:872454 [PubMed: 23936852]
65. Sharpe EJ, Gantz SC, Liu P, Bean BP, Proenza C. 2017 Characteristics of ivabradine-sensitive currents in mouse sinoatrial node myocytes. *Biophys. J* 112(3):35a–36a
66. Sharpe EJ, Larson ED, Proenza C. 2017 Cyclic AMP reverses the effects of aging on pacemaker activity and *I<sub>f</sub>* in sinoatrial node myocytes. *J. Gen. Physiol* 149(2):237–47 [PubMed: 28057842] Study showing cAMP can restore *I<sub>f</sub>* and the firing rate of aged SAMs to youthful levels.
67. Robertson GA. 2017 It's not funny: how changes in *I<sub>f</sub>* limit maximum heart rate with aging. *J. Gen. Physiol* 149(2):177–79 [PubMed: 28057841]

68. Yanni J, Tellez JO, Sutyagin PV, Boyett MR, Dobrzynski H. 2010 Structural remodelling of the sinoatrial node in obese old rats. *J. Mol. Cell. Cardiol* 48(4):653–62 [PubMed: 19729016]
69. Huang X, Yang P, Du Y, Zhang J, Ma A. 2007 Age-related down-regulation of HCN channels in rat sinoatrial node. *Basic Res. Cardiol* 102(5):429–35 [PubMed: 17572839]
70. Li Y-D, Hong Y-F, Yusufuaji Y, Tang B-P, Zhou X-H, et al. 2015 Altered expression of hyperpolarization-activated cyclic nucleotide-gated channels and microRNA-1 and -133 in patients with age-associated atrial fibrillation. *Mol. Med. Rep* 12(3):3243–48 [PubMed: 26005035]
71. Ragueneau I, Laveille C, Jochemsen R, Resplandy G, Funck-Brentano C, Jaillon P. 1998 Pharmacokinetic-pharmacodynamic modeling of the effects of ivabradine, a direct sinus node inhibitor, on heart rate in healthy volunteers. *Clin. Pharmacol. Ther* 64(2):192–203 [PubMed: 9728900]
72. Baruscotti M, Bucchi A, Viscomi C, Mandelli G, Consalez G, et al. 2011 Deep bradycardia and heart block caused by inducible cardiac-specific knockout of the pacemaker channel gene *Hcn4*. *PNAS* 108(4):1705–10 [PubMed: 21220308]
73. Milano A, Vermeer AMC, Lodder EM, Barc J, Verkerk AO, et al. 2014 *HCN4* mutations in multiple families with bradycardia and left ventricular noncompaction cardiomyopathy. *J. Am. Coll. Cardiol* 64(8):745–56 [PubMed: 25145517]
74. Pilcher JD. 1912 The action of caffeine on the mammalian heart. *J. Pharmacol. Exp. Ther* 3(6):609–24
75. Tsien RW, Kass RS, Weingart R. 1979 Cellular and subcellular mechanisms of cardiac pacemaker oscillations. *J. Exp. Biol* 81:205–15 [PubMed: 512578]
76. Rapp PE, Berridge MJ. 1977 Oscillations in calcium-cyclic AMP control loops form the basis of pacemaker activity and other high frequency biological rhythms. *J. Theor. Biol* 66(3):497–525 [PubMed: 196141]
77. Rubenstein DS, Lipsius SL. 1989 Mechanisms of automaticity in subsidiary pacemakers from cat right atrium. *Circ. Res* 64(4):648–57 [PubMed: 2467760]
78. Rigg L, Terrar DA. 1996 Possible role of calcium release from the sarcoplasmic reticulum in pacemaking in guinea-pig sino-atrial node. *Exp. Physiol* 81(5):877–80 [PubMed: 8889484]
79. Chen B, Wu Y, Mohler PJ, Anderson ME, Song LS. 2009 Local control of  $Ca^{2+}$ -induced  $Ca^{2+}$  release in mouse sinoatrial node cells. *J. Mol. Cell. Cardiol* 47(5):706–15 [PubMed: 19615376]
80. Bogdanov KY, Vinogradova TM, Lakatta EG. 2001 Sinoatrial nodal cell ryanodine receptor and  $Na^+$ - $Ca^{2+}$  exchanger. *Circ. Res* 88(12):1254–58 [PubMed: 11420301]
81. Li J, Qu J, Nathan RD. 1997 Ionic basis of ryanodine's negative chronotropic effect on pacemaker cells isolated from the sinoatrial node. *Am. J. Physiol. Heart Circ. Physiol* 273(5):H2481–89
82. Vinogradova TM, Brochet D, Sirenko S, Li Y, Spurgeon H, Lakatta EG. 2010 Sarcoplasmic reticulum  $Ca^{2+}$  pumping kinetics regulates timing of local  $Ca^{2+}$  releases and spontaneous beating rate of rabbit sinoatrial node pacemaker cells. *Circ. Res* 107(6):767–75 [PubMed: 20651285]
83. Dun W, Yagi T, Rosen MR, Boyden PA. 2003 Calcium and potassium currents in cells from adult and aged canine right atria. *Cardiovasc. Res* 58(3):526–34 [PubMed: 12798425]
84. Xu G-J, Gan T-Y, Tang B-P, Chen Z-H, Jiang T, et al. 2013 Age-related changes in cellular electrophysiology and calcium handling for atrial fibrillation. *J. Cell. Mol. Med* 17(9):1109–18 [PubMed: 23837844]
85. Herraiz-Martínez A, Álvarez-García J, Llach A, Molina CE, Fernandes J, et al. 2015 Ageing is associated with deterioration of calcium homeostasis in isolated human right atrial myocytes. *Cardiovasc. Res* 106(1):76–86 [PubMed: 25712961]
86. Jones SA, Boyett MR, Lancaster MK. 2007 Declining into failure the age-dependent loss of the L-type calcium channel within the sinoatrial node. *Circulation* 115(10):1183–90 [PubMed: 17339548]
87. Lipsius SL, Vassalle M. 1978 Dual excitatory channels in the sinus node. *J. Mol. Cell. Cardiol* 10(8):753–67 [PubMed: 702536]
88. Muramatsu H, Zou AR, Berkowitz GA, Nathan RD. 1996 Characterization of a TTX-sensitive  $Na^+$  current in pacemaker cells isolated from rabbit sinoatrial node. *Am. J. Physiol. Heart Circ. Physiol* 270(6):H2108–19

89. Maier SKG, Westenbroek RE, Yamanushi TT, Dobrzynski H, Boyett MR, et al. 2003 An unexpected requirement for brain-type sodium channels for control of heart rate in the mouse sinoatrial node. *PNAS* 100(6):3507–12 [PubMed: 12631690]
90. Ruan Y, Liu N, Priori SG. 2009 Sodium channel mutations and arrhythmias. *Nat. Rev. Cardiol* 6(5):337–48 [PubMed: 19377496]
91. Wu J, Zhang Y, Zhang X, Cheng L, Lammers WJ, et al. 2012 Altered sinoatrial node function and intra-atrial conduction in murine gain-of-function *Scn5a*<sup>+/KPO</sup> hearts suggest an overlap syndrome. *Am. J. Physiol. Heart Circ. Physiol* 302(7):H1510–23 [PubMed: 22287583]
92. Protas L, Oren RV, Clancy CE, Robinson RB. 2010 Age-dependent changes in Na current magnitude and TTX-sensitivity in the canine sinoatrial node. *J. Mol. Cell. Cardiol* 48(1):172–80 [PubMed: 19665465]
93. Bettahi I, Marker CL, Roman MI, Wickman K. 2002 Contribution of the Kir3.1 subunit to the muscarinic-gated atrial potassium channel  $I_{K_{ACh}}$ . *J. Biol. Chem* 277(50):48282–88 [PubMed: 12374786]
94. Weisbrod D, Peretz A, Ziskind A, Menaker N, Oz S, et al. 2013 SK4  $Ca^{2+}$  activated  $K^{+}$  channel is a critical player in cardiac pacemaker derived from human embryonic stem cells. *PNAS* 110(18):E1685–94 [PubMed: 23589888]
95. Lai MH, Wu Y, Gao Z, Anderson ME, Dalziel JE, Meredith AL. 2014 BK channels regulate sinoatrial node firing rate and cardiac pacing in vivo. *Am. J. Physiol. Heart Circ. Physiol* 307(9):H1327–38 [PubMed: 25172903]
96. Weisbrod D, Khun SH, Bueno H, Peretz A, Attali B. 2016 Mechanisms underlying the cardiac pacemaker: the role of SK4 calcium-activated potassium channels. *Acta Pharmacol. Sin* 37(1):82–97 [PubMed: 26725737]
97. Aziz Q, Li Y, Tinker A. 2018 Potassium channels in the sinoatrial node and their role in heart rate control. *Channels* 12(1):356–66 [PubMed: 30301404]
98. Thomas D, Wimmer A-B, Karle CA, Licka M, Alter M, et al. 2005 Dominant-negative  $I_{K_s}$  suppression by KCNQ1- F339 potassium channels linked to Romano-Ward syndrome. *Cardiovasc. Res* 67(3):487–97 [PubMed: 15950200]
99. Horigome H, Nagashima M, Sumitomo N, Yoshinaga M, Ushinohama H, et al. 2010 Clinical characteristics and genetic background of congenital long-QT syndrome diagnosed in fetal, neonatal, and infantile life: a nationwide questionnaire survey in Japan. *Circ. Arrhythm. Electrophysiol* 3(1):10–17 [PubMed: 19996378]
100. Wilders R, Verkerk AO. 2018 Long QT syndrome and sinus bradycardia—a mini review. *Front. Cardiovasc. Med* 5:106 [PubMed: 30123799]
101. Wang L, Swirp S, Duff H. 2000 Age-dependent response of the electrocardiogram to  $K^{+}$  channel blockers in mice. *Am. J. Physiol. Cell Physiol* 278(1):C73–80 [PubMed: 10644514]
102. Furukawa Y, Miyashita Y, Nakajima K, Hirose M, Kurogouchi F, Chiba S. 1999 Effects of verapamil, zatebradine, and E-4031 on the pacemaker location and rate in response to sympathetic stimulation in dog hearts. *J. Pharmacol. Exp. Ther* 289(3):1334–42 [PubMed: 10336524]
103. Kuß J, Stallmeyer B, Goldstein M, Rinné S, Pees C, et al. 2019 Familial sinus node disease caused by a gain of GIRK (G-protein activated inwardly rectifying  $K^{+}$  channel) channel function. *Circ. Genomic Precis. Med* 12(1):e002238
104. Wickman K, Nemeč J, Gendler SJ, Clapham DE. 1998 Abnormal heart rate regulation in *GIRK4* knockout mice. *Neuron* 20(1):103–14 [PubMed: 9459446]
105. Liu Y, Beyer A, Aebersold R. 2016 On the dependency of cellular protein levels on mRNA abundance. *Cell* 165(3):535–50 [PubMed: 27104977]
106. Shiraishi I, Takamatsu T, Minamikawa T, Onouchi Z, Fujita S. 1992 Quantitative histological analysis of the human sinoatrial node during growth and aging. *Circulation* 85(6):2176–84 [PubMed: 1591834]
107. Thery C, Gosselin B, Lekieffre J, Warembourg H. 1977 Pathology of sinoatrial node. Correlations with electrocardiographic findings in 111 patients. *Am. Heart J* 93(6):735–40 [PubMed: 871100]
108. Inoue S, Shinohara F, Niitani H, Gotoh K. 1986 A new method for the histological study of aging changes in the sinoatrial node. *Jpn. Heart J* 27(5):653–60 [PubMed: 3546778]


109. Kistler PM, Sanders P, Fynn SP, Stevenson IH, Spence SJ, et al. 2004 Electrophysiologic and electroanatomic changes in the human atrium associated with age. *J. Am. Coll. Cardiol* 44(1):109–16 [PubMed: 15234418]
110. Sanders P, Morton JB, Kistler PM, Spence SJ, Davidson NC, et al. 2004 Electrophysiological and electroanatomic characterization of the atria in sinus node disease. *Circulation* 109(12):1514–22 [PubMed: 15007004]
111. Monfredi O, Boyett MR. 2015 Sick sinus syndrome and atrial fibrillation in older persons—a view from the sinoatrial nodal myocyte. *J. Mol. Cell. Cardiol* 83:88–100 [PubMed: 25668431]
112. Hao X, Zhang Y, Zhang X, Nirmalan M, Davies L, et al. 2011 TGF- $\beta_1$ -mediated fibrosis and ion channel remodeling are key mechanisms in producing the sinus node dysfunction associated with *SCN5A* deficiency and aging. *Circ. Arrhythm. Electrophysiol* 4(3):397–406 [PubMed: 21493874]
113. Glukhov AV, Kalyanasundaram A, Lou Q, Hage LT, Hansen BJ, et al. 2015 Calsequestrin 2 deletion causes sinoatrial node dysfunction and atrial arrhythmias associated with altered sarcoplasmic reticulum calcium cycling and degenerative fibrosis within the mouse atrial pacemaker complex. *Eur. Heart J* 36(11):686–97 [PubMed: 24216388]
114. de Melo SR, de Souza RR, Mandarim-de-Lacerda CA. 2002 Stereologic study of the sinoatrial node of rats—age related changes. *Biogerontology* 3(6):383–90 [PubMed: 12510177]
115. Swaminathan PD, Purohit A, Soni S, Voigt N, Singh MV, et al. 2011 Oxidized CaMKII causes cardiac sinus node dysfunction in mice. *J. Clin. Investig* 121(8):3277–88 [PubMed: 21785215]
116. Kharche S, Beling J, Biktasheva IV, Zhang H, Biktashev VN. 2013 Simulating cell apoptosis induced sinus node dysfunction. Paper presented at the 35th Annual International Conference of the IEEE Engineering in Medicine and Biology Society, Osaka, Japan, July 3–7
117. Lakatta EG, Gerstenblith G, Angell CS, Shock NW, Weisfeldt ML. 1975 Diminished inotropic response of aged myocardium to catecholamines. *Circ. Res* 36(2):262–69 [PubMed: 1116236]
118. Stratton JR, Cerqueira MD, Schwartz RS, Levy WC, Veith RC, et al. 1992 Differences in cardiovascular responses to isoproterenol in relation to age and exercise training in healthy men. *Circulation* 86(2):504–12 [PubMed: 1638718]
119. Fleg JL, Schulman S, O'Connor F, Becker LC, Gerstenblith G, et al. 1994 Effects of acute beta-adrenergic receptor blockade on age-associated changes in cardiovascular performance during dynamic exercise. *Circulation* 90(5):2333–41 [PubMed: 7955191]
120. Abhishekh HA, Nisarga P, Kisan R, Meghana A, Chandran S, et al. 2013 Influence of age and gender on autonomic regulation of heart. *J. Clin. Monit. Comput* 27(3):259–64 [PubMed: 23297094]
121. Freeling JL, Li Y. 2015 Age-related attenuation of parasympathetic control of the heart in mice. *Int. J. Physiol. Pathophysiol. Pharmacol* 7(3):126–35 [PubMed: 26823961]
122. Baker SE, Limberg JK, Dillon GA, Curry TB, Joyner MJ, Nicholson WT. 2018 Aging alters the relative contributions of the sympathetic and parasympathetic nervous system to blood pressure control in women. *Hypertension* 72(5):1236–42 [PubMed: 30354803]
123. Vestal RE, Wood AJJ, Shand DG. 1979 Reduced  $\beta$ -adrenoceptor sensitivity in the elderly. *Clin. Pharmacol. Ther* 26(2):181–86 [PubMed: 37014]
124. Alboni P, Malcarne C, Pedroni P, Masoni A, Narula OS. 1982 Electrophysiology of normal sinus node with and without autonomic blockade. *Circulation* 65(6):1236–42 [PubMed: 7074781]
125. St. Clair JR, Liao Z, Larson ED, Proenza C. 2013 PKA-independent activation of  $I_f$  by cAMP in mouse sinoatrial myocytes. *Channels* 7(4):318–21 [PubMed: 23756695]
126. Striessnig J, Pinggera A, Kaur G, Bock G, Tuluc P. 2014 L-type  $Ca^{2+}$  channels in heart and brain. *Wiley Interdiscip. Rev. Membr. Transp. Signal* 3(2):15–38 [PubMed: 24683526]
127. Vinogradova TM, Bogdanov KY, Lakatta EG. 2002  $\beta$ -Adrenergic stimulation modulates ryanodine receptor  $Ca^{2+}$  release during diastolic depolarization to accelerate pacemaker activity in rabbit sinoatrial nodal cells. *Circ. Res* 90(1):73–79 [PubMed: 11786521]
128. Shen J. 2006 Isoprenaline enhances local  $Ca^{2+}$  release in cardiac myocytes. *Acta Pharmacol. Sin* 27(7):927–32 [PubMed: 16787578]

129. Yoshida A, Takahashi M, Imagawa T, Shigekawa M, Takisawa H, Nakamura T. 1992 Phosphorylation of ryanodine receptors in rat myocytes during beta-adrenergic stimulation. *J. Biochem* 111(2):186–90 [PubMed: 1349013]
130. Uehara A, Yasukochi M, Mejía-Alvarez R, Fill M, Imanaga I. 2002 Gating kinetics and ligand sensitivity modified by phosphorylation of cardiac ryanodine receptors. *Pflügers Arch.* 444(1–2):202–12 [PubMed: 11976933]
131. Kaese S, Bögeholz N, Pauls P, Dechering D, Olligs J, et al. 2017 Increased sodium/calcium exchanger activity enhances beta-adrenergic-mediated increase in heart rate: whole-heart study in a homozygous sodium/calcium exchanger overexpressor mouse model. *Heart Rhythm* 14(8):1247–53 [PubMed: 28495655]
132. Sanders L, Rakovic S, Lowe M, Mattick PAD, Terrar DA. 2006 Fundamental importance of Na<sup>+</sup>–Ca<sup>2+</sup> exchange for the pacemaking mechanism in guinea-pig sino-atrial node. *J. Physiol* 571(Part 3):639–49 [PubMed: 16423859]
133. Liao Z, Lockhead D, Larson E, Proenza C. 2010 Phosphorylation and modulation of hyperpolarization-activated HCN4 channels by protein kinase A in the mouse sinoatrial node. *J. Gen. Physiol* 136(3):247–58 [PubMed: 20713547]
134. Simmerman HKB, Jones LR. 1998 Phospholamban: protein structure, mechanism of action, and role in cardiac function. *Physiol. Rev* 78(4):921–47 [PubMed: 9790566]
135. O'Connor SW, Scarpace PJ, Abrass IB. 1981 Age-associated decrease of adenylate cyclase activity in rat myocardium. *Mech. Ageing Dev* 16(1):91–95 [PubMed: 6265716]
136. Hardouin S, Bourgeois F, Toraasson M, Oubenaissa A, Elalouf JM, et al. 1998  $\beta$ -Adrenergic and muscarinic receptor mRNA accumulation in the sinoatrial node area of adult and senescent rat hearts. *Mech. Ageing Dev* 100(3):277–97 [PubMed: 9578116]
137. Yaniv Y, Ahmet I, Liu J, Lyashkov AE, Guiriba T-R, et al. 2014 Synchronization of sinoatrial node pacemaker cell clocks and its autonomic modulation impart complexity to heart beating intervals. *Heart Rhythm* 11(7):1210–19 [PubMed: 24713624]
138. Behar J, Yaniv Y. 2017 Age-related pacemaker deterioration is due to impaired intracellular and membrane mechanisms: insights from numerical modeling. *J. Gen. Physiol* 149(10):935–49 [PubMed: 28887411]
139. Ortman JM, Velkoff VA, Hogan H. 2014 An aging nation: the older population in the United States *Curr. Pop. Rep* P25–1140, US Dep. Commer., Washington, DC <https://www.census.gov/prod/2014pubs/p25-1140.pdf>
140. Semelka M, Gera J, Usman S. 2013 Sick sinus syndrome: a review. *Am. Fam. Physician* 87(10):691–96 [PubMed: 23939447]
141. Sideris S, Archontakis S, Dilaveris P, Gatzoulis KA, Trachanas K, et al. 2017 Leadless cardiac pacemakers: current status of a modern approach in pacing. *Hellenic J. Cardiol* 58(6):403–10 [PubMed: 28529181]
142. Stieber J, Wieland K, Stöckl G, Ludwig A, Hofmann F. 2006 Bradycardic and proarrhythmic properties of sinus node inhibitors. *Mol. Pharmacol* 69(4):1328–37 [PubMed: 16387796]
143. Lees-Miller JP, Guo J, Wang Y, Perissinotti LL, Noskov SY, Duff HJ. 2015 Ivabradine prolongs phase 3 of cardiac repolarization and blocks the HERG1 (KCNH2) current over a concentration-range overlapping with that required to block HCN4. *J. Mol. Cell. Cardiol* 85:71–78 [PubMed: 25986146]
144. Melgari D, Brack KE, Zhang C, Zhang Y, El Harchi A, et al. 2015 hERG potassium channel blockade by the HCN channel inhibitor bradycardic agent ivabradine. *J. Am. Heart Assoc* 4(4):e001813 [PubMed: 25911606]
145. Kragie L, Sekovski B. 1992 Theophylline—an alternative therapy for bradyarrhythmia in the elderly. *Pharmacotherapy* 12(4):324–30 [PubMed: 1518730]
146. Pasnoori VR, Leesar MA. 2004 Use of aminophylline in the treatment of severe symptomatic bradycardia resistant to atropine. *Cardiol. Rev* 12(2):65–68 [PubMed: 14766019]
147. Proenza C. 2018 Ion channels: exploiting natural regulation. *eLife* 7:e39664 [PubMed: 30080138]
148. Kane AE, Howlett SE. 2018 Differences in cardiovascular aging in men and women. *Adv. Exp. Med. Biol* 1065:389–411 [PubMed: 30051398]

149. Burke JH, Goldberger JJ, Ehlert FA, Kruse JT, Parker MA, Kadish AH. 1996 Gender differences in heart rate before and after autonomic blockade: evidence against an intrinsic gender effect. *Am. J. Med* 100(5):537–43 [PubMed: 8644766]
150. Nowak B, Misselwitz B, Erdogan A, Funck R, et al. 2010 Do gender differences exist in pacemaker implantation?—results of an obligatory external quality control program. *Europace* 12(2):210–15 [PubMed: 19864309]
151. Tamargo J, Rosano G, Walther T, Duarte J, Niessner A, et al. 2017 Gender differences in the effects of cardiovascular drugs. *Eur. Heart J. Cardiovasc. Pharmacother* 3(3):163–82 [PubMed: 28329228]
152. Park DS, Fishman GI. 2017 Development and function of the cardiac conduction system in health and disease. *J. Cardiovasc. Dev. Dis* 4(2):7 [PubMed: 29098150]
153. Kharche SR, Vigmond E, Efimov IR, Dobrzynski H. 2017 Computational assessment of the functional role of sinoatrial node exit pathways in the human heart. *PLOS ONE* 12(9):e0183727 [PubMed: 28873427]


### SUMMARY POINTS

1. Aging reduces mHR by reducing iHR and thus by reducing the pacemaker activity of the sinoatrial node of the heart.
2. The progressive decrease in iHR with age necessitates a corresponding increase in sympathetic tone in order to maintain rHR.
3. The age-dependent reduction in mHR is crucial for individuals and society because it causes an inevitable reduction in  $V_{O2max}$ , which not only limits athletic performance but can also limit functional independence and quality of life for the growing population of elderly people.
4. The mechanisms by which aging slows the heart include electrical and structural remodeling as well as a reduced response to  $\beta$ AR stimulation. Of these factors, the reductions in cellular mechanisms that are responsible for iHR predominate and represent the best potential for development of drugs that can selectively increase HR and  $V_{O2max}$  in elderly individuals.
5. There is an urgent need for additional functional data to identify molecular mechanisms that contribute to decreased SAN function with age. Important areas for future work include acquisition of expanded data sets to improve mathematical models, description of gender differences in SAN aging, and additional information about electrical coupling in the SAN to improve understanding of tissue-level properties.
6. Given that aging affects all individuals, the clinical implications and benefits of future work deciphering the underlying mechanisms of cardiac pacemaker decline are clear.




**Figure 1.** Similar declines in maximum and intrinsic heart rates with no change in resting heart rate necessitate an increase in sympathetic tone with age. Maximum heart rate (mHR; *red line*) is plotted as  $208 - (0.7 \times \text{age})$ ; data from Reference 5. Intrinsic heart rate (*green line*) is plotted as  $118 - (0.57 \times \text{age})$ ; data from Reference 20. Resting heart rate (*dashed black line*) is plotted as a constant 70 beats per minute (bpm) throughout life. The red arrowhead indicates an average transition age (84.2 years) beyond which sympathetic tone at rest (*pink shading*) predominates over parasympathetic tone at rest (*green shading*) in order to maintain resting heart rate. Sympathetic capacity (*black double arrows*), the difference between intrinsic and maximum heart rate, decreases slightly with age; however, the majority of the age-dependent decline in mHR is caused by the decline in intrinsic heart rate.





**Figure 2.**

The cardiac conduction system and sinoatrial node myocytes. (a) The cardiac conduction system visualized in a heart from a *contactin2-eGFP* reporter mouse. The pacemaker depolarization initiated by the sinus node propagates through the atria to the atrioventricular (AV) node and then on to the ventricles via the His-Purkinje network. Adapted from Reference 152 under the terms of the Creative Commons Attribution (CC BY) License, <http://creativecommons.org/licenses/by/4.0>. (b) Sinoatrial node myocyte (SAM) morphology. SAMs are long, thin cells with few striations. Shown is a mouse SAM in the recording chamber of an electrophysiology rig. Note the patch pipette entering from the right that contacts the cell near its center. Adapted from Reference 133 under the terms of the Creative Commons Attribution-ShareAlike (CC BY-SA) License, <https://creativecommons.org/licenses/by-sa/4.0>. (c) Rich sympathetic innervation of SAMs. Co-immunolabeling of SAMs and sympathetic nerve fibers with antibodies against HCN4 (*red*) and tyrosine hydroxylase (*green*), respectively. Image on right shows enlargement of boxed area from image on left. E. Larson and C. Proenza, unpublished data.




b

| Abbreviation      | Parameter                         | Unit  | Definition  |
|-------------------|-----------------------------------|-------|---|
| DD                | Diastolic duration                | ms    | EDD plus LDD  |
| EDD               | Early DD                          | ms    | Initial phase of DD, including DDR                      |
| LDD               | Late DD                           | ms    | Late phase of DD, from DDR to THR                       |
| APD               | AP duration                       | ms    | Time from THR to MDP                                    |
| APD <sub>50</sub> | AP duration at 50% repolarization | ms    | Time interval at 50% of maximum repolarization          |
| APD <sub>90</sub> | AP duration at 90% repolarization | ms    | Time interval at 90% of maximum repolarization          |
| MDP               | Maximum diastolic potential       | mV    | Minimum membrane potential                              |
| THR               | Threshold potential               | mV    | Potential of significant increase in membrane potential |
| PP                | Peak potential                    | mV    | Maximum potential of AP                                 |
| APA               | AP amplitude                      | mV    | Potential difference of PP minus MDP                    |
| DDR               | Diastolic depolarization rate     | mV/ms | Potential change rate at end of EDD                     |
| MUV               | Maximum upstroke velocity         | mV/ms | Maximum of derivative between MDP and PP                |
| MRR               | Maximum repolarization rate       | mV/ms | Minimum of derivative between PP and MDP                |

**Figure 3.**

Characteristic waveform of sinoatrial node action potentials. (a) Schematic illustration of a sinoatrial action potential (*top*) and its first derivative (*bottom*) indicating key features of the action potential. (b) Summary of action potential waveform parameter descriptions. Adapted from Reference 50. Copyright 2017, Elsevier.



**Figure 4.**

Mechanisms of the age-dependent decline in maximum heart rate (mHR). Shown are known (*black boxes and arrows*, color-coded to corresponding subpanels) and hypothesized (*gray boxes and arrows*) mechanisms associated with reduced pacemaker function in aging as described in this review. mHR is decreased primarily due to a reduction in intrinsic heart rate (iHR), as well as a reduction in the response to  $\beta$ -adrenergic receptor ( $\beta$ AR) stimulation. (a) iHR is decreased in part as a result of reduced action potential (AP) firing rate in sinoatrial myocytes (SAMs) from aged mice (*red*) compared to young mice (*black*). (b) The decreased SAM AP firing rate is caused in turn by a reduced diastolic depolarization rate (DDR), a hyperpolarized maximum diastolic potential (MDP), and an increased AP duration. The decrease in iHR is also accompanied by a reduction in the number of SAMs and fibrosis and structural remodeling of the sinoatrial node (SAN). Panels *a* and *b* adapted from Reference 13. (c) Increased diffuse fibrosis in a SAN section from an aged human heart (*right*), as

indicated by increased blue-green staining compared to a section from a young heart (*left*). Adapted from Reference 153 under the terms of the Creative Commons Attribution (CC BY) License, <http://creativecommons.org/licenses/by/4.0>.  $I_{Ca,L}$ ,  $I_{Ca,T}$ , and  $I_f$  currents have been demonstrated to be reduced in aged mouse SAMs. (*d*) Hyperpolarized voltage dependence of activation of  $I_f$  in excised inside-out membrane patches from SAMs from young (*black*) or old (*red*) mice. Adapted from Reference 66 under the terms of the Creative Commons Attribution-ShareAlike (CC BY-SA) License, <https://creativecommons.org/licenses/by-sa/4.0>. (*e*) Reduced  $I_{Ca,L}$  current density in SAMs from aged mice. Adapted from Reference 13. We speculate that  $I_{Na}$ ,  $I_{NCX}$ , and  $I_K$  may also be decreased with age (*gray boxes*). Reduced expression of NCX1, RYR2, SERCA2, and Cx43 transcripts and protein has been observed in a variety of species. (*f*) Reduced relative mRNA abundance of RYR2 and SERCA2 in SAN of old (*red*) compared to young (*black*) rats, normalized to expression in atrial muscle. Adapted with permission from Reference 48. Copyright 2011, John Wiley & Sons. We believe that aging may also reduce expression of many other channels and signaling molecules (*gray boxes*). Asterisks in panels *e* and *f* indicate significant age-dependent differences ( $p < 0.05$ ).