


ARTICLE

Open Access

Gynoecy instability in cucumber (*Cucumis sativus* L.) is due to unequal crossover at the copy number variation-dependent *Femaleness* (*F*) locus

Zheng Li^{1,2}, Yonghua Han^{1,3}, Huanhuan Niu², Yuhui Wang¹, Biao Jiang^{1,4} and Yiqun Weng^{1,5} 

Abstract

Cucumber, *Cucumis sativus* is an important vegetable crop, and gynoecy has played a critical role in yield increase of hybrid cucumber production. Cucumber has a unique genetic system for gynoecious sex expression, which is determined by the copy number variation (CNV)-based, dominant, and dosage-dependent *femaleness* (*F*) locus. However, this gynoecy expression system seems unstable since monoecious plants could often be found in *F*-dependent gynoecious cucumber inbreds. We hypothesized that gynoecy instability (gynoecy loss) may be due to unequal crossing over (UCO) during meiosis among repeat units of the CNV. In this study, using high throughput genome resequencing, fiber-FISH and genomic qPCR analyses, we first confirmed and refined the structure of the *F* locus, which was a CNV of a 30.2-kb tandem repeat. Gynoecious plants contained three genes: *CsACS1*, *CsACS1G*, and *CsMYB*, of which *CsACS1G* is a duplication of *CsACS1* but with a recombinant distal promoter that may contribute to gynoecy sex expression. In two large populations from self-pollinated gynoecious inbred lines, 'gynoecy loss' mutants were identified with similar mutation rates (~0.12%). We show that these monoecious mutants have lost *CsACS1G*. In addition, we identified gynoecious lines in natural populations that carry two copies of *CSACS1G*. We proposed a model to explain gynoecy instability in *F*-dependent cucumbers, which is caused by UCO among *CSACS1G* units during meiosis. The findings present a convincing case that the phenotypic variation of an economically important trait is associated with the dynamic changes of copy numbers at the *F* locus. This work also has important implications in cucumber breeding.

Introduction

The common ancestor of all angiosperms was likely hermaphroditic with perfect flowers that have both stamens and pistils¹. During the evolution of flowering plants, sex determination is an important mechanism to increase genetic fitness by promoting outcrossing, and decreasing inbreeding². One such strategy taken by species in the Cucurbitaceae family is monoecy in which plants produce unisexual flowers on the same individual. Among the ~950 species in this family, ~50% are

monoecious^{3,4} including the economically important vegetable crop cucumber, *Cucumis sativus* L. that has long been a favorite model for study of sex determination in plants.

Three types of flowers can be present in a cucumber plant: staminate (male), pistillate (female) and hermaphrodite (bisexual/perfect). By default, all floral buds contain staminate and pistillate primordia at early stages of development; selective arrest of either staminate or pistillate flower development results in female or male flowers, respectively, and no abortion of either staminate or pistillate primordia allows development of hermaphroditic flowers^{5–9}. Extensive studies have been conducted on the genetic and physiological bases on sex determination in cucumber. Early studies have established three

Correspondence: Yiqun Weng (yiqun.weng@wisc.edu)

¹Horticulture Department, University of Wisconsin, Madison, WI 53706, USA

²College of Horticulture, Northwest A&F University, Yangling, Shaanxi 712100, China

Full list of author information is available at the end of the article.

© The Author(s) 2020



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

genes, *F* (femaleness), *m* (andromonoecy), and *a* (androecy), and their interplays in sex determination in cucumber^{5,6,10–17}. Thus, a cucumber plant may be monoecious (*MMffAA*, with both male and female flowers), gynoeious (*MMFFAA* or *MMFFaa*, with only female flowers), subgynoeious (*MMFfAA* or *MMFfaa*, with few male flowers in the beginning nodes and all female flowers later), andromonoecious (*mmffAA* with bisexual flowers and male flowers), hermaphroditic (*mmFFAA*, or *mmFFaa*, with only perfect flowers), or androecious (*MMffaa* or *mmffaa* with only male flowers). Sex expression in cucumber could also be modified by other genes or quantitative trait loci (QTL), as well as environmental factors such as temperature, and photoperiod^{6,14–16,18–22}.

The wild (*C. s. var. hardwickii*), semi-wild (*C. s. var. xishuangbannensis*) cucumbers and most landraces of cultivated cucumber (*C. s. var. sativus*) are monoecious. However, from the cucumber breeding perspective, gynoeious sex expression has the obvious advantage for the increase of fruit yield. Tkachenko¹² was probably the first to study the inheritance of gynoeious sex expression in cucumber who reported that ‘femaleness’ in a Japanese variety was governed by a pair of genes with ‘femaleness’ being dominant to ‘maleness’. The monogenic segregation was confirmed by Galun⁶ and Shifriss¹³ who designated the genetic factor as *Acr* for accelerating the differentiation of pistillate flowers in gynoeious lines, which was later renamed as the *F* (*femaleness*) locus. The dominant allele (*F*) functions to increase the percentages of female flowers in a dosage-dependent manner^{13,23}.

In 1950s, the identification of gynoeious lines from Japanese and Korean materials (for example, PI 220860), and discovery of methods for sex conversion^{24–26} expedited the incorporation of gynoeicy into commercial F_1 hybrid cucumber production^{27,28}, and understanding of the genetic and physiological basis of gynoeious sex expression in cucumber. Before the cloning of the *F* locus, it has been well established that the phytohormone ethylene is the major regulator of cucumber sex determination^{29–32}. There is a high level of correlation between *F* locus-dependent femaleness and elevated endogenous levels of ethylene; inhibition of ethylene biosynthesis increases the staminate tendency^{30,33}. Indeed, it was found that all three main sex determination genes, *M*, *F* and *A*, are members of the *aminocyclopropane-1-carboxylic acid synthase* (*ACS*) gene families (*CsACSIG* for *F*; *CsACS2* for *M*, and *CsACS11* for *A*) catalyzing the rate-limiting step in ethylene biosynthesis^{18,34–37}. Trebish et al.³⁴ were the first to link *CsACS1* with the *F* locus: they found that while monoecious (*ff*) cucumbers possess a single *CsACS1* gene, gynoeious ones (*FF*) have an additional copy of *CsACS1*, *CsACSIG*, which co-segregates with the *F* locus; *CsACS1* and *CsACSIG* have different

distal promoter sequences that may be responsible for the exclusive expression of *CsACSIG* in *FF* (gynoeious) and *Ff* (subgynoeious) plants³⁵. Knopf and Trebitsh³⁸ further found that the promoter region of *CsACSIG* was the result of a recombination between *CsACS1* and a *branched-chain amino acid transaminase* (*BCAT*) gene (exons 8, 9, and 10). Recently, during a genome-wide survey of the cucumber genome for structural variation, it was found that the *F* locus was due to copy number variation (CNV) of a 30.2-kb region with monoecious and gynoeious lines carrying one and two copies, respectively³⁹. However, a detailed description of the CNV and the structure of the *F* locus is still lacking.

CNV refers to the rearrangements of DNA segments which typically are larger than 1 kb, resulting in the loss or gain of these genomic sequences⁴⁰. CNVs are widely distributed in the genomes of many organisms and have been extensively studied in the human genome due to their association with numerous diseases^{40–43}. CNV is also ubiquitously present in plant genomes and associated with many traits of biological and agricultural importance, such as flowering time in *Arabidopsis* and wheat^{44–46}, *Rhg1*-conferred cyst nematode resistance in soybean (*SCN*)^{47–50}, grain size in rice⁵¹, aluminum tolerance in maize or boron tolerance in barley^{52,53}, and herbicide resistance in the weed palmer amaranth (*Amaranthus palmeri*)^{54,55}.

CNV is a key contributor to genetic variation. Tandem gene clusters of multigene families in particular are rearrangement hotspots and a major source of novel gene formation^{56–58}. CNVs seem to form at a faster rate than other types of mutation, and abiotic stresses may increase the speed CNV⁴⁰. For example, in *Arabidopsis*, CNVs could be observed among individuals separated by only one generation; numerous CNVs affecting hundreds of genes had already originated after only five generations^{59–61}. In cucumber, Osipowski et al.⁶² reported 626 CNVs among three individual plants derived from a common ancestor after 21 or 22 generations of continuous self-pollination. CNVs could be generated through different genetic mechanisms including nonallelic homologous recombination (NAHR) or unequal crossing over (UCO), which results from aberrant homology recognition during homology-based DNA repair or meiosis^{61,63}.

In cucumber, gynoeicy instability is a common phenomenon in which monoecious plants are found at low frequencies in highly inbred, *F* locus-dependent gynoeious populations. Varying numbers of *CsACSIG* copies could also be found in the same gynoeious inbred line³⁹ (see below). We hypothesize that the monoecious lines found in gynoeious progeny (‘gynoeicy loss’) may be caused by loss of the *CsACSIG* copy at the *F* locus due to UCO during meiosis. Thus, the objectives of this study were triple folds. (1) Clarify the CNV structure of the

F locus governing gynoecious sex expression in cucumber; (2) Examine the scope of CNV at the *F* locus in natural cucumber populations; and (3) Investigate the association of ‘gynoecy loss’/instability with UCO at the *F* locus. We first verified the CNV associated with the *F* locus with fiber-FISH, genome re-sequencing, and quantitative real-time PCR (qPCR) using genomic DNA as templates. In large populations of two gynoecious inbred lines (Gy14 and G06), we identified monocious ‘gynoecy loss’ mutants and confirmed the gynoecy instability was due to elimination of the *CsACSI*G copy through UCO during meiosis. The flow chart and reasoning of our work is summarized in Supplemental file 1 (Fig. S1)

Results

F locus-dependent gynoecy in Gy14 cucumber is associated with a 30-kb tandem repeat

The gynoecious Gy14 is a US pickling type inbred line while the monocious 9930 is a North China type (Chinese Long). Draft genomes of both lines have been developed. Previous studies found that, as compared with monocious cucumbers, gynoecious ones possess an extra copy of *CsACSI* with a recombinant distal promoter (*CsACSI*G), and the 30.2 kb sequences were duplicated which constitutes the *F* locus^{38,39}. We manually annotated this 30.2 kb region in the 9930v2.0 assembly and predicted three genes: *CsACSI*, *CsMYB*, and *CsBACT* (Fig. 1a). Then, using 50 kb of this region in 9930v2.0 (30 kb plus 10 kb up-stream and down-stream sequences) as the reference, we mapped Illumina re-sequencing reads from six cucumber lines including three gynoecious cucumbers Gy14, G421, and WI2757 that are known carrying the *F* locus, and three monocious ones, PI 183967 (wild cucumber, *C. s. var. hardwickii*), WI7167 (semi-wild cucumber, *C. s. var. xishuangbannensis*), and PI 190788 (a landrace from India). Each genome was sequenced at 15–30 × depth of coverage. As compared with the flanking 10 kb regions, the average number of reads was nearly twice as many in the 30-kb region in all three gynoecious lines (Fig. 1B1–B3), but no such difference was observed in the three monocious lines (Fig. 1B4–B6) supporting early notion that the 30.2-kb region was duplicated in gynoecious cucumbers.

To examine the orientation of this duplication, we conducted fiber FISH using single copy DNA sequences from ~40 kb region of 9930 V2.0 as probes. Among 19 probes designed [one probe every 2 kb; Fig. 2a; Supplementary File 2 (Table S1)], two (FISH-2 and FISH-3) were located in the *CsACSI* region, and the rest were in up-stream (one probe) or down-stream (16 probes) of *CsACSI*. The two *CsACSI* probes were pooled and labeled with the green dye, and the remaining 17 probes were pooled and labeled with the red dye in FISH of both 9930 and Gy14 genomic DNA fibers. The distribution of


FISH signals on the DNA fibers at the *F*-locus region is shown in Fig. 2a. As expected, in 9930, the relative positions of red and green signals were consistent with its genome structure, while in the gynoecious Gy14, the physical length of labeled DNA fiber was twice that of 9930, and the green and red signal patterns suggested a tandem duplication of the 30.2 kb region in Gy14.

Genomic structure of *F* locus in Gy14 and 9930 cucumbers

Previous studies have shown that in gynoecious cucumber plants, in addition to the non-specific *CsACSI*, the additional copy of this gene, *CsACSI*G, had a new distal promoter region (upstream from –410 bp) containing genomic DNA sequences of exons 9 and 10 of the *CsBCAT* gene due to recombination between *CsACSI* and *CsBCAT*^{34,35,38}. We annotated this region in Gy14 V1.0 (~60 kb), and confirmed the existence of *CsACSI*G (*BCAT* exons 9 + 10 + *CsACSI*) and *CsMYB* in the tandem repeat. Thus, in Gy14, the first copy of the 30.2 kb region was exactly the same as in 9930 (*CsACSI* + *CsMYB*), and the second copy was *CsACSI*G (*BCAT* exons 9 + 10 + *CsACSI* + *CsMYB*), which was followed by a complete *CsBACT* (Fig. 2b).

When this project was initiated, only Gy14 V1.0 and 9930 V2.0 assemblies were available, both of which had some gaps or N's in this region. Thus, it was not clear if additional sequences may present in the CNV region of Gy14. We employed anchor PCR to estimate the size in target regions of the CNV in Gy14. Primers were designed at three regions for PCR including the 5' region (upstream of *CsACSI*), the ‘junction point’ of *BACT* exons 9–10 and *CsACSI* (upstream of *CsACSI*G), and the 3' region (*CsBCAT* gene) (Fig. 2c). At each location, one primer was anchored, and other primers were designed at locations such that the PCR amplicons were increased in size progressively (Table S1). The results indicated that, in each case, the amplicon sizes were consistent with expected sizes estimated from genomic sequences in Gy14 V1.0 and 9930 V2.0 (Fig. 2c) suggesting no additional large sequences are present in the CNV region in the Gy14 assembly.

We further verified the copy numbers of individual genes in the 30.2 kb region with real-time qPCR using genomic DNA as the templates. Four pairs of gene-specific primers were designated from sequences targeting the *CsACSI*, and *CsMYB* genes, exons 9 and 10 (*BACT*9–10), as well as exons 1–8 (*BACT*1–8) of the *BCAT* gene (Fig. 3a; Table S1). The effectiveness of all primers was first confirmed with consistent electrophoretic bands from regular PCR. The qPCR with genomic DNA was performed in four representative cucumber lines: the gynoecious Gy14, and WI2757, and the monocious 9930, and WI7167. In qPCR, the level of *BCAT* exons 1–8 that are not involved in the CNV at the *F* locus



was used as the standard to normalize the level of other three genes. The results are presented in Fig. 3b. Indeed, as compared with the adjacent genomic region, there were two copies each of the *CsACS1*, *CsMYB*, and *BCAT9–10* in the 30.2 kb region of the two gynococious lines (Gy14 and WI2757), and one copy each in 9930 and WI7167.

Previous studies^{38,39} and the results reported herein all supported ~30.2 kb duplication at the *F* locus in gynococious cucumbers that contains two copies of *CsACS1* and *BCAT* exons 9 and 10. However, the exact sequences and annotation of this region have never been explicitly reported. By the time we were preparing the manuscript, the improved versions of the 9930 V3.0⁶⁴ and Gy14 V2.0 (our

unpublished data) draft genomes were just released to the public (available at <https://www.cucurbitgenomics.org/>). No gaps or N's were found in the CNV region in either draft genome, which allowed verification and refinement of the results obtained in the present study. We manually annotated the CNV regions in both Gy14 and 9930, which are graphically presented in Supplementary File 3 (Fig. S2). In 9930 V3.0, the sequences from transcription start of *CsACS1* to 3' end of *CsBCAT* is 35,572 bp. In the Gy14 V2.0 assembly, unexpectedly, two copies of *CsACS1G* (*BACT9–10* + *CSACS1* + *CsMYB*) were annotated with a total length of 94,849 bp (Fig. S2). Since all Gy14 materials used in various studies (genome sequencing for V1.0/




Fig. 2 Structure of *F* locus in gynocercous Gy14 cucumber. **a** Fiber FISH analysis reveals a tandem duplicate of 30-kb region in Gy14 as compared with the monocious 9930 cucumber. The green signals are from probes of *CsACS1* genomic DNA sequence, and the red signals are from pooled 17 probes from other sequences in the 30-kb region (up). The approximate targeting positions of 19 probes are shown as green and red oval spots. Three representative DNA fibers of Gy14 and 9930 are shown (down). Bar = 10 μ m. **b** Genomic structure of the *F* locus in Gy14. The exons 9 and 10 of the *CsBCAT* gene is part of the second copy of *CsACS1*, which are collectively called *CsACS1G*. **c** One-end anchored PCR is used to analyze the continuity of the *F* locus in the 5' end (green), 3' end (blue), and junction point (brown). M = DNA size ladder

V2.0, and fiber FISH) were from the same source (but different generations from self-pollinations), it may hint that the triple tandem repeats of the ~30 kb region in Gy14 V2.0 may be the result of UCO (see discussion below).

We examined DNA nucleotide diversity in the *CsACS1*/*CsACS1G* gene region. Sequence alignment among four sequences (~4.6 kb each) including *CsACS1* from 9930 V3.0 and Gy14 V2.0, *CsACS1G* from Gy14 V2.0, and the NCBI accession DQ839406 (*CsACS1G* from gynocercous cucumber Marketmore 76F)³⁸. The results are illustrated in Supplementary File 4 (Fig. S3). Sequence features for *CsACS1* (TATA box, transcription start/end, and intron/exon junctions) and *CsACS1G* (proximal and distal promoters) are also highlighted. The ~4.6 kb Gy14 and 9930 *CsACS1* genomic DNA sequences were almost identical. In addition, the sequences from -410 bp to 3' UTR in both *CsACS1* and *CsACS1G* were also largely the same except for two SNPs (Fig. S3). These observations were

consistent with previous findings with regard to sequence variation at the *F* locus in gynocercous and monocious lines³⁸. The data also suggested that the duplication of *CsACS1* resulting in the *F* locus was a very recent event, and the *F* locus in the two gynocercous lines Gy14 and Marketmore 76F (DQ839406) have the same origin (see "Discussion" section below).

Knopf and Trebitsh³⁸ suggested the promoter of *CsACS1G* was derived from recombination between *CsBCAT* and *CsACS1* genes. Indeed, significant sequence diversity was found in the distal promoter regions (upstream from -411 to -1700) between *CsACS1* and *CsACS1G* (Fig. S3). The sequences in *CsACS1G* from -411 upstream of transcription initiation are part of the *CsBCAT* gene which included (reverse complementary strand, in order of presence): the 8th intron (928 bp), Exon 9 (113 bp), the 9th intron (350 bp), and Exon 10 (113 bp) with a total of 1504 bp (up to -1914 from transcription start). Interestingly, the whole length of the




Fig. 3 Evaluation of copy numbers of genes at the *F* locus with qPCR of genomic DNAs in cucumber. **a** Targets regions for qPCR. Green, red, blue and black horizontal bars represent genomic region of *CsACS1*, *CsMYB*, exons 9–10, and 1–8 of the *CsBACT* genes, respectively. Copy numbers of different regions revealed by qPCR among four representative gynoecious and monocious lines, as well as 30 additional cucumber varieties are shown in **b** and **c**, respectively. Each data point in **c** is the mean of seven replications that was normalized to that for exons 1–8 of the *CsBACT* gene. In **c**, asterisk indicates the gynoecious line AM297 (G14) that contains three copies of the 30-kb region at *F* locus. In each line name, H = hermaphroditic, G = gynoecious, M = monocious. Asterisk indicates the gynoecious line AM297, of which CNVs value is 3


8th intron is 984 bp, and 928 bp was present in the distal promoter with the last 56 bp missing.

CNV at *F* locus is associated with gynoecy expression in natural cucumber populations

Next we sought to investigate the association of CNV at the *F* locus with sex expression in natural cucumber populations. Previous studies^{52,53} and our work described above (Fig. 3b) showed that qPCR could be used to estimate copy numbers of genomic DNA sequences. We employed this method to examine the number of copies of the four sequences at the *F* locus (Fig. 3a) among 30 lines of different sex morphs including 5, 14, 8, and 3 of hermaphroditic (*FF*), gynoecious (*FF*), monocious (*ff*), and subgynoecious (SubG, *Ff*) cucumber lines, respectively (Supplementary File 5 or Table S2). The three SubG lines were F_1 from gynoecious \times monocious crosses that were

heterozygous at the *F* locus. The qPCR results among the 30 samples are illustrated in Fig. 3c. All monocious lines had one copy each of the three sequences (*CsACS1*, *CsMYB*, and *BACT9–10*), and all hermaphroditic and gynoecious lines except AM297 (G14 in Fig. 3c) had two, and the three SubG F_1 hybrids had 1.5 copies. These results were consistent with two copies of the 30-kb unit in gynoecious plants containing the *F* locus.

Among the 14 gynoecious varieties examined, AM297 was an interesting exception that had three copies of (*CsACS1* + *CsMYB* + *BACT9–10*) sequences. To clarify the structure of the *F* locus in AM297, we conducted genomic DNA qPCR among Gy14, 9930 and AM297 using primers from *BCAT9–10*, *BCAT1–8*, and a new primer pair, F-gDNA, which targeted the ‘junction point’ in the unique promoter region of *CsACS1G* (Fig. 3a). The left and right primers of F-gDNA were near the 10th exon



of *BCAT*, and the start of *CsACS1*, respectively. Because of the unique design of the primer pair, there would be no amplicon in the monocious 9930 cucumber due to opposite directions of the two primers while the expected amplicon size would be 111 bp in Gy14 (Table S1). The qPCR results (Fig. 4a) suggested that AM297 carries triple tandem repeats of the ~30 kb region with two copies of *CsACS1G* (Fig. 4b). The structures of the *F* locus in the three varieties are summarized in Fig. 4c, in which AM297 was predicted to have the same triple repeat as in Gy14 V2.0 (Fig. S2).

All the gynocious lines carried at least one copy of *CsACS1G* suggesting they may have a single origin. Indeed, all the 14 gynocious lines we used (Table S2) belong to US pickling or slicing market groups or the mini (Beit Alpha) cucumbers with rather recent breeding histories. To confirm this, we examined sequence diversity in the 50-kb genomic region around *CsACS1* region among 48 re-sequenced cucumber varieties including 10 gynocious lines, which are listed in Supplementary File 6 (Table S3). From the re-sequencing data, 68 SNPs and 3 InDels were detected (Table S3), and were used to construct a phylogenetic tree (Supplementary File 7/ Fig. S4). All 10 gynocious varieties were in the same clade, and the grouping of monocious lines was based largely on their taxonomic status or geographic origins. This

observation supported the common origin of the *F* gene in all gynocious lines.

Gynocy instability may be due to loss of *CsACS1G* through UCO among tandem repeats of the CNV at the *F* locus

In cucumber, monocious plants could be found at low frequency among self-pollinated progeny of highly inbred gynocious lines ('gynocy loss'). Varying copies of *CsACS1G* may present in gynocious lines (Fig. 3c). We hypothesize that 'gynocy loss' may be the result of UCO among the 30-kb repeat units at the *F* locus. To prove this, in 2015 field season, we grew 2236 gynocious Gy14 plants in an open field. All these plants were derived from a single Gy14 plant through two generations of self-pollinations. Throughout the growing season, plants with at least one male flower were tagged. By end of the season, 66 such plants were identified, but only three exhibited consistent subgynocious sex expression, which were designated as No. 17, 42, and 63, respectively (Fig. 5a). Thus, the mutation ('gynocy loss') rate was 0.13% in Gy14 (3/2236). In 2017, we conducted a similar experiment using another gynocious line G06 that belongs to the Mediterranean (Beit alpha) type cucumber. Among 2573 gynocious G06 plants grown in greenhouse conditions, three subgynocious plants (G06M-B, G06M-C, and G06M-E) were identified. The mutant frequency of




Table 1 Segregation of sex expression in cucumber lines or populations

Lines/populations	# <i>CsACS1G</i> copies	Expected sex expression ^a	# Plants observed	# G	# of SubG	# of M	<i>P</i> for χ^2 test
Gy14	2	G	2236	2170	0	3 ^b	–
G06	2	G	2573	2537	0	3 ^b	–
9930	0	M	20	0	0	20	–
S52	0	M	20	0	0	20	–
XTMC	0	M	20	0	0	20	–
AM297	4	G	20	20	0	0	–
(Gy14 × 9930) F1	1	SubG	16	1	15	0	–
(Gy14 × XTMC) F1	1	SubG	16	2	14	0	–
(Gy14 × S52) F1	1	SubG	16	0	16	0	–
(Gy14 × 9930) F2	0, 1, or 2	Segregating, 1G: 2 SubG: 1M	200	47	98	55	0.6976
(AM297 × 9930) F1	2	G	16	16	0	0	–
(AM297 × XTMC) F1	2	G	16	16	0	0	–
(AM297 × S52) F1	2	G	16	16	0	0	–
(AM297 × 9930) F2	0, 1, or 2	Segregating, 3G: 1M	1000	735	0	265	0.2733
(Gy14M-17 × 9930) F1	0	M	14	0	0	9	–

^aG gynoeious, SubG Subgynoeious, M Monoeious

^b'Gynoeicy loss' monoeious mutants

'gynoeicy loss' in G06 was 0.11%, which was very close to 0.13% observed in Gy14 suggesting gynoeicy instability is a common phenomenon in *F*-dependent gynoeious cucumbers regardless of the genetic backgrounds.

The three subgynoeious mutant plants from Gy14 were self-pollinated, and the progeny were observed for sex expression. The segregating sex phenotypes (gynoeious, subgynoeious, and monoeious) in their progeny populations suggested the three lines were true 'gynoeicy loss' mutations. We further self-pollinated the monoeious individuals, and the offspring were all monoeious in the following six generations of self-pollinations, which were designated as Gy14M-17, Gy14M-42, and Gy14M-63, respectively (Fig. 5b), indicating these 'gynoeicy loss' mutations were stable. We re-sequenced the three mutant lines with >15× coverage, and the reads were aligned against the 30-kb *F* locus region of the 9930 V2.0 draft genome. We found the same depth of coverage of reads in the 30-kb region as the flanking regions (Supplementary File 8/ Fig. S5B2-B4) suggesting the extra copy of *CsACS1* in Gy14 (i.e., *CsACS1G*; Fig. S5B1) have lost in the three monoeious mutant lines. We also conducted qPCR using primers in the *F* locus to examine copy numbers of target genes at the *F* locus in the three 'gynoeicy loss' lines, which further supported the loss of *CsACS1G* in these mutants (Supplementary File 9/ Fig. S6).

UCO among repeat units at the *F* locus occurs in gynoeious (G) plants would generate two types of gametes: gametes with duplication (1 *CsACS1* + 2

CsACS1G alleles) and gametes with deletion (1 *CsACS1*). The three 'gynoeicy loss' mutants (Gy14M-17, –42 and –63) might represent the products from deletion of *CsACS1G*. While we did not try to identify products from duplication events because the large amount of work to screen gynoeious plants. Alternatively, AM297 with two copies of *CsACS1G* might represent such product from duplication of the 30-kb repeat of UCO. Thus, we investigated the inheritance of sex expression by examining lines and derived segregating populations which carry different copies of *CsACS1G* (Table 1). It is known that the *F* gene in cucumber has a dosage effect, and the F₁ from gynoeious (*FF*) × monoeious (*ff*) cross is subgynoeious (*Ff*, SubG). Indeed, of 16 Gy14 × 9930 F₁ plants observed, 15 were SubG (Table 1; Supplementary File 10/ Fig. S7A, B). Similar results were obtained in F₁ of Gy14 × XTMC (M), and Gy14 × S52 (M) crosses. In contrast, all 16 F₁ plants from AM297 × 9930 cross were gynoeious. In addition, all the Gy14M-17/–42/–63 × 9930 F₁ plants were monoeious (Table 1; Fig. S7C, D). In the Gy14 × 9930 F₂ population, of the 220 plants, the segregation of G:SubG:M plants was consistent with the expected 1:2:1 ratio. But among over 1000 AM297 × 9930 F₂ plants, no SubG plants were present, and the ratio of G to M plants exhibited roughly 3:1 segregation suggesting at least two copies of the *F* gene in the AM297 haploid genome (Table 1). The no show of SubG plants in the large population also indicated that the two copies of the *F* gene were co-segregating (no recombination), which

was consistent with the model structure of tandem duplication of the three 30-kb regions in AM297 predicted in Fig. 4c. These data support the hypothesis that AM297 carries two copies of *CsACSI*G, which might be derived from UCO.

From previous studies, it is well known that a high correlation exists between ethylene accumulation and female sex expression (see section “Introduction”). We examined ethylene releases in wild type gynoeious Gy14 and the ‘gynoecy loss’ mutants, and found significant decrease of ethylene in the three monoeious mutant lines (Fig. 5c). We further examined the expression levels of *CsACSI*, *CsACSI*2 (*andromonoecious* or *m* locus), and *CsMYB* at the *F* locus with mRNA qPCR (Fig. 5d). Among all three mutant lines, as compared with Gy14, the expression of both *CsACSI* and *CsACSI*2 was significantly down regulated, whereas no significant difference in expression was found for *CsMYB* among the four lines. In our previous study, transcriptome profiling for sex expression revealed no expression difference of the *CsBCAT* gene among the three mutant lines and Gy14⁶⁵. These data suggest that expression level of both *CsMYB* and *CsBCAT* genes at the *F* locus may have no direct link with gynoecy expression in cucumber.

Discussion

Structure and function of the *F* locus in cucumber

Cucumber has a unique system for gynoeious sex expression which is based on CNV of a 30.2 kb region containing *CsACSI*G, a duplicated copy of *CsACSI* but with recombinant promoter^{38,39}. In melon and watermelon, the gynoeious sex expression is controlled by *WIP1* homologs encoding a C2H2 zinc-finger transcription factor, which is a negative regulator of femaleness^{66,67}. No gynoeious *WIP1* mutant (*CsWIP1*) have been identified in natural cucumber populations. *CsWIP1* may perform a similar function in cucumber because mutations obtained by CRISPR/Cas9 (clustered regularly interspaced short palindromic repeats/CRISPR-associated protein 9) assay in this gene significantly increase the degree of femaleness⁶⁸.

In the present study, data from fiber-FISH, analysis of resequencing reads, as well as qPCR (Figs. 1–3) all confirmed that the *F* locus for gynoecy in cucumber is a duplication of 30.2 kb region. These results were further validated with analysis of the newest 9930 (V3.0) and Gy14 (V2.0) draft genome assemblies which also, for the first time, presented a more complete fine structure of the *F* locus (Figs. S2 and S3). From previous studies^{35,38,39,69–71} and work reported herein, the following conclusions could be made. (1) The *F* locus is a duplication of 30.2 kb repeat unit. In monoeious lines (*ff*), this region contains three genes: *CsACSI*, *CsMYB*, and *CsBCAT*; in gynoeious lines (*FF*), the structure of the *F* locus is

CsACSI–*CsMYB*–*CsACSI*G–*CsMYB*–*CsBACT* (Fig. 2b). (2) There is no sequence difference between *CsACSI* and *CsACSI*G in the proximal region (+1 to –410 bp) and CDS in the gynoeious plant tested. (3) The 5'-distal promoter (upstream from –410 up to –1900) of *CsACSI*G is composed of sequences from intronic (8th and 9th) and exonic (exons 9 and 10) sequences of the *CsBCAT* gene (Fig. S3), which is very different from that in *CsACSI* supporting its origin from recombination between the *CsACSI* and *CsBCAT* genes³⁸.

The *CsACSI*G sequences from Gy14 (a US pickling cucumber) and Marketmore 76F (A US slicing cucumber) are identical (Fig. S3). This is expected because the *F* locus in both gynoeious lines were probably from the same source, MSU713–5, which, in turn, was introduced from the Korean cucumber line ‘Shogoin’ (PI 220860)²⁵. In fact, most modern gynoeious cucumbers in commercial production may share the same source of *F* locus-dependent gynoecy from MSU713–5^{28,72,73}. This can explain the clustering of all gynoeious lines in one clade (Fig. S4) which would otherwise be clustered based on their geographic origins if genome-wide molecular markers were used in phylogenetic analysis^{74,75}.

The effects of the duplication at the *F* locus on the expression of *CsACSI* and *CsACSI*G in gynoeious and monoeious cucumbers have been investigated in several studies. First, the recombination between *CsBACT* and *CsACSI* resulting in the distal promoter of *CsACSI*G, which introduced a putative open-reading frame (ORF) (from position 528 of Gy14 *CsACSI*G or position 666 in DQ839406_ACS1G.1, Fig. S2), but Shiber et al.⁷¹ did not detect any transcripts from this putative ORF thus ruling out the possibility that its product may be the *F* locus. In gynoeious plants (*FF*), the transcript level of *CsACSI*/G is much higher^{38,65}. On the other hand, knock-down of *CsACSI*G expression in gynoeious cucumber resulted in monoeocy⁷¹. We found that, in ‘gynoecy loss’ mutants (loss of *CsACSI*G), both ethylene level and expression of *CsACSI* were significantly reduced as compared with the wild type (Fig. 5d). These observations suggested the higher expression of *CsACSI*/G in gynoeious plants is directly linked with gynoecium development.

Among the three genes involved in the CNV at this *F* locus (Figs. 3b and S2), the expression of *CsMYB* and *CsBACT* did not seem to be affected by the duplication event in the generation of the *F* locus. No significant differences in expression of the two genes were identified between monoeious and gynoeious cucumbers⁶⁵ (also Fig. 5d). This is in contrast with the *RhgI* locus of soybean. *RhgI*-conferred cyst nematode resistance in soybean^{47–50} in which the SCN resistance level is positively correlated with copy numbers of a 31.2 kb repeat unit. The simultaneous upregulated expression of all three genes in the repeat unit is required for the resistance⁴⁷. It

is interesting to see why *CsMYB* expression remains unchanged in gynoecious cucumbers.

Gynoecy instability and CNV dynamics in gynoecious cucumbers

In cucumber, the *F* locus-dependent gynoecy is relatively stable. However, it is common that monocious plants are found at low frequency in the self-pollinated progeny of gynoecious inbred lines which we call ‘gynoecy loss’. Indeed, in two gynoecious lines, the US pickling cucumber Gy14 (Fig. 5) and the mini cucumber inbred G06, we found 0.13% and 0.11% plants that were monocious ‘gynoecy loss’ mutants. We show that these mutant lines have lost the *CsACSI*G copy at the *F* locus (Table 1, Fig. S5). Since the *F* locus contains two copies of the 30.2 kb repeat, these data suggested that gynoecy expression is not stable due probably to the dynamic change of copy numbers of the 30-kb unit.

Such dynamic change of copy numbers is common at CNV loci in plant and animal genomes, which is most likely caused by NAHR or UCO^{63,76}. UCO depends on misalignment of adjacent homologous sequences. The result of UCO is deletion or duplication/triplication of a genomic sequence, which is likely to form a CNV locus. A CNV locus with tandem repeat structure, or located in adjacent position in the same chromosome, may have the potential to take place further UCO^{57,76}. One good example for phenotype-associated CNV instability due to UCO is the *Rhg1* locus for SCN resistance in soybean. The level of *Rhg1*-conferred SCN resistance level in soybean is positively correlated with copy numbers of a 31.2 kb repeat unit⁴⁷. While there is an extensive diversity in copy numbers among different resistant lines, significant variation of copy numbers and disease resistance level was also observed within the same variety⁵⁰. It has been shown that copy number at *Rhg1* is unstable within a released variety over a relatively small number of generations due to NAHR or UCO^{47–50}. NAHR has also been shown to be responsible for CNV of different alleles at the *I* (inhibitor) locus in soybean for the colors of seed coat⁷⁷, and many other CNV-based phenotypic variation^{78–80}.

UCO may also explain the gynoecy instability or ‘gynoecy loss’ observed in the present study. For easy description, a model is provided in Fig. 6 to explain UCO events resulting in ‘gynoecy loss’ monocious mutations in Gy14 cucumber. During meiosis, the two 30.2 kb repeat units containing three genes misalign and subsequent UCO (Fig. 6a) would result in gametes with deletion and duplication of *CsACSI*G sequences (Fig. 6b). In self-pollination, the union of these gametes with normal *F* gametes will result in gynoecious (G) and subgynoecious (SubG) plants (Fig. 6c). Self-pollination of SubG individuals will segregate for G, SubG, and monocious (M) plants (Fig. 6d). The M plants were confirmed by the

observation of no sex expression segregation in the following self-pollination (Fig. 5b). The mutation rate for the M plants from UCO is likely low (0.13% in Gy14). We did not genotype G plants carrying two or more *CsACSI*G copies in the field due to the sheer large numbers. The presence of CNV = 3 (two copies of *CsACSI*G) plants could be indirectly evidenced from the identification of AM297 in natural cucumber populations (Fig. 4), and in two copies of *CsACSI*G in Gy14 V2.0 (Supplementary Fig. S2). In fiber FISH, the hybridization patterns in multiple gynoecious lines also indicated possible varying copies # in the *F* locus (data not shown).


It is puzzling why the CNV has 30.2 kb as the repeat unit in which two of the three genes (*CsMYB* and *BCAT*) have no difference in expression level before (in monocious) and after (in gynoecious) the duplication. Interestingly, after examining several draft genomes (<https://cucurbitgenomics.org/> and <https://www.ncbi.nlm.nih.gov/>), we found high degree of micro-synteny of the *ACSI*–*MYB*–*BCAT* trio in both gene order and content across multiple cucurbit crops, which also extends to as far as Arabidopsis. It is not known if the triplet genes are required for gamete fertility or any other fitness advantages during evolution.

Materials and methods

Identification and phenotypic characterization of ‘gynoecy loss’ mutants

Two gynoecious cucumber inbred lines Gy14 and G06 (genotype *FF* at the *F* locus) were used to test our hypothesis on ‘gynoecy loss’, which belong to the North American pickling and Mediterranean beta alpha (mini) market group, respectively. Both lines have undergone at least six generations of self-pollination. In 2015 field season, 2500 Gy14 plants were grown in the field plot at the University of Wisconsin Madison Hancock Agriculture Research Station (HARS, Hancock, WI, USA). In 2017 spring, a similar work was conducted with G06, in which sex expression was examined among over 3000 plants in greenhouses at the Jingyang Agriculture Research Station (Xianyang, Shaanxi, China). At reproductive stage, each plant in both experiments was examined for sex expression. All plants with one or more male flowers were tagged. Subgynoecious plants were self-pollinated. Multiple plants from the selfed progeny were planted in the glasshouses for observation of sex expression. Selected mutants were self-pollinated to observe segregation among the progeny.

The gynoecious cucumber inbred line AM297 (aka, ‘Telegraph Improved’) was found to contain three copies of *CsACSI*/G gene the *F* locus in haploid genome. Both Gy14 and AM297 were crossed with three monocious lines 9930, XinTaiMiCi (North China type), and S52 (South China type). In addition, the three ‘gynoecy loss’



(monocious) mutant lines (Gy14M-17/-42/-63) were crossed with 9930. Sex expression of these F_1 plants were recorded in the greenhouses. In addition, F_2 plants were produced from AM297 \times 9930 F_1 and Gy14 \times 9930 F_1 for observation of segregation of sex expression.

Selected cucumber varieties (total 37, Supplementary Table S2) were also grown at HARS in multiple seasons for observation of sex expression according to Li et al.⁸¹. Leave samples of these lines were collected for analysis of copy numbers for genes at the *F* locus.

Copy number estimation of F locus with fiber fluorescence in situ hybridization (FISH) and resequencing reads alignment

For visual access of the copy numbers at the *F* locus, fiber FISH was conducted with Gy14 and 9930 inbred lines. Based on the 9930 V2.0 genome assembly (<https://cucurbitgenomics.org/>), 19 primer pairs were designed in the targeting ~40.0 kb region of single copy DNA sequences. The locations of each probe in different versions of Gy14 and 9930 genome assemblies are listed in

Supplementary Table S1. The resulting PCR amplicons (average length ~2 kb) were used as probes.

FISH procedures on genomic DNA fibers were described previously^{82,83}. The probe labeling was performed according to published protocols⁸³. DNA probes were labeled with digoxigenin-dUTP or biotin-dUTP via nick translation and detected with anti-dioxigenin antibody coupled with Rhodamine (Roche) or anti-avidin antibody conjugated with FITC (Vector Laboratories), respectively. Hybridization signals in two-color fiber-FISH were detected with a three-layer antibody detection system as described⁸⁴. Images were captured digitally using a CCD camera, and the final image adjustments were done with Adobe Photoshop v6.0.

We estimated CNV at the *F* locus from resequencing data of nine cucumber lines with different sex morphs including the gynoecious Gy14, G421, and WI2757, monocious lines WI7167, PI 183967, PI 197088, and the three Gy14 ‘gynoecy loss’ mutant lines (Gy14M-17/-42/-63). The genome of each line was re-sequenced with Illumina Hi-Seq 2000 sequencing platform at >15× coverage. Sequencing reads of different mutant lines were aligned against the 40 kb *F* gene region of the 9930 V2.0 draft genome using BWA pipeline following Pan et al.⁸⁵.

Genome structure analysis at *F* locus

The draft genomes of *F* locus region in both 9930 and Gy14 were manually annotated. To verify the structure of *CsACS1G*, we amplified the 5′ and 3′ regions as well as the *CsACS1*–*BCAT* ‘junction point’ with progressive PCR in which one end of the amplicon was anchored and the other end was at different locations. The primer sequences and their positions are listed in Table S1. All amplicons in the junction point were confirmed with sub-cloning and Sanger sequencing.

The structure of the *F* locus by evaluating copy numbers of genes in the CNV region was confirmed with real-time quantitative PCR (qPCR) using genomic DNA as the template. The primers were designed from different regions of selected genes (Table S1). All PCR reactions were conducted in a 96-well plate using the ABI 7500 Fast Real-Time PCR System (Applied Biosystems, CA, USA) with the SYBR green qPCR master mix (Bio-Rad, Hercules, CA, USA). The amplification was initiated by heating to 95 °C for 10 min, followed by 40 cycles at 95 °C for 20 s and 65 °C for 30 s. Each result was derived from seven independent replications.

The genomic DNA sequence diversity of the 50 kb *F* locus was examined in 48 re-sequenced cucumbers lines reported previously^{86,87} and the three ‘gynoecy loss’ mutants. SNPs within the target region were called with the BWA-GATK4.0 workflow using 9930 V2.0 as the reference following Wang et al.⁸⁸. All detected polymorphisms (SNPs or InDels) in the 30-kb CNV sequence

in the ‘gynoecy loss’ mutants were verified by PCR and Sanger sequencing. These polymorphic markers in the 50 kb region were employed to construct a phylogenetic tree among the 48 cucumber lines to understand the origin of the *F* locus. Sequence alignment and clustering were performed with MEGA 7.0 (<http://www.megasoftware.net/>) using the maximum likelihood method with 1000 bootstrap replications.

The new versions of draft genomes for both Gy14 (V2.0) (our unpublished data) and 9930 (V3.0)⁶⁴ were recently released (<https://cucurbitgenomics.org/>). We compared the 30-kb CNV region in 9930 with that of Gy14 to verify results from other experiments in this study.

Expression of sex determination-related genes with mRNA qPCR

We examined expression level of *CsACS1*, *CsMYB* genes in the CNV region as well as the andromonoecy (*m*) gene *CsACS2* in relation to sex expression. The apices of Gy14, 9930 and three ‘gynoecy loss’ mutants were harvested for RNA extraction and first-strand cDNA synthesis⁸⁹ and qPCR using the procedure described above with the cucumber *actin2* as the reference to normalize the expression data (primer info is provided in Table S1). Each sample was run with three biological and at least three technical replicates with appropriate statistical (*t* tests) analysis of the data.

Measurement of ethylene release

To measure the ethylene production rate, shoot apices were excised from seedlings at the fourth true leaf stage. Equal amount of samples of the three mutant lines was mixed as ‘Gy14-M’. There were three biological replicates from both Gy14 and Gy14-M with each replicate containing shoot apices from 15 plants. Each sample was placed in a 12 ml container and sealed with a rubber stopper. After incubation at 25 °C for 14 h in the dark, 1 ml of gas was withdrawn using a gas-tight syringe from the headspace, which was analyzed using the Trace GC Ultra gas chromatography system (Thermo Scientific, USA) that was equipped with a flame-ionization detector and a capillary column for ethylene measurement. The instrument was calibrated with an ethylene gas standard, and the amount of ethylene released from shoot apices per 1 kg fresh weight and per hour was calculated. All determinations were made in triplicate.

Funding

The work in Z.L.’s lab was supported by the National Natural Science Foundation of China (Nos. 31672150 and 31872111), the Fundamental Research Fund for the Central Universities (2452016004), and the Key Research and Development Plan (2018NY-034) of Shaanxi Province. Work in Y.Q.W.’s lab was supported by USDA National Institute of Food and Agriculture under following award numbers 2015-51181-24285 and 2017-67013-26195. USDA is an equal opportunity provider and employer.

Author details

¹Horticulture Department, University of Wisconsin, Madison, WI 53706, USA. ²College of Horticulture, Northwest A&F University, Yangling, Shaanxi 712100, China. ³Institute of Integrative Plant Biology, School of Life Sciences, Jiangsu Normal University, Xuzhou 221116, China. ⁴Vegetable Research Institute, Guangdong Academy of Agricultural Science, Guangzhou, Guangdong 510640, China. ⁵USDA-ARS, Vegetable Crops Research Unit, Madison, WI 53706, USA

Author contributions

Y.Q.W. and Z.L. designed and supervised the experiments. Z.L., H.N., and B.J. performed majority of the laboratory work. Y.H. conducted fiber FISH. Y.H.W. conducted bioinformatics analysis of the CNV region. Z.L. and Y.Q.W. wrote the manuscript with inputs from all authors.

Conflict of interest

The authors declare that they have no conflict of interest.

Supplementary Information accompanies this paper at (<https://doi.org/10.1038/s41438-020-0251-2>).

Received: 28 October 2019 Revised: 6 January 2020 Accepted: 15 January 2020

Published online: 15 March 2020

References

- Cronquist, A. *The Evolution and Classification of Flowering Plants* (New York Botanical Gardens, Bronx, New York, 1988).
- Chuck, G. Molecular mechanisms of sex determination in monoecious and dioecious plants. *Adv. Bot. Res.* **54**, 54–83 (2010).
- Schaefer, H. & Renner, S. Cucurbitaceae. In *Families and Genera of Flowering Plants*, Vol. 10 (ed. Kubitzki, K) 112–174 (Springer Verlag, Berlin, Germany, 2011).
- Renner, S. S. Pathways for making unisexual flowers and unisexual plants: moving beyond the “two mutations linked on one chromosome” model. *Am. J. Bot.* **103**, 587–589 (2016).
- Correns, C. Bestimmung. *Vererbung und Verteilung des Geschlechtes bei den höheren Pflanzen. Handbuch der Vererbungswissenschaft*, 3(II, C). Band II (English: *Determination, Inheritance and Distribution of Sex in Higher Plants. Handbook of Inheritance Science*, 3 (II, C) Vol. II (Bomtraeger, 1928).
- Galun, E. Study of the inheritance of sex expression in the cucumber: the interaction of major genes with modifying genetic and non-genetic factors. *Genetica* **32**, 134–163 (1961).
- Kater, M. M., Franken, J., Carney, K. J., Colombo, L. & Angenent, G. C. Sex determination in the monoecious species cucumber is confined to specific floral whorls. *Plant Cell* **12**, 481–494 (2001).
- Hao, Y. et al. DNA damage in the early primordial anther is closely correlated with stamen arrest in the female flower of cucumber (*Cucumis sativus* L.). *Planta* **217**, 888–895 (2003).
- Bai, S. et al. Developmental analyses reveal early arrests of the spore-bearing parts of reproductive organs in unisexual flowers of cucumber (*Cucumis sativus* L.). *Planta* **220**, 230–240 (2004).
- Rosa, J. T. The inheritance of flower types in *Cucumis* and *Citrullus*. *Hilgardia* **3**, 235–250 (1928).
- Whitaker, T. W. Sex ratio and sex expression in the cultivated cucurbits. *Am. J. Bot.* **18**, 359–366 (1930).
- Tkachenko, N. N. Preliminary results of a genetic investigation of the cucumber, *Cucumis sativus* L. *Bull. Appl. Plant Breed.* **9**, 311–356 (1935).
- Shifriss, O. Sex control in cucumber. *J. Hered.* **52**, 5–12 (1961).
- Kubicki, B. Investigations of sex determination in cucumber (*Cucumis sativus* L.). IV. Multiple alleles of locus. *Genet. Pol.* **10**, 23–68 (1969).
- Kubicki, B. Investigations of sex determination in cucumber (*Cucumis sativus* L.). V. Genes controlling intensity of femaleness. *Genet. Pol.* **10**, 69–86 (1969).
- Kubicki, B. Investigations on sex determination in cucumbers (*Cucumis sativus* L.). VI. Androecism. *Genet. Pol.* **10**, 87–99 (1969).
- Malepszy, S. & Niemirowicz-Szczytt, K. Sex determination in cucumber (*Cucumis sativus*) as a model system for molecular biology. *Plant Sci.* **80**, 39–47 (1991).
- Chen, H. et al. An ACC oxidase gene essential for cucumber carpel development. *Mol. Plant* **9**, 1315–1327 (2016).
- Bu, F. et al. A major quantitative trait locus conferring subgynoecy in cucumber. *Theor. Appl. Genet.* **129**, 97–104 (2016).
- Win, K. T. et al. Identification of quantitative trait loci governing subgynoecy in cucumber. *Theor. Appl. Genet.* **132**, 1505–1521 (2019).
- Pawelkowicz, M. E., Skarzyńska, A., Płader, W. & Przybecki, Z. Genetic and molecular bases of cucumber (*Cucumis sativus* L.) sex determination. *Mol. Breed.* **39**, 50 (2019).
- Li, D., Sheng, Y., Niu, H. & Li, Z. Gene interactions regulating sex determination in Cucurbits. *Front. Plant Sci.* **10**, 1231 (2019).
- Shifriss, O., George, W. L. & Quinones, J. A. Gynodioecism in cucumber. *Genetics* **49**, 285–291 (1964).
- Ito, H. & Kato, T. Factors responsible for the sex expression of Japanese cucumbers. *Hort. Assoc. Jpn. J.* **22**, 138–144. 1953 (1953).
- Peterson, C. E. A gynoecious inbred line of cucumber. *Quart. Bull. Mich. Agric. Exp. Stat.* **3**, 40–42 (1960).
- Peterson, C. E. & Anhder, L. D. Induction of staminate flowers on gynoecious cucumbers with gibberellin A3. *Science* **131**, 1673–1674 (1960).
- Wehner, T. C. & Miller, C. H. Effect of gynoecious expression on yield and earliness of a fresh-market cucumber hybrid. *J. Am. Soc. Hort. Sci.* **110**, 464–466 (1985).
- Staub, J. E., Robbins, M. D. & Wehner, T. C. Cucumber. In *Vegetables I. Handbook of Plant Breeding*, Vol. 1 (eds. Prohens, J. & Nuez, F.) (Springer, New York, NY, 2008).
- Rudich, J., Halevy, A. & Kedar, N. Ethylene evolution from cucumber plants as related to sex expression. *Plant Physiol.* **49**, 998–999 (1972).
- Takahashi, H. & Jaffe, M. Further studies of auxin and ACC induced feminization in the cucumber plant using ethylene inhibitors. *Phyton* **44**, 81–86 (1984).
- Yin, T. J. & Quinn, J. A. Tests of a mechanistic model of one hormone regulating both Sexes in *Cucumis sativus* (Cucurbitaceae). *Am. J. Bot.* **82**, 1537–1546 (1995).
- Yamasaki, S., Fujii, N. & Takahashi, H. Characterization of ethylene effects on sex determination in cucumber plants. *Sex. Plant Reprod.* **16**, 103–111 (2003).
- Atsmon, D. & Tabbak, C. Comparative effects of gibberellin, silver nitrate and aminoethoxyvinyl glycine on sexual tendency and ethylene evolution in cucumber plant (*Cucumis sativus* L.). *Plant Cell Physiol.* **20**, 1547–1555 (1979).
- Treibitsh, T., Staub, J. E. & O'Neill, S. D. Identification of a 1-aminocyclopropane-1-carboxylic acid synthase gene linked to the female (F) locus that enhances female sex expression in cucumber. *Plant Physiol.* **113**, 987–995 (1997).
- Mibus, H. & Tatlioglu, E. T. Molecular characterization and isolation of the F/f gene for femaleness in cucumber (*Cucumis sativus* L.). *Theor. Appl. Genet.* **109**, 1669–1676 (2004).
- Li, Z. et al. Molecular isolation of the M gene suggests that a conserved-residue conversion induces the formation of bisexual flowers in cucumber plants. *Genetics* **182**, 1381–1385 (2009).
- Boualem, A. et al. A cucurbit androecy gene reveals how unisexual flowers develop and dioecy emerges. *Science* **350**, 688–691 (2015).
- Knopf, R. R. & Treibitsh, T. The female-specific CS-ACS1G gene of cucumber. A case of gene duplication and recombination between the non-sex-specific 1-aminocyclopropane-1-carboxylate synthase gene and a branched-chain amino acid transaminase gene. *Plant Cell Physiol.* **47**, 1217–1228 (2006).
- Zhang, Z. et al. Genome-wide mapping of structural variations reveals a copy number variant that determines reproductive morphology in cucumber. *Plant Cell* **27**, 1595–1604 (2015).
- Zmienko, A., Samelak, A., Kozłowski, P. & Figlerowicz, M. Copy number polymorphism in plant genomes. *Theor. Appl. Genet.* **127**, 1–18 (2014).
- Hollox, E. J. et al. Psoriasis is associated with increased beta-defensin genomic copy number. *Nat. Genet.* **40**, 23–25 (2008).
- Conrad, D. F. et al. Origins and functional impact of copy number variation in the human genome. *Nature* **464**, 704–712 (2010).
- Weischenfeldt, J., Symmons, O., Spitz, F. & Korbel, J. O. Phenotypic impact of genomic structural variation: insights from and for human disease. *Nat. Rev. Genet.* **14**, 125–138 (2013).
- Diaz, A., Zikhali, M., Turner, A. S., Isaac, P. & Laurie, D. A. Copy number variation affecting the *photoperiod-B1* and *vernalization-A1* genes is associated with altered flowering time in wheat (*Triticum aestivum*). *PLoS ONE* **7**, e33234 (2012).
- Würschum, T., Boeven, P. H., Langer, S. M., Longin, C. F. & Leiser, W. L. Multiply to conquer: copy number variations at *Ppd-B1* and *Vrn-A1* facilitate global adaptation in wheat. *BMC Genet.* **16**, 96 (2015).

46. Theißen, G., Rümpler, F. & Gramzow, L. Array of MADS-Box genes: facilitator for rapid adaptation? *Trends Plant Sci.* **23**, 563–576 (2018).
47. Cook, D. E. et al. Copy number variation of multiple genes at *Rhg1* mediates nematode resistance in soybean. *Science* **338**, 1206–1209 (2012).
48. Cook, D. E. et al. Distinct copy number, coding sequence and locus methylation patterns underlie *Rhg1*-mediated soybean resistance to soybean cyst nematode. *Plant Physiol.* **165**, 630–647 (2014).
49. Lee, T. G., Kumar, I., Diers, B. W. & Hudson, M. E. Evolution and selection of *Rhg1*, a copy-number variant nematode-resistance locus. *Mol. Ecol.* **24**, 1774–1791 (2015).
50. Lee, T. G., Diers, B. W. & Hudson, M. E. An efficient method for measuring copy number variation applied to improvement of nematode resistance in soybean. *Plant J.* **88**, 143–153 (2016).
51. Wang, Y. et al. Copy number variation at the *GL7* locus contributes to grain size diversity in rice. *Nat. Genet.* **47**, 944–948 (2015).
52. Sutton, T. et al. Boron-toxicity tolerance in barley arising from efflux transporter amplification. *Science* **318**, 1446–1449 (2007).
53. Maron, L. G. et al. Aluminum tolerance in maize is associated with higher *MATE1* gene copy number. *Proc. Natl Acad. Sci. USA* **110**, 5241–5246 (2013).
54. Gaines, T. A. et al. Mechanism of resistance of evolved glyphosate-resistant palmer amaranth (*Amaranthus palmeri*). *J. Agric. Food Chem.* **59**, 5886–5889 (2011).
55. Gaines, T. A. et al. Gene amplification confers glyphosate resistance in *Amaranthus palmeri*. *Proc. Natl Acad. Sci. USA* **107**, 1029–1034 (2010).
56. Caicedo, A. L., Richards, C., Ehrenreich, I. M. & Purugganan, M. D. Complex rearrangements lead to novel chimeric gene fusion polymorphisms at the *Arabidopsis thaliana* MAF2-5 flowering time gene cluster. *Mol. Biol. Evol.* **26**, 699–711 (2009).
57. Magadum, S., Banerjee, U., Murugan, P., Gangapur, D. & Ravikesavan, R. Gene duplication as a major force in evolution. *J. Genet.* **92**, 155–161 (2013).
58. Lye, Z. N. & Purugganan, M. D. Copy number variation in domestication. *Trends Plant Sci.* **24**, 352–366 (2019).
59. de Bolt, S. Copy number variation shapes genome diversity in *Arabidopsis* over immediate family generational scales. *Genome Biol. Evol.* **2**, 441–453 (2010).
60. Kondrashov, F. A. Gene duplication as a mechanism of genomic adaptation to a changing environment. *Proc. R. Soc. B* **279**, 5048–5057 (2012).
61. Lu, P. et al. Analysis of *Arabidopsis* genome-wide variations before and after meiosis and meiotic recombination by resequencing Landsberg erecta and all four products of a single meiosis. *Genome Res.* **22**, 508–518 (2012).
62. Osipowski, P. et al. A high-quality cucumber genome assembly enhances computational comparative genomics. *Mol. Genet. Genomics* <https://doi.org/10.1007/s00438-019-01614-3> (2019).
63. Hastings, P. J., Lupski, J. R., Rosenberg, S. M. & Ira, G. Mechanisms of change in gene copy number. *Nat. Rev. Genet.* **10**, 551–564 (2009).
64. Li, Q. et al. A chromosome-scale genome assembly of cucumber (*Cucumis sativus* L.). *Gigascience* **8**, 1–10 (2019).
65. Tao, Q. et al. Ethylene responsive factor ERF110 mediates ethylene-regulated transcription of a sex determination-related orthologous gene in two *Cucumis* species. *J. Exp. Bot.* **69**, 2953–2965 (2018).
66. Martin, A. et al. A transposon-induced epigenetic change leads to sex determination in melon. *Nature* **461**, 1135–1138 (2009).
67. Zhang, J. et al. A unique chromosome translocation disrupting *CIWIP1* leads to gynoeicy in watermelon. *Plant J.* <https://doi.org/10.1111/tpj.14537> (2019).
68. Hu, B. et al. Engineering non-transgenic gynoeicy cucumber using an improved transformation protocol and optimized CRISPR/Cas9 system. *Mol. Plant* **10**, 1575–1578 (2017).
69. Shiomi, S. et al. cDNA cloning of ACC synthase and ACC oxidase genes in cucumber fruit and their differential expression by wounding and auxin. *J. Jpn. Soc. Hort. Sci.* **67**, 685–692 (1998).
70. Mathooko, F. M. et al. Expression characteristics of *CSACS1*, *CSACS2* and *CSACS3*, three members of the 1-aminocyclopropane-1-carboxylate synthase gene family in cucumber (*Cucumis sativus* L.) fruit under carbon dioxide stress. *Plant Cell Physiol.* **40**, 164–172 (1999).
71. Shiber et al. The origin and mode of function of the *Female* locus in cucumber. In *Cucurbitaceae 2008, Proc. 9th EUCARPIA Meeting on Genetics and Breeding of Cucurbitaceae*, May 21–24, 2008, France (ed. Pitrat, M.) 263–270 (INRA, Avignon, 2008).
72. Kooistra, K. Femaleness in breeding glasshouse cucumbers. *Euphytica* **6**, 1–17 (1967).
73. Davidi, H. A historical survey of cucumber breeding in Israel. Proc. 4th Balkan Symposium on Vegetables and Potatoes (eds. Krasteva, L. and Panayotov, N.) *Acta Hort.* **830**, 33–36 (2009).
74. Lv, J. et al. Genetic diversity and population structure of cucumber (*Cucumis sativus* L.). *PLoS ONE* **7**, e46919 (2012).
75. Wang, X. et al. The USDA cucumber (*Cucumis sativus* L.) collection: genetic diversity, population structure, genome-wide association studies and core collection development. *Hortic. Res.* **5**, 64 (2018).
76. Zmienko, A., Samelak-Czajka, A., Kozłowski, P., Szymanska, M. & Figlerowicz, M. *Arabidopsis thaliana* population analysis reveals high plasticity of the genomic region spanning *MSH2*, *AT3G18530* and *AT3G18535* genes and provides evidence for NAHR-driven recurrent CNV events occurring in this location. *BMC Genomics* **17**, 893 (2016).
77. Cho, Y. B., Jones, I. & Vodkin, L. O. Nonallelic homologous recombination events responsible for copy number variation within an RNA silencing locus. *Plant Direct* **3**, e00162 (2019).
78. Leister, D. Tandem and segmental gene duplication and recombination in the evolution of plant disease resistance genes. *Trends Genet.* **20**, 116–122 (2004).
79. Oling, D., Masoom, R. & Kvint, K. Loss of *Ubp3* increases silencing, decreases unequal recombination in rDNA, and shortens the replicative life span in *Saccharomyces cerevisiae*. *Mol. Biol. Cell* **25**, 1916–1924 (2014).
80. Pettersson, M., Eisfeldt, J., Syk Lundberg, E., Lundin, J. & Lindstrand, A. Flanking complex copy number variants in the same family formed through unequal crossing-over during meiosis. *Mutat. Res.* **812**, 1–4 (2018).
81. Li, Z. et al. Development and fine mapping of three co-dominant SCAR markers linked to the *M/m* gene in the cucumber plant (*Cucumis sativus* L.). *Theor. Appl. Genet.* **117**, 1253–1260 (2008).
82. Jackson, S. A., Wang, M. L., Goodman, H. M. & Jiang, J. Application of fiber-FISH in genome analysis of *Arabidopsis thaliana*. *Genome* **41**, 566–572 (1998).
83. Han, Y. et al. Distribution of the tandem repeat sequences and karyotyping in cucumber (*Cucumis sativus* L.) by fluorescence in situ hybridization. *Cytogenet. Genome Res.* **122**, 80–88 (2008).
84. Jiang, J., Gill, B. S., Wang, G., Ronald, P. C. & Ward, D. C. Metaphase and interphase fluorescence in situ hybridization mapping of the rice genome with bacterial artificial chromosomes. *Proc. Natl Acad. Sci. USA* **92**, 4487–4491 (1995).
85. Pan, Y., Bo, K., Cheng, Z. & Weng, Y. The loss-of-function GLABROUS 3 mutation in cucumber is due to LTR-retrotransposon insertion in a class IV HD-ZIP transcription factor gene *CsGL3* that is epistatic over *CsGL1*. *BMC Plant Biol.* **15**, 302 (2015).
86. Yang, L. et al. Chromosome rearrangements during domestication of cucumber as revealed by high-density genetic mapping and draft genome assembly. *Plant J.* **71**, 895–906 (2012).
87. Qi, J. et al. A genomic variation map provides insights into the genetic basis of cucumber domestication and diversity. *Nat. Genet.* **45**, 1510–1515 (2013).
88. Wang, Y. et al. STAYGREEN, STAY HEALTHY: a loss-of-susceptibility mutation in the STAYGREEN gene provides durable, broad-spectrum disease resistances for over 50 years of US cucumber production. *New Phytol.* **221**, 415–430 (2019).
89. Li, Z. et al. A putative positive feedback regulation mechanism in *CsACS2* expression suggests a modified model for sex determination in cucumber (*Cucumis sativus* L.). *J. Exp. Bot.* **63**, 4475–4484 (2012).