


Soil fauna diversity increases CO₂ but suppresses N₂O emissions from soil

Ingrid M. Lubbers^{1,2}  | Matty P. Berg^{3,4} | Gerlinde B. De Deyn¹ |
Wim H. van der Putten^{5,6} | Jan Willem van Groenigen¹

¹Soil Biology Group, Wageningen University, Wageningen, The Netherlands

²Soil Geography and Landscape Group, Wageningen University, Wageningen, The Netherlands

³Department of Ecological Science, Animal Ecology Group, Vrije Universiteit, Amsterdam, The Netherlands

⁴Groningen Institute of Evolutionary Life Science, Community and Conservation Ecology Group, University of Groningen, Groningen, The Netherlands

⁵Department of Terrestrial Ecology, Netherlands Institute of Ecology (NIOO-KNAW), Wageningen, The Netherlands

⁶Laboratory of Nematology, Wageningen University, Wageningen, The Netherlands

Correspondence

Ingrid M. Lubbers, Soil Biology Group, Wageningen University, PO Box 47, 6700AA Wageningen, The Netherlands.
Email: Ingrid.Lubbers@wur.nl

Funding information

Netherlands Organization for Scientific Research/Earth and Life Sciences, Grant/Award Number: 823.01.016

Abstract

Soil faunal activity can be a major control of greenhouse gas (GHG) emissions from soil. Effects of single faunal species, genera or families have been investigated, but it is unknown how soil fauna diversity may influence emissions of both carbon dioxide (CO₂, end product of decomposition of organic matter) and nitrous oxide (N₂O, an intermediate product of N transformation processes, in particular denitrification). Here, we studied how CO₂ and N₂O emissions are affected by species and species mixtures of up to eight species of detritivorous/fungivorous soil fauna from four different taxonomic groups (earthworms, potworms, mites, springtails) using a microcosm set-up. We found that higher species richness and increased functional dissimilarity of species mixtures led to increased faunal-induced CO₂ emission (up to 10%), but decreased N₂O emission (up to 62%). Large ecosystem engineers such as earthworms were key drivers of both CO₂ and N₂O emissions. Interestingly, increased biodiversity of other soil fauna in the presence of earthworms decreased faunal-induced N₂O emission despite enhanced C cycling. We conclude that higher soil fauna functional diversity enhanced the intensity of belowground processes, leading to more complete litter decomposition and increased CO₂ emission, but concurrently also resulting in more complete denitrification and reduced N₂O emission. Our results suggest that increased soil fauna species diversity has the potential to mitigate emissions of N₂O from soil ecosystems. Given the loss of soil biodiversity in managed soils, our findings call for adoption of management practices that enhance soil biodiversity and stimulate a functionally diverse faunal community to reduce N₂O emissions from managed soils.

KEYWORDS

community composition, functional dissimilarity, GHG mitigation, net diversity effect, soil-derived GHG emission, species richness

1 | INTRODUCTION

There is considerable interest in how soils influence global climate change, and through which feedback mechanisms soil quality may be deteriorating. The biodiversity within an active soil biome is a large contributor to the soil function–climate feedback (Wall, 2012). However, empirical research on the connection between soil biodiversity and climate regulation is lacking. Specifically, the close relationship between soil-derived greenhouse gas (GHG) emissions and soil processes such as biogeochemical cycling of C and N may result in feedback mechanisms that can either amplify or diminish the initial climate forcing (Crowther et al., 2016; Van Nes et al., 2015). As part of the soil biodiversity, soil fauna species can have a major impact on this relationship through their influence on biogeochemical cycling of nutrients (Coulis et al., 2015; Filser et al., 2016; Sauvadet, Chauvat, Brunet, & Bertrand, 2017; Wall et al., 2008), plant productivity (Liiri, Setälä, Haimi, Pennanen, & Fritze, 2002; Van Groenigen et al., 2014), and soil-derived GHG emissions (Lubbers, Van Groenigen, et al., 2013).

Agroecosystems are a known focal point of problems related to soil biodiversity, soil-derived GHG emissions and the biogeochemical cycling of nutrients. Feedback mechanisms that may influence global climate change are especially relevant in these agroecosystems, where soil biodiversity is mostly negatively affected by agricultural intensification (Ponge et al., 2013; Postma-Blaauw, Goede, Bloem, Faber, & Brussaard, 2012; Tsiafouli et al., 2015), where inputs of nitrogen (N) fertilizers are increasing globally (FAO, 2016), and where soil-derived GHG emissions are high (IPCC, 2014). Agroecosystems also have the best potential for C storage in the soil by restoring soil organic C that was previously lost due to agricultural practices (Lal, 2004; Paustian, Cole, Sauerbeck, & Sampson, 1998; Smith, 2016; Stockmann et al., 2013). Yet, increasing soil C stocks can be negated by elevated soil-derived emissions of the potent GHG N₂O (Lubbers, Groenigen, Brussaard, & Groenigen, 2015; Powlson et al., 2014; Powlson, Whitmore, & Goulding, 2011). Global change models that underpin predictions of future climate scenarios still lack the potentially intricate and non-linear interactions between soil fauna and microbes regulating soil biogeochemical cycles (Grandy, Wieder, Wickings, & Kyker-Snowman, 2016). The lack of sufficient empirical data addressing ecological feedbacks to ecosystem processes such as GHG emissions precludes the parameterization necessary for further optimization of such models (Blagodatsky & Smith, 2012; Crowther et al., 2015; Grandy et al., 2016). To further unravel how soils and soil disturbance contribute to GHG emissions, mechanistic research is needed on functional diversity of soil fauna affecting the soil processes underlying climate regulation.

Belowground, gas fluxes of CO₂, methane (CH₄) and N₂O are the result of a variety of (micro)biotic processes, occurring in 'hotspots' and during 'hot moments' (Kuzakov & Blagodatskaya, 2015): CO₂ is produced by soil respiration (including root, faunal and microbial respiration; Rastogi, Singh, & Pathak, 2002), CH₄ through methanogenesis (Le Mer & Roger, 2001), while N₂O is produced by a combination of microbial transformation processes,

most importantly denitrification (Opdyke, Ostrom, & Ostrom, 2009), as well as nitrification and nitrifier–denitrification (Wrage-Monnig et al., 2018). All these soil (micro)biotic processes resulting in the production of CO₂, CH₄ and N₂O are controlled by numerous interacting conditions, characterized by key physico-chemical factors such as the availability of labile C, mineral N and oxygen (O₂), soil pH, soil moisture, temperature and diffusivity (Granli & Bøckman, 1994). Whereas CO₂ is an end product of soil respiration, the net efflux of CH₄ is the balance between anaerobic decomposition by methanogens and oxidation by methanotrophic bacteria (Dalal & Allen, 2008). Nitrous oxide can be both an intermediate and a by-product of N transformation processes, and N₂O gas can therefore also be consumed during (upward) diffusion through the soil profile (Klefoth, Oenema, & Willem Van Groenigen, 2012). This makes the net emission rates of N₂O from soils highly variable and notoriously difficult to predict: N₂O emission is dependent on the prevailing soil physico-chemical factors, as well as on soil faunal activity (Kuiper, Deyn, Thakur, & Groenigen, 2013; Lubbers, Van Groenigen, et al., 2013).

The only soil faunal group so far acknowledged to have a pronounced influence on soil-derived N₂O emissions are earthworms (Lubbers, Van Groenigen, et al., 2013). These macro-detrivores have been well studied with respect to N₂O emissions and they can increase N₂O emissions considerably through interacting with the microbial community (Drake & Horn, 2006; Nebert, Bloem, Lubbers, & Groenigen, 2011), by increasing the mineralization of nutrients and altering the soil structure (Lubbers, Brussaard, Otten, & Groenigen, 2011; Lubbers, Van Groenigen, et al., 2013). Other macrofauna, such as isopods and grubs, and mesofaunal groups, such as potworms, fungivorous mites and springtails, can also influence N₂O emissions (Kuiper et al., 2013; Majeed et al., 2014; Tianxiang, Huixin, Tong, & Feng, 2008; Van Vliet, Beare, Coleman, & Hendrix, 2004). However, their (mostly enhancing) effects on N₂O emission have mainly been determined as single-species effects. The few studies on species combinations of different soil fauna feeding groups with respect to N₂O emissions show diverging results: from a strong increase of soil-derived N₂O emissions for a combination of mesofaunal species (Thakur, Groenigen, Kuiper, & Deyn, 2014) to the absence of any interactive effects between earthworms and mesofauna (Schorpp et al., 2016; Wu, Lu, Lu, Guan, & He, 2015). Furthermore, most soil fauna–GHG emission studies to date have focussed on combinations of very few (up to four) different species under controlled laboratory conditions. Based on modelling studies (De Vries et al., 2013), it has been argued that changes in soil biodiversity in the field have substantial consequences for ecosystem processes, such as soil-derived GHG emissions. However, no controlled laboratory studies so far have looked into the potential role of community composition and functional diversity of soil fauna in increasing or decreasing net GHG emissions from soil. In order to establish that changes in soil biodiversity can indeed influence the production and emission of GHG emissions, a proof-of-principle incubation study is needed to focus on causal relationships between soil fauna diversity and GHG emissions from soil.

Here, we quantified biodiversity effects on soil-derived emissions of N_2O and CO_2 under controlled laboratory conditions. Methane was not measured because the experimental conditions representative of temperate managed soils (aerobic, relatively low ambient temperature) were not favourable to significant CH_4 production (Levy et al., 2012). We experimentally varied species composition in hay-amended soil microcosms with an increasing number of detritivorous/fungivorous soil mesofauna and macrofauna species: none, one, two, four and eight species per microcosm. We randomly assigned eight species belonging to four different taxonomic groups (earthworms, potworms, mites, springtails) to multispecies treatments following a 'constrained random sampling design' (Heemsbergen et al., 2004). Given the collective evidence that a higher species richness enhances fundamental ecological soil processes (Cardinale et al., 2006; Nielsen, Ayres, Wall, & Bardgett, 2011; Tresch et al., 2019; Wall, 2012), we hypothesized increasing soil fauna biodiversity to enhance the microbial (transformation) processes that underlie soil-derived emissions of N_2O and CO_2 . To investigate this hypothesis, we measured N_2O and CO_2 emissions over an experimental period of 120 days and calculated *net biodiversity effects* (Heemsbergen et al., 2004; Loreau & Hector, 2001). We also determined soil biogeochemical properties following the destructive sampling of microcosm soil and calculated the *functional dissimilarity* among species (Heemsbergen et al., 2004).

2 | MATERIALS AND METHODS

2.1 | Experimental design

We manipulated faunal species composition in hay-amended soil microcosms at two levels: species and taxa level. We set the number of soil mesofauna and macrofauna at none, one, two, four and eight species per microcosm treatment (Table 1). The eight selected detritivorous/fungivorous soil fauna species belonged to four different taxonomic groups (earthworms, potworms, mites, springtails; Table 1) and were either obtained from cultures (Vrije Universiteit, Amsterdam, the Netherlands; Wageningen University, Wageningen, the Netherlands) or from grass fields on campus at Wageningen University. To determine compositional effects on soil-derived GHG fluxes (N_2O and CO_2), different two- and four-species combinations were included following a 'constrained random sampling design' (Heemsbergen et al., 2004). This means that each species was assigned randomly to multispecies treatments with two constraints: (a) each species to be represented equally (six times) and (b) two- and four-species combinations containing similar taxonomic group diversity, that is, one versus two taxonomic groups and two versus four taxonomic groups respectively. Single-species treatments of all eight species were included to quantify their per capita (mg^{-1} dry weight animal) effect on N_2O and CO_2 emissions, and these were used for *net biodiversity effect* and *functional dissimilarity* calculations (see Section 2.6). Two control treatments, 'Control Soil' and 'Control Hay', completed the number of treatments to 23 (Table 1).

TABLE 1 Overview of the fauna treatments, including treatment codes, # species, # taxa and faunal density (# individuals microcosm⁻¹) per treatment

Abbr.	Fauna	Species	# Species																	
			1	2	2	4	4	4	8											
			# Taxa																	
Pw1	Potworm	<i>Enchytraeus albidus</i>	C	Pw1	50	K	Pw1	25	O	Pw2	25	S	Mi1	100	U	Pw1	12	W	Pw1	6
Pw2	Potworm	<i>Enchytraeus crypticus</i>	D	Pw2	50	L	Pw2	25	P	Ew1	2	T	Mi2	100	V	Sp2	65		Pw2	6
Mi1	Mite	<i>Rhizoglyphus robini</i>	E	Mi1	400	M	Mi1	200	Q	Pw1	25		Ew1	1		Ew1	1		Mi1	50
Mi2	Mite	<i>Oppia nitens</i>	F	Mi2	400	N	Mi2	200	R	Sp2	130		Ew2	1		Mi1	100		Mi2	50
Sp1	Springtail	<i>Sinella curviseta</i>	G	Sp1	260	J	Sp1	130		Mi1	200		Pw1	13		Ew2	1		Sp1	33
Sp2	Springtail	<i>Folsomia candida</i>	H	Sp2	260		Sp2	130		Sp1	130		Pw2	13		Mi2	100		Sp2	33
Ew1	Earthworm	<i>Aporrectodea caliginosa</i>	I	Ew1	4		Ew1	2		Mi2	200		Sp1	65		Sp1	65		Ew1	1
Ew2	Earthworm	<i>Lumbricus rubellus</i>	J	Ew2	2		Ew2	1		Ew2	1		Sp2	65		Pw2	12		Ew2	1

Note: Codes for treatments are alphabetically: A up to W, for, respectively, Control Soil, Control Hay, Pw1, Mi1, Mi2, Sp1, Sp2, Ew1, Ew2, Pw1 & Pw2, etc. Two control treatments were included in the experimental design: Control Soil (treatment code = A) and Control Hay (treatment code = B), completing the number of treatments to 23.

2.2 | Microcosm set-up

Polypropylene microcosms (diameter = 6.7 cm, height = 15 cm, volume = 500 cm³) were filled with 250 g dry soil. The sandy soil (*typic Endoaquoll*, 75% sand, 23% silt and 2% clay) was collected from 'Droevendaal Agricultural Farm' (51°59'N, 5°39'E) from 0 to 25 cm soil depth and contained 14.6 g total C/kg, 1.3 g total N/kg and had a pH (measured in 0.01 M CaCl₂) of 5.4. The soil was sieved over an 8 mm mesh and was heated for 24 hr at 70°C to eliminate mesofauna and macrofauna while maintaining a viable soil microbial population (Kaneda & Kaneko, 2011). Distilled water was added at a rate of 220 ml/kg to reach a water-filled pore space (WFPS) of 62% after gently pressing the soil to a bulk density of 1.4 g/cm³. The soil depth was 6 cm.

Hay residue (32.8 g N/kg dry matter, 448.5 g C/kg dry matter, C:N ratio = 13.8) was chopped in pieces of ~1 cm and wet autoclaved. To represent a realistic N fertilization of 125 kg N/ha, 1.34 g (of dry hay equivalent) was placed on top of the soil surface. After 2 weeks of pre-incubation, the experiment was started by introducing the soil fauna species to the microcosms according to densities shown in Table 1. Both potworm species were added in an equal ratio of adults to juveniles. Mites (*Rhizoglyphus robini*) could only be added with some yeast flakes from their cultures; therefore, all microcosms received some yeast flakes. Earthworms were prepared following the wet filter paper method, that is, earthworms were moved to damp filter paper to void gut contents for 48 hr before weighing (Dalby, Baker, & Smith, 1996). All added earthworms, potworms, mites and springtails were added individually by hand with tweezers or a brush.

To avoid intraspecific competition (Dias et al., 2013), the total abundance of the soil mesofauna was kept constant across all treatments, according to established soil food web numbers (De Ruiter, Van Veen, Moore, Brussaard, & Hunt, 1993). For individual mesofauna species, this meant that we added abundances corresponding to their share in the treatment (100% for single-species treatment, 50% for two-species combinations, 25% for the four-species combination and 12.5% for the eight-species combination, as indicated in Table 1). The total abundance of the macrofauna, that is, the earthworms, was also kept constant across treatments. However, because of their large body size and low abundance, we assigned earthworms on the basis of their body mass rather than their abundance (Table 1, Table S1).

The microcosms were covered tightly with black cloth to prevent fauna from escaping, facilitate gaseous exchange, but minimize moisture loss. Soil moisture was resupplied gravimetrically every 2–3 days after taking gas flux measurements.

The microcosms were arranged in a randomized block design consisting of five replicates divided over five separate blocks; replicates six, seven and eight were placed in block one, two and three, respectively, to be destructively sampled earlier in the experimental period than replicates 1–5. During the experimental period of 120 days, there were two destructive sampling moments: on day 47 ($n = 3$, replicates six, seven and eight) and on day 120 ($n = 5$,

replicates 1–5). The microcosms were placed in the dark at 16°C and a relative humidity of 60%.

2.3 | N₂O and CO₂ flux measurements

Nitrous oxide and CO₂ fluxes were measured 34 times over the experimental period of 120 days. Using a static closed chamber technique, gaseous concentrations of N₂O and CO₂ in the microcosm headspace were measured with an Innova 1302 photoacoustic infrared multi gas analyser (Brüel & Kjaer; Velthof, Kuikman, & Oenema, 2002). For N₂O flux measurements, a soda-lime filter was used to minimize interference by CO₂; CO₂ headspace concentration was therefore measured separately from N₂O headspace concentration.

Cumulative N₂O and CO₂ emissions were calculated assuming a linear change in emission rates between subsequent flux measurements, which was checked occasionally. Values for N₂O and CO₂ were transformed to CO₂ equivalents (CO₂-eq; IPCC, 2013), using a 100 year time horizon as in the Kyoto Protocol. The conversion of N₂O emission rates to CO₂ equivalents was done as follows: 1 kg N₂O = 298 kg CO₂ (Lubbers, Van Groenigen, et al., 2013).

Hay decomposition rates were determined by calculating the difference in CO₂ production (mg CO₂-C/microcosm) between the control treatment with (Control Hay: 'B') and without hay residues (Control Soil: 'A'). The cumulative hay-derived decomposition was calculated to confirm that the experiment had proceeded within realistic respiration rates for laboratory conditions (Heemsbergen et al., 2004; Kuiper et al., 2013; Porre, Groenigen, Deyn, Goede, & Lubbers, 2016; Thakur et al., 2014). Hay decomposition rates have not been used in further analyses of faunal effects, since soil fauna diversity effects on these decomposition rates would be similar to effects on cumulative CO₂ emissions (see Section 2.6 for the determination of soil fauna diversity effects).

2.4 | Soil analyses

Destructive sampling for soil property analyses was done on day 1 ($n = 4$), day 47 ($n = 3$) and on day 120 ($n = 4$, randomly block 2–5). Each microcosm was divided vertically in two halves, one of which was used for faunal extractions and the other for soil chemical and microbial analyses. Because of the extraction of the soil fauna from the hay-soil interface, we did not remove and weigh the leftover hay residues from the microcosms on day 47 or day 120. Since in many cases the residues were much stuck to the soil and were partly or entirely mixed in, we carefully removed any left-over residues from the vertical halves that were kept aside for soil analyses. After thoroughly mixing this latter half, a subsample was used to determine microbial biomass nitrogen (MBN), following the chloroform fumigation and extraction technique (using a correction factor of 0.54; Brookes, Landman, Pruden, & Jenkinson, 1985). Subsequently, total dissolved N (Nts), ammonia (NH₄⁺), nitrate and nitrite (NO₃⁻ + NO₂⁻) concentrations were

measured colorimetrically in a 0.5 M K_2SO_4 extract, at 1:4 w/v, by segmented flow analysis (3x Segmented Flow Analyzer; Klug, 1999); the dissolved organic N (DON) content was calculated by subtracting $NH_4^+ + (NO_3^- + NO_2^-)$ from Nts. The pH and dissolved organic C (DOC) were measured in a 0.01 M $CaCl_2$ extract, at 1:10 w/v, by segmented flow analysis (3x Segmented Flow Analyzer; Houba, Temminghoff, Gaikhorst, & Vark, 2000).

2.5 | Fauna abundances

Soil fauna was extracted from the other vertical half of the microcosm on day 47 ($n = 3$) and on day 120 ($n = 5$). When both potworms and mites were present in the microcosms, we subdivided one microcosm halve further in two, following Thakur et al. (2014). This was necessary because of the different extraction techniques for potworms (wet extraction with a Beermann funnel; Petersen & Luxton, 1982) and mites (dry extraction with a Berlese funnel; Tullgren funnel; Petersen & Luxton, 1982). Earthworms were carefully collected by hand from the entire microcosm. The numbers of live earthworms were recorded and fresh weights were determined following the wet filter paper method (Dalby et al., 1996). Microcosms belonging to the 'Control Hay' treatment were treated in the same way as when potworms and mites were present, to ascertain that no faunal infestations had occurred. Fauna abundances at two destructive sampling dates are shown in Table S1.

More information on the determination of fauna dry weight is in Supporting Information.

2.6 | Calculations of net biodiversity and functional dissimilarity

The net biodiversity effect was calculated following Heemsbergen et al. (2004) and Loreau and Hector (2001). First, for each single species, its per capita effect (g/DW animal) on the cumulative N_2O and CO_2 emissions after 120 days was determined by subtracting the Control Hay from the respective single-species treatments; the average dry weight animal over 120 days was used, that is, the average dry weight at the start of the experiment and at the final harvest after 120 days (Table S1). Second, the expected effect of a multispecies treatment (two-, four- and eight-species combinations) on either N_2O or CO_2 was calculated as the sum of the per capita effects of its component species. The difference between the expected effect (Y_E) and the observed GHG flux (Y_O) of a multispecies treatment was then the net biodiversity effect (ΔY): $\Delta Y = Y_O - Y_E$. Positive net biodiversity effects point to facilitative interactions and negative effects point to inhibitive interactions.

Apart from the influence of species richness on net biodiversity effects, it is known that communities composed of functionally dissimilar species are likely to have stronger effects on process rates (Heemsbergen et al., 2004). Therefore, we also calculated net biodiversity effects in relation to mean functional dissimilarity of species in multispecies mixtures in terms of the key controlling factors of

N_2O and CO_2 production. To analyse the role of biodiversity in soil-derived GHG emissions, we considered the key driving factors for N_2O production and emission (concentrations of NH_4^+ , $NO_3^- + NO_2^-$, DOC and pH; Granli & Bøckman, 1994; these factors also impact CO_2 production and emission) to shape an 'attribute space' in which the functional dissimilarity between species could be quantified as the standardized 'ecological distance' (Walker, Kinzig, & Langridge, 1999). The variables used to define the 'attribute space' were the per capita concentrations of NH_4^+ , $NO_3^- + NO_2^-$, DOC and pH in the single-species treatments (Table S2). Since these variables do not necessarily have the same units, we defined a linear scale between the smallest and largest value for each variable, and the single-species treatments were placed along the non-dimensionalized scales. The use of these normalized scales to shape our 'four-dimensional attribute space' allowed an estimate of functional dissimilarity (sum of pairwise 'ecological distances' between species in the four-dimensional space, divided by the number of interspecific distances between fauna species present in a combination) in which all variables had an equal weight in the calculation.

When including the mean pore size as a soil structural element in a new 'attribute space', we decided to interchange pH for mean pore size. This was done for two reasons: (a) keep the 'attribute space' four-dimensional because using four variables (instead of more) is optimal for our number of samples; and (b) pH was selected over the other variables because pH was significantly correlated with most of the other variables used for the 'attribute space'.

2.7 | Statistical analyses

All statistical analyses were conducted in SPSS, version 22.0 (SPSS Inc.). Our data of the soil-derived GHG fluxes were much skewed by the presence of earthworms, and therefore, treatment imposed differences were first established by a nonparametric statistical test: the Kruskal–Wallis test was used to establish the difference between treatments with and without earthworms. Subsequently, the with- and without-earthworm data sets were checked for normality and homoscedasticity using the Shapiro–Wilk test and Levene's test respectively. If needed, data were transformed to normality (with the inverse square root in the case of cumulative N_2O fluxes) before using standard one-way analysis of variance (ANOVA) with blocking and post hoc Tukey HSD to quantify the significance of the effects of soil fauna. Our data of the soil properties (NH_4^+ , $NO_3^- + NO_2^-$, MBN, DOC and pH) were also much skewed. The same procedure was repeated; when no transformations to normality appeared to be possible on day 120, the Kruskal–Wallis Test was again used with subsequent pairwise comparisons (Mann–Whitney U Test with a Bonferroni correction) to test for differences between fauna treatments. The treatment 'Control Soil' was never part of statistical analysis.

To test for differences in the net biodiversity effect (on cumulative N_2O and CO_2 emissions after 120 days) between two-species, four-species and eight-species combinations, an independent samples t test assuming unequal variances was used. A one-sample t test

was used to test the difference between net biodiversity effects and the constant zero; not significantly different from zero indicates a neutral net biodiversity effect.

To identify the relationship between the mean functional dissimilarity (predictor variable) and the net biodiversity effect on cumulative N_2O and CO_2 (response variables), we used a fitted linear regression model.

For all analyses, a p -value of .05 or smaller was considered significant.

3 | RESULTS

3.1 | Net biodiversity effects on faunal-induced GHG emissions

Figure 1 shows the net biodiversity effect on per capita soil-derived emissions of N_2O and CO_2 in relation to species number. The 0-line indicates a neutral net biodiversity effect. Deviations from this 0-line (i.e. positive or negative interactions among species) could be used to explain the observed variation in soil-derived emissions of N_2O and CO_2 within multispecies treatments. The net biodiversity effects on per capita soil-derived emissions of N_2O and CO_2 showed contrasting patterns: on average, species richness reduced cumulative faunal-induced N_2O emissions, whereas it increased cumulative faunal-induced CO_2 emissions (Figure 1). For N_2O , we found significant differences between the two- and eight-species treatments ($p = .013$) and the four- and eight-species treatments ($p = .001$). For CO_2 , there were no significant differences between the two-, four- and eight-species groups. However, for both N_2O and CO_2 , the net biodiversity effects in the four- and eight-species groups were significantly different from the constant 0: net biodiversity effects for N_2O in the four- and eight-species treatments were negative, pointing to inhibitive interactions; net biodiversity effects for CO_2 in the four- and eight-species treatments were positive, pointing to facilitative interactions (Figure 1).

Figure 2 shows the net biodiversity effect on per capita soil-derived emissions of N_2O and CO_2 in relation to mean functional dissimilarity. Similar to the effects of net biodiversity in relation to species number (Figure 1), we found contrasting patterns for net biodiversity effects in relation to mean functional dissimilarity: a negative relationship for soil-derived N_2O emission and a positive relationship for soil-derived CO_2 emission with mean functional dissimilarity (both significant; Figure 2). For N_2O , we found a significant negative regression between mean functional dissimilarity and the net biodiversity effect ($p = .030$). For CO_2 , we found a significant positive regression between mean functional dissimilarity and the biodiversity effect ($p < .001$). Since the functional dissimilarity of species mixtures was neither related to species number nor number of taxa, we could assume that it is the divergent influences of the different detritivorous/fungivorous species on the controlling factors of N_2O and CO_2 production that explain the negative net biodiversity effects for N_2O and the positive net biodiversity effects for CO_2 emission.




FIGURE 1 Net biodiversity effect on per capita soil-derived emissions of N_2O and CO_2 ($mg\ N_2O-N\ m^{-2}\ mg^{-1}\ DW$ and $g\ CO_2-C\ m^{-2}\ mg^{-1}\ DW$, respectively) in relation to species number. The 0-line indicates a neutral net biodiversity effect. Each box plot shows the 5th and 95th percentiles and the mean (red line) of all treatments for two species ($n = 40$), four species ($n = 20$) and eight species ($n = 5$), respectively. Data points that lie outside the 5th and 95th percentiles are shown as dots. For N_2O , significant differences between the two- and eight-species treatments (two-sample t test assuming unequal variances, t statistic = 2.61, p two-tail = .013) and the four- and eight-species treatments (two-sample t test assuming unequal variances, t statistic = 3.70, p two-tail = .001) are indicated by different letters. For CO_2 , there are no significant differences between the two-, four- and eight-species groups. For both N_2O and CO_2 , the four- and eight-species groups are significantly different from the constant 0 (established by a one-sample t test); levels of significance are * $<.05$; ** $<.01$; *** $<.001$; ns = not significant

3.2 | Community composition effects on faunal-induced GHG emissions

Figure 3 shows the observed cumulative emissions of N_2O and CO_2 after 120 days. Faunal-induced N_2O and CO_2 emissions in our experiment were dominantly triggered by the presence of large ecosystem engineering detritivores, the two earthworm species, *Aporrectodea caliginosa* and *Lumbricus rubellus*. We observed considerably increased rates of N_2O emissions for treatments that included earthworms, especially the endogeic earthworm *A. caliginosa* (Figure 3, the black bars; and Table S3). Increased rates of CO_2 emission in single-species earthworm treatments (treatments 'I' and 'N' for *A. caliginosa* and *L. rubellus*, respectively) were similar for both earthworm species, and much smaller than the increased rates of N_2O . The extremely increased N_2O emission corresponded with equally increased concentrations of mineral N (NO_3^- and/or NH_4^+); after the experimental period of 120 days, there were no significant earthworm (or any other faunal species) effects on dissolved organic C (Table S2).




FIGURE 2 Net biodiversity effect on per capita soil-derived emissions of N_2O and CO_2 ($\text{mg N}_2\text{O-N m}^{-2} \text{mg}^{-1} \text{DW}$ and $\text{g CO}_2\text{-C m}^{-2} \text{mg}^{-1} \text{DW}$, respectively) in relation to mean functional dissimilarity (sum of effects on NH_4 , NO_3 , DOC and pH (Granli & Bøckman, 1994) of species in the community). The 0-line indicates a neutral net biodiversity effect. Each vertical series of dots represent a treatment ($n = 5$ replicates per treatment; some dots overlap). For N_2O , a significant negative regression between mean functional dissimilarity and the net biodiversity effect (linear regression, $F_{64,63} = 4.95$, $p = .030$) indicates that communities with functionally dissimilar species are more likely to have negative net biodiversity effects. For CO_2 , a significant positive regression between mean functional dissimilarity and the net biodiversity effect (linear regression, $F_{64,63} = 14.5$, $p < .001$) indicates that communities with functionally dissimilar species are more likely to have positive biodiversity effects. Uppercase letters at the top of the figure refer to the species combination given in Table 1. Colours indicate the number of species present: blue = 2 species present, pink = 4 species present and green = 8 species present. Underlined uppercase letters refer to species combinations that include an earthworm species




FIGURE 3 Observed cumulative emissions of N_2O and CO_2 after 120 days (in $\text{mg CO}_2\text{-eq per microcosm}$). A nonparametric test (Independent-samples Mann-Whitney U test) established the difference in cumulative emissions from treatments with earthworms present (black bars '+EW') compared to treatments without earthworms present (grey bars '-EW'; black bars vs. grey bars: $p < .001$ for both N_2O and CO_2). Lowercase letter denotes differences between treatments with earthworms; any other differences occurring are reported in Table S3. Capital letters placed underneath each bar refer to the species combination given in Table 1. Colours indicate the number of species present: blue = 2 species present, pink = 4 species present and green = 8 species present. Underlined uppercase letters refer to species combinations that include an earthworm species

4 | DISCUSSION

4.1 | Increased species richness and functional dissimilarity result in contrasting faunal-induced emissions of N_2O and CO_2

Considering the major contribution of soil fauna species to below-ground processes, our findings emphasize the importance of soil fauna functional diversity in the soil function-climate feedback. We found an unexpected contrasting pattern for net biodiversity effects on soil-derived emissions of N_2O and CO_2 : N_2O emissions became smaller than expected when species number increased, whereas CO_2 emissions became larger. Consequences of soil fauna diversity for total soil-derived GHG emissions, therefore, can only be evaluated when understanding the net effects of reduced N_2O and enhanced CO_2 emission.

For CO_2 emissions, the results are corresponding to what is generally known about the effects of soil fauna species richness on ecosystem functioning: increasing species richness generally speeds up the rates of ecosystem processes, such as litter decomposition and/or soil respiration and nutrient cycling (Bardgett & Van der Putten, 2014; Bender, Wagg, & Heijden, 2016; Handa et al., 2014; Nielsen et al., 2011), especially at the low end of the diversity spectrum, as is the case in our controlled experimental set-up. However, the decreasing N_2O emissions with increasing species richness are remarkably opposite to this well-studied pattern. The explanation might lie in the fundamentally different nature of N_2O and CO_2 emissions. Whereas CO_2 is produced as the end product of respiration and decomposition processes, N_2O is produced as an intermediate or by-product in heterotrophic microbial N transformation processes (Wrage-Monnig et al., 2018). Therefore, any produced N_2O gas could be consumed again and further converted to elemental N_2 (Klefoth et al., 2012).

Mesofauna and macrofauna activity could potentially have both facilitative and inhibiting effects on the soil microbial transformation processes underlying N_2O emission. These constitute direct effects by, for example, stimulating the soil denitrifier community (Drake & Horn, 2006), or indirect effects by, for example, altering the in situ (an)aerobic conditions and influencing the C and N substrate availability (De Ruiter, Moore, et al., 1993; Schroter, Wolters, & Ruiter, 2003). For instance, net biodiversity effects on per capita MBN in relation to species number showed a similar pattern compared to the net biodiversity effects on N_2O emission (Figure S1); this could mean that the changes in MBN resulting from increasing species numbers are directly underlying the biodiversity effects found for the per capita N_2O emissions. Whether the changes in MBN can be related to increased/decreased activity of N-related microbes, or whether communities of N-related microbes were altered by increasing species richness is impossible to deduce from Figure S1.

Soil engineering species such as earthworms and potworms may also cause indirect effects by changing soil structural properties that affect gas diffusion in and out of the soil (Lee & Foster, 1991; Lubbers et al., 2011; Porre et al., 2016). All these faunal interactions with multiple microbial transformations converge in 'hotspots' and 'hot moments' of microbial activity (Kuzaykov & Blagodatskaya, 2015), and may shift the complex balance of N_2O production and consumption towards less N_2O production or more complete denitrification, and therefore to reduced rates of faunal-induced N_2O emission.

Next to establishing the influence of species richness on faunal-induced emissions of CO_2 and N_2O , we determined the effect of functional dissimilarity of species on CO_2 and N_2O in our multispecies mixtures. Production of CO_2 was most dominant in combinations including ecosystem engineering species (earthworms and potworms; Figure 2 combinations K, N, O, R, S, U, V, W) and therewith corresponded with previous findings by Heemsbergen et al. (2004). The observed inhibition of N_2O was likewise mostly influenced by species combinations that included ecosystem engineers, foremost the two earthworm species. As in our experiment the net biodiversity effects were explained most by soil ecosystem engineering traits (i.e. measurable effects on soil structural properties, Capowiez, Sammartino, & Michel, 2011; Lubbers et al., 2011; Porre et al., 2016), we included a soil structural element in our calculations of mean functional dissimilarity. Within the functional dissimilarity matrix, we interchanged pH for mean pore size (as measured by soil X-ray micro computed tomography; see Porre et al., 2016 for information about the X-ray tomography procedures and used software and programming, and see Sections 2 and 2.6 for information on calculations of mean *functional dissimilarity*). Especially the regression between the net biodiversity effect on N_2O and the mean functional dissimilarity of species in the community improved when mean pore size was included ($F_{64,63} = 8,56, p = .005$, Figure S2). For CO_2 , the regression model did not improve. This indicates that ecosystem engineering traits in soil animals are more important for N_2O emission than for CO_2 emission.

A similar point can be made when designing the dissimilarity matrix with MBN instead of pH and DON instead of NO_3 , thereby shifting the matrix more towards a focus on N cycling: the regression

between the net biodiversity effect on N_2O and the mean functional dissimilarity of species in the community became statistically stronger for N_2O when including MBN and DON in the dissimilarity matrix ($F_{64,63} = 8.79, p = .004$, Figure S3). For CO_2 , the regression model did not improve. The effects of soil fauna on N cycling properties thus also add explanatory power to the relationship between net biodiversity and mean functional dissimilarity. The reason why this relationship became stronger for N_2O and not for CO_2 is likely because the MBN varied mostly between the single-taxum earthworm treatments and the other non-earthworm fauna treatments (see Table S2), and this was also especially true for N_2O emissions (and to a lesser degree for CO_2 ; see Figure 3; Table S3). However, in what ways (size and activity of the N-related microbial community, as well as its community composition) faunal effects on MBN in the bulk soil influences N_2O emission remains elusive in this study.

To establish the influence of species richness on MBN as a response variable, we determined the effect of functional dissimilarity of species on MBN in our multispecies mixtures (see Figure S4 with a dissimilarity matrix of NH_4 , NO_3 , DOC and pH; and Figure S5 with a dissimilarity matrix of NH_4 , NO_3 , DOC and pore size). The net biodiversity effect on MBN in relation to mean functional dissimilarity shows a significant negative regression in both cases ($F_{51,50} = 4.81, p = .033$ in Figure S4 and $F_{51,50} = 4.48, p = .039$ in Figure S5). Like the net biodiversity effect on MBN in relation to species number, this pattern of reduced MBN with increasing functional dissimilarity follows the responses of N_2O emission in relation to species number and functional dissimilarity. However, when the soil structural element 'pore size' is included in the dissimilarity matrix, the statistical relationship between net biodiversity effect on MBN in relation to mean functional dissimilarity does not become stronger. This corroborates our expectation that ecosystem engineering traits (i.e. effects on soil structure such as pore size) in soil animals are especially important to consider in N_2O emissions.

4.2 | Community composition is key to explaining faunal-induced GHG emissions

Increased GHG emissions (N_2O and CO_2) by earthworm activity from intensively managed soils have been well established (Lubbers, Van Groenigen, et al., 2013). Mechanistic explanations for earthworm-induced N_2O emission by previous studies focus on increased N mineralization, nitrification and denitrification, as well as on a changed pore size distribution and an improved connectivity of soil pores with the atmosphere in the presence of earthworms (Bradley, Whalen, Chagnon, Lanoix, & Alves, 2011; Lubbers et al., 2011; Lubbers, González, Hummelink, & Groenigen, 2013; Nebert et al., 2011; Postma-Blaauw et al., 2006). Here, we show that earthworms increased N mineralization and activated N transformation processes, while C substrate to facilitate heterotrophic microbial processes did not become limiting. However, when we increased biodiversity of other fauna in the presence of earthworms (by increasing species richness as well as soil functional dissimilarity), faunal-induced N_2O emissions decreased despite enhanced C cycling.

4.3 | Implications for biodiversity effects on soil GHG mitigation

Different hierarchical levels of soil biodiversity may have differential influences on key soil ecosystem processes (Kardol, Throop, Adkins, & Graaff, 2016). However, body size of soil fauna may span the entire hierarchical biodiversity effect, and large soil ecosystem engineers such as earthworm species can dominate soil processes and have disproportional impact on GHG emissions. Therefore, we considered the earthworm effects on soil-derived N_2O and CO_2 apart from biodiversity effects by relating observed cumulative GHG emissions to earthworm share (in %, based on earthworm biomass) in the species combinations (Figure S6). The earthworm share is positively related to the effect of earthworm activity on CO_2 emissions (linear regression, $p < .001$, see Figure S7). This is in accordance with the mass ratio hypothesis (Grime, 1998), which states that a given ecosystem process is strongly influenced by the characteristics of a dominant species—proportional to its relative dominance in the community. The earthworm share is also positively related to the effect of earthworm activity on N_2O emissions (quadratic regression, $p < .001$, see Figure S7); however, in the case of N_2O emissions, the effects of earthworm presence are nonadditive and are not predictable from the sum of single species. This is indicative of a dominant effect of a unique trait. In our study, it could mean that the diminishing of earthworm-induced N_2O emissions is enhanced by increased variation in species trait values in the community; so in fact by increased functional biodiversity.

Bílá et al. (2014) concluded that dominant trait values in the community play a chief role in driving ecosystem processes such as C cycling processes (so corroborating the mass ratio hypothesis); it is yet to be determined whether the generality of their findings also hold true for other ecosystem processes. Here, we find that ecosystem engineers like our two earthworm species indeed possess dominant trait values concerning emissions of N_2O and CO_2 . However, while for CO_2 , the mass ratio hypothesis was supported; for N_2O , we have shown that functional diversity (indicative of the variation in species trait values in the community) promotes a non-linear decrease of faunal-induced N_2O emissions.

5 | POTENTIAL CAVEATS AND OUTLOOK

The extreme earthworm- N_2O effects observed in our experiment may partly be a consequence of our microcosm set-up. Earthworm numbers in the microcosms ranged from 1,140/m² to 570/m² for *A. caliginosa* and *L. rubellus*, respectively, in single-species treatments, to 285/m² for both earthworm species in the eight-species treatment. It is possible to have very high abundances of earthworms in both natural and cultivated environments; almost as high as in the single-species treatments of our microcosm set-up (Schon, Mackay, Gray, Kotten, & Dodd, 2017). However, especially when *A. caliginosa* was present in a single-species treatment (treatment 'I'), or combined with *L. rubellus* (treatment 'N'), its presence could result in

the collapse of soil structure. The microcosm soil could subsequently become compacted throughout, and partly anaerobic (likely going hand-in-hand with dramatic increases in the WFPS locally), but with the earthworm burrows mostly conserved. With this severe impact on soil structure, *A. caliginosa* may have caused its extreme N_2O effect by creating perfect in situ conditions for N_2O production, and simultaneously, a surface-connected pore network by which N_2O could easily escape to the atmosphere.

High earthworm abundances are ultimately associated with grassland systems (Schon et al., 2017). However, our microcosm set-up did not allow for plant growth (in the dark at 16°C and covered with cloth to keep the introduced soil animals alive and inside) and our results are therefore valid for bulk soil without an active rhizosphere. When an active rhizosphere is present, it is plausible that it has a substantial influence on earthworm-induced N_2O and CO_2 emissions. On the one hand, active plant roots take up nutrients such as $N-NH_4^+$ and $N-NO_3^-$ for growth (Kraiser, Gras, Gutiérrez, González, & Gutiérrez, 2011), and thereby (partly) remove the substrate for the manifold nitrate-reducing bacteria and denitrifiers that carry out the N transformation processes underpinning N_2O production. On the other hand, it is also to be expected that more easily available C is excreted in root exudates that are produced by active plant roots in the rhizosphere (Jones, Nguyen, & Finlay, 2009). Related consequences of an active rhizosphere may be enhanced microbial activity (increased C substrate/increased C availability), changing soil water and O_2 dynamics (give-and-take of water by roots as well as O_2 diffusion from the soil-atmosphere interface), and likely an interchanging use of the (bio)pore network by roots and soil fauna (perhaps slowing down diffusion rates of N_2O to the atmosphere and thereby creating more possibilities for N_2O to be reduced to N_2). Such consequences may especially cascade down in soil fauna diversity effects on the microbial processes (nitrification, denitrification, nitrifier-denitrification, decomposition) that underlie N_2O and CO_2 emissions. It is then to be expected that fauna diversity effects on N_2O and CO_2 are smaller or even occluded in an active rhizosphere. Yet, previous experimental work by Lubbers et al. (2011; Lubbers, González, et al., 2013) confirmed the occurrence of earthworm-induced effects on N_2O emissions in the presence of growing plants (*Lolium perenne*). These previously found effects could vary enormously, especially under field conditions: depending on the season and on earthworm species, earthworms could increase N_2O by 0%–394% (Lubbers, González, et al., 2013).

A potential unexpected source of crucial substrates for N_2O production is the extra N and C that entered the microcosm soil following the death of a number of earthworms. However, their dead biomass or emaciation during the experimental period (Table S1) was unlikely to have been the cause of either extremely increased or less increased rates of N_2O emission. Not only did few earthworms die but also there was no relationship between replicate microcosms in which earthworms had died and the N_2O or CO_2 emissions from the corresponding microcosms. In a previous study, Lubbers et al. (2011) calculated the highest possible contribution of N released from

decaying earthworm tissue and concluded that earthworm-induced N_2O emissions could not have resulted from dead earthworms.

These considerations in relation to potential caveats accompanying microcosm studies such as ours call for experimental work under realistic conditions including growing plants, prevailing weather conditions, and up-to-date agricultural management practices. Microcosm incubation studies that confirm 'proof of principle', namely, in our study effects of soil fauna diversity on emissions of CO_2 and N_2O , are a prerequisite for extended investigations that can be successfully extrapolated to realistic field conditions.

To conclude, we found that higher species richness and functional dissimilarity led to increased faunal-induced CO_2 emission but decreased N_2O emission. These contrasting dynamics are best explained by the fundamentally different processes underlying CO_2 and N_2O emissions: CO_2 is produced as the end product of respiration and decomposition processes, while N_2O is produced as an intermediate or by-product of microbial N transformation processes; during upward diffusion, any produced N_2O can be consumed again and be converted to elemental N_2 . Community composition appeared to determine the observed cumulative GHG emissions over an experimental period of 120 days. The presence of keystone species such as earthworms greatly enhanced soil-derived emissions of N_2O , but increasing species richness and functional dissimilarity reduced these elevated earthworm-induced N_2O emissions. These effects in our study call for further investigations that may contribute to the adoption of management practices that enhance soil biodiversity and stimulate a functionally diverse faunal community (rather than stimulating e.g. only target species such as earthworms) as to reduce N_2O emissions from managed soils.

Taken together, our findings recognize the importance of soil biodiversity conservation and call for incorporation of methods to maintain and stimulate the diversity of the living biota that provide critical processes and ecosystem services. New research may be directed towards a better mechanistic understanding of biodiversity effects on in situ soil structure including an active rhizosphere, with implications for gas diffusion and microbial diversity and activity. Within the contexts of 'soil ecological engineering' (Bender et al., 2016) and 'climate-smart soils' (Paustian et al., 2016), we need knowledge on faunal diversity effects connected to the potential of soils to mitigate GHG emissions that can be applied to achieve sustainable agriculture and viable agricultural systems.

ACKNOWLEDGEMENTS

We thank Tamás Salánki for extracting the potworms, mites and springtails from our soil samples, and Remco Hamoen for his assistance by obtaining the X-ray micro tomography scans (XRT) using CAT-AgroFood equipment. In addition, we would like to thank Jaap Nelemans, Willeke van Tintelen and Gerlinde Vink for their assistance in the laboratory. This research was supported by a grant from the Netherlands Organization for Scientific Research/Earth and Life Sciences (NWO-ALW, grant number 823.01.016).

AUTHOR CONTRIBUTIONS

IML and JWG conceived and designed the study, with suggestions and input from MPB and GBDD; IML and GBDD set up and destructively sampled the experiment; IML measured all GHG fluxes, maintained the experiment, did laboratory analyses and performed the statistical analyses; IML, JWG, MPB, WHP and GBDD interpreted and discussed the results; IML, JWG and WHP wrote the paper, with substantial contributions from all co-authors; JWG had the overall supervision of the project.

ORCID

Ingrid M. Lubbers  <https://orcid.org/0000-0002-2168-104X>

DATA AVAILABILITY STATEMENT

Raw data were generated at the Soil Physics Laboratory and the Chemical Biological Soil Laboratory of the Wageningen University, the Netherlands. Derived data supporting the findings of this study are available from the corresponding author upon request.

REFERENCES

- Bardgett, R. D., & Van Der Putten, W. H. (2014). Belowground biodiversity and ecosystem functioning. *Nature*, *515*, 505–511. <https://doi.org/10.1038/nature13855>
- Bender, S. F., Wagg, C., & Van Der Heijden, M. G. A. (2016). An underground revolution: Biodiversity and soil ecological engineering for agricultural sustainability. *Trends in Ecology & Evolution*, *31*, 440–452. <https://doi.org/10.1016/j.tree.2016.02.016>
- Bílá, K., Moretti, M., De Bello, F., Dias, A. T. C., Pezzatti, G. B., Van Oosten, A. R., & Berg, M. P. (2014). Disentangling community functional components in a litter-macrodetrivore model system reveals the predominance of the mass ratio hypothesis. *Ecology and Evolution*, *4*, 408–416. <https://doi.org/10.1002/ece3.941>
- Blagodatsky, S., & Smith, P. (2012). Soil physics meets soil biology: Towards better mechanistic prediction of greenhouse gas emissions from soil. *Soil Biology and Biochemistry*, *47*, 78–92. <https://doi.org/10.1016/j.soilbio.2011.12.015>
- Bradley, R. L., Whalen, J., Chagnon, P. L., Lanoix, M., & Alves, M. C. (2011). Nitrous oxide production and potential denitrification in soils from riparian buffer strips: Influence of earthworms and plant litter. *Applied Soil Ecology*, *47*, 6–13. <https://doi.org/10.1016/j.apsoil.2010.11.007>
- Brookes, P. C., Landman, A., Pruden, G., & Jenkinson, D. S. (1985). Chloroform fumigation and the release of soil nitrogen: A rapid direct extraction method to measure microbial biomass nitrogen in soil. *Soil Biology and Biochemistry*, *17*, 837–842. [https://doi.org/10.1016/0038-0717\(85\)90144-0](https://doi.org/10.1016/0038-0717(85)90144-0)
- Capowiez, Y., Sammartino, S., & Michel, E. (2011). Using X-ray tomography to quantify earthworm bioturbation non-destructively in repacked soil cores. *Geoderma*, *162*, 124–131. <https://doi.org/10.1016/j.geoderma.2011.01.011>
- Cardinale, B. J., Srivastava, D. S., Emmett Duffy, J., Wright, J. P., Downing, A. L., Sankaran, M., & Jouseau, C. (2006). Effects of biodiversity on the functioning of trophic groups and ecosystems. *Nature*, *443*, 989–992. <https://doi.org/10.1038/nature05202>
- Coulis, M., Fromin, N., David, J.-F., Gavinet, J., Clet, A., Devidal, S., ... Hättenschwiler, S. (2015). Functional dissimilarity across trophic levels as a driver of soil processes in a Mediterranean decomposer system

- exposed to two moisture levels. *Oikos*, 124, 1304–1316. <https://doi.org/10.1111/oik.01917>
- Crowther, T. W., Thomas, S. M., Maynard, D. S., Baldrian, P., Covey, K., Frey, S. D., ... Bradford, M. A. (2015). Biotic interactions mediate soil microbial feedbacks to climate change. *Proceedings of the National Academy of Sciences of the United States of America*, 112, 7033–7038. <https://doi.org/10.1073/pnas.1502956112>
- Crowther, T. W., Todd-Brown, K. E. O., Rowe, C. W., Wieder, W. R., Carey, J. C., Machmuller, M. B., ... Bradford, M. A. (2016). Quantifying global soil carbon losses in response to warming. *Nature*, 540, 104–108. <https://doi.org/10.1038/nature20150>
- Dalal, R. C., & Allen, D. E. (2008). Greenhouse gas fluxes from natural ecosystems. *Australian Journal of Botany*, 56, 369–407. <https://doi.org/10.1071/BT07128>
- Dalby, P. R., Baker, G. H., & Smith, S. E. (1996). "Filter paper method" to remove soil from earthworm intestines and to standardise the water content of earthworm tissue. *Soil Biology and Biochemistry*, 28, 685–687.
- De Ruiter, P. C., Van Veen, J. A., Moore, J. C., Brussaard, L., & Hunt, H. W. (1993). Calculation of nitrogen mineralization in soil food webs. *Plant and Soil*, 157, 263–273. <https://doi.org/10.1007/BF00011055>
- De Ruiter, P. C. D., Moore, J. C., Zwart, K. B., Bouwman, L. A., Hassink, J., Bloem, J., ... Brussaard, L. (1993). Simulation of nitrogen mineralization in the below-ground food webs of two winter wheat fields. *Journal of Applied Ecology*, 30, 95–106. <https://doi.org/10.2307/2404274>
- de Vries, F. T., Thebault, E., Liiri, M., Birkhofer, K., Tsiafouli, M. A., Bjornlund, L., ... Bardgett, R. D. (2013). Soil food web properties explain ecosystem services across European land use systems. *Proceedings of the National Academy of Sciences of the United States of America*, 110, 14296–14301. <https://doi.org/10.1073/pnas.1305198110>
- Dias, A. T. C., Berg, M. P., De Bello, F., Van Oosten, A. R., Bilá, K., & Moretti, M. (2013). An experimental framework to identify community functional components driving ecosystem processes and services delivery. *Journal of Ecology*, 101, 29–37. <https://doi.org/10.1111/1365-2745.12024>
- Drake, H. L., & Horn, M. A. (2006). Earthworms as a transient heaven for terrestrial denitrifying microbes: A review. *Engineering in Life Sciences*, 6, 261–265. <https://doi.org/10.1002/elsc.200620126>
- FAO. (2016). *FAO statistical yearbooks; FAO statistical pocketbooks*. Rome, Italy: FAO. Retrieved from <https://www.fao.org/economic/ess/esspublications/en/#.WD1c-7lrJhE>
- Filser, J., Faber, J. H., Tiunov, A. V., Brussaard, L., Frouz, J., De Deyn, G., ... Jiménez, J. J. (2016). Soil fauna: Key to new carbon models. *Soil*, 2, 565–582. <https://doi.org/10.5194/soil-2-565-2016>
- Grandy, A. S., Wieder, W. R., Wickings, K., & Kyker-Snowman, E. (2016). Beyond microbes: Are fauna the next frontier in soil biogeochemical models? *Soil Biology and Biochemistry*, 102, 40–44. <https://doi.org/10.1016/j.soilbio.2016.08.008>
- Granli, T., & Bøckman, O. C. (1994). Nitrous oxide from agriculture. *Norwegian Journal of Agricultural Sciences*, Supplement No. 12, 1–128.
- Grime, J. P. (1998). Benefits of plant diversity to ecosystems: Immediate, filter and founder effects. *Journal of Ecology*, 86, 902–910. <https://doi.org/10.1046/j.1365-2745.1998.00306.x>
- Handa, I. T., Aerts, R., Berendse, F., Berg, M. P., Bruder, A., Butenschoten, O., ... Hättenschwiler, S. (2014). Consequences of biodiversity loss for litter decomposition across biomes. *Nature*, 509, 218–221. <https://doi.org/10.1038/nature13247>
- Heemsbergen, D. A., Berg, M. P., Loreau, M., Van Hal, J. R., Faber, J. H., & Verhoef, H. A. (2004). Biodiversity effects on soil processes explained by interspecific functional dissimilarity. *Science*, 306, 1019–1020. <https://doi.org/10.1126/science.1101865>
- Houba, V. J. G., Temminghoff, E. J. M., Gaikhorst, G. A., & Van Vark, W. (2000). Soil analysis procedures using 0.01 M calcium chloride as extraction reagent. *Communications in Soil Science and Plant Analysis*, 31, 1299–1396.
- IPCC. (2013). Climate change 2013: The physical science basis. In T. F. Stocker, D. Qin, G.-K. Plattner, M. Tignor, S. K. Allen, J. Boschung, ... P. M. Midgley (Eds.), *Contribution of working group I to the fifth assessment report of the Intergovernmental Panel on Climate Change*. Cambridge, UK: Cambridge University Press, 1585 pp.
- IPCC. (2014). Climate change 2014: Mitigation of climate change. In O. Edenhofer, R. Pichs-Madruga, Y. Sokona, E. Farahani, S. Kadner, K. Seyboth, ... J. C. Minx (Eds.), *Contribution of working group III to the fifth assessment report of the Intergovernmental Panel on Climate Change*. Cambridge, UK: Cambridge University Press, 31 pp.
- Jones, D., Nguyen, C., & Finlay, D. R. (2009). Carbon flow in the rhizosphere: Carbon trading at the soil–root interface. *Plant and Soil*, 321, 5–33. <https://doi.org/10.1007/s11104-009-9925-0>
- Kaneda, S., & Kaneko, N. (2011). Influence of Collembola on nitrogen mineralization varies with soil moisture content. *Soil Science and Plant Nutrition*, 57, 40–49. <https://doi.org/10.1080/00380768.2010.551107>
- Kardol, P., Throop, H. L., Adkins, J., & De Graaff, M.-A. (2016). A hierarchical framework for studying the role of biodiversity in soil food web processes and ecosystem services. *Soil Biology and Biochemistry*, 102, 33–36. <https://doi.org/10.1016/j.soilbio.2016.05.002>
- Klefoth, R., Oenema, O., & Willem Van Groenigen, J. (2012). A novel method for quantifying nitrous oxide reduction in soil. *Vadose Zone Journal*, 11(4). <https://doi.org/10.2136/vzj2011.0107>
- Klug, M. J. (1999). The determination of microbial biomass. From: Standard soil methods for long-term ecological research. In G. Robertson, D. C. Coleman, C. S. Bledsoe, & P. Sollins (Eds.), *Standard soil methods for long-term ecological research* (p. 291). Oxford, UK: Oxford University Press.
- Kraiser, T., Gras, D. E., Gutiérrez, A. G., González, B., & Gutiérrez, R. A. (2011). A holistic view of nitrogen acquisition in plants. *Journal of Experimental Botany*, 62, 1455–1466. <https://doi.org/10.1093/jxb/erq425>
- Kuiper, I., De Deyn, G. B., Thakur, M. P., & Van Groenigen, J. W. (2013). Soil invertebrate fauna affect N₂O emissions from soil. *Global Change Biology*, 19, 2814–2825.
- Kuzyakov, Y., & Blagodatskaya, E. (2015). Microbial hotspots and hot moments in soil: Concept & review. *Soil Biology and Biochemistry*, 83, 184–199.
- Lal, R. (2004). Soil carbon sequestration impacts on global climate change and food security. *Science*, 304, 1623–1627. <https://doi.org/10.1126/science.1097396>
- Le Mer, J., & Roger, P. (2001). Production, oxidation, emission and consumption of methane by soils: A review. *European Journal of Soil Biology*, 37, 25–50. [https://doi.org/10.1016/S1164-5563\(01\)01067-6](https://doi.org/10.1016/S1164-5563(01)01067-6)
- Lee, S. Y., & Foster, R. C. (1991). Soil fauna and soil structure. *Australian Journal of Soil Research*, 29, 745–775. <https://doi.org/10.1071/SR9910745>
- Levy, P. E., Burden, A., Cooper, M. D. A., Dinsmore, K. J., Drewer, J., Evans, C., ... Zieliński, P. (2012). Methane emissions from soils: Synthesis and analysis of a large UK data set. *Global Change Biology*, 18, 1657–1669. <https://doi.org/10.1111/j.1365-2486.2011.02616.x>
- Liiri, M., Setälä, H., Haimi, J., Pennanen, T., & Fritze, H. (2002). Relationship between soil microarthropod species diversity and plant growth does not change when the system is disturbed. *Oikos*, 96, 137–149. <https://doi.org/10.1034/j.1600-0706.2002.960115.x>
- Loreau, M., & Hector, A. (2001). Partitioning selection and complementarity in biodiversity experiments. *Nature*, 412, 72–76. <https://doi.org/10.1038/35083573>
- Lubbers, I. M., Brussaard, L., Otten, W., & Van Groenigen, J. W. (2011). Earthworm-induced N mineralization in fertilized grassland increases both N₂O emission and crop-N uptake. *European Journal of Soil Science*, 62, 152–161. <https://doi.org/10.1111/j.1365-2389.2010.01313.x>

- Lubbers, I. M., González, E. L., Hummelink, E. W. J., & Van Groenigen, J. W. (2013). Earthworms can increase nitrous oxide emissions from managed grassland: A field study. *Agriculture, Ecosystems & Environment*, 174, 40–48. <https://doi.org/10.1016/j.agee.2013.05.001>
- Lubbers, I. M., Van Groenigen, K. J., Brussaard, L., & Van Groenigen, J. W. (2015). Reduced greenhouse gas mitigation potential of no-tillage soils through earthworm activity. *Scientific Reports*, 5, 13787. <https://doi.org/10.1038/srep13787>
- Lubbers, I. M., Van Groenigen, K. J., Fonte, S. J., Six, J., Brussaard, L., & Van Groenigen, J. W. (2013). Greenhouse-gas emissions from soils increased by earthworms. *Nature Climate Change*, 3, 187–194. <https://doi.org/10.1038/nclimate1692>
- Majeed, M. Z., Miambi, E., Barois, I., Randriamanantsoa, R., Blanchart, E., & Brauman, A. (2014). Contribution of white grubs (Scarabaeidae: Coleoptera) to N₂O emissions from tropical soils. *Soil Biology and Biochemistry*, 75, 37–44. <https://doi.org/10.1016/j.soilbio.2014.03.025>
- Nebert, L., Bloem, J., Lubbers, I., & Van Groenigen, J. (2011). Association of earthworm – Denitrifier interactions with increased emissions of nitrous oxide from soil mesocosms amended with crop residue. *Applied Environmental Microbiology*, 77, 4097–4104.
- Nielsen, U. N., Ayres, E., Wall, D. H., & Bardgett, R. D. (2011). Soil biodiversity and carbon cycling: A review and synthesis of studies examining diversity-function relationships. *European Journal of Soil Science*, 62, 105–116. <https://doi.org/10.1111/j.1365-2389.2010.01314.x>
- Opdyke, M. R., Ostrom, N. E., & Ostrom, P. H. (2009). Evidence for the predominance of denitrification as a source of N₂O in temperate agricultural soils based on isotopologue measurements. *Global Biogeochemical Cycles*, 23, GB4018.
- Paustian, K., Cole, C. V., Sauerbeck, D., & Sampson, N. (1998). CO₂ mitigation by agriculture: An overview. *Climatic Change*, 40, 135–162.
- Paustian, K., Lehmann, J., Ogle, S., Reay, D., Robertson, G. P., & Smith, P. (2016). Climate-smart soils. *Nature*, 532, 49–57. <https://doi.org/10.1038/nature17174>
- Petersen, H., & Luxton, M. (1982). A comparative analysis of soil fauna populations and their role in decomposition processes. *Oikos*, 39, 288–388. <https://doi.org/10.2307/3544689>
- Ponge, J.-F., Pérès, G., Guernion, M., Ruiz-Camacho, N., Cortet, J., Pernin, C., ... Cluzeau, D. (2013). The impact of agricultural practices on soil biota: A regional study. *Soil Biology & Biochemistry*, 67, 271–284. <https://doi.org/10.1016/j.soilbio.2013.08.026>
- Porre, R. J., Van Groenigen, J. W., De Deyn, G. B., De Goede, R. G. M., & Lubbers, I. M. (2016). Exploring the relationship between soil mesofauna, soil structure and N₂O emissions. *Soil Biology and Biochemistry*, 96, 55–64. <https://doi.org/10.1016/j.soilbio.2016.01.018>
- Postma-Blaauw, M. B., Bloem, J., Faber, J. H., Van Groenigen, J. W., De Goede, R. G. M., & Brussaard, L. (2006). Earthworm species composition affects the soil bacterial community and net nitrogen mineralization. *Pedobiologia*, 50, 243–256. <https://doi.org/10.1016/j.pedobi.2006.02.001>
- Postma-Blaauw, M. B., De Goede, R. G. M., Bloem, J., Faber, J. H., & Brussaard, L. (2012). Agricultural intensification and de-intensification differentially affect taxonomic diversity of predatory mites, earthworms, enchytraeids, nematodes and bacteria. *Applied Soil Ecology*, 57, 39–49. <https://doi.org/10.1016/j.apsoil.2012.02.011>
- Powlson, D. S., Stirling, C. M., Jat, M. L., Gerard, B. G., Palm, C. A., Sanchez, P. A., & Cassman, K. G. (2014). Limited potential of no-till agriculture for climate change mitigation. *Nature Climate Change*, 4, 678–683. <https://doi.org/10.1038/nclimate2292>
- Powlson, D. S., Whitmore, A. P., & Goulding, K. W. T. (2011). Soil carbon sequestration to mitigate climate change: A critical re-examination to identify the true and the false. *European Journal of Soil Science*, 62, 42–55. <https://doi.org/10.1111/j.1365-2389.2010.01342.x>
- Rastogi, M., Singh, S., & Pathak, H. (2002). Emission of carbon dioxide from soil. *Current Science*, 82, 510–518.
- Sauvadet, M., Chauvat, M., Brunet, N., & Bertrand, I. (2017). Can changes in litter quality drive soil fauna structure and functions? *Soil Biology & Biochemistry*, 107, 94–103. <https://doi.org/10.1016/j.soilbio.2016.12.018>
- Schon, N. L., Mackay, A. D., Gray, R. A., Van Koten, C., & Dodd, M. B. (2017). Influence of earthworm abundance and diversity on soil structure and the implications for soil services throughout the season. *Pedobiologia*, 62, 41–47. <https://doi.org/10.1016/j.pedobi.2017.05.001>
- Schorpp, Q., Riggers, C., Lewicka-Szczepak, D., Giesemann, A., Well, R., & Schrader, S. C. R. C. M. R. (2016). Influence of *Lumbricus terrestris* and *Folsomia candida* on N₂O formation pathways in two different soils – With particular focus on N₂ emissions. *Rapid Communications in Mass Spectrometry*, 30, 2301–2314.
- Schroter, D., Wolters, V., & De Ruiter, P. C. (2003). C and N mineralisation in the decomposer food webs of a European forest transect. *Oikos*, 102, 294–308. <https://doi.org/10.1034/j.1600-0579.2003.12064.x>
- Smith, P. (2016). Soil carbon sequestration and biochar as negative emission technologies. *Global Change Biology*, 22, 1315–1324. <https://doi.org/10.1111/gcb.13178>
- Stockmann, U., Adams, M. A., Crawford, J. W., Field, D. J., Henakaarchchi, N., Jenkins, M., ... Zimmermann, M. (2013). The knowns, known unknowns and unknowns of sequestration of soil organic carbon. *Agriculture, Ecosystems & Environment*, 164, 80–99. <https://doi.org/10.1016/j.agee.2012.10.001>
- Thakur, M. P., Van Groenigen, J. W., Kuiper, I., & De Deyn, G. B. (2014). Interactions between microbial-feeding and predatory soil fauna trigger N₂O emissions. *Soil Biology and Biochemistry*, 70, 256–262. <https://doi.org/10.1016/j.soilbio.2013.12.020>
- Tianxiang, L., Huixin, L., Tong, W., & Feng, H. (2008). Influence of nematodes and earthworms on the emissions of soil trace gases (CO₂, N₂O). *Acta Ecologica Sinica*, 28, 993–999. [https://doi.org/10.1016/S1872-2032\(08\)60033-5](https://doi.org/10.1016/S1872-2032(08)60033-5)
- Tresch, S., Frey, D., Le Bayon, R. C., Zanetta, A., Rasche, F., Fliessbach, A., & Moretti, M. (2019). Litter decomposition driven by soil fauna, plant diversity and soil management in urban gardens. *Science of the Total Environment*, 658, 1614–1629. <https://doi.org/10.1016/j.scitotenv.2018.12.235>
- Tsiafouli, M. A., Thébault, E., Sgardelis, S. P., de Ruiter, P. C., van der Putten, W. H., Birkhofer, K., ... Hedlund, K. (2015). Intensive agriculture reduces soil biodiversity across Europe. *Global Change Biology*, 21, 973–985. <https://doi.org/10.1111/gcb.12752>
- Van Nes, E. H., Scheffer, M., Brovkin, V., Lenton, T. M., Ye, H., Deyle, E., & Sugihara, G. (2015). Causal feedbacks in climate change. *Nature Climate Change*, 5, 445–448. <https://doi.org/10.1038/nclimate2568>
- Van Vliet, P. C. J., Beare, M. H., Coleman, D. C., & Hendrix, P. F. (2004). Effects of enchytraeids (Annelida: Oligochaeta) on soil carbon and nitrogen dynamics in laboratory incubations. *Applied Soil Ecology*, 25, 147–160. <https://doi.org/10.1016/j.apsoil.2003.08.004>
- Van Groenigen, J. W., Lubbers, I. M., Vos, H. M. J., Brown, G. G., De Deyn, G. B., & van Groenigen, K. J. (2014). Earthworms increase plant production: A meta-analysis. *Scientific Reports*, 4(1), <https://doi.org/10.1038/srep06365>
- Velthof, G., Kuikman, P., & Oenema, O. (2002). Nitrous oxide emission from soils amended with crop residues. *Nutrient Cycling in Agroecosystems*, 62, 249–261.
- Walker, B., Kinzig, A., & Langridge, J. (1999). Plant attribute diversity, resilience, and ecosystem function: The nature and significance of dominant and minor species. *Ecosystems*, 2, 95–113.
- Wall, D. (2012). *Soil ecology and ecosystem services*. Oxford, UK: Oxford University Press.

- Wall, D. H., Bradford, M. A., St. John, M. G., Trofymow, J. A., Behan-Pelletier, V., Bignell, D. E., ... Zou, X. (2008). Global decomposition experiment shows soil animal impacts on decomposition are climate-dependent. *Global Change Biology*, *14*, 2661–2677. <https://doi.org/10.1111/j.1365-2486.2008.01672.x>
- Wrage-Monnig, N., Horn, M. A., Well, R., Muller, C., Velthof, G., & Oenema, O. (2018). The role of nitrifier denitrification in the production of nitrous oxide revisited. *Soil Biology & Biochemistry*, *123*, A3–A16. <https://doi.org/10.1016/j.soilbio.2018.03.020>
- Wu, H., Lu, M., Lu, X., Guan, Q., & He, X. (2015). Interactions between earthworms and mesofauna has no significant effect on emissions of CO₂ and N₂O from soil. *Soil Biology and Biochemistry*, *88*, 294–297. <https://doi.org/10.1016/j.soilbio.2015.06.005>

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section.

How to cite this article: Lubbers IM, Berg MP, De Deyn GB, van der Putten WH, van Groenigen JW. Soil fauna diversity increases CO₂ but suppresses N₂O emissions from soil. *Glob Change Biol*. 2020;26:1886–1898. <https://doi.org/10.1111/gcb.14860>