



Correction to: Japanese Clinical Practice Guideline for Diabetes 2016

Masakazu Haneda¹ · Mitsuhiro Noda² · Hideki Origasa³ · Hiroshi Noto⁴ · Daisuke Yabe⁵ · Yukihiro Fujita¹ · Atsushi Goto⁶ · Tatsuya Kondo⁷ · Eiichi Araki⁷

Published online: 20 December 2019
© The Japan Diabetes Society 2019

Correction to: Diabetology International (2018) 9:1–45
<https://doi.org/10.1007/s13340-018-0345-3>

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

In the original version of the article, Table 5 was published incorrectly. The correct version of Table 5 is given below.

Table 5 The lipid control target values in patients with diabetes. Edited by Japan Atherosclerosis Society: Guidelines for Prevention of Atherosclerotic Cardiovascular Diseases, published 2012, p. 42

Coronary artery disease	Lipid control target values (mg/dL)			
	LDL-C	HDL-C	TG	Non-HDL-C
Absent	< 120			< 150
		≥ 40	< 150	
Present	< 100			< 130

The original article can be found online at <https://doi.org/10.1007/s13340-018-0345-3>.

✉ Masakazu Haneda
jds-journal@jds.or.jp

- ¹ Asahikawa Medical University, Asahikawa, Japan
- ² Saitama Medical University, Saitama, Japan
- ³ University of Toyama, Toyama, Japan
- ⁴ St Luke's International Hospital, Tokyo, Japan
- ⁵ Department of Diabetes, Endocrinology and Nutrition, Kyoto University Graduate School of Medicine, Kyoto, Japan
- ⁶ Center for Public Health Sciences, National Cancer Center, Tokyo, Japan
- ⁷ Department of Metabolic Medicine, Kumamoto University, Kumamoto, Japan